Integrated phenotypic and mutational approach defines EBF3-related HADD syndrome genotype-phenotype relationships

Cole A. Deisseroth,[1,2,14,16] Aarushi Nayak,[3] Nathan D. Bliss,[4] Vanesa Lerma,[2,5] Ashley W. LeMaire,[6] Vinodh Narayanan,[7] Christopher Balak,[8] Ginevra Zanni,[9] Enza Maria Valente,[10,11] Enrico Bertini,[9] Paul J. Benke,[12] Michael F. Wangler,[2,13,16] Hsiao-Tuan Chao[1,2,5,13,14,15,17]

[1]Medical Scientist Training Program, Baylor College of Medicine, Houston, TX, 77030, USA

[2]Jan and Dan Duncan Neurological Research Institute, Texas Children's Hospital, Houston, TX, 77030, USA

[3]Case Western Reserve University, Cleveland, OH, 44106, USA

[4]Department of Biomedical Engineering, Texas A&M University, College Station, TX, 77843, USA

[5]Department of Pediatrics, Division of Neurology and Developmental Neuroscience, Baylor College of Medicine, Houston, TX, 77030, USA

[6]Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, TX, 77030, USA

[7]Center for Rare Childhood Disorders, Translational Genomics Research Institute, Phoenix, AZ, 85004, USA

[8]Biomedical Sciences Graduate Program, University of California at San Diego, San Diego, CA, 92093, USA

[9]Department of Neurosciences, Unit of Neuromuscular and Neurodegenerative Disorders, Bambino Gesu Children's Research Hospital IRCCS, Rome, 00146, Italy

[10]Department of Molecular Medicine, University of Pavia, Pavia, 27100, Italy

[11]Neurogenetics Research Centre, IRCCS Mondino Foundation, Pavia, 27100, Italy

[12]Joe DiMaggio Children's Hospital, Hollywood, FL, 33021, USA

[13]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, 77030, USA

[14]Department of Neuroscience, Baylor College of Medicine, Houston, TX, 77030, USA

[15]McNair Medical Institute, The Robert and Janice McNair Foundation, Houston, TX, 77030, USA

[16]Co-corresponding authors: Cole.Deisseroth@bcm.edu (C.A.D.), mw147467@bcm.edu (M.F.W.)

[17]Corresponding author: hc140077@bcm.edu (H.T.C.), 832-826-0461, 1250 Moursund St, Suite 925, Room 930.01, Houston, TX, 77030

Abstract

Hypotonia, Ataxia, and Delayed Development syndrome is a neurodevelopmental disorder caused by heterozygous *Early B-Cell Factor 3* (*EBF3*) loss-of-function variants. Identified in 2016, the full spectrum of clinical findings and the relationship between the *EBF3* genotype and clinical outcomes has not been determined beyond its namesake features. We combined a phenotypic assessment of 33 individuals molecularly diagnosed with *EBF3* pathogenic variants with a meta-analysis of 34 previously reported individuals. The combined 62 unique individuals enabled comparative cross-sectional phenotype and genotype analysis in the largest cohort to date of affected individuals. Cardinal distinguishing features were identified that facilitate phenotypic stratification for clinical diagnosis. We developed assessment scales to ascertain individuals at risk for pathogenic *EBF3* variants, stratify the clinical severity, and connect variant-specific molecular phenotypes to clinical outcomes. Our findings show that a specific class of *EBF3* variants affecting the evolutionarily conserved Zinc Finger (ZNF) motif, which is critical for stabilizing the protein interaction with the DNA target sequence, is associated with an increased risk of persistent motor and language impairments. These findings highlight the impact of combining variant-specific molecular phenotypes with comprehensive clinical data to predict neurodevelopmental outcomes and potentially guide personalized decisions for therapeutic interventions.

Introduction

Early B-Cell Factor 3 (*EBF3* [MIM: 607407] HGNC: 19087) encodes a Collier/Olf/EBF (COE) transcription factor located on chromosome 10q26. COE transcription factors control gene expression via DNA methylation, nucleosome remodeling, and active chromatin modifications[1–3], and regulate nervous system, immune system, bone, and muscle development[4–7]. Loss-of-function manipulations of *EBF3* homologs in worms (*unc-3*, *CeO/E*), flies (*knot, collier*), frogs (*Xcoe2*), and mice (*Ebf3*) are deleterious to survival and impair neurodevelopmental processes[8]. In mice, *Ebf3* has been shown to regulate neuronal proliferation, sub-type specification, and migration[6,9–11]. Furthermore, cellular studies revealed that *EBF3* binds to the promoters of many genes associated with neurodevelopmental and autism spectrum disorders[12], suggesting that EBF3 dysfunction could result in a wider phenotypic spectrum than currently recognized.

Over 30 years ago, genomic deletions containing *EBF3* were identified in the 10q26-deletion syndrome, which is characterized by delayed development, hypotonia, ataxia, growth problems, neuropsychiatric comorbidities, and genitourinary abnormalities [MIM: 609625][13–17]. However, the principal genetic alterations mediating pathogenesis in 10q26-deletion syndrome was unknown until recently. In 2016, *de novo EBF3* variants and a heterozygous *Ebf3* single gene deletion were discovered to cause the Hypotonia, Ataxia, and Delayed Development Syndrome (HADDS [MIM: 617330]), which is characterized by the namesake features and includes impaired expressive language, neuropsychiatric comorbidities including autism, and genitourinary abnormalities[12,18–20]. The strong overlap in features between HADDS and 10q26-deletion syndrome suggests that *EBF3* haploinsufficiency may be the principal pathogenic factor in 10q26-deletion syndrome.

Although the discovery of a monogenic disorder associated with pathogenic *EBF3* variants is

fairly recent, there is increasing evidence that *EBF3*-related disorders may be more common than

previously recognized because of a wide range of clinical severity[20–25]. This is consistent with a

prior statistical model of *de novo* variants for autism spectrum disorder (ASD) and

developmental and intellectual disability (DD/ID) that identified *EBF3* as one of ~1,000 genes

with significant enrichment of *de novo* loss-of-function variants in affected individuals[26].

Therefore, expanding our understanding of the HADDS phenotypic spectrum and co-morbid

conditions will advance diagnosis, facilitate prognosis, and improve neurodevelopmental

outcomes for individuals with HADDS, 10q26-deletion syndrome, and other prevalent

neurodevelopmental disorders with similar features.

A common challenge faced with newly recognized or relatively rare neurodevelopmental

disorders is that incomplete knowledge about distinguishing features limits the stratification of

affected individuals for traditional phenotypically driven diagnostics. To expand our

understanding of the HADDS phenotypic spectrum and identify individuals at risk for

pathogenic *EBF3* variants, we conducted a comprehensive phenotypic and mutational analysis of

33 individuals diagnosed with HADDS and correlated the findings with a literature meta-analysis

for a combined assessment of 62 unique individuals. Our findings expand our understanding of

the HADDS phenotypic spectrum and reveal that *EBF3* variant type and location are associated

with quantifiable clinical severity.

MATERIALS AND METHODS

BCM investigators systematically evaluated 22 individuals in person at Texas Children's

Hospital and 11 through medical record reviews for a total of 33 individuals. Individuals were

enrolled in a study approved by the Institutional Review Board of Baylor College of Medicine

(BCM), following its ethical standards. Written and informed consent was obtained from all

individuals before the study was conducted. All procedures followed national standards. All data

were de-identified. Three individuals opted out of publishing genomic information. We

compared our cohort to a meta-analysis of nine studies containing a total of 34 individuals with

pathogenic *EBF3* variants[12,18–25]. Five of the individuals enrolled in our study were previously

reported. Probands 7, 15, 19, 25, and 26 from our study (**Table S1**) are the same as meta-analysis

probands M-11, M-1, M-25, M-31, and M-32, respectively (**Table S2**). We assessed a total of 62

unique individuals and compared demographics, clinical features, and molecular findings in our

cohort to the meta-analysis (**Tables S1-S5**). Variant nomenclature is according to gene transcript

(GenBank: NM_001005463.2, GRCh37). Odds ratios (OR) = AD/BC, where A is the number of

probands from our cohort with the feature, B is the number of probands from our cohort without

the feature, C is the number of probands from the meta-analysis with the feature, and D is the

number of probands from the meta-analysis without the feature. 95% confidence intervals (CI)

were calculated for a sample size *n* and proportion *p*:

$$CI = p \pm 1.96 \sqrt{\frac{p(1-p)}{n}}$$

For height, weight, and occipito-frontal circumference (OFC) measurements, we calculated LMS

(lambda-mu-sigma)-based Z-scores[27], where for a given L (degree of skewness), M (median),

and S (variation coefficient) values, and an experimental value X, the z-score is $= [(X/M)^L -$

1]/(LS). L, M, and S values for height and weight based on sex and age (birth to 20 years) are

from the Centers for Disease Control and Prevention (CDC)[28]. For OFC, the L, M, and S values

were taken from the World Health Organization (WHO, birth to five years)[29]. Delayed

development was determined by failure to achieve developmental milestones within the CDC's

standard age range (**Tables S4, S5**)[30].

HADDS Developmental and Behavioral Scale (HADDS-DBS)

The HADDS-DBS assesses for seven major and 14 minor features (**Table 1**). Major features

include decreased pain sensitivity (HP: 0007328), weak/absent cry (HP: 0001612), motor delay

(HP: 0001270), hypotonia (HP: 0001252), speech delay (HP: 0000750), ataxia (HP: 0001251),

and hypomimia (HP: 0000338). Minor features include poor eye contact, strabismus, repetitive

scratching in any location, repetitive scratching on back or neck, attraction to lights, aversion to

crowds, aversion to loud noises, constipation, frontal bossing, recurrent urinary tract infections,

feeding difficulties, cerebellar anomalies, gait ataxia, and short stature. Based on the number of

diagnostic features identified, the individuals are stratified into four tiers (**Table S6**). Higher-

numbered tiers indicate a higher risk for a pathogenic *EBF3* variant.

Positive (PPV) and negative (NPV) predictive values were calculated by combining the

HADDS-DBS tier assignments with the molecularly confirmed genetic diagnoses. True positives

(TP) are individuals with HADDS and a Tier-3 or -4 assignment. False positives (FP) are

individuals with non-*EBF3* related genetic syndromes and a Tier-3 or -4 assignment. True

negatives (TN) are individuals with non-*EBF3* related genetic syndromes and a Tier-1 or -2

assignment. False negatives (FN) are individuals with HADDS and a Tier-1 or -2 assignment.

Sensitivity = TP/(TP+FN). Specificity = TN/(TN+FP). PPV = TP/(TP+FP). NPV =

TN/(TN+FN).

For each diagnostic tier, OR = AD/BC, where A is the number of probands from our cohort at

the specified tier, B is the number of probands from our cohort not at the specified tier, C is the

number of probands from the meta-analysis at the specified tier, and D is the number of probands

from the meta-analysis not at the specified tier. Statistically significant differences at the 0.05

level were determined by one-way ANOVA with Tukey's post-hoc analysis.

HADDS Developmental Delay Severity Scale (HADDS-DDSS)

The four-tier HADDS-DDSS assesses for delayed development and neuroanatomical findings in

individuals at or greater than three years of age (**Table S7**). Higher-numbered tiers indicates

increasingly delayed development, persistent limitations in motor and language abilities, and

neuroanatomical alterations. Tier 1 includes individuals who achieved first word (12 months)

and independent walking (18 months) milestones appropriately for age and without regression.

Tier 2 includes individuals who achieved independent walking and first word milestones by three

years of age. Tier 3 includes individuals who achieved independent walking or first word by

three years of age with subsequent loss of the acquired skills or did not achieve the milestones by

three years of age. Tier 4 includes individuals with neither speech nor independent walking

achieved after three years of age. If information for the developmental milestones is limited, then

the presence or absence of cerebellar anomalies is considered a secondary factor. If either motor

or language developmental milestone is unavailable and brain imaging results are not available,

then the proband is excluded from the analysis. Statistically significant differences at the 0.05

level were determined by one-way ANOVA with Tukey's post-hoc analysis. OR for each

severity tier were calculated using the same formula as the HADDS-DBS.

RESULTS

General features, demographics, and inheritance in HADDS

Demographics, growth parameters, genotype, and phenotype findings in our cohort were

evaluated with a comparative literature meta-analysis (**Fig.1A-D, Table S3**). Growth parameters

reveal below-average height and weight for individuals with HADDS (**Table S3**). However,

head circumference both at birth and recent assessment remained within average parameters.

Gender distribution was equivalent between our cohort and the meta-analysis (OR=1.19)

(**Fig.1A**). More than 90% of affected individuals were identified at <18 years old (**Fig.1B**). Self-

reported ethnicity in our study included European (76%), African (3%), East Asian (3%), Middle

Eastern (6%), and multiethnic (12%) descent. Ethnicity was reported in <45% of individuals in

the meta-analysis, but there was overall a similar distribution (**Fig.1C**).

The inheritance pattern for the majority of *EBF3* pathogenic variants was *de novo* in 79% of our

study and 88% of the meta-analysis (OR=0.50, **Fig.1D**). Mendelian inheritance of a pathogenic

EBF3 variant was previously reported in one affected mother and son pair[23], and inheritance of a

pathogenic *EBF3* variant from an unaffected parent with mosaicism was reported in an

unaffected mother and two affected children[12]. In our study, we identified Mendelian inheritance

in four affected individuals from three unrelated families (OR=4.55) and mosaicism in the

parents of two affected and unrelated individuals (OR=1.03) (**Fig.1D**).

Pathogenic *EBF3* variants and genomic deletions

We identified 13 individuals with noncoding *EBF3* variants and genomic deletions (**Fig.1E**) and

46 individuals with coding variants (**Fig.1F**). The majority of *EBF3* variants cluster within the

N-terminal DNA binding domain with 27% of pathogenic variants located within five amino

acids of the zinc finger motif (ZNF, amino acid 157-170, OR=2.2), a region critical for

stabilizing the interaction of EBF3 with the DNA target[31]. A prior *in vitro* study of the

paralogous EBF1 showed that disrupting conserved residues in the ZNF, including p.Arg163,

abolished DNA-binding activity[31]. Intriguingly, we identified the most frequent recurrent *EBF3*

variants affected the p.Arg163 codon in eight unrelated individuals with p.Arg163Gln

(c.488G>A) in three, p.Arg163Pro (c.488G>C) in two, p.Arg163Trp (c.487C>T) in two, and

p.Arg163Leu (c.488G>T) in one individual. Other recurrent variants affected the p.Arg209

codon in six, p.Arg206 codon in five, p.Cys161 codon in three, and p.His157 codon in three

individuals. The recurrent variants are likely due to the position of the affected nucleotides in

CpG-dinucleotide islands, which are mutational hotspots underlying over one-third of *de novo*

missense variants associated with human diseases[32,33]. Other potential mechanisms include the

selfish spermatogonial selection process[34], the error-prone replication hypothesis[35], or a selection

bias for similar phenotypes.

Common neurologic features in HADDS

203   Prevalent neurologic features identified in the cohort include delayed development, motor

204   incoordination, perturbed expressive language, altered sensory processing, neuropsychiatric

205   comorbidities, and cerebellar alterations. Consistent between our cohort and the meta-analysis,

206   we found that more than 95% of affected individuals had delayed motor development and more

than 80% had delayed expressive language development (OR=3.32, **Fig.2A**). Intriguingly, we

found discordance between expressive and receptive language development with 36% (CI[16,

56]) of individuals diagnosed with HADDS able to follow one-step verbal commands on time

and 37% (CI[15, 58]) able to follow two-step verbal commands on time. This finding suggests

that individuals with HADDS have a greater prevalence of delayed expressive language

development, potentially related to motor incoordination and speech apraxia due to cerebellar

dysfunction[36,37], but maintain age appropriate receptive language development. Another common

language finding was dysarthria (30%, CI[15, 46]), which was identified in a similar frequency

in the meta-analysis (OR=1.68, **Fig.2B**). We also identified speech apraxia in 27% (CI[12, 42],

OR=6.00, **Fig.2B**) and weak or absent cry in 97% (CI[91, 103]) of affected individuals in our

study, a higher prevalence than previously reported (OR 512, **Fig.2B**).

Consistent with the namesake features, we found a high prevalence of hypotonia (100%) and

ataxia (84%, CI[73, 97], OR=3.92) in our cohort (**Fig.2C**). The phenotypic analysis in our cohort

also revealed a high frequency of motor incoordination (70%, CI[54, 85], OR=10.73),

hypomimia (73%, CI[57, 88], OR=12.44), and hypertonia (42%, CI[26, 59], OR=24.32), but

these features were reported in less than 20% of affected individuals in the meta-analysis

(**Fig.2C**). Additional neurologic findings include an increased risk of seizures (36%, CI[20, 53],

OR=18.86), febrile seizures (27%, CI[12, 42], OR=3.875), and abnormalities on

electroencephalogram (EEG) recordings (18%, CI[5, 31] OR=1.04) (**Fig.2D**). Intriguingly, we

found that altered sensory processing is a cardinal feature of HADDS. All individuals in our

study had chronic decreased pain response, but a smaller prevalence was identified in the meta-

analysis (18%, CI[5, 30]), potentially due to incomplete ascertainment (**Fig.2E**). Other sensory

findings in our cohort include repetitive scratching behaviors (58%, CI[41, 74]), hyperacusis

(48%, CI[31, 66]), and discomfort with loud noises (45%, CI [28, 62]) (**Fig.2E**), none of which

were previously reported.

Neuropsychiatric comorbidities are commonly observed in affected individuals with HADDS.

We found that these features occur at a higher frequency than previously reported. Prevalent

neuropsychiatric comorbidities in HADDS include autistic features (70%, CI [54, 85],

OR=10.73), formal autism diagnosis (27%, CI [12, 42], OR=8.82), attention deficit and

hyperactivity disorder (ADHD, 18%, CI [5, 31], OR=1.03), and emotional lability (18%, CI [5,

31], OR=3.56) (**Fig.2F**). Common autistic features identified in our study include stereotypies

(67%, CI [51, 83], OR=15.0), poor eye contact (48%, CI [31, 66]), attraction to lights (24%, CI

[10, 39]), and aversion to crowds (21%, CI [7, 35]) (**Fig.2G**). In contrast, the meta-analysis

identified autistic features, autism, or autism spectrum disorder in 8-18% of individuals (**Fig.2F**).

Consistent with the neurodevelopmental impairments, neuroimaging studies in our cohort

revealed altered cerebellar structure in 33% (CI [17, 49]) of individuals, compared to 24% (CI [9,

38]) in the meta-analysis (OR=1.63, **Fig.2H**). Cerebellar vermian hypoplasia was observed in

30% (CI [15, 46]) of individuals in our study, compared to 18% (CI [5, 30]) in the meta-analysis

(OR=2.03), and is the most common neuroanatomical finding. Less frequent findings in our

cohort include prominent ventricles (9%, CI [-1, 19]), abnormal corpus callosum (two

individuals), delayed myelination (one individual), hydrocephalus (one individual), and partial

rhomboencephalosynapsis (one individual) (**Fig.2H**).

Non-neurological findings in HADDS

Other features commonly seen in HADDS include craniofacial, gastrointestinal, ophthalmologic, genitourinary, and musculoskeletal involvement. In our study, 88% (CI[77, 99]) of individuals had at least one craniofacial feature (OR=0.97). Recurrent features included deep-set eyes or prominent forehead (64%, CI[47, 80]), ear abnormalities (39%, CI[22, 56]), mouth anomalies (42%, CI[26, 59]), straight eyebrows (27%, CI[12, 42]), tall forehead (21%, CI[7, 35]), long face (18%, CI[5, 31]), pointed chin (21%, CI[7, 35]), myopathic facies (21%, CI[7, 35]), triangular facies (9%, CI[-1, 19]), anteverted nares (9%, C [-1, 19]), and short, broad chin (6%, CI[-2, 14]) (**Fig.3A**). Ophthalmologic findings also occur frequently in HADDS with strabismus reported in 88% (CI[77, 99]) of individuals in our cohort compared to 62% (CI[45, 78]) in the meta-analysis (OR=4.49) and nystagmus reported in 6% (CI[2, 14], OR=1.03) of individuals in both our cohort and the meta-analysis (**Fig.3B**).

In our cohort, we identified a high frequency of constipation (88%, CI[77, 99], OR=54.38), feeding difficulties (88%, CI[77, 99], OR=24) and gastroesophageal reflux (58%, CI[41, 74], OR=10), compared to meta-analysis frequencies of 12-21% for these features (**Fig.3C**). Genitourinary complications are frequently observed in both males and females. In our cohort, we found that recurrent urinary tract infections (UTIs) occurred in 48% (CI[31, 66]) of individuals compared to 24% (CI[9, 38]) in the meta-analysis (OR=3.06), with a higher prevalence in females (**Fig.3D**). Anatomical findings in our study included abnormal kidney or bladder anatomy (36%, CI[20, 53], OR=3.31), vesicoureteric reflux (18%, CI[5, 31], OR=0.62), abnormal external genitalia (9%, CI[-1, 19], OR=0.47), and cryptorchidism (6%, CI[-2, 14], OR=0.48) (**Fig.3D**), which had a similar frequency in the meta-analysis. Female reproductive

organ anomalies were seen in 6% of the meta-analysis (**Table S2**) but were not observed in our

cohort (**Table S1**). The most prevalent musculoskeletal finding was short stature, which we

identified in 39% (CI[23, 56]) of individuals in our cohort and was similarly reported in 41%

(CI[24, 58] of the meta-analysis (OR=0.71). Other musculoskeletal dysmorphisms occurred in

less than 10% of individuals in our cohort and in the meta-analysis (OR≤3). Dysmorphisms

included clinodactyly (9%, CI[-1,19]), pectus excavatum (6%, CI[-2,14]), and low muscle mass

(6%, CI[-2,14]) (**Fig.3E**). Short digits were also reported in two individuals in the literature

(**Table S2**).

Diagnostic utility of the HADDS developmental and behavioral scale

To aid clinical recognition and expedited diagnosis of HADDS, we developed a diagnostic

assessment scale (HADDS-DBS) to identify individuals "at risk" for pathogenic *EBF3* variants

and prioritize *EBF3*-specific gene testing. We designed the HADDS-DBS to address the

challenges of assessing children in various age groups and developed the scale for use by

caregivers and healthcare providers. The HADDS-DBS can be used to screen for *EBF3*

pathogenic variant "risk" in children as young as 15-months old and takes the form of a symptom

questionnaire comprised of 21 clinical features (**Table 1**). Based on the number of major and

minor clinical features identified in the questionnaire, the HADDS-DBS assigns one of four tiers

based on increasing risk of carrying a pathogenic *EBF3* variant with recommendations for

molecular testing (**Table S6**). When applied to our cohort, the HADDS-DBS identified 30

individuals as Tier-4, two individuals as Tier-3, and one individual as Tier-2 (**Fig.4A, Table S8**).

When applied to the 34 individuals in the meta-analysis, the HADDS-DBS identified three

individuals as Tier-4 (OR=103), seven individuals as Tier-3 (OR=0.25), 12 individuals as Tier-2

(OR=0.06), and 12 individuals as Tier-1 (**Fig.4A**, **Table S9**). These differences in phenotypic

stratification between our cohort and the meta-analysis reflects the variability in phenotypic

ascertainment across different clinical sites and highlights the value of comprehensive clinical

phenotyping (**Fig.4A**, $p<0.001$, F=102.04).

To determine the specificity of the HADDS-DBS, we applied the diagnostic scale to three

neurodevelopmental disorders with similar clinical features (hypotonia, delayed development,

ataxia, and variable cognitive impairment), but do not have *EBF3* gene alterations (**Table S10**).

Three studies were selected that described eight individuals with Harel-Yoon syndrome (HYS

[MIM: 617183]) caused by *ATAD3A* variants[38], 20 individuals with Mental Retardation and

Microcephaly with Pontine and Cerebellar Hypoplasia (MICPCH [MIM: 300749]) caused by

CASK variants[39], and 15 individuals with Takenouchi-Kosaki syndrome (TKS [MIM: 616737])

caused by *CDC42* variants[40]. Of the eight individuals with HYS[38], two were classified Tier-1

(25%) and six as Tier-2 (75%). For the 20 individuals with MICPCH[39], 10 were classified as

Tier-1 (50%) and 10 as Tier-2 (50%). Finally, of the 15 individuals with TKS[40], all were

classified as Tier-1 (100%). Statistical comparisons with one-way ANOVA reveal a significant

difference for the phenotypic stratification of the HADDS meta-analysis compared to individuals

with either HYS, MICPCH, or TKS (**Fig.4A**, $p<0.051$, F=8.34) with sensitivity=0.29,

specificity=1.0, PPV=1.0, and NPV=0.64. In contrast, comparison of our HADDS cohort to

individuals with either HYS, MICPCH, or TKS reveal sensitivity=0.97, specificity=1.0,

PPV=1.0, and NPV=0.97, consistent with the comprehensive phenotypic ascertainment in our

study cohort.

Correlating *EBF3* variant type with severity of motor and language delays

We used the HADDS-DDSS to identify molecular phenotype groups with differential risk of

motor and language impairments (**Table S7**). Of the combined 42 individuals diagnosed with

HADDS who were more than three years old, we classified seven individuals as Tier-1, 18 as

Tier-2, eight as Tier-3, and nine as Tier-4. We examined the HADDS-DDSS classification

relative to the *EBF3* variant type (noncoding, gene deletion, or coding) and the location of

coding variants relative to the ZNF. For the seven individuals with *EBF3* noncoding variants or

gene deletions, one was classified as Tier-1, three as Tier-2, one as Tier-3, and two as Tier-4

severity (**Fig.4B**). For the 21 individuals with *EBF3* coding variants located outside of the ZNF,

six were classified as Tier-1, 11 as Tier-2, three as Tier-3, and one as Tier-4. Finally, for the 14

individuals with *EBF3* coding variants located in the ZNF region, the HADDS-DDSS classified

four individuals as Tier-2, four as Tier-3, and six as Tier-4 (**Fig.4B**). A significant correlation

was identified between increased clinical severity and coding variants within five amino acids of

the ZNF (**Fig.4B**, $p<0.01$, F=7.72). Intriguingly, more than 25% of the individuals with HADDS

have *EBF3* variants located within the ZNF, which may reflect a selection bias for consistent

phenotypes or increased clinical testing due to more pronounced phenotypic features.

DISCUSSION

Despite increased information available regarding *EBF3*-related HADDS, the need remains for a

clear understanding of the phenotypic variability and how the *EBF3* genotype affects specific

clinical findings. Our comprehensive phenotyping in a large cohort of individuals molecularly

diagnosed with pathogenic *EBF3* variants allow for an expanded understanding of the variation

in clinical outcomes and improved clinical management (**Fig.S1**). Neuroanatomical studies

reveal that the cerebellum is the most common structure perturbed in individuals with *EBF3*

pathogenic variants. Extensive anatomical and functional studies revealed critical roles for the

cerebellum in motor and non-motor functions, including language, cognition, and social

behavior[37]. Thus, many of the features observed in individuals with HADDS may be consistent

with cerebellar dysfunction. To further understand the relationship between the *EBF3* molecular

phenotype and clinical outcome, we designed two assessment scales that determine an

individual's risk for having a pathogenic *EBF3* variant (HADDS-DBS) and their degree of

functional impairments (HADDS-DDSS). Our findings validated the clinical utility of the scales,

revealed critical insights into the *EBF3* genotype-phenotype map, and demonstrated that

molecular phenotypes correlate with quantitative clinical findings (**Fig.4**).

The ability to broadly predict the risk of a pathogenic *EBF3* variant advances clinical diagnosis,

facilitates prognosis, and aids in resolving *EBF3* variants of uncertain significance (VUS).

Moreover, we were interested in determining whether the *EBF3* molecular phenotype could

provide insights into the variation of clinical outcomes. By integrating the HADDS-DDSS

assessment with the *EBF3* molecular phenotype, we found that missense variants within the ZNF

define a subgroup of individuals at risk for persistent motor and language impairments. This

finding may be related to a stronger dominant-negative effect resulting from *EBF3* missense

variants located affecting the ZNF. Haploinsufficient non-coding *EBF3* variants resulting in

nonsense mediated decay or *EBF3* gene deletion appears to correlate with intermediate clinical

outcomes. However, this interpretation is limited by the small sample size. Furthermore, the

variation in clinical outcomes associated with *EBF3* variants raises the possibility that subtle

impairments in EBF3 function or protein levels may result in a limited subset of HADDS

features or manifest primarily as a non-syndromic ASD or non-specific developmental delay.

This hypothesis is supported by prior work identifying *EBF3* as one of ~1,000 genes with

significant enrichment of *de novo* loss-of-function variants in individuals with ASD and

DD/ID[26].

By using the largest set of clinically annotated *EBF3* variants examined to date, we strengthen

prior findings by showing that HADDS-associated *EBF3* variants are related to a wide spectrum

of clinical outcomes, identified Mendelian transmission of potentially hypomorphic variants, and

expanded our understanding of cardinal diagnostic features. Furthermore, we found that the

distribution of pathogenic *EBF3* variants remains predominantly in the DNA-binding domain,

suggesting that variants in the C-terminal domains are less damaging to EBF3 function. Of note,

we found that *EBF3* variants affecting the ZNF are associated with an increased risk of persistent

motor and/or language impairments. These differential risk profiles may account for the

increased frequency and severity of *de novo* missense variants in the ZNF and surrounding

nucleotides, whereas milder variants resulting in subtle impairments in EBF3 function may not

be identified primarily on the basis of a clinical HADDS diagnosis.

Despite advances in our recognition of *EBF3*-related HADDS, the neurobiological basis of

differential risk profiles based on *EBF3* variant type and location remains to be elucidated. As

EBF3 functions through homo- or heterodimerization, the missense variants may result in a

dominant-negative effect that alters the function of the wildtype EBF3 protein and reduces the

overall amount of functional EBF3 dimers. This would result in increased perturbations to the

EBF3-dependent transcriptional regulation in the developing nervous system. In contrast, *EBF3*

single gene deletion or non-coding variants may preserve functional EBF3 dimers but result in an

overall reduction of EBF3 available to regulate developmental processes. Ideally, a

comprehensive analysis would include the effect of variation on EBF3's protein-protein

interactions, subcellular localizations, regulation of target gene expression profiles, and nuclear

function. The increasing recognition of individuals at risk for pathogenic *EBF3* variants and new

high-throughput assays might make such datasets available for expanded analysis of the

neurobiological consequences of the molecular phenotype.

In conclusion, we show that pathogenic *EBF3* variants cause a broad spectrum of phenotypic

severity. This expands our understanding of cardinal clinical features in HADDS. Through the

integration of the HADDS-DBS and HADDS-DDSS assessments with the molecular phenotype,

we found that a subset of individuals with *EBF3* missense variants affecting the ZNF have a

higher risk of persistent functional impairments. Furthermore, the phenotypic overlap between

HADDS with ASD and DD/ID suggest that subtle EBF3 dysfunction may result in non-

syndromic features and that HADDS may share common pathogenic mechanisms with ASD and

DD/ID. Longitudinal follow-up of these individuals in conjunction with new prospective

recruitment efforts will be needed to comprehensively delineate the *EBF3* genotype-phenotype

map, which may lead to promising avenues for therapeutic interventions.

DECLARATION OF INTERESTS

The authors declare no competing interests.

ACKNOWLEDGEMENTS

We thank the families for participation in this study, Keri Ramsey with clinical assistance, and Mingshan Xue, Sahana Murthy, Harim Delgado-Seo, Maimuna Paul, and Joshi Stephen for manuscript feedback. There is no targeted funding for this study. H.T.C. is partially supported from NIH-1DP5OD026428. C.D. is supported by the BCM Medical Scientist Training Program. N.D.B. was supported by the Autism Science Foundation. V.L. is supported by The Gordon and Mary Cain Foundation and Annie and Bob Graham. EB and GZ are members of the European Reference Network for Rare Neurological Diseases-Project #739510. EB, EMV, and GZ are supported by the Ministry of Health (Ricerca Corrente 2020, Ricerca Finalizzata NET-2013-02356160).

WEB RESOURCES

OMIM, https://www.omim.org

GenBank, https://www.ncbi.nlm.nih.gov/genbank

Centers for Disease Control and Prevention growth charts,

https://www.cdc.gov/growthcharts/index.htm

World Health Organization growth charts, https://www.who.int/childgrowth/standards/en/

gnomAD-Genome Aggregation Database, https://gnomad.broadinstitute.org/

CADD-Combined Annotation Dependent Depletion, https://cadd.gs.washington.edu/snv

UCSC Genome Browser (GRCh37/hg19), https://genome.ucsc.edu/cgi-bin/hgGateway

DATA AND CODE AVAILABILITY

The de-identified data supporting the current study have not been deposited in a public repository but are available from the corresponding author on request.

REFERENCES

1. Decker, T., Pasca di Magliano, M., McManus, S., Sun, Q., Bonifer, C., Tagoh, H., and

Busslinger, M. (2009). Stepwise activation of enhancer and promoter regions of the B cell

commitment gene Pax5 in early lymphopoiesis. Immunity *30*, 508–520.

2. Maier, H., Ostraat, R., Gao, H., Fields, S., Shinton, S.A., Medina, K.L., Ikawa, T., Murre, C.,

Singh, H., Hardy, R.R., et al. (2004). Early B cell factor cooperates with Runx1 and mediates

epigenetic changes associated with mb-1 transcription. Nat. Immunol. *5*, 1069–1077.

3. Treiber, T., Mandel, E.M., Pott, S., Györy, I., Firner, S., Liu, E.T., and Grosschedl, R. (2010).

Early B cell factor 1 regulates B cell gene networks by activation, repression, and transcription-

independent poising of chromatin. Immunity *32*, 714–725.

4. Green, Y.S., and Vetter, M.L. (2011). EBF proteins participate in transcriptional regulation of

Xenopus muscle development. Dev. Biol. *358*, 240–250.

5. Kuriki, M., Sato, F., Arai, H.N., Sogabe, M., Kaneko, M., Kiyonari, H., Kawakami, K.,

Yoshimoto, Y., Shukunami, C., and Sehara-Fujisawa, A. (2020). Transient and lineage-restricted

requirement of Ebf3 for sternum ossification. Development *147*,.

6. Wang, S.S., Lewcock, J.W., Feinstein, P., Mombaerts, P., and Reed, R.R. (2004). Genetic

disruptions of O/E2 and O/E3 genes reveal involvement in olfactory receptor neuron projection.

Development *131*, 1377–1388.

7. Sugiyama, T., Omatsu, Y., and Nagasawa, T. (2019). Niches for hematopoietic stem cells and

immune cell progenitors. Int. Immunol. *31*, 5–11.

8. Liberg, D., Sigvardsson, M., and Akerblad, P. (2002). The EBF/Olf/Collier family of

transcription factors: regulators of differentiation in cells originating from all three embryonal

germ layers. Mol. Cell. Biol. *22*, 8389–8397.

9. Chiara, F., Badaloni, A., Croci, L., Yeh, M.L., Cariboni, A., Hoerder-Suabedissen, A.,

Consalez, G.G., Eickholt, B., Shimogori, T., Parnavelas, J.G., et al. (2012). Early B-cell factors 2

and 3 (EBF2/3) regulate early migration of Cajal-Retzius cells from the cortical hem. Dev. Biol.

365, 277–289.

10. Fulp, C.T., Cho, G., Marsh, E.D., Nasrallah, I.M., Labosky, P.A., and Golden, J.A. (2008).

Identification of Arx transcriptional targets in the developing basal forebrain. Hum. Mol. Genet.

17, 3740–3760.

11. Garel, S., Garcia-Dominguez, M., and Charnay, P. (2000). Control of the migratory pathway

of facial branchiomotor neurones. Development *127*, 5297–5307.

12. Harms, F.L., Girisha, K.M., Hardigan, A.A., Kortüm, F., Shukla, A., Alawi, M., Dalal, A.,

Brady, L., Tarnopolsky, M., Bird, L.M., et al. (2017). Mutations in EBF3 Disturb Transcriptional

Profiles and Cause Intellectual Disability, Ataxia, and Facial Dysmorphism. Am. J. Hum. Genet.

100, 117–127.

13. Lewandowski, R.C., Kukolich, M.K., Sears, J.W., and Mankinen, C.B. (1978). Partial

deletion 10q. Hum Genet *42*, 339–343.

14. Shapiro, S.D., Hansen, K.L., Pasztor, L.M., DiLiberti, J.H., Jorgenson, R.J., Young, R.S., and

Moore, C.M. (1985). Deletions of the long arm of chromosome 10. Am J Med Genet *20*, 181–

196.

15. Mehta, L., Duckett, D.P., and Young, I.D. (1987). Behaviour disorder in monosomy 10qter.

J. Med. Genet. *24*, 185–186.

16. Tanabe, S., Akiba, T., Katoh, M., and Satoh, T. (1999). Terminal deletion of chromosome

10q: clinical features and literature review. Pediatr Int *41*, 565–567.

17. Irving, M., Hanson, H., Turnpenny, P., Brewer, C., Ogilvie, C.M., Davies, A., and Berg, J.

(2003). Deletion of the distal long arm of chromosome 10; is there a characteristic phenotype? A

report of 15 de novo and familial cases. Am. J. Med. Genet. A *123A*, 153–163.

18. Chao, H.-T., Davids, M., Burke, E., Pappas, J.G., Rosenfeld, J.A., McCarty, A.J., Davis, T.,

Wolfe, L., Toro, C., Tifft, C., et al. (2017). A Syndromic Neurodevelopmental Disorder Caused

by De Novo Variants in EBF3. Am J Hum Genet *100*, 128–137.

19. Sleven, H., Welsh, S.J., Yu, J., Churchill, M.E.A., Wright, C.F., Henderson, A., Horvath, R.,

Rankin, J., Vogt, J., Magee, A., et al. (2017). De Novo Mutations in EBF3 Cause a

Neurodevelopmental Syndrome. Am. J. Hum. Genet. *100*, 138–150.

20. Lopes, F., Soares, G., Gonçalves-Rocha, M., Pinto-Basto, J., and Maciel, P. (2017). Whole

Gene Deletion of EBF3 Supporting Haploinsufficiency of This Gene as a Mechanism of

Neurodevelopmental Disease. Front Genet *8*, 143.

21. Blackburn, P.R., Barnett, S.S., Zimmermann, M.T., Cousin, M.A., Kaiwar, C., Pinto E Vairo,

F., Niu, Z., Ferber, M.J., Urrutia, R.A., Selcen, D., et al. (2017). Novel de novo variant in EBF3

is likely to impact DNA binding in a patient with a neurodevelopmental disorder and expanded

phenotypes: patient report, in silico functional assessment, and review of published cases. Cold

Spring Harb Mol Case Stud *3*, a001743.

22. Tanaka, A.J., Cho, M.T., Willaert, R., Retterer, K., Zarate, Y.A., Bosanko, K., Stefans, V.,

Oishi, K., Williamson, A., Wilson, G.N., et al. (2017). De novo variants in EBF3 are associated

with hypotonia, developmental delay, intellectual disability, and autism. Cold Spring Harb Mol

Case Stud *3*,.

23. Beecroft, S.J., Olive, M., Quereda, L.G., Gallano, M.P., Ojanguren, I., McLean, C.,

McCombe, P., Laing, N.G., and Ravenscroft, G. (2019). Cylindrical spirals in two families:

Clinical and genetic investigations. Neuromuscul. Disord.

24. Harkness, J.R., Beaman, G.M., Teik, K.W., Sidhu, S., Sayer, J.A., Cordell, H.J., Thomas,

H.B., Wood, K., Stuart, H.M., Woolf, A.S., et al. (2020). Early B-cell Factor 3-Related Genetic

Disease Can Mimic Urofacial Syndrome. Kidney Int Rep *5*, 1823–1827.

25. Hildebrand, M.S., Jackson, V.E., Scerri, T.S., Van Reyk, O., Coleman, M., Braden, R.O.,

Turner, S., Rigbye, K.A., Boys, A., Barton, S., et al. (2020). Severe childhood speech disorder.

Neurology *94*, e2148.

26. Samocha, K.E., Robinson, E.B., Sanders, S.J., Stevens, C., Sabo, A., McGrath, L.M.,

Kosmicki, J.A., Rehnström, K., Mallick, S., Kirby, A., et al. (2014). A framework for the

interpretation of de novo mutation in human disease. Nat. Genet. *46*, 944–950.

27. Flegal, K.M., and Cole, T.J. (2013). Construction of LMS parameters for the Centers for

Disease Control and Prevention 2000 growth charts (US Department of Health and Human

Services, Centers for Disease Control and …).

28. Kuczmarski, R.J. (2002). 2000 CDC Growth Charts for the United States: methods and

development (Department of Health and Human Services, Centers for Disease Control and …).

29. World Health Organization (2007). WHO child growth standards: methods and development

(World Health Organization).

30. Milestone Checklist.

31. Hagman, J., Gutch, M.J., Lin, H., and Grosschedl, R. (1995). EBF contains a novel zinc

coordination motif and multiple dimerization and transcriptional activation domains. EMBO J.

14, 2907–2916.

32. Magewu, A.N., and Jones, P.A. (1994). Ubiquitous and tenacious methylation of the CpG

site in codon 248 of the p53 gene may explain its frequent appearance as a mutational hot spot in

human cancer. Mol. Cell. Biol. *14*, 4225–4232.

33. Mancini, D., Singh, S., Ainsworth, P., and Rodenhiser, D. (1997). Constitutively methylated

CpG dinucleotides as mutation hot spots in the retinoblastoma gene (RB1). Am. J. Hum. Genet.

61, 80–87.

34. Goriely, A., McGrath, J.J., Hultman, C.M., Wilkie, A.O.M., and Malaspina, D. (2013).

"Selfish spermatogonial selection": a novel mechanism for the association between advanced

paternal age and neurodevelopmental disorders. Am J Psychiatry *170*, 599–608.

35. Dhokarh, D., and Abyzov, A. (2016). Elevated variant density around SV breakpoints in

germline lineage lends support to error-prone replication hypothesis. Genome Res *26*, 874–881.

36. Bolceková, E., Mojzeš, M., Van Tran, Q., Kukal, J., Ostrý, S., Kulišťák, P., and Rusina, R.

(2017). Cognitive impairment in cerebellar lesions: a logit model based on neuropsychological

testing. Cerebellum Ataxias *4*, 13.

37. Strick, P.L., Dum, R.P., and Fiez, J.A. (2009). Cerebellum and nonmotor function. Annu.

Rev. Neurosci. *32*, 413–434.

38. Harel, T., Yoon, W.H., Garone, C., Gu, S., Coban-Akdemir, Z., Eldomery, M.K., Posey, J.E.,

Jhangiani, S.N., Rosenfeld, J.A., Cho, M.T., et al. (2016). Recurrent De Novo and Biallelic

Variation of ATAD3A, Encoding a Mitochondrial Membrane Protein, Results in Distinct

Neurological Syndromes. Am. J. Hum. Genet. *99*, 831–845.

39. Moog, U., Kutsche, K., Kortüm, F., Chilian, B., Bierhals, T., Apeshiotis, N., Balg, S.,

Chassaing, N., Coubes, C., Das, S., et al. (2011). Phenotypic spectrum associated with CASK

loss-of-function mutations. J. Med. Genet. *48*, 741–751.

40. Martinelli, S., Krumbach, O.H.F., Pantaleoni, F., Coppola, S., Amin, E., Pannone, L., Nouri,

K., Farina, L., Dvorsky, R., Lepri, F., et al. (2018). Functional Dysregulation of CDC42 Causes

Diverse Developmental Phenotypes. Am. J. Hum. Genet. *102*, 309–320.

 TABLE AND FIGURE LEGENDS

Figure 1: Demographics and molecular findings in *EBF3*-related HADDS

Comparison of demographic and *EBF3* variant findings from 33 individuals in this study and 34

individuals in the meta-analysis. A total of 62 unique individuals were evaluated. **(A)** Gender

distribution was equivalent between our cohort and the meta-analysis. **(B)** The molecularly

confirmed diagnosis of *EBF3*-related HADDS occurred predominantly in the pediatric

population, with more than 90% of affected individuals identified at less than 18 years old. **(C)**

Ethnicity distribution in our study compared to meta-analysis. UNK = Unknown, EUR =

European, AFR = African, EAS = East Asian, SAS = South Asian, MEA = Middle Eastern,

MUL = Multiethnic. **(D)** Inheritance pattern for the majority of identified *EBF3* pathogenic

variants was *de novo*. **(E, F)** There are 13 individuals with noncoding *EBF3* variants and

genomic deletions **(E)** and 46 individuals with coding variants **(F)**. Published variants are shown

in magenta, variants unique to this study are in purple, and variants shared between our cohort

and the literature are in black. DBD is the DNA binding domain, ZNF is the zinc finger motif,

IPT/TIG stands for immunoglobin-like, plexins, transcription factors/transcription factor

immunoglobin, fold structure, and HLH is the helix-loop-helix region.

Figure 2: Frequency of neurological features in *EBF3*-related HADDS

From the combined analysis of 62 unique individuals, we identified distinct common neurologic

features associated with pathogenic *EBF3* variants. Findings from our cohort are shown in solid

bars and findings from the meta-analysis are shown in the outlined bars. Data shown as percent

of individuals with the feature. Sample sizes are shown above each bar. Females are shown in

red and males in blue. **(A)** Percent of individuals reporting language and gross motor

developmental milestones as either on time (green) or delayed (purple) reveal prevalent motor

and language delays. **(B)** Other language findings in our cohort include speech delay, weak or

absent crying during infancy, speech apraxia, and dysarthria. **(C)** Frequent motor findings

observed in affected individuals include hypotonia, motor delay, ataxia, motor incoordination,

hypomimia, and hypertonia. **(D)** A risk of seizures, febrile seizures, and abnormal EEG findings

are observed in our cohort. **(E)** Sensory alterations are prevalent in HADDS with frequent

findings of decreased pain response, repetitive itching behaviors, hyperacusis, and aversion to

loud noise. **(F)** Prevalent neuropsychiatric comorbidities in HADDS include autistic features,

autism, attention deficit and hyperactivity disorder (ADHD), and emotional lability. **(G)**

Common autistic features identified in our study include stereotypies, poor eye contact, attraction

to bright lights, and aversion to crowded environments. **(H)** Neuroanatomical findings observed

in our HADDS cohort include cerebellar-vermian hypoplasia (CVH), prominent ventricles,

abnormal corpus callosum, delayed myelination, hydrocephalus, and partial

rhomboencephalosynapsis (RES).

Figure 3: Frequency of non-neurological features in *EBF3*-related HADDS

Other features commonly seen in individuals with HADDS include craniofacial, gastrointestinal,

ophthalmologic, genitourinary, and musculoskeletal involvement. Findings from the current

study are shown in solid bars and findings from the meta-analysis are shown in the outlined bars.

Data is shown as a percentage of individuals with the feature. Sample sizes are shown above

each bar. Females are shown in red and males in blue. **(A)** Craniofacial dysmorphisms are

frequently observed with 84% of affected individuals in our cohort found to have at least one

craniofacial finding. **(B)** Strabismus is the most common ophthalmology finding in individuals

with HADDS. Nystagmus was reported in a subset of individuals. **(C)** Constipation, feeding

difficulties, and gastroesophageal reflux disease (GERD) are common gastrointestinal (GI)

findings identified in affected individuals. **(D)** Genitourinary complications are frequently

observed in both genders with recurrent urinary tract infections (UTIs) and bladder control

problems as the most prevalent findings. Other genitourinary findings include vesicoureteral

reflux (VUR), abnormal bladder and/or kidney anatomy, abnormal external genitalia, and

cryptorchidism **(E)** Musculoskeletal findings include short stature, clinodactyly, pectus

excavatum, and low muscle mass.

Figure 4: Diagnostic assessment scale and genotype-phenotype mapping in *EBF3*-related

HADDS

The HADDS-DBS is a symptom questionnaire designed to identify individuals at risk for

pathogenic *EBF3* variants. **(A)** The HADDS-DBS diagnostic scale was applied to individuals

with molecularly confirmed HADDS enrolled in our study and previously reported in the

literature. The majority of affected individuals enrolled in our study are classified as Tier-4,

whereas the majority of affected individuals in the meta-analysis are classified as Tiers-2 and -3.

For the five probands in our study who were previously published, we used the classification

determined from phenotypic information collected for the study. When applied to three non-

EBF3 related disorders [MIM: 617183, 300749, 616737], the HADDS-DBS primarily classified

the reported individuals as Tier-1. Data is shown as a percentage of each cohort classified in

Tiers-1 through -4. Sample sizes are shown above each bar. Tier-1 is shown in black, Tier-2 is

shown in purple, Tier-3 is shown in blue, and Tier-4 is shown in red. Statistically significant

differences were determined by one-way ANOVA with Tukey's post-hoc analysis. Significance

shown as * *p*<0.05 and *** *p*<0.001. **(B)** The HADDS-DDSS is a symptom assessment scale

designed to determine the severity of neurological limitations. The scale is based on readily

ascertained developmental milestones (age of first word and age of independent ambulation) and

the presence or absence of MRI brain findings. **(i)** The HADDS-DDSS tier assignment is

examined relative to the *EBF3* variant type and location relative to the ZNF motif for genotype-

phenotype mapping. A significant correlation was identified between symptom severity and

coding variants within the ZNF motif (amino acids 157-170) and the flanking five amino acids

(total range 152-175). Data is shown as the percentage of each cohort classified in Tiers-1

through -4. Sample sizes are shown above each bar. Tier-1 is in black, Tier-2 is in purple, Tier-3

is in blue, and Tier-4 is in red. Statistically significant differences were determined by one-way

ANOVA with Tukey's post-hoc analysis. Significance shown as ** $p < 0.01$. **(ii)** Illustration

shows the distribution of *EBF3* missense variants within five amino acids flanking the ZNF

motif.

Table 1: HADDS-Diagnostic Behavioral Scale Questionnaire

MAJOR FEATURE	QUESTION (YES OR NO)
Weak or absent cry	As a baby, does your child seem to rarely cry or cry weakly?
Decreased pain sensitivity	Does your child have a reduced response to painful stimuli, such as needle sticks or falls?
Hypomimia	Does your child have reduced facial expressions, such as difficulty with smiling?
Hypotonia	Does your child have low muscle tone or appear to be "floppy" or more "flexible" as a baby?
Ataxia	Does your child have trouble keeping their balance and coordinating their movements?
Speech delay	Is your child taking longer than 12-15 months to say their first word?
Motor delay	Is your child taking longer than expected to sit, stand, and/or walk with assistance?
MINOR FEATURE	QUESTION (YES OR NO)
Poor eye contact	Is your child less likely to look directly at another person's eyes?
Strabismus	Does your child's eyes have difficulty aligning with each other when looking at an object or person?
Repetitive scratching	Does your child exhibit repetitive scratching when tired or stressed and there is no clear evidence of another trigger (ie, insect bites, allergy)?
Repetitive scratching on back or neck	Does your child exhibit repetitive scratching on the back or neck with no clear evidence of another trigger?
Attraction to lights	Is your child drawn to bright lights?
Disturbed by loud noises	Does your child exhibit discomfort with loud noises?
Aversion to crowds	Does your child exhibit an aversion to crowded environments?
Constipation	Does your child have difficulty emptying their bowels?
Frontal bossing	Does your child have a prominent forehead?
Recurrent urinary tract infections	Does your child have frequent urinary infections?
Difficulty feeding	As a baby, did your child have trouble nursing?
Cerebellar anomaly	Does your child have abnormal cerebellum findings on MRI brain imaging studies?
Gait ataxia	Does your child have trouble keeping their balance while walking?
Short stature	Is your child's height below expected based on parental heights?

642 The HADDS-DBS scale is comprised of seven major and 14 minor features and designed for use

643 by parents and clinicians. Seven of the clinical features are considered "major" due to their high

644 prevalence among the patients in the current cohort. The remaining 14 clinical features are

645 considered "minor".

A. Diagnostic tiers assigned

B. Greater severity associated with Zinc Finger variants

i.

ii.

*p < 0.05
***p < 0.001

**p < 0.01

ZNF = amino acids 157-171