Plasma S1P links to hypertension and biomarkers of inflammation and cardiovascular disease – findings from a translational investigation

Short running title: Plasma S1P and hypertension.

Amra Jujic [a,b,c,d]*, Frank Matthes [d,e]*, Lotte Vanherle [d,e], Henning Petzka [f], Marju Orho-Melander [a], Peter M Nilsson [a,g], Martin Magnusson [a,b,d,h] & Anja Meissner [d,e]

[a] Department of Clinical Sciences, Lund University, Malmö, Sweden

[b] Department of Cardiology, Skåne University Hospital, Malmö, Sweden

[c] Lund University Diabetes Centre, Lund University, Malmö, Sweden

[d] Wallenberg Centre for Molecular Medicine, Lund University, Sweden

[e] Department of Experimental Medical Sciences, Lund University, Sweden

[f] Department of Mathematics, Lund Technical University (LTH), Sweden

[g] Department of Internal Medicine, Clinical Research Unit, Malmö, Sweden

[h] Hypertension in Africa Research Team (HART), North West University Potchefstroom, South Africa

* denotes equal contribution

Correspondence to: Anja Meissner; Klinikgatan 32, SE-22184 Lund, Sweden; anja.meissner@med.lu.se; +46 (0)46 22 20641

Manuscript category: Original Article

Word count: 6117

Abstract

Background: Recent studies identified sphingosine-1-phosphate (S1P) as an important player in immune cell trafficking and vascular dysfunction contributing to the development and progression of overt hypertension. Although targeting S1P signaling revealed therapeutic potential in different experimental hypertension studies, validations of S1P-blood pressure associations in humans are lacking.

Methods and Results: In a translational approach, we explored the associations between plasma S1P, quantified using LC-MS/MS, and blood pressure in a family-based study cohort (Malmö Offspring (MOS) study; N=1026), and in a longitudinally conducted murine hypertension cohort. In MOS, linear multivariate regression analyses showed that plasma S1P associates with increased systolic blood pressure (β=1.06, P=0.015). Study subjects with systolic blood pressure ≥140 mmHg presented with significantly higher S1P plasma concentrations compared to subjects with blood pressure <120 mmHg independent of age and sex. The S1P-blood pressure association was validated in a murine model where plasma S1P increased with systolic blood pressure (r=0.7018, R^2=0.4925; P<0.0001). In a sub-sample of the human study population (N=444), proteomic profiling for markers of inflammation, metabolism and cardiovascular disease was carried out using proximity Extension Assays. Testing S1P associations revealed multiple significant interactions, some of them with marked sex-specificity. Amongst them, interleukin-18, which exerts apparent vascular and immune responses during hypertension and associates to adverse cardiovascular events, strongly correlates with plasma S1P concentrations in females but not males in both humans and mice. *In vitro* and *ex vivo* validation of S1P effects on endothelial and monocytic cells of murine or human origin and resistance arteries isolated from mice disclosed augmented expression of different vascular dysfunction and inflammation markers in response to exogenously added S1P.

Conclusion: Taken together, our translational findings strongly suggest a link between plasma S1P and systolic blood pressure as well as several inflammation and cardiovascular disease biomarkers in humans, encouraging further studies to investigate S1P's potential as a therapeutic target in hypertensive disease.

Introduction

Hypertension is the most common preventable risk factor for cardiovascular disease (CVD) with detrimental implications for target organs including the heart, the kidneys, and the brain. Thus, it is the leading single contributor to all-cause mortality and disability worldwide and a major global cause of premature death.[1] Blood pressure control is therefore considered the gold standard approach to reducing the proportion of population burden of blood pressure-induced CVD. Epidemiological studies reported an existing link between blood pressure and cardiovascular risks at systolic blood pressure levels of as low as 110–115 mmHg and diastolic blood pressure levels of 70–75 mmHg.[1] Besides, increasing blood pressure thresholds of about 20 mmHg systolic or 10 mmHg diastolic blood pressure were shown to double the risk of stroke and heart disease mortality,[2] highlighting the need to intervene early. Following the results of the SPRINT study,[3] both the American College of Cardiology (ACC)/American Heart Association (AHA) and the European Society of Cardiology/European Society of Hypertension (ESC/ESH) have gradually updated their advice on diagnosis and treatment of arterial hypertension. The ACC/AHA guidelines replaced "prehypertension" with "elevated blood pressure" for blood pressure levels 120-129/80-89 mm Hg while the ESC/ESH still defines this category as "normal" blood pressure. However, when comparing the guidelines, the definition of normal (as used in the ACC/AHA guidelines) or optimal blood pressure (as used in the ESC/ESH guidelines) is the same for systolic (<120 mm Hg) and diastolic blood pressure (<80 mmHg).[4, 5] This implementation of lower blood pressure thresholds[5] increases the prevalence of hypertension substantially,[6] with a large young low-risk population that presents with intermediate blood pressure who are recommended healthy lifestyle and antihypertensive treatment according to the new guidelines. However, blood pressure lowering interventions in these low blood pressure spans are most likely only beneficial in parts of the population and broad therapy would without a doubt result in treatment of a substantial amount of people with

no or little benefit. Thus, the incorporation of biomarkers to aid in the selection of patients that would actually respond to antihypertensive therapy (personalized medicine)[7] and for the overall assessment of cardiovascular risk has been proposed.[8-11] Selecting from the numerous inflammation, metabolism and CVD biomarkers that associate with a higher cardiovascular risk might inflict bias, hence the use of biomarkers with association to both blood pressure and established CVD, inflammation and metabolism markers might be advantageous.

A major regulator of vascular functions, inflammatory and metabolic processes relevant to the pathology of hypertension and associated cardiovascular events is the bioactive sphingophospholipid sphingosine-1-phosphate (S1P) and its signaling axis.[12-18] Besides holding cell type- and receptor-specific vasomodulator potency,[12, 14] S1P's chemotactic properties critically regulate immune cell trafficking, specifically T-cell egress and homing.[19] S1P signaling is furthermore involved in immune cell differentiation and polarization[20-22] and in barrier function control,[23, 24] largely affecting lipid and glucose metabolism.[25-27] In preclinical experimental studies, augmented S1P signaling has been linked to several cardiovascular condition including stroke, heart failure, atherosclerosis and hypertension.[12, 13, 15, 28] Small scale biomarker studies associated plasma S1P levels with impaired left ventricular ejection fraction (N=74) and all-cause mortality (N=210) in systolic heart failure patients.[29, 30] In another study, plasma S1P was used to predict the severity of coronary artery atherosclerosis (N=59).[31] With respect to hypertension, S1P's role in disease development and propagation is mainly elusive,[13] despite a few reports that indicate altered sphingolipid metabolism in different forms of experimental hypertension.[15, 28, 32-35] To date, human cohort-based studies investigating associations between S1P plasma concentrations and blood pressure are lacking. Therefore, this study explored the relationship between plasma S1P and blood pressure in a large family-based cohort study, clinically validating results obtained in an experimental murine hypertension model. Additionally, evaluation of S1P associations with established

CVD, inflammation and metabolism biomarkers using multiplex proteomic profiling and *in vitro* and *ex vivo* validation approaches were performed to study novel biomarker associations and to test the suitability of S1P as predictive marker for hypertensive disease.

Methods

Animal Study

The investigations using research animals conform to the EU Directive 2010/63/EU for animal experiments and were conducted in accordance with European Animal Protection laws. All protocols were approved by the institutional ethical committee of Lund University (Dnr. 5.8.18/12637/2017). Commercially available male and female wild-type (WT) C57BL/6N mice were obtained from Taconic (Ejby, Denmark) and housed under standard 12h:12h light-dark cycle with access to food and water *ad libitum*. Mice with a body weight (BW) ≥ 25g were housed in groups of four to five mice per cage. Experimental groups were designed in a way to minimize stress for the animals and to guarantee maximal information using the lowest group size possible using a power calculation with Type I error $\alpha = 0.05$ and Power of $1-\beta > 0.8$ (80%) based on previous studies.[15] Hypertension was induced using angiotensin II (AngII)-releasing osmotic pumps (Alzet-2006, AgnTho's, Sweden). In brief, animals were anaesthetized with isoflurane (IsoFlo® vet 100%, Sweden; 2.5% at 1.5L/min in room air) for subcutaneous pump implantation containing AngII (infusion rate 20ng/kg/min over four weeks). Blood pressure was measured bi-weekly in conscious mice using non-invasive tail-cuff plethysmography (CODA, EMKA, France), starting one week before pump implantation after a training period of 7 days. Weekly blood draws from the vena saphena were performed starting prior to pump implantation (=baseline) at seven days intervals and blood was collected in EDTA-coated tubes (Sarstedt, Germany). At termination, mice were anaesthetized (isoflurane 2.5% at 1.5L/min in

room air) before euthanasia through cervical dislocation. Mesenteric arteries were dissected and immediately processed for RNA isolation using the Trizol method as per manufacturer's instructions and subsequent quantitative real-time PCR.

Human study population

The Malmö Offspring Study (MOS) is an on-going family-based cohort study initiated in March of 2013 with the general aim to map risk factor patterns for family traits of chronic diseases.[36] The study population consists of adult children (MOS-G2) and grandchildren (MOS-G3) of participants from the Malmö Diet and Cancer Study – Cardiovascular Cohort (MDCS-CC) with a mean attendance rate of 47%.[37] Participants were recruited using official register information from the Swedish Tax Agency. Not understanding information in Swedish was the only exclusion criteria. Study half-time was reached in May 2017, with 2645 participants included (1326 children, 1321 grandchildren). The study population used here consisted of children to MDCS-CC participants (N=1326). Subjects with missing data in any of the co-variates used in analyses of associations between S1P and blood pressure were excluded, resulting in 1046 eligible subject (**Supplemental Figure S1**). Ethical approval has been obtained for MOS at the Regional Ethics Committee in Lund (Dnr. 2012/594).

Clinical assessment

Participants height (cm) and weight (kg) were measured in light indoor clothing. Resting blood pressure (mmHg) was measured after 10 minutes of rest in the supine position using an automatic device (Omron). A mean of two readings with one minute apart was calculated. Smoking (yes/no), anti-hypertensive treatment (AHT) and alcohol usage were self-reported in a web-based questionnaire. Diabetes was defined as either self-reported diabetes diagnosis, use of anti-diabetic medication or fasting plasma glucose levels ≥7.0 mmol/L at two separate visits at the research facility.

Laboratory assays

Blood samples were drawn after an overnight fast and analyzed for plasma glucose, creatinine and cystatin C at the Department of Clinical Chemistry, Malmö, which is part of a national standardization and quality control system. Plasma glucose was measured using the HemoCue Glucose System (HemoCue, Ängelholm, Sweden). Plasma creatinine was measured using an enzymatic colorimetric assay with an IDMS-traceable calibrator on the Hitachi Modular P analysis system (Roche, Switzerland). Plasma levels of cystatin C were determined by an automated particle-based immunoassay using Hitachi Modular P analysis system and reagents from DAKO (Dako A/S, Denmark). Relative estimated glomerular filtration rate (eGFR) was calculated as a mean of eGFR derived from creatinine and eGFR derived from cystatin C and reported as $mL/min/1.73\ m^2$.

Proteomic profiling

Proximity Extension Assay (PEA) technique was applied to analyze plasma levels of proteins (Olink Bioscience, Uppsala, Sweden) in a sub-sample of the population (N=444; consecutive subjects from 6th of March 2013 until 17th of June 2015 with complete data on all examinations). Four OLINK panels were analyzed: (1) Inflammation Panel (inflammation panel specifics), (2) Metabolism Panel (metabolism panel specifics), (3) CVD II panel (CVDII panel specifics) and (4) CVD III panel (CVDIII panel specifics), all comprising 92 proteins each within different domains (N = 92x4 = 368). Proteins with ≥15% samples below limit of detection were excluded (N=20 for the Metabolism panel; N=27 for the Inflammation panel; N=7 for the CVDII panel; and N=4 for the CVDIII panel; **Supplemental Table S1**). Additionally, 11 proteins were overlapping between panels thus, only one of each marker was included in linear regression analyses (**Supplemental Table S2**), resulting in 299 proteins that were taken forward to analyses. Validation data and coefficients of variance for all panels is available on the Olink homepage (http://www.olink.com).

S1P plasma quantitation

S1P was extracted by mixing 10 µl of plasma with 90 µl of ice-cold methanol containing 22.2 nM S1P-D7 (Avanti Polar Lipids / Merck, Darmstadt, Germany) as internal standard. After incubation on ice for 30 min precipitate was removed by centrifugation (20,000 g for 10 min at 4 °C). The supernatant was analyzed by LC-MS/MS on a 6495 QQQ instrument (Agilent Technologies, Sweden) essentially as described.[38] Extracts were separated on a 2.1 x 50 mm Acquity UPLC Peptide HSS T3 C18 column (Waters, Sweden) at a flow rate of 0.5 ml/min using eluents (A) water / 0.1 % formic acid / 1 mM ammonium formate, and (B) methanol / 0.1 % formic acid / 1 mM ammonium formate, with a gradient of 20 % A and 80 % B to 100 % B over 2 min, followed by 100 % B for 6 min. By multiple reaction monitoring, MS/MS transitions of m/z 380 to 264 (with 380 to 82 as qualifier) for S1P, and 387 to 271 (with 387 to 82 as qualifier) for S1P-D7, were measured. A calibration curve consisting of 7 concentrations in the range of 0.1 to 2.4 µM S1P in 4 % fatty acid-free BSA was generated in triplicates and measured during each session. Additionally, 3 plasma samples were used as quality control samples and measured during each session to ensure reproducibility.

Ex vivo **and** *in vitro* **testing**

Endothelial cells of murine (bEND.3; ATCC® CRL-2299™) and human (HMEC-1; ATCC® CRL-3243™) origin were cultured in DMEM containing 10% FBS and 1% Penicillin-Streptomycin and MCDB131 supplemented with 10% FBS, 1% Penicillin-Streptomycin, 1% Glutamine (2mM), 1% NEAA, 1% sodium pyruvate, 0.1% Amphotericin B and 0.1% hEGF (10ng/ml), respectively. Cell cultures were maintained at 37°C with 5 % CO_2 and split 1:4 at a seeding density of 10^6 cells. Cells were incubated with 1 µM S1P or vehicle (4% fatty acid-free BSA) for 6, 12 or 24 hrs prior to processing for RNA isolation using the Trizol method as per manufacturer's instructions and subsequent quantitative real-time PCR. Mesenteric arteries isolated from male WT C57Bl/6N mice were cultured in DMEM (Gibco Life Technology,

Sweden) containing 10% FBS (Gibco Life Technology, Sweden) and 1% Penicillin-Streptomycin (Sigma- Aldrich, Sweden) for 24 hrs in the presence of 1 µM S1P (Cayman Chemicals, BioNordika, Sweden) or vehicle (4% fatty acid-free BSA) prior to processing for RNA isolation and subsequent quantitative real-time PCR. Standard biochemical procedures were utilized for experiments involving reverse transcription polymerase chain reaction, quantitative PCR, and ELISA. Methodological details and primer sequences are provided in the data supplement.

Statistical methods

Human study: Comparisons between blood pressure groups were carried out using one-way ANOVA for continuous variables, and $\chi 2$ tests for binary variables. For linear regression analyses of associations between S1P and blood pressure, S1P was z-transformed. For associations between S1P and systolic blood pressure and diastolic blood pressure, linear regression analyses were carried out unadjusted, adjusted for age and sex (*Model 1*), and further adjusted for BMI, eGFR, diabetes status, AHT, smoking status and alcohol usage (*Model 2*). Interaction analyses between age, eGFR, BMI, sex, and S1P were carried out in linear regression analyses using a moderator variable. For associations between S1P and systolic blood pressure ≥120 mmHg; ≥130 mmHg; and ≥140 mmHg, logistic regressions were carried out unadjusted, adjusted for age and sex (*Model 1*), and further adjusted for BMI, eGFR, diabetes status, AHT, smoking status and alcohol usage (*Model 2*). For both linear and logistic regressions, P<0.05 was considered statistically significant.

For associations between proteins from the OLINK-panels and S1P, NPX (log2) data were used and S1P was log2-transformed accordingly as recommended for OLINK data (Measuring protein biomarkers with OLINK). Linear regression analyses were carried out unadjusted and adjusted for age and sex. In order to adjust for multiple testing, a Bonferroni-corrected P-value of 0.05/299 ($P<1.67x10^{-4}$) was applied. Pearson's correlation analyses were carried out with

two-tailed significance testing and computation of exact correlation coefficients (Pearson's r). With correlation matrices being calculated for four different panels, multiple comparison adjustment was performed using Bonferroni-corrected P-values for each correlation matrix individually; ($P \leq 7.81 \times 10^{-4} = 0.05/64$ for the inflammation panel, $P \leq 7.04 \times 10^{-4} = 0.05/71$ for the metabolism panel, $P \leq 5.95 \times 10^{-4} = 0.05/84$ for the CVDII panel and $P \leq 5.75 \times 10^{-4} = 0.05/87$ for the CVDIII panel). The panel-wise correction for multiple comparison avoids a (possible incorrect) rejection of significant correlations when correction is instead based on strict P-values from the large amount of overall 299 markers. For methodological correctness, we always compare to the Bonferroni-corrected significance of all 299 markers used in the linear regression. Both, Pearson's correlation and linear regression analyses were carried out using SPSS 26.0 (IBM).

Animal study: All data are expressed as mean ± SEM, where N is the number of animals. For assessment of differences in blood pressure and plasma S1P levels over time, two-way repeated measure ANOVA followed by Dunnett post hoc testing was performed. For comparison of multiple independent groups, parametric one-way ANOVA test was used, followed by Tukey's post hoc test with exact P value computation. For comparison of two groups a two-tailed unpaired t-test was utilized. Differences were considered significant at error probabilities of $P \leq 0.05$. All statistical analyses were carried out using GraphPad software (Version 8.4.2).

Ex vivo and in vitro studies: All data are expressed as mean ± SEM, where N is the number of animals or independent *in vitro* experiments. For comparison of two groups a two-tailed unpaired t-test was utilized. Differences were considered significant at error probabilities of $P \leq 0.05$. All statistical analyses were carried out using GraphPad software (Version 8.4.2).

Visualization of correlation data: The matrix of pairwise correlations was used to construct a network graph with nodes given by the individual markers and edges drawn between nodes if their corresponding markers show a correlation of absolute value r ≥ 0.3. Edge width was weighted by correlation in absolute values. The network was grouped into clusters using Gephi 0.9.2,[39] which implements clustering by modularity taking edge weight into consideration.[40] Resolution was chosen around the default value of 1.0 so that a maximal modularity score is obtained.

A separate t-SNE [41] analysis on the correlation data considers each variable as a data point with the correlation to all other variables as its features, defining a high-dimensional representation of the variables. We utilized the scikit-learn 0.23.1-implementation of t-SNE in Python (3.8.3) to learn a two-dimensional representation suitable for visualization, which reflects the relation of similar correlations encoded as proximity in the high-dimensional data.

Search strategy for biomarker associations with CVD and inflammatory disease: PubMed searches for relevant studies published within the past five years, using the following combination of Medical Subject Heading (MeSH) terms and text words, with no language limitations: "biomarker name"[MeSH Terms] OR "biomarker name"[All Fields] OR "biomarker name"[All Fields]) AND "CVD"[All Fields] for associations with CVD, and "biomarker name"[MeSH Terms] OR "biomarker name"[All Fields] OR "biomarker name"[All Fields]) AND "inflammatory disease"[All Fields] for associations with inflammatory disease. All observational and clinical studies as well as journal articles were included in the search.

Results

S1P plasma levels associate with systolic blood pressure. S1P plasma levels were quantified in MOS-G2 participants with recorded systolic and diastolic blood pressure information. Characteristics of the study population are presented in **Table 1**. There was a steady significant

increase in S1P plasma levels as systolic blood pressure values increased within cut offs defined as <120 mmHg; ≥120 mmHg; ≥130 mmHg, and ≥140 mmHg (P=0.024; **Table 1**). Subjects with systolic blood pressure ≥140 mmHg were older and presented with significantly higher S1P plasma concentrations, higher BMI, higher diastolic blood pressure, more frequent anti-hypertensive treatment and lower eGFR compared to subjects with blood pressure <120 mmHg. Further, one-way ANOVA analysis revealed no statistically significant differences in plasma levels of S1P between the sexes (P=0.163).

To explore associations between S1P and blood pressure, linear regression analyses were performed revealing that each one standard deviation (1SD) increment of S1P was associated with increasing systolic blood pressure but not diastolic blood pressure when adjusted for age, sex, BMI, eGFR, diabetes status, anti-hypertensive treatment, smoking and alcohol usage (**Supplemental Table S3**). The relationships between S1P and blood pressure were not mediated by age (P=0.975), sex (P=0.495), eGFR (P=0.980), or BMI (P=0.834) as determined by interaction analyses. Further, logistic regression analyses revealed that each 1SD increment of S1P was associated with systolic blood pressure ≥120 mmHg, ≥130 mmHg and ≥140mmHg in adjusted models, with similar odds ratios (**Table 2**).

In order to verify a possible positive correlation between plasma S1P and systolic blood pressure, we longitudinally assessed plasma S1P concentrations in a murine model of slowly developing hypertension (induced by a low-pressor dose of 20ng/kg/min AngII over the course of four weeks). In this model, blood pressure steadily increased and established significance at four weeks after AngII pump implantation (**Figure 1A**). Similar to systolic blood pressure, plasma S1P concentrations were significantly elevated compared to baseline after four weeks of AngII perfusion (**Figure 1B**). Thus, S1P plasma levels presented with a positive linear

relationship to systolic blood pressure in our murine model ($r=0.7018$, $R^2=0.4925$; $P<0.0001$; **Figure 1C**).

Proteomic profiling reveals significant associations between S1P plasma levels and inflammation, metabolism and CVD markers. As hypertension is a major modifiable risk factor for CVD,[12, 13, 15, 28] relates to metabolic disease,[25-27] and its pathophysiology strongly links to immune system activation and inflammation,[19] [20-22] we tested if S1P plasma levels associate to biomarkers of CVD, metabolism and inflammation in a subset of our study population (N=444). We categorized all markers with values above detection limit based on reported associations to cardiovascular or inflammatory diseases. **Figure 2** illustrates an overview of biomarkers detected in each OLINK panel, incorporating all significant S1P correlations after single comparison (grey dots) and multiple comparison testing (black dots). Pearson correlation analysis revealed 23 significant associations between plasma S1P and inflammation panel markers (panel-specific Bonferroni-correction), 21 of which were validated with Bonferroni-correction for testing all 299 markers (**Figure 2A** and **Supplemental Figure 2A**). Altogether 23 metabolism panel markers significantly correlated with plasma S1P (significant after both for panel specific Bonferroni-correction and for testing all 299 markers **Figure 2B** and **Supplemental Figure 2B**). CVDII and CVDIII panels presented with 22 and 24 proteins that correlated significantly with S1P levels (Bonferroni-corrected for individual panels), of which 21 and 16 remained significant after Bonferroni-correction for testing all 299 markers (**Figure 2C/D** and **Supplemental Figure S2C/D**). Overall, Pearson correlations revealed a total of 185 correlations with S1P levels across all panels, of which 92 and 81 remained significant after Bonferroni-correction for the number of panel-specific tests and all 299 tests, respectively (**Supplemental Figure S2**). Correction for age and sex in the linear regression analyses resulted in the same number of significant associations for the individual biomarker panels. A

full list of all unadjusted and age- and sex-adjusted associations in linear regression analyses between S1P and 299 proteins from all four panels (Inflammation, Metabolism, CVDII and CVDIII) is presented in **Supplemental Table S4**.

When applying a cut-off defined as $r \geq 0.3$ or $r \leq -0.3$ to determine the strongest significant Pearson's correlations[42, 43] with plasma S1P in all four panels, a total of 29 markers were extracted that all revealed significant associations with S1P in linear regression analyses. 16 out of 29 strongly correlated markers belong to CVDII and CVDIII panels, 9 to the metabolism panel and 4 to the inflammation panel. **Figure 3** illustrates individual correlation networks of all four OLINK panels. Modularity clustering was applied to all networks as an algorithmic approach to detect communities of markers with strong pairwise correlations within a community and less frequent inter-correlations to markers of other communities. For the inflammation panel, marker clusters can be linked to T-cell homeostasis, immune cell homeostasis and chemotaxis (*cluster A*), neutrophil chemotaxis and angiogenesis (*cluster B*), or regulation of immune responses and association to CVD (*cluster C*), respectively (**Figure 3A**). S1P associations belong to *cluster B* and link to T-cell metabolism and apoptosis. Similarly, modularity clustering of metabolism panel markers detected 3 groups, separating markers associated to cell adhesion (*cluster A*), apoptosis and cellular stress response (*cluster B*), or cellular metabolism (*cluster C*). S1P associations link to markers representing *cluster B* (**Figure 3B**). CVDII panel clusters represent features linking to inflammation and metabolism associated to atherosclerosis, plaque development and cardiovascular events (*cluster A*), angiogenesis (*cluster B*), vascular dysfunction and inflammation (*cluster C*), the latter including all S1P associations of this panel (**Figure 3C**). For the CVDIII panel, marker clusters link to vascular dysfunction, remodeling and inflammation (*cluster A*), endothelial activation and inflammation (*cluster B*) or development of atherosclerosis and cardiovascular events (*cluster C*), respectively (**Figure 3D**). Strongest S1P associations belong to *cluster B*.

Modularity clustering applied to a network constructed from correlation factors of markers of all panels combined, separates markers into five distinct clusters, which can be linked to different processes involved in inflammation and CVD previously reported by clinical and pre-clinical studies (**Figure 4)**. All significant S1P correlations are highlighted in red and associate to clusters 2, 3 and 4, representing markers linked to vascular inflammation, cell adhesion, immune cell metabolism and chemotaxis as well as atherosclerosis and cardiovascular events. To support the significance and stability of the found clusters, an application of a t-distributed stochastic neighbor embedding (t-SNE) visualizes the high dimensional correlation data in two dimensions, revealing an underlying data structure that can be compared to the clusters (**Supplemental Figure 3**). While the positioning of markers within a modularity cluster is arbitrary, the t-SNE locations of markers are meaningful where proximity is explained by similarity in correlations with other markers. The t-SNE arranges the markers into 9 visual groups that were linked to different processes involved in inflammation and CVD. When comparing tSNE groups with the modularity clusters, 81-92% marker overlap was observed between 5 tSNE groups (groups 1- 3, 7, 9; **Supplemental Figure S3**) and clusters 1 to 5 (**Figure S4**). All S1P correlations are associated to groups 7 and 8, resembling marker groups linked to vascular inflammation, immune cell homeostasis and cell adhesion or endothelial inflammation and thrombosis, respectively. Marker overlap with cluster 3 (92%), where most markers can be linked to vascular inflammation, immune cell homeostasis and cell adhesion, largely confirms the meaningfulness of the modularity clusters. Correlation data used for all visualizations are presented as correlation matrices (**Supplemental Figure S4**).

To experimentally test the effect of S1P on marker expression profiles, we performed *in vitro* and *ex vivo* experiments utilizing endothelial cells of murine and human origin that were treated with 1 μM S1P. As illustrated in **Figure 4B-D**, exposure to S1P significantly increased the expression of markers characteristic for angiogenesis and endothelial activation such as PAI

(plasminogen activation inhibitor-1; **Figure 4B**), PECAM-1 (Platelet endothelial cell adhesion molecule 1; **Figure 4C**) and SELP (P-selectin; **Figure 4D**) in cultured human or mouse endothelial cells. All markers presented strongest positive associations with plasma S1P in our study cohort (r=0.409, β=0.409 for PAI-1; r=0.389, β=0.378 for PECAM-1 and r=0.383, β=0.384 for SELP). Remarkably, mesenteric arteries isolated from hypertensive mice presented with augmented PECAM-1 and SELP mRNA expression levels when compared to normotensive controls (**Figure 4E** and **Figure 4F**), strongly suggesting a link between plasma S1P and hypertension-associated vascular dysfunction and inflammation as systolic blood pressure significantly correlated with plasma S1P levels in this model (see **Figure 1C**). Although cross-sectional associations do not allow conclusions on causality, several different markers increased expression after exposure to high S1P concentrations, including small resistance artery Ang-1 (angiopoietin 1), endothelial cell Casp3 (caspase 3), endothelial cell IL18 and monocytic cell CD40 (**Supplemental Figure S5**).

Proteomic profiling discloses sex-specific associated effects of S1P for a subset of markers. As sex-dependent differences have been discussed for various S1P responses,[44] we investigated the possibility of sex-specific S1P associations of all protein markers tested in our study. In single comparisons, we identified 66 significant sex dependent S1P correlations, out of which 32 showed stronger associations with male sex and 34 with female sex (**Supplemental Table S5**). After panel-specific correction for multiple comparisons, the majority of sex-specific differences were dictated by female sex (3 of 4 in the inflammation panel, 3 of 3 in the CVDII panel and 1 of 2 in the metabolism panel), while only one male-specific association remained significant after correction for multiple comparisons in the CVDIII panel. Eight out of ten sex-specific associations were confirmed with strict Bonferroni-correction for all 299 markers (**Table 3**). Amongst them IL18, which exerts apparent vascular and immune responses during hypertension and associates to adverse cardiovascular events, presented with a significantly

stronger correlation in females (r = 0.278 and β = 0.538; P = 1.80×10^{-5}) compared to males (r = 0.157 and β = 0.283; P = 0.021). Investigating the relation between IL18 and S1P in a controlled experimental setting revealed a similar sex-specific difference as evident by higher IL18 plasma levels in female hypertensive mice compared to their male counterparts (**Figure 5A**). Testing a linear relationship between IL18 and plasma S1P disclosed an extremely strong association in female mice (r = 0.9260, R^2 = 0.8574; P < 0.0001) but not in male mice (r = 0.2175, R^2 = 0.0473; P = 0.436) in our model (**Figure 5B**). Similar to observations in our human cohort, plasma S1P responses to blood pressure increases did not differ sex-specifically (**Figure 5C**). In neither mice nor humans did IL18 associate to blood pressure in either sex (male mice: r = 0.257, R^2 = 0.066, P = 0.354; female mice r = -0.222, R^2 = 0.491, P = 0.427; men: r = 0.070, P = 0.308 and β = 1.27; P = 0.308; women: r = 0.076, P = 0.257 and β = 2.90, P = 0.257).

Discussion

For the first time, we show significant association between plasma S1P and systolic blood pressure levels in a large human cohort study (N=1046) and validate these findings in a longitudinally conducted pre-clinical murine hypertension study where S1P plasma levels positively correlate with systolic blood pressure. Additionally, our data provide first evidence of significant associations between plasma S1P and multiple cardiovascular, inflammation and metabolism biomarkers assessed by proteomic profiling of 444 MOS participants. Some of these markers present with marked sex-specific interactions. Amongst them IL18, that has been associated to hypertension and adverse cardiovascular events, strongly correlates with plasma S1P concentrations in women but not in men. This sex-specific interaction is independent of blood pressure and was verified in a pre-clinical murine hypertension model. Taken together, our novel translational findings strongly suggest a link between S1P plasma concentrations and

systolic blood pressure and encourages the consideration of S1P as potential therapeutic target or risk marker in hypertensive disease.

The pathogenesis of hypertension is profoundly complex, involving several mechanisms, ranging from vascular and sympathetic involvement to immune system activation. Given the large number of affected patients and the subsequent risk of cardiovascular events and mortality, efforts trying to better understand the pathogenesis of hypertension are of utmost clinical importance. The bioactive phosphosphingolipid S1P, as one of the most potent regulators of leukocyte migration, with critical roles in both the vascular and the immune system and proven involvement in experimental hypertension[12, 15, 45, 46], might link inflammation, vascular function, blood pressure regulation and CVD. S1P's critical role in the pathogenesis of experimental hypertension was previously shown. Specifically, one of the enzyme generating S1P, sphingosine kinase 2 (SphK2), presented as a key player in mediating plasma S1P responses to AngII, immune cell egress from secondary lymphoid tissue and vascular dysfunction and thereby, contributing to the development of overt hypertension.[15] Thus far, human-based studies investigating S1P associations with blood pressure were lacking. As the risk of CVD mortality doubles by each 20 mmHg increment in systolic blood pressure,[2] additional tools that help identifying those at risk are needed. Our study reveals associations between increasing plasma levels of S1P and systolic blood pressure cut-offs of <120mmHg, ≥120mmHg, ≥130mmHg, and ≥140mmHg. Concurrently, the effect size of S1P association with systolic blood pressure in humans is comparable to those of well-established hypertension-associated biomarkers such as renin, soluble interleukin 1 receptor-like 1 (ST2) and IL6. Together, these promising results call for further prospective studies to investigate S1P's potential suitability as predictive marker for identifying subjects at high risk of incident hypertension in the population. As the first study showing significant associations between

plasma S1P and systolic blood pressure in a large population cohort, it is reassuring that the mean S1P plasma levels measured in our human study population validate those recently published for a study group comprising 174 healthy participants with a median age of 45.5 years,[47] which further substantiates our findings. Similar to our study, neither S1P plasma nor serum levels were affected by sex, age or BMI.[47, 48]

Previously, S1P was assigned with marker capacity for coronary artery disease.[49] The authors speculated whether S1P might be a marker for inflammatory processes associated with coronary disease or whether sphingolipids are ischemic markers. Here, we provide first evidence of significant associations between plasma S1P and multiple CVD, inflammation and metabolism markers assessed by proteomic profiling. Modularity clustering that detects communities of markers with strong pairwise correlations within a community arranged the strongest S1P associations into marker communities linked to vascular inflammation, immune cell homeostasis, cell adhesion, endothelial inflammation and thrombosis. PECAM-1 as multi-functional vascular cell adhesion molecule, that is highly expressed at endothelial cell-cell junctions with critical functions in endothelial cell permeability,[50] presented with strong positive associations to S1P in humans. A tSNE visualization, where locations of markers are meaningful as proximity is explained by similarity in correlations with other markers, grouped PECAM-1 in close proximity to other makers related to endothelial inflammation, thrombosis, atherosclerosis and cardiovascular events, indicating S1P-PECAM-1 involvement in endothelial damage and deregulation of inflammatory responses at the blood-endothelial interface.[51] In light of this, we observe augmentation of PECAM-1 expression in response to excess exogenous S1P *in vitro*, *in vivo* and *ex vivo*, suggesting a pivotal relationship that supports previous studies demonstrating PECAM-1 as a downstream target for S1P signaling in human endothelial cells.[52] Together with our findings that show elevated resistance artery PECAM-1 expression and increased plasma S1P in hypertensive mice, these data are

supportive of potential contributions of S1P-PECAM-1 deregulations in hypertensive disease. Similarly, SELP that plays an essential role in the recruitment of leukocytes during inflammation[53] presented with strong S1P associations in our human cohort and was upregulated in human and murine endothelial cells following exogenous S1P treatment. In our visualization approaches, SELP clusters together with markers of vascular inflammation and cell adhesion and in close proximity to PECAM-1, respectively. This is particularly intriguing since soluble SELP levels are elevated among healthy women at risk for future vascular events[54] thus, encouraging the consideration of S1P as therapeutic target or risk marker for hypertensive disease.

A potential sex-specificity for S1P plasma concentrations is controversially discussed with equally many studies suggesting sex-dependent differences as those reporting similar S1P plasma levels between the sexes.[44, 47, 48, 55-58] In our study, no significant sex-specific statistical differences in human or murine plasma S1P concentrations were observed. When investigating the association between S1P levels, sex and pre/postmenopausal status in 108 healthy participants, a previous study demonstrated a significant association between plasma S1P and estrogen, with higher S1P plasma concentration in pre-menopausal than post-menopausal women.[44] Evaluating a much smaller study population, the authors assessed plasma S1P levels in individuals in an age range between 16 to 55 years as compared to our study population with a narrow age range, 51.7 ($\pm$7.9) years, and presumably a predominance of post-menopausal women. Nonetheless, the herein observed sex-independence of S1P concentrations, specifically in respect to disease, requires verification in even larger cohorts.

In light of known sex-specific differences in blood pressure control,[59] experimental studies highlighted the critical role of inflammation (i.e. regulatory T-cells and T helper 17 cells) in mediating such differences.[60] Despite the relative sex-independence of plasma S1P in our

study, some of the herein identified S1P associations with proteomic markers presented with sex-specific interactions. Amongst them IL18, which has been suggested as independent predictor of cardiovascular events in subjects with metabolic syndrome[61] and chronic kidney disease,[62] is involved in destabilization of atherosclerotic plaques[63] and was linked to essential hypertension.[64] The herein observed sex-specific differences in S1P-IL18 associations are interesting as they are independent of blood pressure (i.e. correlation and linear regression analyses revealed no significant associations between IL18 and blood pressure in either sex). Thus far, only one experimental study reported significantly higher IL18 mRNA expression in the renal cortex of female spontaneously hypertensive rats compared to age-matched males[65] despite 10-15 mmHg higher blood pressure in male compared to female rats.[66] This is particularly interesting since blood pressure in men significantly differs from women in our study cohort (125 mmHg ($\pm$14) and 119 ($\pm$17) for men and women respectively; $P=7.5 \times 10^{-10}$). Similar female-driven increases in the renal cortex of spontaneously hypertensive rats were observed for CD40 expression,[65] which positively associates with S1P plasma levels in women but not men in our study. Thus, our findings show that blood pressure-independent sex-specific S1P associations to different inflammation, metabolism and CVD markers exist, suggestive of sex-specific differences in immune and vascular responses during hypertension or its development with potential significance for cardiovascular health that might only be unveiled when directly comparing to S1P plasma levels.

Conclusion

Our translational data provide evidence of associations between plasma S1P and systolic blood pressure in both men and mice. We provide first evidence of significant associations between plasma S1P and multiple CVD, inflammation and metabolism biomarkers in humans and sex-specific interactions for some of these markers. Taken together, our novel translational findings

encourage the consideration of S1P as potential therapeutic target or risk marker for hypertensive disease.

Strengths and limitations

By studying a general population and adjusting for risk factors for hypertension, we demonstrated that S1P plasma level are associated with increased systolic blood pressure in humans. Further, we substantiated these findings in an animal model of hypertension. However, since blood pressure regulation is multifactorial, drawing conclusions about associations should be done with caution. The present study shares limitations common to all cross-sectional studies as no conclusion of causality can be drawn. Moreover, the MOS study was carried out in subjects of mainly Swedish descent, and the conclusions may not be generalizable to all populations or ethnicities. Also, since the MOS study is ongoing and participants still fairly young, we were unable to investigate possible association of S1P with incident disease (e.g., incident hypertension or CVD).

The use of mass spectrometry to determine S1P concentrations may be seen as limitation due to the specific tissue preparation requirements and the limited availability of this technological platform. Thus far, however, no feasible alternative for processing large sample sizes with such accuracy is available.

Proteomic profiling of four different OLINK panels (inflammation, metabolism, CVDII and CVDIII) was performed using inter-plate controls for markers within the individual panels but no standardization was performed to compare between the different panels. By analyzing markers that appeared in more than one panel, a high degree of similarity (i.e. correlation of the same marker between two panels) was observed for inflammation, metabolism and CVDII panels ($r \sim 0.8$). In a few cases of the CVDIII panel, however, lower correlation coefficients were observed ($r \sim 0.45$ for two extreme cases). We identified a technical variation associated

to the OLINK data as reason for low correlation between duplicate markers, which has been reported before[67] and which became substantial for some markers with a biological variance below 0.4 NPX. Thus, correlations to markers outside the panels, in particular S1P, should not be expected to markedly surpass correlation factors of duplicate markers and some associations may remain undetected due to the commonly used cut-off r > 0.3. Additionally, we observed a small bias on correlation factors within a panel. This induces a higher chance for markers within one panel to fall into the same cluster when combining all four panels. To mitigate the role of the technical error in the cluster formations, we performed additional individual panel clustering for data interpretation.

Acknowledgement: The authors thank Dr. Nicholas Don-Doncow for longitudinal blood sampling in the murine hypertension model.

Sources of Funding: This work was supported by the following funding sources: The Knut and Alice Wallenberg foundation [F 2015/2112; AM; MM]; Swedish Research Council [VR; 2017-01243; AM]; German Research Foundation [DFG; ME 4667/2-1; AM]; Åke Wibergs Stiftelse [M19-0380; AM]; Inger Bendix Stiftelse [AM-2019-10; AM] and Stohnes Stiftelse [AM]; Lund University [AJ, AM] and Region Skåne [AJ; MM], Swedish Medical Research Council [MM, MOM], the Swedish Heart and Lung Foundation [MM, MOM], the Albert Påhlsson Research Foundation [MM], the Crafoord Foundation [MM], the Ernhold Lundströms Research Foundation [MM], the Hulda and Conrad Mossfelt Foundation [MM], the Southwest Skanes Diabetes foundation [MM], the King Gustaf V and Queen Victoria Foundation [MM], the Lennart Hanssons Memorial Fund [MM], the Marianne and Marcus Wallenberg Foundation [MM], Novo Nordic Foundation (MOM and the European Research Council Consolidator grant (MOM). The Malmö Offspring Study (MOS) has been funded by

the Research Council of Sweden (grant 521-2013-2756), the Heart and Lung Foundation (grant 20150427), and by funds obtained from the local Region Skane County Council (ALF) [PMN and MOM].

Disclosures: None declared.

Data availability: The data underlying this article are available in the article and in its online supplementary material. Study population-specific data that support the findings of this study are available upon reasonable request from the Steering Committee of Malmö Diet and Cancer study by contacting its chair, Professor Olle Melander (olle.melander@med.lu.se), but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly available due to ethical and legal restrictions related to the Swedish Biobanks in Medical Care Act (2002:297) and the Personal Data Act (1998:204).

References

1.   Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, Alexander L, Estep K, Hassen Abate K, Akinyemiju TF, Ali R, Alvis-Guzman N, Azzopardi P, Banerjee A, Barnighausen T, Basu A, Bekele T, Bennett DA, Biadgilign S, Catala-Lopez F, Feigin VL, Fernandes JC, Fischer F, Gebru AA, Gona P, Gupta R, Hankey GJ, Jonas JB, Judd SE, Khang YH, Khosravi A, Kim YJ, Kimokoti RW, Kokubo Y, Kolte D, Lopez A, Lotufo PA, Malekzadeh R, Melaku YA, Mensah GA, Misganaw A, Mokdad AH, Moran AE, Nawaz H, Neal B, Ngalesoni FN, Ohkubo T, Pourmalek F, Rafay A, Rai RK, Rojas-Rueda D, Sampson UK, Santos IS, Sawhney M, Schutte AE, Sepanlou SG, Shifa GT, Shiue I, Tedla BA, Thrift AG, Tonelli M, Truelsen T, Tsilimparis N, Ukwaja KN, Uthman OA, Vasankari T, Venketasubramanian N, Vlassov VV, Vos T, Westerman R, Yan LL, Yano Y, Yonemoto N, Zaki ME and Murray CJ. Global Burden of Hypertension and Systolic Blood Pressure of at Least 110 to 115 mm Hg, 1990-2015. *JAMA*. 2017;317:165-182.

2.   Lewington S, Clarke R, Qizilbash N, Peto R, Collins R and Prospective Studies C. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903-13.

3.   Group SR, Wright JT, Jr., Williamson JD, Whelton PK, Snyder JK, Sink KM, Rocco MV, Reboussin DM, Rahman M, Oparil S, Lewis CE, Kimmel PL, Johnson KC, Goff DC, Jr., Fine LJ, Cutler JA, Cushman WC, Cheung AK and Ambrosius WT. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med*. 2015;373:2103-16.

4.   Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, de Simone G, Dominiczak A, Kahan T, Mahfoud F, Redon J, Ruilope L, Zanchetti A, Kerins M, Kjeldsen SE, Kreutz R, Laurent S, Lip GYH, McManus R, Narkiewicz K, Ruschitzka F, Schmieder RE, Shlyakhto E, Tsioufis C, Aboyans V, Desormais I and Group ESCSD. 2018 ESC/ESH Guidelines for the management of arterial hypertension. *Eur Heart J*. 2018;39:3021-3104.

5.   Whelton PK, Carey RM, Aronow WS, Casey DE, Jr., Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbiagele B, Smith SC, Jr., Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Williams KA, Sr., Williamson JD and Wright JT, Jr. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2018;71:e127-e248.

6.   Muntner P, Carey RM, Gidding S, Jones DW, Taler SJ, Wright JT, Jr. and Whelton PK. Potential US Population Impact of the 2017 ACC/AHA High Blood Pressure Guideline. *Circulation*. 2018;137:109-118.

7.   Byrd JB. Personalized medicine and treatment approaches in hypertension: current perspectives. *Integr Blood Press C*. 2016;9:59-66.

8.   Pokharel Y, Sun W, de Lemos JA, Taffet GE, Virani SS, Ndumele CE, Mosley TH, Hoogeveen RC, Coresh J, Wright JD, Heiss G, Boerwinkle EA, Bozkurt B, Solomon SD, Ballantyne CM and Nambi V. High-sensitivity troponin T and cardiovascular events in systolic

blood pressure categories: atherosclerosis risk in communities study. *Hypertension*. 2015;65:78-84.

9. Uddin SMI, Mirbolouk M, Kianoush S, Orimoloye OA, Dardari Z, Whelton SP, Miedema MD, Nasir K, Rumberger JA, Shaw LJ, Berman DS, Budoff MJ, McEvoy JW, Matsushita K, Blaha MJ and Graham G. Role of Coronary Artery Calcium for Stratifying Cardiovascular Risk in Adults With Hypertension. *Hypertension*. 2019;73:983-989.

10. Patel KV, Pandey A and de Lemos JA. Conceptual Framework for Addressing Residual Atherosclerotic Cardiovascular Disease Risk in the Era of Precision Medicine. *Circulation*. 2018;137:2551-2553.

11. Pandey A, Patel KV, Vongpatanasin W, Ayers C, Berry JD, Mentz RJ, Blaha MJ, McEvoy JW, Muntner P, Vaduganathan M, Correa A, Butler J, Shimbo D, Nambi V, deFilippi C, Seliger SL, Ballantyne CM, Selvin E, de Lemos JA and Joshi PH. Incorporation of Biomarkers Into Risk Assessment for Allocation of Antihypertensive Medication According to the 2017 ACC/AHA High Blood Pressure Guideline: A Pooled Cohort Analysis. *Circulation*. 2019;140:2076-2088.

12. Cantalupo A, Gargiulo A, Dautaj E, Liu C, Zhang Y, Hla T and Di Lorenzo A. S1PR1 (Sphingosine-1-Phosphate Receptor 1) Signaling Regulates Blood Flow and Pressure. *Hypertension*. 2017;70:426-434.

13. Don-Doncow N, Zhang Y, Matuskova H and Meissner A. The emerging alliance of sphingosine-1-phosphate signalling and immune cells: from basic mechanisms to implications in hypertension. *Br J Pharmacol*. 2019;176:1989-2001.

14. Hoefer J, Azam MA, Kroetsch JT, Leong-Poi H, Momen MA, Voigtlaender-Bolz J, Scherer EQ, Meissner A, Bolz SS and Husain M. Sphingosine-1-phosphate-dependent activation of p38 MAPK maintains elevated peripheral resistance in heart failure through increased myogenic vasoconstriction. *Circulation research*. 2010;107:923-33.

15. Meissner A, Miro F, Jimenez-Altayo F, Jurado A, Vila E and Planas AM. Sphingosine-1-phosphate signalling-a key player in the pathogenesis of Angiotensin II-induced hypertension. *Cardiovascular research*. 2017;113:123-133.

16. Rivera J, Proia RL and Olivera A. The alliance of sphingosine-1-phosphate and its receptors in immunity. *Nature reviews Immunology*. 2008;8:753-63.

17. Yagi K, Lidington D, Wan H, Fares JC, Meissner A, Sumiyoshi M, Ai J, Foltz WD, Nedospasov SA, Offermanns S, Nagahiro S, Macdonald RL and Bolz SS. Therapeutically Targeting Tumor Necrosis Factor-alpha/Sphingosine-1-Phosphate Signaling Corrects Myogenic Reactivity in Subarachnoid Hemorrhage. *Stroke; a journal of cerebral circulation*. 2015;46:2260-70.

18. Yang J, Noyan-Ashraf MH, Meissner A, Voigtlaender-Bolz J, Kroetsch JT, Foltz W, Jaffray D, Kapoor A, Momen A, Heximer SP, Zhang H, van Eede M, Henkelman RM, Matthews SG, Lidington D, Husain M and Bolz SS. Proximal cerebral arteries develop myogenic responsiveness in heart failure via tumor necrosis factor-alpha-dependent activation of sphingosine-1-phosphate signaling. *Circulation*. 2012;126:196-206.

19. Matloubian M, Lo CG, Cinamon G, Lesneski MJ, Xu Y, Brinkmann V, Allende ML, Proia RL and Cyster JG. Lymphocyte egress from thymus and peripheral lymphoid organs is dependent on S1P receptor 1. *Nature*. 2004;427:355-60.

20.   Eken A, Duhen R, Singh AK, Fry M, Buckner JH, Kita M, Bettelli E and Oukka M. S1P1 deletion differentially affects TH17 and Regulatory T cells. *Scientific reports*. 2017;7:12905.

21.   Garris CS, Wu L, Acharya S, Arac A, Blaho VA, Huang Y, Moon BS, Axtell RC, Ho PP, Steinberg GK, Lewis DB, Sobel RA, Han DK, Steinman L, Snyder MP, Hla T and Han MH. Defective sphingosine 1-phosphate receptor 1 (S1P1) phosphorylation exacerbates TH17-mediated autoimmune neuroinflammation. *Nature immunology*. 2013;14:1166-72.

22.   Liao JJ, Huang MC and Goetzl EJ. Cutting edge: Alternative signaling of Th17 cell development by sphingosine 1-phosphate. *Journal of immunology*. 2007;178:5425-8.

23.   Wilkerson BA and Argraves KM. The role of sphingosine-1-phosphate in endothelial barrier function. *Biochim Biophys Acta*. 2014;1841:1403-1412.

24.   Chen T, Lin R, Jin S, Chen R, Xue H, Ye H and Huang Z. The Sphingosine-1-Phosphate/Sphingosine-1-Phosphate Receptor 2 Axis in Intestinal Epithelial Cells Regulates Intestinal Barrier Function During Intestinal Epithelial Cells-CD4+T-Cell Interactions. *Cell Physiol Biochem*. 2018;48:1188-1200.

25.   Fox TE, Bewley MC, Unrath KA, Pedersen MM, Anderson RE, Jung DY, Jefferson LS, Kim JK, Bronson SK, Flanagan JM and Kester M. Circulating sphingolipid biomarkers in models of type 1 diabetes. *J Lipid Res*. 2011;52:509-17.

26.   Frej C, Andersson A, Larsson B, Guo LJ, Norstrom E, Happonen KE and Dahlback B. Quantification of sphingosine 1-phosphate by validated LC-MS/MS method revealing strong correlation with apolipoprotein M in plasma but not in serum due to platelet activation during blood coagulation. *Anal Bioanal Chem*. 2015;407:8533-42.

27.   Fayyaz S, Henkel J, Japtok L, Kramer S, Damm G, Seehofer D, Puschel GP and Kleuser B. Involvement of sphingosine 1-phosphate in palmitate-induced insulin resistance of hepatocytes via the S1P2 receptor subtype. *Diabetologia*. 2014;57:373-82.

28.   Spijkers LJ, van den Akker RF, Janssen BJ, Debets JJ, De Mey JG, Stroes ES, van den Born BJ, Wijesinghe DS, Chalfant CE, MacAleese L, Eijkel GB, Heeren RM, Alewijnse AE and Peters SL. Hypertension is associated with marked alterations in sphingolipid biology: a potential role for ceramide. *PloS one*. 2011;6:e21817.

29.   Xue Y, Jiang W, Ma Q, Wang X, Jia P, Li Q, Chen S, Song B, Wang Y, Zhang J, Liu J, Yang G, Lin Y, Liu J, Wei L, Dong C, Li H, Xie Z, Bai L and Ma A. U-shaped association between plasma sphingosine-1-phosphate levels and mortality in patients with chronic systolic heart failure: a prospective cohort study. *Lipids Health Dis*. 2020;19:125.

30.   Polzin A, Piayda K, Keul P, Dannenberg L, Mohring A, Graler M, Zeus T, Kelm M and Levkau B. Plasma sphingosine-1-phosphate concentrations are associated with systolic heart failure in patients with ischemic heart disease. *J Mol Cell Cardiol*. 2017;110:35-37.

31.   Sattler K, Lehmann I, Graler M, Brocker-Preuss M, Erbel R, Heusch G and Levkau B. HDL-bound sphingosine 1-phosphate (S1P) predicts the severity of coronary artery atherosclerosis. *Cell Physiol Biochem*. 2014;34:172-84.

32.   Chen J, Tang H, Sysol JR, Moreno-Vinasco L, Shioura KM, Chen T, Gorshkova I, Wang L, Huang LS, Usatyuk PV, Sammani S, Zhou G, Raj JU, Garcia JG, Berdyshev E, Yuan JX, Natarajan V and Machado RF. The sphingosine kinase 1/sphingosine-1-phosphate pathway in pulmonary arterial hypertension. *American journal of respiratory and critical care medicine*. 2014;190:1032-43.

33.   Zhao YD, Chu L, Lin K, Granton E, Yin L, Peng J, Hsin M, Wu L, Yu A, Waddell T, Keshavjee S, Granton J and de Perrot M. A Biochemical Approach to Understand the Pathogenesis of Advanced Pulmonary Arterial Hypertension: Metabolomic Profiles of Arginine, Sphingosine-1-Phosphate, and Heme of Human Lung. *PloS one*. 2015;10:e0134958.

34.   Gairhe S, Joshi SR, Bastola MM, McLendon JM, Oka M, Fagan KA and McMurtry IF. Sphingosine-1-phosphate is involved in the occlusive arteriopathy of pulmonary arterial hypertension. *Pulmonary circulation*. 2016;6:369-80.

35.   Tabeling C, Yu H, Wang L, Ranke H, Goldenberg NM, Zabini D, Noe E, Krauszman A, Gutbier B, Yin J, Schaefer M, Arenz C, Hocke AC, Suttorp N, Proia RL, Witzenrath M and Kuebler WM. CFTR and sphingolipids mediate hypoxic pulmonary vasoconstriction. *Proceedings of the National Academy of Sciences of the United States of America*. 2015;112:E1614-23.

36.   Brunkwall L, Jonsson D, Ericson U, Hellstrand S, Kennback C, Ostling G, Jujic A, Melander O, Engstrom G, Nilsson J, Ohlsson B, Klinge B, Orho-Melander M, Persson M and Nilsson PM. The Malmo Offspring Study (MOS): design, methods and first results. *Eur J Epidemiol*. 2020.

37.   ROSVALL M, JANZON L, BERGLUND G, ENGSTRÖM G and HEDBLAD B. Incident coronary events and case fatality in relation to common carotid intima-media thickness. *Journal of Internal Medicine*. 2005;257:430-437.

38.   Chipeaux C, de Person M, Burguet N, Billette de Villemeur T, Rose C, Belmatoug N, Heron S, Le Van Kim C, Franco MC, 2017 #34} and Moussa F. Optimization of ultra-high pressure liquid chromatography - tandem mass spectrometry determination in plasma and red blood cells of four sphingolipids and their evaluation as biomarker candidates of Gaucher's disease. *J Chromatogr A*. 2017;1525:116-125.

39.   Bastian M. HS, Jacomy M. Gephi: an open source software for exploring and manipulating networks. *International AAAI Conference on Weblogs and Social Media*. 2009.

40.   Blondel VD GG, Lambiotte R, Lefebvre E. Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*. 2008;10.

41.   GE vdMLaH. Visualizing High-Dimensional Data Using t-SNE. . *Journal of Machine Learning Research* 2008:2579-2605.

42.   Mukaka MM. Statistics corner: A guide to appropriate use of correlation coefficient in medical research. *Malawi Med J*. 2012;24:69-71.

43.   Schober P, Boer C and Schwarte LA. Correlation Coefficients: Appropriate Use and Interpretation. *Anesth Analg*. 2018;126:1763-1768.

44.   Guo S, Yu Y, Zhang N, Cui Y, Zhai L, Li H, Zhang Y, Li F, Kan Y and Qin S. Higher level of plasma bioactive molecule sphingosine 1-phosphate in women is associated with estrogen. *Biochim Biophys Acta*. 2014;1841:836-46.

45.   Swendeman SL, Xiong Y, Cantalupo A, Yuan H, Burg N, Hisano Y, Cartier A, Liu CH, Engelbrecht E, Blaho V, Zhang Y, Yanagida K, Galvani S, Obinata H, Salmon JE, Sanchez T, Di Lorenzo A and Hla T. An engineered S1P chaperone attenuates hypertension and ischemic injury. *Sci Signal*. 2017;10.

46.   Cantalupo A, Zhang Y, Kothiya M, Galvani S, Obinata H, Bucci M, Giordano FJ, Jiang XC, Hla T and Di Lorenzo A. Nogo-B regulates endothelial sphingolipid homeostasis to control vascular function and blood pressure. *Nat Med*. 2015;21:1028-1037.

47.   Daum G, Winkler M, Moritz E, Muller T, Geffken M, von Lucadou M, Haddad M, Peine S, Boger RH, Larena-Avellaneda A, Debus ES, Graler M and Schwedhelm E. Determinants of Serum- and Plasma Sphingosine-1-Phosphate Concentrations in a Healthy Study Group. *TH Open*. 2020;4:e12-e19.

48.   Moritz E, Wegner D, Gross S, Bahls M, Dorr M, Felix SB, Ittermann T, Oswald S, Nauck M, Friedrich N, Boger RH, Daum G, Schwedhelm E and Rauch BH. Reference intervals for serum sphingosine-1-phosphate in the population-based Study of Health in Pomerania. *Clin Chim Acta*. 2017;468:25-31.

49.   Deutschman DH, Carstens JS, Klepper RL, Smith WS, Page MT, Young TR, Gleason LA, Nakajima N and Sabbadini RA. Predicting obstructive coronary artery disease with serum sphingosine-1-phosphate. *Am Heart J*. 2003;146:62-8.

50.   Lertkiatmongkol P, Liao D, Mei H, Hu Y and Newman PJ. Endothelial functions of platelet/endothelial cell adhesion molecule-1 (CD31). *Curr Opin Hematol*. 2016;23:253-259.

51.   Privratsky JR, Newman DK and Newman PJ. PECAM-1: conflicts of interest in inflammation. *Life Sci*. 2010;87:69-82.

52.   Huang YT, Chen SU, Chou CH and Lee H. Sphingosine 1-phosphate induces platelet/endothelial cell adhesion molecule-1 phosphorylation in human endothelial cells through cSrc and Fyn. *Cell Signal*. 2008;20:1521-7.

53.   Robinson SD, Frenette PS, Rayburn H, Cummiskey M, Ullman-Culleré M, Wagner DD and Hynes RO. Multiple, targeted deficiencies in selectins reveal a predominant role for P-selectin in leukocyte recruitment. *Proceedings of the National Academy of Sciences*. 1999;96:11452-11457.

54.   Ridker PM, Buring JE and Rifai N. Soluble P-selectin and the risk of future cardiovascular events. *Circulation*. 2001;103:491-5.

55.   Yafasova A, Mandrup CM, Egelund J, Nyberg M, Stallknecht B, Hellsten Y, Nielsen LB and Christoffersen C. Effect of menopause and exercise training on plasma apolipoprotein M and sphingosine-1-phosphate. *J Appl Physiol (1985)*. 2019;126:214-220.

56.   Chung MY, Park SY, Chung JO, Cho DH and Chung DJ. Plasma sphingosine 1-phosphate concentrations and cardiovascular autonomic neuropathy in individuals with type 2 diabetes. *Scientific reports*. 2020;10:12768.

57.   Winkler MS, Nierhaus A, Holzmann M, Mudersbach E, Bauer A, Robbe L, Zahrte C, Geffken M, Peine S, Schwedhelm E, Daum G, Kluge S and Zoellner C. Decreased serum concentrations of sphingosine-1-phosphate in sepsis. *Crit Care*. 2015;19:372.

58.   Ohkawa R, Nakamura K, Okubo S, Hosogaya S, Ozaki Y, Tozuka M, Osima N, Yokota H, Ikeda H and Yatomi Y. Plasma sphingosine-1-phosphate measurement in healthy subjects: close correlation with red blood cell parameters. *Ann Clin Biochem*. 2008;45:356-63.

59.   Maranon R and Reckelhoff JF. Sex and gender differences in control of blood pressure. *Clin Sci (Lond)*. 2013;125:311-8.

60.   Tipton AJ, Baban B and Sullivan JC. Female spontaneously hypertensive rats have a compensatory increase in renal regulatory T cells in response to elevations in blood pressure. *Hypertension*. 2014;64:557-64.

61.   Troseid M, Seljeflot I, Hjerkinn EM and Arnesen H. Interleukin-18 is a strong predictor of cardiovascular events in elderly men with the metabolic syndrome: synergistic effect of inflammation and hyperglycemia. *Diabetes Care*. 2009;32:486-92.

62.   Formanowicz D, Wanic-Kossowska M, Pawliczak E, Radom M and Formanowicz P. Usefulness of serum interleukin-18 in predicting cardiovascular mortality in patients with chronic kidney disease--systems and clinical approach. *Scientific reports*. 2015;5:18332.

63.   Mallat Z, Corbaz A, Scoazec A, Besnard S, Leseche G, Chvatchko Y and Tedgui A. Expression of interleukin-18 in human atherosclerotic plaques and relation to plaque instability. *Circulation*. 2001;104:1598-603.

64.   S OZ and Ulucam ZM. Association Between Interleukin-18 Level and Left Ventricular Mass Index in Hypertensive Patients. *Korean Circ J*. 2017;47:238-244.

65.   Tipton AJ and Sullivan JC. Sex differences in T cells in hypertension. *Clin Ther*. 2014;36:1882-1900.

66.   Sullivan JC, Semprun-Prieto L, Boesen EI, Pollock DM and Pollock JS. Sex and sex hormones influence the development of albuminuria and renal macrophage infiltration in spontaneously hypertensive rats. *Am J Physiol Regul Integr Comp Physiol*. 2007;293:R1573-9.

67.   Yeh CY, Adusumilli R, Kullolli M, Mallick P, John EM and Pitteri SJ. Assessing biological and technological variability in protein levels measured in pre-diagnostic plasma samples of women with breast cancer. *Biomark Res*. 2017;5:30.

Figures

Figure 1 – S1P plasma concentrations increase with systolic blood pressure in a mouse model of AngII-induced hypertension. |A| Longitudinal assessment of systolic blood pressure (BP_{sys} in mmHg) in responses to a low pressor dose of AngII (20ng/kg/min) in WT mice over the course of four weeks. **|B|** Longitudinal plasma S1P quantification in WT mice developing hypertension over the course of four weeks. **|C|** Linear regression analysis of systolic blood pressure (BP_{sys} in mmHg) and plasma S1P concentrations of WT mice developing hypertension in response to a low pressor dose of AngII (20ng/kg/min). In **A** and **B**, N=7; * denotes $P \leq 0.05$ compared to baseline after two-way repeated measure ANOVA followed by Dunnett's post hoc testing. In **C**, N=7 per group and time point; linear regression analysis of association between plasma S1P and systolic blood pressure levels (BP_{sys} in mmHg) with a calculated goodness of fit measurement (R^2), Pearson's r and exact P-value computation ($P<0.0001$).

Figure 2 – S1P significantly associates with various biomarkers in inflammation, metabolism and CVD OLINK panels. Overview of significant S1P correlations and associations amongst 299 proteins from |**A**| inflammation, |**B**| metabolism, |**C**| CDVII and |**D**| CDVIII OLINK panels. The numbers represent all panel markers with previously reported links to cardiovascular (green) or inflammatory disease (yellow), including overlapping markers in each panel. Significant S1P associations to these markers after Pearson's correlation and linear regression analyses are presented for two-tailed test (grey dots), and for multiple comparisons with Bonferroni-correction of P-values (black dots). N=444

Figure 3 – Correlation networks showing significant S1P correlations with biomarkers in the inflammation, metabolism and CVD OLINK panels. Modularity clustering applied to correlation networks for |**A**| inflammation panel (modularity = 0.271), |**B**| metabolism panel (modularity = 0.273), |**C**| CVDII panel (modularity = 0.189) and |**D**| CVDIII panel (modularity = 0.03) where each node represents a marker and edges between nodes a correlation factor of r ≥ 0.3 or r ≤ -0.3. Coloring according to cluster and edge width is weighted by correlation.

Figure 4 – S1P increases various markers of inflammation and vascular dysfunction. |A| Modularity clustering applied to a network constructed from correlation factors of markers ($r \geq 0.3$ or $r \leq -0.3$) from inflammation, metabolism, CVDII and CVDIII OLINK panels combined. Significant S1P correlations highlighted in red. Clusters were categorized based on previously reported marker involvement in inflammation and CVD. **|B|** Augmentation of endothelial activation marker mRNA expression in human endothelial cells in response to 1 μM S1P (6hrs). **|C|** and **|D|** Augmentation of endothelial activation marker mRNA expression in murine endothelial cells in response to 1 μM S1P (12hrs). **|E|** and **|F|** Augmentation of endothelial activation marker mRNA expression in mesenteric arteries isolated from normotensive and hypertensive mice. In **A**, N=444. In **B-D**, N=3 per group in triplicates; * denotes P≤0.05 after single unpaired comparisons. In **E-F**, N=5 per group; * denotes P≤0.05 after single unpaired comparisons.

Figure 5 – Plasma S1P-IL18 associations differ sex-dependently. |**A**| Plasma IL18 concentrations in normotensive and hypertensive female and male WT mice. |**B**| Linear regression showing associations between plasma S1P and IL18 in normotensive and hypertensive female and male WT mice. |**C**| Plasma S1P concentrations in normotensive and hypertensive female and male WT mice. In **A** and **C**, N=7-8 per group; * denotes P≤0.05 compared to same sex control and & denotes P≤0.05 compared to female hypertension group after one-way ANOVA followed by Tukey's post hoc testing. In **B**, N=6-8 per group; Pearson correlation with goodness of fit and r computation; significance from zero was calculated for female (P<0.0001) but not male mice (P=0.4361).

Tables

Table 1 - Characteristics of the study population. Values are means (± standard deviations) or n (%). Statistical significance is given by exact P-value computation after multiple comparisons between different blood pressure groups. AHT=anti-hypertensive treatment. BMI=body mass index; DBP=diastolic blood pressure; eGFR=estimated glomerular filtration rate; SBP=systolic blood pressure.

	Total	<120mmHg	120-129 mmHg	130-139 mmHg	≥140mmHg	P
	N=1046	N=537	N=229	N=153	N=127	
S1P (nmol/L)	770.7 (±173.7)	757.4 (±163.5)	771.6 (±172.8)	788.5 (±195.2)	804.1 (±184.9)	0.024
Age (years)	51.7 (±7.9)	49.4 (±7.8)	52.7 (±7.5)	54.7 (±7.2)	56.6 (±6.2)	1.0×10^{-27}
Sex (women, N (%))	554 (53)	338 (62.9)	89 (38.9)	58 (37.9)	69 (54.3)	1.3×10^{-11}
BMI (kg/m^2)	26.6 (±4.5)	25.6 (±4.2)	27.3 (±4.7)	27.8 (±4.6)	28.2 (±4.3)	4.3×10^{-13}
SBP (mmHg)	121 (±16)	109 (±7)	124 (±3)	134 (±3)	152 (±10)	-
DBP (mmHg)	76 (±10)	70 (±6)	77 (±6)	83 (±6)	90 (±8)	1.3×10^{-195}
eGFR (mL/min/1.73m^2)	81.4 (±9.7)	82.9 (±9)	82 (±10)	79 (±10)	76 (±11)	8.0×10^{-11}
Diabetes status (yes, N (%))	43 (4.1)	11 (2.0)	10 (4.4)	15 (9.8)	7 (5.5)	2.7×10^{-4}
AHT (yes, N (%))	137 (13.1)	28 (5.2)	37 (16.2)	40 (26.1)	32 (25.2)	3.5×10^{-15}
Smoking (yes, N (%))	94 (9.0)	42 (7.8)	20 (8.7)	17 (11.1)	15 (11.8)	0.393
Alcohol usage (N (%))						0.016
Never	69 (6.6)	41 (7.6)	8 (3.5)	11 (7.2)	9 (7.1)	
Monthly or less	169 (16.2)	104 (19.4)	30 (13.1)	14 (9.2)	21 (16.5)	
2-4 times a month	377 (36.0)	192 (35.8)	84 (36.7)	63 (41.2)	38 (29.9)	
2-3 times a week	368 (35.2)	169 (31.5)	93 (40.6)	52 (34.0)	54 (42.5)	
≥ 4 times a week	63 (6.0)	31 (5.8)	14 (6.1)	13 (8.5)	5 (3.9)	

Table 2 - **Associations between S1P and systolic blood pressure ≥120mmHg; ≥130mmHg and ≥140mmHg.** Values are odds ratios (OR) and confidence intervals (CI95%) and significance values (P) for logistic regression analyses. AHT=anti-hypertensive treatment. BMI=body mass index; DBP=diastolic blood pressure; eGFR=estimated glomerular filtration rate; SBP=systolic blood pressure.

	Logistic regression analyses					
	≥120 mmHg		≥130 mmHg		≥140 mmHg	
	OR (CI95%)	P	OR (CI95%)	P	OR (CI95%)	P
Unadjusted						
S1P	1.17 (1.04-1.32)	0.011	1.21 (1.06-1.38)	0.005	1.23 (1.03-1.46)	0.021
Model 1						
S1P	1.14 (1.01-1.30)	0.047	1.19 (1.03-1.37)	0.016	1.21 (1.01-1.46)	0.037
Age	1.10 (1.08-1.12)	1.0×10^{-21}	1.10 (1.08-1.13)	1.7×10^{-19}	1.11 (1.08-1.15)	5.4×10^{-13}
Sex	0.41 (0.31-0.53)	1.9×10^{-11}	0.66 (0.49-0.88)	0.004	1.12 (0.76-1.65)	0.563
Model 2						
S1P	1.15 (1.01-1.32)	0.046	1.21 (1.04-1.40)	0.012	1.23 (1.02-1.48)	0.034
Age	1.09 (1.06-1.11)	9.0×10^{-14}	1.08 (1.06-1.11)	6.7×10^{-11}	1.10 (1.06-1.13)	6.2×10^{-8}
Sex	0.48 (0.36-0.64)	2.6×10^{-7}	0.77 (0.57-1.05)	0.098	1.20 (0.80-1.81)	0.387
BMI	1.09 (1.06-1.13)	1.0×10^{-7}	1.07 (1.04-1.11)	6.3×10^{-6}	1.07 (1.03-1.12)	0.002
eGFR	1.01 (0.99-1.02)	0.643	0.99 (0.97-1.01)	0.202	0.98 (0.96-1.00)	0.048
Diabetes status	1.00 (0.46-2.20)	0.993	0.86 (0.43-1.72)	0.676	1.71 (0.69-4.23)	0.249
AHT	2.89 (1.80-4.63)	1.1×10^{-5}	2.12 (1.41-3.20)	3.1×10^{-4}	1.47 (0.89-2.43)	0.130
Smoking	1.25 (0.77-2.02)	0.377	1.27 (0.77-2.11)	0.348	1.14 (0.60-2.16)	0.686
Alcohol						
Never	1.09 (0.50-2.40)	0.828	1.48 (0.63-3-44)	0.366	2.11 (0.62-7-17)	0.232
Monthly or less	0.98 (0.51-1.91)	0.961	0.94 (0.45-1.97)	0.877	2.24 (0.75-6.72)	0.149

2-4 times a month	1.43 (0.79-1.91)	0.238	1.30 (0.68-2.47)	0.433	1.74 (0.63-4.83)	0.286
2-3 times a week	1.53 (0.84-2.76)	0.164	1.27 (0.67-2.41)	0.466	2.54 (0.94-6.92)	0.067
≥ 4 times a week	0.92 (0.42-2.02)	0.828	0.79 (0.41-1.50)	0.466	0.57 (0.21-1.59)	0.286

Table 3 - List of proteins from four proteomic panels that show significant sex-dependent associations with plasma S1P with association-determining sex highlighted in orange. Multiple comparison adjustment was performed using Bonferroni-corrected P-values of 0.05/299. For the inflammation panel, two additional significant sex-specific association with plasma S1P are found significant according to panel-specific Bonferroni correction of 0.05/64 (P≤7.81x10^{-4}) that are highlighted in yellow. N=444.

Inflammation panel	Male (N=216)			Female (N=228)		
	Pearson's r	β	P	Pearson's r	β	P
MMP-1	0.258	0.911	1.28x10^{-4}	0.202	0.663	0.002
CD40	0.200	0.345	0.003	0.254	0.382	1.02x10^{-4}
TWEAK	0.039	0.050	0.571	0.246	0.283	1.78x10^{-4}
CD244	0.160	0.235	0.019	0.245	0.308	1.84x10^{-4}
Metabolism panel	Pearson's r	β	P	Pearson's r	β	P
ENO2	0.250	0.601	1.97x10^{-4}	0.316	0.677	7.97x10^{-7}
SUMF2	0.284	0.654	2.10x10^{-5}	0.196	0.435	0.003
CVDII panel	Pearson's r	β	P	Pearson's r	β	P
LOX-1	0.194	0.305	0.004	0.250	0.423	1.21x10^{-4}
IL18	0.157	0.283	0.021	0.278	0.538	1.80x10^{-5}
GLO1	0.241	0.619	3.41x10^{-4}	0.389	0.929	8.93x10^{-10}
CVDIII panel	Pearson's r	β	P	Pearson's r	β	P
CTSD	0.276	0.567	4.0x10^{-5}	0.189	0.271	0.009