Through The Back Door: Expiratory Accumulation Of SARS-Cov-2 In The Olfactory Mucosa As Mechanism For CNS Penetration

Carlotta Pipolo[1], Antonio Mario Bulfamante[1], Andrea Schillaci[2], Jacopo Banchetti[2], Luca Castellani[1], Alberto Maria Saibene[1], Giovanni Felisati[1] * & Maurizio Quadrio[2] *

*These authors are both last co-authors

[1] Unit of Otolaryngology, ASST Santi Paolo e Carlo, Department of Health Sciences, Università degli Studi di Milano, Milan, Italy

[2] Dept. of Aerospace Science and Technologies, Politecnico di Milano, Italy

Corresponding author
Antonio Mario Bulfamante, MD
Università degli Studi di Milano
Otolaryngology Department
San Paolo Hospital
Via A. di Rudinì, 8
20142 Milan – Italy
Phone: +39 02 8184 4249 Fax: +39 02 5032 3166 Mail: antonio.bulfamante@unimi.it

Keywords

Computational Fluid Dynamics, SARS-Cov-2, Olfactory Mucosa, Nose

Abstract

Introduction

SARS-CoV-2 is a respiratory virus supposed to enter the organism through aerosol or fomite transmission to the nose, eyes and oropharynx. It is responsible for various clinical symptoms, including hyposmia and other neurological ones. Current literature suggests the olfactory mucosa as a port of entry to the CNS, but how the virus reaches the olfactory groove is still unknown. Because the first neurological symptoms of invasion (hyposmia) do not correspond to first signs of infection, the hypothesis of direct contact through airborne droplets during primary infection and therefore during inspiration is not plausible. The aim of this study is to evaluate if a secondary spread to the olfactory groove in a retrograde manner during expiration could be more probable.

Methods

Four three-dimensional virtual models were obtained from real CT scans and used to simulate expiratory droplets. The volume mesh consists of 25 million of cells, the simulated condition is a steady expiration, driving a flow rate of 270 ml/s, for a duration of 0.6 seconds. Droplet diameter is of 5 μm.

Results

The analysis of the simulations shows the virus to have a high probability to be deployed in the rhinopharynx, on the tail of medium and upper turbinates. The possibility for droplets to access the olfactory mucosa during the expiratory phase is lower than other nasal areas, but consistent.

Discussion

The data obtained from these simulations demonstrates that virus can be deployed in the olfactory groove during expiration. Even if the total amount in a single act is very low, it must be considered that it is repeated tens of thousands of times a day, and the source of contamination continuously acts on a timescale of several days. The present results also imply CNS penetration of SARS-CoV-2 through olfactory mucosa might be considered a complication and, consequently ,prevention strategies should be considered in diseased patients.

Abbreviations

SARS-CoV-2: severe acute respiratory syndrome coronavirus 2; CT: computed tomography; CNS: Central nervous system; LES: Large eddy simulation; CFD: Computational fluid dynamics; ACE-2: Angiotensin converting enzyme; COVID-19: Coronavirus disease 2019

Introduction

SARS-CoV-2 is a respiratory virus, still widely spreading throughout the globe, that is thought to enter the organism through aerosol or fomite transmission to the nose, eyes and oropharynx (1,2). Presentation ranges from respiratory symptoms such as cough and fever to neurological symptoms such as headache, dizziness and hyposmia, showing different target organs of the virus (3,4). Current literature has recently started to study access points into the CNS and the anatomical proximity between neurons, nerve fibers and the mucosa within the olfactory groove (5); the reported clinical-neurological signs related to alteration in smell suggest that SARS-CoV-2 exploits this neuro-mucosal interface as port of entry. Even though early reports (6) are indeed supporting this hypothesis through autopsy sampling, no literature exists as to how the SARS-CoV-2 reaches the mucosa at the level of the olfactory cleft, and whether the olfactory mucosa involvement is a direct consequence of viral particle deposition or due to a secondary viral invasion of these tissues during the course of the infection.

From other respiratory viruses we know that aerosols, which are responsible for the transmission of airborne microorganisms, consist of small droplet nuclei (1–5µm) or droplets (>5µm) (7); these have specific characteristics regarding their distribution inside the nose and respiratory tract. Considering that the first neurological symptoms of invasion (hyposmia) do not correspond to first signs of presentation of infection, the hypothesis of direct contact through airborne at the stage of primary infection and therefore during inspiration is not plausible.

The second hypothesis of a secondary spread to the olfactory groove in a retrograde manner during for example expiration in an already challenged organism seems to be more likely. This would make CNS penetration a complication secondary to e.g. pulmonary infection, thus opening the field to so far unconsidered preventative measures.

Our group has therefore used computational fluid dynamics to study distribution of airflow and deposition of supposed infectious sub-micron droplets during breathing, to better understand the possible routes of infection and penetration inside the nasal cavities and the olfactory mucosa.

Materials and Methods

This study was granted exemption from the Institutional Review Board of the San Paolo Hospital, Milano, Italy, due to its retrospective nature and is based on a set of CFD simulations of breathing,

where only expiration is considered. The Large Eddy Simulation (LES) technique on CT scan

reconstructions of nasal anatomies is used. LES is a high-cost and high-fidelity CFD approach, which

allows fine control over the modelling error in dealing with complex and possibly turbulent flows.

LES numerical simulations were performed starting from a set of four CT scans, whose sinonasal

anatomy was defined by consensus by all authors as devoid of any appreciable anatomic anomaly

(i.e. a straight septum, normotrophic turbinates with orthodox bending, symmetrical distribution of

anatomical features among the two sinonasal emi-systems).

CT scans have a 512 × 512 matrix with a 0.49 mm × 0.49 mm spatial resolution in the sagittal-coronal

plane and a 0.625 mm gap between consecutive axial slices, with 250-350 native images for each

case. More details on CT images processing, choice of the threshold value and 3D reconstruction,

carried out via the software 3D Slicer (9), have been already reported in the literature documenting

the entire procedure (10–14). The CFD simulations were carried out with the finite-volumes

OpenFOAM software package (15).

The CFD analysis of each of the four cases (patients from P1 to P4) was conducted on finely

discretized volume mesh consisting of 25 million of cells, yielding extreme accuracy. The simulated

condition is a steady expiration driving a flow rate of 270 ml/s, which corresponds to low to medium

intensity breathing (16), for a duration of 0.6 seconds. A large number of droplets with diameter of

5 μm, in accordance with the expected droplet size described above (7), was placed at the lower

boundary of the computational domain and allowed to enter as time progresses. Droplets are

transported by the airflow, and most of them are exhaled after travelling through the nasal

chamber, becoming responsible for the potential airborne contagion. Because of their inertia,

however, a fraction of the droplets deposits on the mucosal lining of the nose. The simulations

identify the deposited droplets, and therefore provide a representation of the deposition pattern,

highlighting areas of preferential deposition during expiration. Particular attention is given to the

particles that reach the olfactory slit, qualitatively sketched in figure 1. This study provides

information on the preferential site of adhesion of expiratory droplets to the olfactory mucosa and

computes the spatially varying degree of probability for a droplet to deposit in a specific location

instead of being convected to the external ambient.

Results

The simulations portrait the preferential sites of droplet deposition on the nasal mucosa during

expiration. It is clearly visible in Figure 2, 3 and 4 that, although virus deposition is prevalent in the

nasal vestibule and rhinopharynx. some droplets indeed do deposit in the area corresponding to the

olfactory mucosa. Moreover, as expected, interindividual differences are visible. Droplets have been

emphasized with a red dot.

The analysis of sagittal sections (fig. 2, where only the left nasal fossa is shown for clarity) shows the

virus to have a high probability to be deployed in the rhinopharynx, on the tail of medium and upper

turbinates. The possibility for droplets to access the olfactory mucosa during the expiratory phase

is of primary interest. The evaluation of axial projections (Fig. 3) confirms a high concentration of

particles in the posterior segments in addition to a better visualisation of particle distribution

between medial and lateral compartments. Although heterogeneously, it can be observed how the

particles, and consequently the virus, are more likely to settle in the medial quadrants of the nasal

cavities than in lateral ones. During the expiratory phase, the particles have a significant probability

to impact and adhere to the mucosa of the ethmoidal rostrum. Finally, the analysis of coronal

projections (Fig. 4) confirms previous observations, although it demonstrates a better visualisation

of the septal rostrum region and of both portions of the rhinopharynx. Coronal projections confirm

that indeed some particles reach of the olfactory slit.

Discussion

Regardless of the mechanism for viral transmission (direct respiratory, aerosol or fomite), first

access to the nose must happen through inspiration. Once the virus has gained entry to the sinonasal

cavities, however, many different potential mechanisms concur to further diffusion, among which

transport, local replication, and invasion of proximal structures. The ability of SARS-CoV-2 to bind

the ACE-2 receptor, enter the respiratory epithelium cells and thereby initiate its replication has

been thoroughly demonstrated (17,18).

Respiratory droplets containing viral particles are unable to massively reach the olfactory cleft,

which should not be therefore considered as a primary target for COVID-19 infection. The droplet

ability to deposit on the olfactory cleft is a direct function of the particle size, given that the olfactory

cleft is anatomically developed to receive smaller particles such as odorants, while droplets carrying

the viral load can be larger (8). Such an ineffective viral deposition on to the olfactory mucosa,

coupled with the known defensive mechanisms employed by the olfactory mucosa to protect from

environmental noxae (17,19), make the direct infection of the olfactory cleft by SARS-CoV-2 at the

time of primary entry into the organism unlikely at best. On the other hand, cumulative exposure of

the olfactory cleft to expiratory droplets from the lower respiratory tract in an already diseased

organism may be more likely. In fact, the viral load in the lung is much higher than a one-time aerosol

reaching the nose, the act of expiration is repeated tens of thousands of times a day, and the source

of contamination continuously acts on a timescale of several days. This route to the olfactory cleft

and maybe to CNS may also explain the time lag between first symptoms and first neurological

impairments such as hyposmia (20,21).

The present results also imply that CNS penetration of SARS-CoV-2 through olfactory mucosa might

be a complication of an already present infection of the lower respiratory tract. Hence, prevention

of the olfactory mucosa penetration by the virus should be considered in diseased patients. High

volumes nasal washes, usually performed with saline, can be used to reduce the adherence of viral

parts emitted from the lower respiratory tract towards the nasal cavities, thus weakening the virus

ability to spread to the olfactory mucosa. Indeed, other authors have advocated the use of nasal

lavages in SARS-CoV-2 infection as a preventive measure (22), and prior studies on viral upper

respiratory tract infections with hypertonic saline showed reduced viral shedding and patients

infectivity (23) in the already diseased. Prior studies on viral upper respiratory tract infections with

hypertonic saline showed decreased viral shedding and reduced patients infectivity (23) in in the

already diseased. Others propose other types of medications: in (24) inhalation of acetic acid is

suggested to be effective in shortening duration of symptoms. The present study -- besides

suggesting the olfactory region as a target for inhibition of the secondary viral infection which

endangers the CNS -- provides further support for the effectiveness of such preventive measures,

since a diffuse droplet deposition takes place in the nasal fossae, that can be easily reached by

washing or by other nasal medications.

Further studies are thus required to focus on nasal washes not only as a preventive measure to

infection but also as a means to inhibit the secondary spreading of the virus to the olfactory mucosa

and therefore to the CNS for COVID-19 patients. These studies should clinically quantify the ability

of nasal washes, i.e. a simple and non-invasive treatment, to halt the progression of the disease by

containing its complications involving the CNS.

Competing interests

The authors have declared that no competing interest exists

References

1. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. New England Journal of Medicine. 2020 Apr 16;382(16):1564–7.

2. Kwok YLA, Gralton J, McLaws M-L. Face touching: a frequent habit that has implications for hand hygiene. Am J Infect Control. 2015 Feb;43(2):112–4.

3. Cui C, Yao Q, Zhang D, Zhao Y, Zhang K, Nisenbaum E, et al. Approaching Otolaryngology Patients During the COVID-19 Pandemic. Otolaryngol Head Neck Surg. 2020 Jul 1;163(1):121–31.

4. Tong JY, Wong A, Zhu D, Fastenberg JH, Tham T. The Prevalence of Olfactory and Gustatory Dysfunction in COVID-19 Patients: A Systematic Review and Meta-analysis. Otolaryngol Head Neck Surg. 2020;163(1):3–11.

5. Doty RL. The olfactory vector hypothesis of neurodegenerative disease: is it viable? Ann Neurol. 2008 Jan;63(1):7–15.

6. Bulfamante G, Chiumello D, Canevini MP, Priori A, Mazzanti M, Centanni S, et al. First ultrastructural autoptic findings of SARS -Cov-2 in olfactory pathways and brainstem. Minerva Anestesiol. 2020;86(6):678–9.

7. Zemouri C, de Soet H, Crielaard W, Laheij A. A scoping review on bio-aerosols in healthcare and the dental environment. PLoS One. 2017;12(5):e0178007.

8. Zhao K, Scherer PW, Hajiloo SA, Dalton P. Effect of anatomy on human nasal air flow and odorant transport patterns: implications for olfaction. Chem Senses. 2004 Jun;29(5):365–79.

9. 3D Slicer: A Platform for Subject-Specific Image Analysis, Visualization, and Clinical Support | SpringerLink [Internet]. [cited 2020 Nov 3]. Available from: https://link.springer.com/chapter/10.1007/978-1-4614-7657-3_19

10. Saibene AM, Felisati G, Pipolo C, Bulfamante AM, Quadrio M, Covello V. Partial Preservation of the Inferior Turbinate in Endoscopic Medial Maxillectomy: A Computational Fluid Dynamics Study. Am J Rhinol Allergy. 2020 May;34(3):409–16.

11. Buijs EFM, Covello V, Pipolo C, Saibene AM, Felisati G, Quadrio M. Thermal water delivery in the nose: experimental results describing droplet deposition through computational fluid dynamics. Acta Otorhinolaryngol Ital. 2019 Dec;39(6):396–403.

12. Covello V, Pipolo C, Saibene A, Felisati G, Quadrio M. Numerical simulation of thermal water delivery in the human nasal cavity. Comput Biol Med. 2018 01;100:62–73.

13. Quadrio M, Pipolo C, Corti S, Lenzi R, Messina F, Pesci C, et al. Review of computational fluid dynamics in the assessment of nasal air flow and analysis of its limitations. Eur Arch Otorhinolaryngol. 2014 Sep;271(9):2349–54.

14. Quadrio M, Pipolo C, Corti S, Messina F, Pesci C, Saibene AM, et al. Effects of CT resolution and radiodensity threshold on the CFD evaluation of nasal airflow. Med Biol Eng Comput. 2016 Mar;54(2–3):411–9.

15. Weller HG, Tabor G, Jasak H, Fureby C. A tensorial approach to computational continuum mechanics using object-oriented techniques. Computers in Physics. 1998 Nov 1;12(6):620–31.

16. Calmet H, Gambaruto AM, Bates AJ, Vázquez M, Houzeaux G, Doorly DJ. Large-scale CFD simulations of the transitional and turbulent regime for the large human airways during rapid inhalation. Computers in Biology and Medicine. 2016 Feb 1;69:166–80.

17. Hui KPY, Cheung M-C, Perera RAPM, Ng K-C, Bui CHT, Ho JCW, et al. Tropism, replication
competence, and innate immune responses of the coronavirus SARS-CoV-2 in human respiratory tract
and conjunctiva: an analysis in ex-vivo and in-vitro cultures. The Lancet Respiratory Medicine. 2020 Jul
1;8(7):687–95.

18. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al. SARS-CoV-2 Cell
Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell.
2020 Apr;181(2):271-280.e8.

19. Herbert RP, Harris J, Chong KP, Chapman J, West AK, Chuah MI. Cytokines and olfactory bulb
microglia in response to bacterial challenge in the compromised primary olfactory pathway. J
Neuroinflammation. 2012 May 29;9:109.

20. Ahmad I, Rathore FA. Neurological manifestations and complications of COVID-19: A literature review. J
Clin Neurosci. 2020 Jul;77:8–12.

21. Niazkar HR, Zibaee B, Nasimi A, Bahri N. The neurological manifestations of COVID-19: a review
article. Neurol Sci. 2020 Jul;41(7):1667–71.

22. Singh S, Sharma N, Singh U, Singh T, Mangal DK, Singh V. Nasopharyngeal wash in preventing and
treating upper respiratory tract infections: Could it prevent COVID-19? Lung India. 2020 Jun;37(3):246–
51.

23. Ramalingam S, Graham C, Dove J, Morrice L, Sheikh A. A pilot, open labelled, randomised controlled
trial of hypertonic saline nasal irrigation and gargling for the common cold. Sci Rep. 2019 31;9(1):1015.

24. Pianta L, Vinciguerra A, Bertazzoni G, Morello R, Mangiatordi F, Lund VJ, et al. Acetic acid disinfection
as a potential adjunctive therapy for non-severe COVID-19. Eur Arch Otorhinolaryngol. 2020
Oct;277(10):2921–4.

Figure 1 Sagittal view of the left nostril. The olfactory slit is highlighted in red.

P1

P2

P3

P4

Figure 2 Distribution of droplet deposition during expiration, in sagittal projection. 3D models of the nasal fossae (left column), obtained from the CT, which are shown in the right panel. Particle size is increased to improve clarity.

Figure 3: As in figure 1, but axial projection.

467
468
469

Figure 4: As in figure 1, but coronal projection.