Main Manuscript for

Post Hoc Evaluation of Probabilistic Model Forecasts: A COVID-19 Case
Study

Kyle J. Colonna[a]*, Roger M. Cooke[b,c], and John S. Evans[a]

[a] Environmental Health Department, Harvard T.H. Chan School of Public Health, Harvard University,
Boston, MA 02115
[b] Resources for the Future, Washington, DC 20036
[c] Department of Mathematics, Delft University of Technology, Delft, The Netherlands 2628 XE

*Corresponding Author: Kyle J. Colonna

Address: 42 Beach St., Unit 9C, Boston, MA, 02111 USA

Phone Number: 484-832-4003

Email: kcolonna@g.harvard.edu

Author Contributions: *Concept and design:* All authors
Acquisition, analysis, or interpretation of data: Colonna, Cooke
Drafting of the manuscript: All authors
Critical revision of the manuscript for important intellectual content: All authors
Supervision: Evans

Competing Interest Statement: The authors declare they have no competing interests that might be
perceived to influence the results and/or discussion reported in this manuscript. There has also been no
prior discussion with a PNAS editor.

Classification: Biological, Health, and Medical Sciences – Biostatistics; Public Health and Epidemiology

Keywords: COVID-19; COVID-19 Decision-Making; Forecasting; Uncertainty Analysis; Cooke's Classical
Model

This PDF file includes:

Main Text
Figure 1 and 2 Legends
Tables 1 and 2

Abstract

To combat the spread of coronavirus disease 2019 (COVID-19), decision-makers and the public may
desire forecasts of the cases, hospitalizations, and deaths that are likely to occur. Thankfully, dozens of
COVID-19 forecasting models exist and many of their forecasts have been made publicly available.
However, there has been little published peer-reviewed information regarding the performance of these
models and what is available has focused mostly on the performance of their central estimates (i.e.,
predictive performance). There has been little reported on the accuracy of their uncertainty estimates (i.e.,
probabilistic performance), which could inform users how often they would be surprised by observations
outside forecasted confidence intervals. To address this gap in knowledge, we borrow from the literature
on formally elicited expert judgment to demonstrate one commonly used approach for resolving this
issue. For two distinct periods of the pandemic, we applied the Classical Model (CM) to evaluate
probabilistic model performance and constructed a performance-weighted ensemble based on this
evaluation. Some models which exhibited good predictive performance were found to have poor
probabilistic performance, and vice versa. Only two of the nine models considered exhibited superior
predictive and probabilistic performance. Additionally, the CM-weighted ensemble outperformed the
equal-weighted and predictive-weighted ensembles. With its limited scope, this study does not provide
definitive conclusions on model performance. Rather, it highlights the evaluation methodology and
indicates the utility associated with using the CM when assessing probabilistic performance and
constructing high performing ensembles, not only for COVID-19 modeling but other applications as well.

Significance Statement

Coronavirus disease 2019 (COVID-19) forecasting models can provide critical information for decision-
makers and the public. Unfortunately, little information on their performance has been published,
particularly regarding the accuracy of their uncertainty estimates (i.e., probabilistic performance). To
address this research gap, we demonstrate the Classical Model (CM), a commonly used approach from
the literature on formally elicited expert judgment, which considers both the tightness of forecast
confidence intervals and frequency in which confidence intervals contain the observation. Two models
exhibited superior performance and the CM-based ensemble consistently outperformed the other
constructed ensembles. While these results are not definitive, they highlight the evaluation methodology
and indicate the value associated with using the CM when assessing probabilistic performance and
constructing high performing ensembles.

60
Main Text

1. Introduction

Effective non-pharmaceutical interventions (NPIs), or community mitigation strategies, are crucial in
combating the spread of contagious illnesses like coronavirus disease 2019 (COVID-19) (1). Community
NPIs, such as social distancing guidelines, restrictions, closures, and lockdowns, can effectively delay
and diminish an epidemic peak (1-3), also known as flattening the epidemic curve (2). However, these
NPIs can also have immediate educational and economic consequences (4-6). To make decisions on the
implementation of community NPIs amidst the COVID-19 pandemic, the relevant decision-makers (e.g.,
government officials, community leaders, school administrators etc.) may desire estimates of the number
of coronavirus cases, hospitalizations, and deaths likely to occur in their region of interest within the
coming weeks.
72
Information about, and forecasts from, dozens of COVID-19 forecasting models have been made
available via the University of Massachusetts Amherst Reich Lab's COVID-19 Forecast Hub (7). All the
modeling groups that have participated have provided predictions (or, central estimates), and most also
quantitatively characterize the uncertainty in their predictions – often giving an interquartile range (25%
LCL to 75% UCL) and a 90% confidence interval (5% LCL to 95% UCL) for each prediction (8). These
models could serve as incredible resources for decision-makers and the public. To understand how much

confidence is warranted based on a model's forecasts, they must have access to evidence about the
performance of these models (i.e., how well do forecasts correspond with observations).

Unfortunately, there has been little published peer-reviewed information regarding the performance of
COVID-19 forecasting models. However, one study has investigated their predictive accuracy (i.e., how
close are predictions to the actual observed values). The authors compared the median absolute percent
error (MAPE) of predictions from seven individual models, stratified across world region, month of model
estimation, and weeks of extrapolation (9). Collectively, the seven models' forecasts over a twelve-week
range had a MAPE of about 27%, with errors tending to increase with longer forecasts and the best
performing model varying by region (9).

While this information is quite useful and provides a sense of the typical bias of model predictions, other
aspects of model performance may also be of interest. Users may also care about a model's precision in
its predictions (i.e., how close are point estimates to each other relative to the actual observed values)
and its probabilistic performance (i.e., the modeler's ability to properly characterize the uncertainty in their
predictions). A model's prediction does not fully capture what the modeling group believes might occur
over a given time range; they are not claiming that their prediction will occur with 100% likelihood. The
probability density provided by a model's forecast is just as important as their prediction, as the range of
this probability density could suggest that a decision-maker takes decisive action for their next step, or
exercises caution. Without such information, decision-makers and the public have no sense of how often
they will be surprised by observations that are well outside the confidence intervals of predictions
provided by the modelers. Frequent surprises may cause the public to lose confidence in the forecasting
process and render a potentially valuable tool in the arsenal of COVID-19 mitigation strategies effectively
useless.

Scientists concerned with the use of formally elicited expert judgment in support of public policy have
faced this issue for decades and have devised approaches for evaluating the probabilistic performance of
expert judgments. In this case, models may serve as the 'experts' and their forecasts may serve as their
'judgments'. One such approach, that can evaluate the probabilistic performance of judgments post hoc,
is the Classical Model (CM) method (10). This approach has been employed in many studies (11-13) and
has been suggested/implemented in other studies concerned with model forecasting (14, 15). Forty-nine
applications since 2006 are referenced and analyzed in the recent Cooke, Marti, & Mazzuchi (2021) study
(16). Other protocols for formally elicited expert judgment studies (e.g., IDEA [investigate, discuss,
estimate, and aggregate] and the Sheffield elicitation framework [SHELF]), require behavioral aggregation
or plenary expert deliberation which are unsuitable for post hoc analysis (17, 18).

Some may also wonder whether better forecasts might be obtained by averaging the forecasts of two or
more individual models. The Reich Lab has created one such ensemble model (19), which has been
prominently featured by the Centers for Disease Control and Prevention (CDC) (20). It involves an equal-
weighted *quantile*-averaged combination of individual model forecasts and is not performance-weighted
(19). While this quantile-averaged equal-weighted ensemble generally outperforms many of the individual
models and has greater performance consistency, (21, 22) performance-weighted combinations have
been demonstrated to consistently outperform equal-weighted combinations in the expert judgment
literature (23, 24). Further, quantile averaging of probabilistic forecasts forms a 90% confidence interval
whose width is the average of forecasters' 90% confidence interval widths regardless of how far apart
these intervals are. Thus, if one forecast has a confidence interval of 0 to 1 with a median of 0.5 while a
second has a confidence interval of 9 to 10 with a median of 9.5, the quantile average yields a confidence
band of 4.5 to 5.5 with a median of 5. This leads to poor statistical accuracy in over half of the cases and
is not reflective of what either forecaster believes (23, 25). Other COVID-19 related studies have
previously commented on the value of using the full distributions of forecasts (14, 15). These issues
deserve consideration in the development of any ensemble forecasting models.

To address these research gaps, our analysis compared model forecasts of COVID-19 mortality for states
within the United States with subsequent observations using several measures that reflect predictive (i.e.,
accuracy and precision) and probabilistic (i.e., statistical accuracy [or, as we call it, calibration] and
information) performance. We also constructed three ensemble models, one equal-weighted and two

performance-weighted, all of which averaged forecast probability densities, and compared their
performance with each other and with the best performing individual models. Lastly, we explored whether
the available models tend to provide better forecasts under certain circumstances (i.e., differing case
rates, racial demographics, and forecast time periods/phases of the pandemic).
139

140 **2. Results**

141 **2.1. Performance Criteria**

142
143 Two aspects of performance were evaluated for each model/ensemble– (i) predictive performance (i.e.,
144 the accuracy and precision of the model/ensemble's central estimates, or predictions); and (ii)
145 probabilistic performance (the modeler's ability to appropriately characterize the uncertainty in their
146 predictions, reflected by their reported confidence intervals).
147
148 First, to evaluate the performance of a model's predictions, each observation, O_i, was divided by the
149 corresponding prediction, $P_{m,i}$, provided by the model, m, to obtain the accuracy ratio, $R_{m,i}$ – where i
150 indicates the question, or item, which reflects the date, state, and time range of interest:
151
152 $$R_{m,i} = O_i / P_{m,i}$$
153
154 The distribution of the resulting accuracy ratios was then examined. For each model, the geometric mean
155 (GM) and geometric standard deviation (GSD), based on the accuracy ratios for all items, N, were
156 computed and used as respective measures of the observed accuracy and precision of the model's
157 predictions.
158
159 $$GM_m = \exp(\text{mean}(\ln(R_i), \ldots, \ln(R_N)))$$
160
161 $$GSD_m = \exp(\text{SD}(\ln(R_i), \ldots, \ln(R_N)))$$
162
Second, to evaluate each model's ability to appropriately characterize the uncertainty in its predictions,
their probabilistic performance was assessed using the Classical Model (CM) method (10). The CM
assesses 'statistical accuracy', or 'calibration', C, using Shannon's relative information statistic, Is, which
compares the forecasted probability distribution for each item with the corresponding observation (10,
26). Each observation may fall within six inter-quantile intervals, based on the five uncertainty distribution
quantiles (i.e., 5%, 25%, 50%, 75%, 95%) provided by each model for every item. The calibration score
for each model, C_m, takes the following formula:
170
171 $$C_m = 1 - X^2(2N * Is), \text{ where } Is = \Sigma_{k=1, \ldots, 6}(s_k * \ln(s_k / p_k))$$
172
where X^2 is the chi-square distribution function with five degrees of freedom (based on the number of
quantiles), s_k is the percentage of observations falling in the specific inter-quantile interval, k, and p_k is the
inherent probability of the k interval (10, 26). As the divergence between s_k and p_k increases, Is increases
and C moves toward 0 (the highest calibration score possible is 1). 2N * Is, where N corresponds to the
total number of items, is the log likelihood ratio which is asymptotically chi-square distributed assuming
the observations are independent (10, 26). C_m is therefore the classical chi-square goodness of fit statistic
for testing multinomial hypotheses (10, 26).
180
The CM assesses 'information', Inf, by comparing the width of the confidence intervals provided by each
model with the 'intrinsic range' of each item (10, 26). The intrinsic range for an item is defined as the
difference between the largest of the forecasted and observed values and the smallest of the forecasted
and observed values (10, 26). This range is then expanded slightly by multiplying it by a user-defined
expansion factor, 1+ F, where F is typically a small fraction (for our data, the standard 10% was used)
(10, 26). Using this framework, the information score for each model and each item is defined as:
187
188 $$Inf_{m,i} = \Sigma_{k=1,\ldots,6}(p_k * \ln(p_k / r_k))$$

189
where r_k is the probability from a uniform (or log-uniform) probability density function over the intrinsic
range (10, 26). Models which concentrate their forecasts in a narrow range will have high information
scores. Therefore, while a precision score of 1 is optimal (i.e., all predictions are equally close to each
other relative to the observation), the higher the information score, the better (i.e., the tighter the forecast
distribution, the more information the forecast provides). See *SI appendix*, **Note 1** for more details on the
CM.
196
197 To illustrate the varying model performances, **Fig. 1** provides a forest plot of forecasts reported by the
198 individual models, as well as the constructed ensembles, during the summer 2020 period. A forest plot for
199 the winter 2021 period is provided by *SI appendix*, **Fig. S1**.
200
201   **2.2. Individual Model Performance**
202
203 The predictive and probabilistic performance results for each individual forecasting model are
204 summarized in **Fig. 2** for both the summer 2020 and winter 2021 time periods. More quantitative detail on
205 the performance results is provided in *SI appendix*, **Table S1**.
206
Focusing initially on the predictive performance of the models during the summer 2020 period, we see
that – (i) all models, except model C, had a GM greater than 1, indicating a systemic underprediction of
COVID-19 mortality, (ii) typically model predictions had little bias when compared with the observation (≤
35%) and precision within a factor of 2, (iii) models A, D, and I had excellent accuracy (GM = 1.01, 1.11,
and 1.09, respectively) and good precision (GSD = 1.57, 1.71, and 1.47, respectively), and (iv) models B,
C, and E had substantial bias (> 0.35%) and models B, E, G, and H had far worse precision (> 2). When
focusing on the probabilistic performance of models during this period, only three (or perhaps, four) of the
nine models considered performed at all well – models A, D, F, and to a lesser extent, G. The models that
had the highest information scores (models B, E, H, and I) all had extremely low calibration scores (<<
0.01), suggesting 'overconfidence' – i.e., that their stated confidence intervals were far too narrow while
simultaneously poorly capturing the true value.
218
Looking at the predictive performance during the winter 2021 period, we see that – (i) models F, G, and H
no longer provide forecasts, (ii) all six remaining models, except model E, had a GM less than 1,
indicating systemic overprediction of mortality, (iii) the remaining models again had little bias (< 30%) with
precision within a factor of 2, except model C, (iv) model D exhibited even better accuracy (GM = 0.99)
and precision (GSD = 1.33) than it did in the summer 2020 period, and model E now had excellent
accuracy (GM = 1.04) and better precision (GSD = 1.52), (v) model A had good precision (GSD = 1.53)
but so-so accuracy (GM = 0.76), and (vi) the models generally had better accuracy and precision than the
summer 2020 period. For the probabilistic model performance, models A and D exhibited the best
calibration scores (0.17 and 0.21, respectively), and again models B, E, and I were informative, but their
calibration was incredibly poor (<<0.01). Interestingly, unlike accuracy and precision, model calibration
and information scores generally seemed to be worse during the winter period.
230
231 The random expert hypothesis was tested to assess whether the putative differences in performance
232 between models was due to noise. This hypothesis was rejected for both periods of assessment. More
233 detail on this analysis can be found in *SI appendix*, **Note 2**.
234
Additional sensitivity analyses were conducted to evaluate whether the pervasive overconfidence in
individual model forecasts was due to our selection of states with extreme case rates (i.e., only states with
high or low case rates; no states with middling case rates). We carried out two sensitivity analyses, where
– (i) forecasts for four states with middling case rates were added to the main dataset, and (ii) forecasts
for six states with middling case rates were analyzed on their own. For these sensitivity analyses, it was
expected that, if the models were sensitive to state selection by case rate, model forecasts would exhibit
more appropriate confidence and substantially improve in performance. However, as illustrated by *SI
appendix*, **Fig. S2-5**, while some models for summer 2020 demonstrated slight improvements in
performance and exhibited slightly less confidence (i.e., lower information scores), overconfidence was
still present in the same models and all models demonstrated declines in performance for the winter 2021

period. In fact, for winter 2021, all models, except Model D, had increased confidence (i.e., higher
information scores) while calibration scores either remained very low (i.e., <0.01) or declined to
insignificance (i.e., <0.05).
248
249 **2.3. Ensemble Model Designs**
250
251 It is possible that better and more robust estimates might be obtained by producing an ensemble based
252 on weighted combinations of the forecasts provided by the individual models. Two approaches of
253 potential interest were – (i) equal-weighted combinations, and (ii) performance-weighted combinations. In
254 addition to a density-averaged equal-weighted ensemble, two density-averaged performance-weighted
255 ensembles were considered, based on – (a) 'predictive-performance' weights and (b) CM weights.
256
257 For 'predictive-performance' weighting, each random variable from each individual model was weighted in
258 inverse proportion to the model's predictive variance (i.e., inverse-variance weighting the distribution of
259 accuracy ratios of observations to predictions for each individual model):
260
261 $$\text{Weight}_m \propto 1 / \text{Var}(R_i, \ldots, R_N)$$
262
263 This weighting method is based on, but is not to be confused with, the inverse-variance weighting method
264 commonly used in meta-analysis (27).
265
For CM weighting, based on the theory of proper scoring rules and in accordance with the CM method,
performance weights were calculated as the product of both calibration and information scores (10, 26).
The CM-weighted ensemble also used a threshold significance level of 0.05 (i.e., a model with a
calibration score that is less than 0.05 was unweighted) for its normalized CM-weights to gain robustness
without significant loss in performance (13, 26). More detail on combining "experts" (or, in this case,
models) and why this cutoff was used is available in *SI appendix*, **Notes 3 and 4**.
272
273 The equal-weighted ensemble's forecasts were established by equally weighting the probability densities
274 of individual model forecasts, not their quantiles, to obtain aggregate forecasts. Meanwhile, for both the
275 predictive-performance-weighted and the CM-weighted ensembles, we applied their normalized model
276 performance weights to the probability densities of the individual model forecasts to obtain aggregate
277 performance-based forecasts.
278
279 **2.4. Ensemble Model Performance**
280
281 The predictive and probabilistic performance results for each of the density-averaged ensemble models
282 are summarized in **Table 1** for both the summer 2020 and winter 2021 time periods.
283
Both performance-weighted ensembles would be expected to outperform the equal-weighted ensemble
(23, 24) and they did fulfill this expectation on almost all measures of performance, however, the
differences in predictive performance were often small. The accuracy of the performance-based
combinations was appreciably better than the equal-weighted combination in the summer 2020 data, but
this advantage was not seen in the winter 2021 data. Very small differences in precision were observed in
both data sets.
290
To explore whether these differences in accuracy between the summer 2020 and winter 2021 periods
were because three of the nine models included in the summer 2020 analysis were not reflected in the
winter 2021 data, we reassessed model performance in the summer of 2020 using only data from the six
models that were available during both seasons. The differences in accuracy seen in the summer of 2020
between the equal-weighted and performance-weighted combinations were even greater (equal weight =
1.23, predictive weight = 0.94, CM unnormalized weight = 1.02) than those observed in the full data set.
More detail on this analysis is provided in *SI appendix*, **Table S2**.
298
Though predictive performance differences were small, there were substantial differences in probabilistic
performance. The calibration scores of the CM-weighted ensemble (0.54 in summer 2020; 0.44 in winter

2021) were far better than those of the equal-weighted ensemble (0.03 during the summer, 0.17 during
the winter) or the predictive-performance-weighted ensemble (0.04 during the summer, 0.25 during the
winter). There were also small differences in information scores, where again the CM-weighted ensemble
slightly outperformed the other two ensembles – resulting in substantially higher CM weights during both
periods.

2.5. Performance by Domain

We also compared the performance of the individual models across three domains of interest – (i) race
(i.e., states which are heavily non-Hispanic White vs. states with relatively large non-Hispanic Black
populations), (ii) COVID-19 case rates (i.e., states with a relatively low amount of weekly cases per
100,000 population vs. states with a relatively high amount of weekly cases per 100,000 population), and
(iii) forecast period (i.e., forecasts of mortality for the upcoming week vs. forecasts of mortality for the
week ending four weeks from the date on which the forecast was made). The performance of the equal-
weighted density-averaged ensemble stratified by these three different domains (i.e., eight forecasts per
domain) is provided in **Table 2**.
315
316 Both the accuracy and precision of predictions for states with relatively high non-Hispanic Black
317 populations were better than for states with a largely non-Hispanic White population. This difference
318 appeared to be stable and was seen in both the summer 2020 and winter 2021 data. However, there is
319 no evidence of any stable differences in the calibration and information scores of forecasts depending on
320 the racial composition of the state.
321
322 The summer 2020 data suggests that predictions were more accurate and precise in states with low case
323 rates, and this remained true in the more recent winter 2021 data, but the difference in accuracy is not as
324 substantial. On the other hand, model calibration scores were somewhat better in states with high case
325 rates. This difference appeared to be stable and was seen with both the summer 2020 and winter 2021
326 data.
327
328 Lastly, as expected, the predictions of deaths in the near-term appear to be more accurate and precise
329 than those in the mid-term, especially for the summer 2020 data. This remained true for the more recent
330 winter 2020 data, but the differences in both accuracy and precision were smaller. There is some
331 evidence that calibration scores were better for near-term forecasts in the summer data, but this did not
332 persist in the winter data.
333

3. Discussion

335 There has been little published peer-reviewed information regarding the performance of COVID-19
336 forecasting models, and what is available has focused mostly on the performance of their central
337 estimates (i.e., predictive performance). Very little has been reported on the performance of their
338 uncertainty estimates (i.e., probabilistic performance), which could inform users how often they would be
339 surprised by observations outside of a model's forecasted confidence intervals.
340
To address this gap in knowledge, we borrowed from the formally elicited expert judgment literature to
demonstrate a commonly used approach for resolving this issue (i.e., the Classical Model; CM). We then
evaluated both the predictive (i.e., accuracy and precision) and probabilistic (i.e., calibration and
information) performance of COVID-19 forecasting models during two independent eight-week time
periods (i.e., summer 2020 and winter 2021). In our analysis, some models exhibited good predictive
performance but were found to have poor probabilistic performance (e.g., model I, and for the winter, E),
and vice versa. Of the nine selected forecasting models, only two models, A and D, did reasonably well in
both predictive and probabilistic performance, with model D demonstrating the most consistent
dominance in all performance measures.
350
351 This study was the first to utilize the CM method with COVID-19 forecasting models to measure their
352 probabilistic performance and these CM metrics provided valuable insights. The fact that all but three of

353 the nine models considered (models A, D, and F) had such low calibration scores for both time periods
354 and that most of those also had high information scores indicates pervasive overconfidence.
355
There is strong evidence that most expert judgments are overconfident, (15, 28, 29) that is why there are
often elaborate procedures to minimize this in expert elicitations (30). It is understandably difficult to
predict changes in human behavior, and, while rare, there are occasional anomalies in reported data
(e.g., large revisions or bulk reporting) that may further bias a model's forecast. This issue has been
highlighted in other assessments of these models (22, 31). While, to the best of our knowledge, there
were no such anomalies demonstrated in the observed data utilized for this study, it is clear from these
results and the literature that many modelers need to carefully re-evaluate how they are quantifying
uncertainty in their predictions.
364
We found that the tendency of models to underpredict in the summer of 2020 was replaced by a tendency
to overpredict in the winter of 2021 – suggesting that model predictions lag the actual changes in disease
rates, which were generally increasing in the summer of 2020 and generally decreasing in the winter of
2021. This relative unresponsiveness to more rapid shifts has been well documented by the COVID-19
Forecast Hub and their studies (7, 22). This also could be the reason why model forecasts generally
performed worse during winter 2021 versus summer 2020, as many states experienced rapid and
substantial declines in weekly mortality counts during late winter 2021.
372
373 All the assessed individual models were variations of a susceptible-exposed-infectious-removed (SEIR)
374 compartmental model (32-40). They all forecasted at the U.S. state level, and some also forecasted at the
375 national and county levels (8). Every model provided forecasts at the daily scale, which could then be
376 aggregated to the weekly scale for at least four weeks ahead from forecast date (8).
377
However, while all models used mortality data to inform compartment transition rates, they varied in how
they incorporated case, hospitalization, demographic, and mobility data (32-40). Both models A and D
utilized case and mortality data, and model A utilized demographic data, but neither model incorporated
hospitalization nor mobility data (32, 35). Using case and mortality data, models A and D extrapolated
deaths from lagged case counts, with a 14-day lag for model A (32) and a varying 15- to 30-day lag for
model D (35). Additionally, while models A and D, like most other models, incorporated time-varying
COVID-19 transmission rates which could indirectly reflect social distancing measures, neither model
directly incorporated changes in social distancing measures for forecasts in the near-term (although,
model A may have incorporated changes in social distancing measures for forecasts past three weeks)
(32, 35).
388
The designs of these forecasting models have also not been static. Modeling groups have been learning
and adapting their models over the course of the pandemic. Thus, a model's forecasts for the summer
and winter are broadly comparable, but its methods to produce the forecasts are not necessarily identical.
This further emphasizes the significance of our integration of expert judgment methods, as we essentially
assessed the performance of 'judgments' from model forecasting groups rather than from the models
themselves, which are ever evolving.
395
For both time periods, we constructed and evaluated a density-averaged predictive-performance and CM-
performance-weighted ensemble, which were respectively based on the predictive and probabilistic
performance of the individual forecasting models. Both the performance-weighted ensembles
outperformed the equal-weighted ensemble, although the differences in predictive performance were
modest. This result is anticipated in the Cooke, Marti, & Mazzuchi (2021) study (16). In contrast, when
evaluated in terms of probabilistic performance there were clear advantages to performance weighting
using the CM weights. While a density-averaged equal-weighted ensemble may still hold merit (e.g.,
improved performance scores over many individual models; does not require prior forecasts/realizations
to train the ensemble; easy to produce), for best results, particularly when probabilistic performance is
considered, CM performance-weighting should be utilized. Additionally, the CM method removes from
aggregation those models that are not able to appropriately characterize the uncertainty in their
predictions, and thus should not be relied upon. In any case, if performance-based modeling is not
desired, ensembles should at least be constructed based on averaged densities and not quantiles bins.

409
Across the three domains and two performance attributes considered, consistent and substantial
performance differences were seen only in a few instances – and these differed depending on whether
one focuses on predictive or probabilistic performance. Predictive performance was generally better for
forecasts in states with relatively high non-Hispanic Black populations and relatively low case rates, but
the observed differences were often modest. Probabilistic performance was consistently better when case
rates were high. This difference in performance based on case rates could be explained by models
having difficulty adapting to rapid shifts (reflected in states with high case rates), but this added
uncertainty may be more appropriately reflected in their forecasts. Finally, as expected based on previous
study findings (9, 31) and the disease progression timeline (i.e., cases now strongly predict deaths in 17-
21 days), (41) near-term predictions (i.e., deaths in the next week) outperformed mid-term predictions
(i.e., deaths in the week ending four weeks from forecast date). However, we would like to emphasize
that these results are not definitive and may not be representative of the results that would be based on
all models and all states.
423
424 One study, currently in preprint, also evaluated the predictive and probabilistic accuracy of individual
425 models (using different metrics) and found that accuracy degraded as models made predictions further
426 into the future and that individual models were frequently overconfident (31). The authors also found that
427 the models with few data inputs were among the most accurate and the same models demonstrated
428 superior performance in their analysis (31). Thus, many of their findings agreed with ours.
429
430 It is also important to note that the CM is not the only protocol that analyzes and aggregates expert
431 judgments (or, in this case, model forecasts). Although other expert judgment protocols that typically
432 involve behavioral aggregation could not be utilized in this post hoc analysis, (17, 18) it would be
433 interesting to elicit from modeling groups a "best" model design and this could be the focus of a separate
434 paper.
435
Our sense is that these results are more suggestive than conclusive, because – (i) due to the selection
criteria, only nine models were examined, and three of these ceased forecasting by the winter 2021
period; (ii) they are based on model performance in one eight-week period in the summer of 2020 and
one eight-week period in the winter of 2021; (iii) they are limited to eight states (although more states
were examined for sensitivity in *SI appendix,* **Fig. S2-S5**); and (iv) our analysis did not attempt to explain
what model design choices led to better or worse performance.
442
If we were to reexamine the data, we would have analyzed all models that met our criteria for the winter
period, not just those that were analyzed for the summer and continued forecasting into the winter.
Additionally, in the expert judgment literature, 8 to 15 calibration variables are standard and any more is
thought to be unnecessary (23). Thus, we believe this analysis to be an adequate demonstration of the
CM for the selected models for those selected time periods and states. However, we understand that,
considering the limited scope of this analysis, and with the situation and the modeling groups' ability to
forecast always changing, for more definitive conclusions on the performance of the specific individual
models, a broader evidence base should be utilized. Excalibur, the program that was used to run the CM,
was designed for the analysis of formally elicited expert judgment with limited calibration questions (42).
In the future, we hope to adapt the Excalibur code so that it can handle the full set of data available for
the evaluation of COVID-19 models.
454
Conversely, our analysis has several strengths – (i) it evaluated the performance of a set of leading
models which have been used to forecast COVID-19 mortality in the US; (ii) it considered *both* predictive
(i.e., accuracy and precision) and probabilistic (i.e., calibration and information) performance, and was the
first study to utilize the CM method to accomplish this; (iii) it assessed several approaches for
constructing *density-averaged* ensemble models, and the CM-weighted ensemble exhibited superior
performance; (iv) it examined performance differences across states selected to reflect differences in
racial composition and variations in COVID-19 case rate; and (v) it considered data from two distinct
phases of the pandemic.
463

We conducted this analysis to demonstrate the limitations of performance evaluations which consider
only predictive performance. While we believe this analysis of individual COVID-19 forecasting models to
be adequate, considering its limited scope and the dynamic model designs, we do not believe that our
results provide definitive conclusions about the performance of the specific models considered. Rather,
we suggest the availability, utility, and importance of an approach that has been used for years in the
expert judgment literature for characterizing the probabilistic performance of such forecasts and
constructing high performing ensembles. We believe this application demonstrates that the CM could be
a useful methodology not just for COVID-19 modeling but other applications as well. We hope that users
will find this approach to be valuable in their efforts to better characterize and communicate forecast
performance.
474

475 **4. Materials and Methods**

476 Our analysis involved a comparison of model forecasts with subsequent observations of weekly deaths
477 from COVID-19 in four states (Idaho, Louisiana, New York, and Maine) over an eight-week period during
478 the summer of 2020 and four other states (Georgia, Maryland, Vermont, and Wyoming) over a
479 subsequent eight-week period during the winter of 2021.
480
The states considered in our analysis were selected based on recent case rates of COVID-19
(cases/100,000 population within the previous week) (43) and racial composition (majority non-Hispanic
Black vs. majority non-Hispanic White) (44). Changes in COVID-19 case rates were of interest as they
precede changes in COVID-19 death rates by a period of about 18.5 days (41). Racial composition was
of interest as the COVID-19 mortality rate for non-Hispanic Black Americans at the time of analysis was
more than twice that of non-Hispanic White Americans, and this disparity persisted even when accounting
for other demographic and socioeconomic factors (45).
488
With these two domains in mind, our goal was to assess two states for each time period with relatively
high case rates (Idaho and Louisiana; Georgia and Vermont); two with relatively low case rates (Maine
and New York; Maryland and Wyoming); two with a relatively high fraction of population reported as non-
Hispanic Black (Louisiana and New York; Georgia and Maryland); and two with a relatively high fraction
of population reported as non-Hispanic White (Idaho and Maine; Vermont and Wyoming). This was done
to assess how models may perform forecasting for states under varying circumstances. More detail on
how the case rates and racial composition for states were determined, as well as how states were
selected, is available in *SI appendix*, **Note 5** and **Tables S3 and S4**.
497
498 We were also interested in the models' ability to forecast COVID-19 deaths in both the near-term and the
499 medium-term. Near-term performance was gauged using projected COVID-19 deaths in the week
500 immediately after the forecast was made. Medium-term performance was gauged using projected COVID-
501 19 deaths in the week ending four weeks after the forecast was made.
502
503 Our evaluations of model performance for the eight states and the two forecast periods of interest (week
504 ending one week in the future and week ending four weeks in the future) were examined twice within
505 each period – once for forecasts made on June 13th, 2020, or January 10th, 2021, and a second time for
506 forecasts made on July 11th, 2020, or February 7th, 2021 (no overlap in forecasts). In total, 16
507 comparisons of model forecasts with observed deaths were made for each time period and each model.
508
Of the many models providing data to the COVID-19 Forecast Hub's data repository (8), only the models
that provided weekly COVID-19 mortality forecasts made on the same day of the week, with forecasts at
the weekly timescale for all four of the initially assessed states and without missing forecasts were
selected for this analysis. These include: OliverWyman-Navigator (Model A) (32), MOBS-GLEAM_COVID
(Model B) (33), JHU_IDD-CovidSP (Model C) (34), UMass-MechBayes (Model D) (35), UCLA-SuEIR
(Model E) (36), YYG-ParamSearch (Model F) (37), UT-Mobility (Model G) (38), USACE-ERDC_SEIR
(Model H) (39), and Covid19Sim-Simulator (Model I) (40). To understand whether model performance
was unique to one phase of the pandemic or whether performance was consistent across the two phases,
these same models were assessed during the winter 2021 period.

518
519
Code Availability

520
521
522 Data was analyzed using Microsoft Excel and EXCALIBUR (a software package for using the Classical
523 Model) (42).
524
525
526 **Data Availability**
527
528 Observed state COVID-19 mortality and case data was gathered from the Centers for Disease Control
529 and Prevention (CDC) (43). State population and racial composition data was collected from one-year
530 estimates from the Census Bureau's 2018 and 2019 American Community Survey (ACS) (44). For racial
531 composition statistics and case rate data, please see *SI appendix*, **Tables S3 and S4.**
532
533 Model forecasting data was gathered from the COVID-19 Forecast Hub's publicly available structured
534 data storage repository on GitHub (8). For the model and ensemble predictions, their uncertainty
535 distributions, and the subsequent observations of COVID-19 mortality, please see *SI appendix*, **Tables
536 S5 and S6**.
537
538
539 **Acknowledgments**
540
541 We want to thank Willy Aspinall, Jouni Tuomisto, and Jacqueline Macdonald for contributing to our
542 thoughts about this and for their feedback on our early drafts of the paper.
543
544
545 **Funding Sources**
546
547 Kyle J. Colonna's involvement was funded by the Harvard Population Health Sciences PhD scholarship.
548 Roger M. Cooke's involvement was pro bono. John S. Evans' involvement was funded by the Department
549 of Environmental Health and the Harvard Cyprus Initiative at the Harvard T.F. Chan School of Public
550 Health.
551
552
553 **References**
554
1. Centers for Disease Control and Prevention (CDC), National Center for Emerging and Zoonotic
Infectious Diseases (NCEZID), Division of Global Migration and Quarantine (DGMQ),
Nonpharmaceutical Interventions (NPIs). *CDC* (2020). https://www.cdc.gov/nonpharmaceutical-
interventions/index.html (accessed 3 May 2021).
2. R. M. Anderson, H. Heesterbeek, D. Klinkenberg, T. D. Hollingsworth, How will country-based
mitigation measures influence the course of the COVID-19 epidemic? *Lancet* **395**, 931–934
(2020).
3. S. Flaxman, *et al.*, Estimating the effects of non-pharmaceutical interventions on COVID-19 in
Europe. *Nature* **584**, 257–261 (2020).
4. International Monetary Fund. (IMF) Research Dept., World Economic Outlook, April 2020: The
Great Lockdown. *IMF* (2020) (accessed 3 May 2021).
5. The United Nations Educational, Scientific and Cultural Organization (UNESCO), Adverse
consequences of school closures. *UNESCO* (2020).
https://en.unesco.org/covid19/educationresponse/consequences (accessed 3 May 2021).
6. M. Nicola, *et al.*, The socio-economic implications of the coronavirus pandemic (COVID-19): A
review. *Int J Surg* **78**, 185–193 (2020).
7. University of Massachusetts Amherst (UMass-Amherst), Reich Lab, The COVID-19 Forecast
Hub. *UMass-Amherst* (2021). *https://covid19forecasthub.org/* (accessed 3 May 2021).

8. University of Massachusetts Amherst (UMass-Amherst), Reich Lab, Data from "reichlab/covid19-forecast-hub." *Github*. Available at https://github.com/reichlab/covid19-forecast-hub. Deposited 3 May 2021.

9. J. Friedman, *et al.*, Predictive performance of international COVID-19 mortality forecasting models. *Nat Commun* **12**, 2609 (2021).

10. R. M. Cooke, *Experts in Uncertainty: Opinion and Subjective Probability in Science* (Oxford University Press, 1991).

11. R. M. Cooke, *et al.*, A Probabilistic Characterization of the Relationship between Fine Particulate Matter and Mortality: Elicitation of European Experts. *Environ. Sci. Technol.* **41**, 6598–6605 (2007).

12. T. Hald, *et al.*, World Health Organization Estimates of the Relative Contributions of Food to the Burden of Disease Due to Selected Foodborne Hazards: A Structured Expert Elicitation. *PLoS One* **11** (2016).

13. J. L. Bamber, M. Oppenheimer, R. E. Kopp, W. P. Aspinall, R. M. Cooke, Ice sheet contributions to future sea-level rise from structured expert judgment. *PNAS* **116**, 11195–11200 (2019).

14. K. Shea, *et al.*, COVID-19 reopening strategies at the county level in the face of uncertainty: Multiple Models for Outbreak Decision Support. *medRxiv* [Preprint] (2020). https://doi.org/10.1101/2020.11.03.20225409 (accessed 24 August 2021)

15. K. Shea, *et al.*, Harnessing multiple models for outbreak management. *Science* **368**, 577–579 (2020).

16. R. M. Cooke, D. Marti, T. Mazzuchi, Expert forecasting with and without uncertainty quantification and weighting: What do the data say? *International Journal of Forecasting* **37**, 378–387 (2021).

17. A. M. Hanea, M. F. McBride, M. A. Burgman, B. C. Wintle, Classical meets modern in the IDEA protocol for structured expert judgement. *Journal of Risk Research* **21**, 417–433 (2018).

18. A. O'Hagan, Expert Knowledge Elicitation: Subjective but Scientific. *The American Statistician* **73**, 69–81 (2019).

19. E. L. Ray, *et al.*, Ensemble Forecasts of Coronavirus Disease 2019 (COVID-19) in the U.S. *medRxiv* [Preprint] (2020). https://doi.org/10.1101/2020.08.19.20177493 (accessed 3 May 2021)

20. Centers for Disease Control and Prevention (CDC), National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases, Forecasts of COVID-19 Deaths. *CDC* (2020). https://www.cdc.gov/coronavirus/2019-ncov/covid-data/forecasting-us.html (accessed 3 May 2021).

21. L. C. Brooks, *et al.*, Comparing ensemble approaches for short-term probabilistic COVID-19 forecasts in the U.S. *International Institute of Forecasters* (2020).

22. E. L. Ray, *et al.*, Challenges in training ensembles to forecast COVID-19 cases and deaths in the United States. *International Institute of Forecasters* (2021).

23. A. R. Colson, R. M. Cooke, Cross validation for the classical model of structured expert judgment. *Reliability Engineering & System Safety* **163**, 109–120 (2017).

24. A. R. Colson, R. M. Cooke, Expert Elicitation: Using the Classical Model to Validate Experts' Judgments. *Rev Environ Econ Policy* **12**, 113–132 (2018).

25. J. Bamber, W. Aspinall, R. Cooke, A commentary on "how to interpret expert judgment assessments of twenty-first century sea-level rise" by Hylke de Vries and Roderik SW van de Wal. *Climatic Change* **137**, 321–328 (2016).

26. R. Cooke, L. Goossens, Procedures Guide for Structured Expert Judgment. *European Communities, Luxembourg, EUR* (2000).

27. J. Hartung, G. Knapp, B. K. Sinha, *Statistical Meta-Analysis with Applications* (Wiley, 2008).

28. P. E. Tetlock, D. Gardner, Superforecasting: The Art and Science of Prediction. *Random House* (2016).

29. M. A. Burgman, Trusting Judgements: How to Get the Best out of Experts. *Cambridge Univ. Press* (2015).

30. M. G. Morgan, Use (and abuse) of expert elicitation in support of decision making for public policy. *PNAS* **111**, 7176–7184 (2014).

31. E. Y. Cramer, *et al.*, Evaluation of individual and ensemble probabilistic forecasts of COVID-19 mortality in the US. medRxiv [Preprint] (2021). https://doi.org/10.1101/2021.02.03.21250974 (accessed 24 August 2021)

32. Oliver Wyman, Oliver Wyman COVID-19 Pandemic Navigator. *Oliver Wyman* (2021).
https://pandemicnavigator.oliverwyman.com/ (accessed 3 May 2021).
33. Laboratory for the Modeling of Biological + Socio-Technical Systems (MOBS), COVID-19
Mobility. *The Gleam Project* (2021). https://covid19.gleamproject.org/mobility (accessed 3 May
2021).
34. Infectious Disease Dynamics (IDD) at Johns Hopkins Bloomberg School of Public Health
(JHSPH), Projects | COVID-19. *IDD* (2021). http://www.iddynamics.jhsph.edu/projects/covid-19
(accessed 3 May 2021).
35. D. Sheldon, C. Gibson, N. Reich, Data from "*dsheldon/covid*." *Github*. Available at
https://github.com/dsheldon/covid . Deposited on 12 April 2021.
36. Statistical Machine Learning Lab (ML) at the University of California, Los Angeles (UCLA),
UCLAML Combating COVID-19. *UCLAML* (2021). https://covid19.uclaml.org/index.html
(accessed 3 May 2021).
37. Y. Gu, COVID-19 Projections Using Machine Learning. *Youyang Gu* (2021). https://covid19-
projections.com/ (accessed on 3 May 2021).
38. COVID-19 Modeling Consortium at the University of Texas at Austin (UT Austin), COVID-19
Mortality Projections for US States. *Texas Advanced Computing Center* (2021). https://covid-
19.tacc.utexas.edu/dashboards/us/ (accessed 3 May 2021).
39. United States Army Corps of Engineers (USACE), Engineer Research and Development Center
(ERDC) , Data from "*erdc-cv19/seir-model*." *Github*. Available at https://github.com/erdc-
cv19/seir-model. Deposited on 3 August 2020.
40. Massachusetts General Hospital (MGH) Institute for Technology Assessment, COVID-19
Simulator. *MGH Institute for Technology Assessment* (2021). https://covid19sim.org/ (accessed 3
May 2021).
41. University of Washington School of Medicine (UWSOM), Institute for Health Metrics and
Evaluation (IHME), Daily Deaths. *IHME* (2021). https://covid19.healthdata.org/united-states-of-
america?view=daily-deaths&tab=trend (accessed 29 August 2021).
42. LightTwist Software, Excalibur. *LightTwist Software.* https://lighttwist-software.com/excalibur/
(accessed 3 May 2021).
43. Centers for Disease Control and Prevention (CDC), COVID-19 Response, COVID-19 Case
Surveillance Public Data Access, Summary, and Limitations. *CDC* (2021).
https://data.cdc.gov/Case-Surveillance/United-States-COVID-19-Cases-and-Deaths-by-State-
o/9mfq-cb36 (accessed 3 May 2021).
44. United States Census Bureau (UCSB), American Community Survey (ACS) Data. *USCB* (2021).
https://www.census.gov/programs-surveys/acs/data.html (accessed 3 May 2021).
45. Centers for Disease Control and Prevention (CDC), National Center for Immunization and
Respiratory Diseases (NCIRD), Division of Viral Diseases, Hospitalization and Death by
Race/Ethnicity. *CDC* (2020). https://www.cdc.gov/coronavirus/2019-ncov/covid-
data/investigations-discovery/hospitalization-death-by-race-ethnicity.html (accessed 18 August
2020).

Figures Legends & Tables

Fig. 1: A forest plot of forecasts provided by the individual models and constructed ensembles for each of the variables of interest for summer 2020. The horizontal axis represents deaths during the week in question, the true values are given by the red vertical lines, the error bars indicate the 5th and 95th percentiles, and the dots represent the model's predictions. A forest plot of forecasts for winter 2021 is provided in *SI appendix*, **Fig. S1.**

Fig. 2. Accuracy (GM) vs. precision (GSD) of the central estimates provided by the individual models for summer 2020 (*panel A*) and winter 2021 (*panel B*) periods. Calibration vs. information scores based on the forecast densities provided by the individual models for the summer 2020 (*panel C*) and winter 2021 (*panel D*) periods. The optimal accuracy, precision, and calibration scores are 1, while the higher the information score the better.

681 **Table 1. Ensemble model performance for summer 2020 (*Top*) and winter 2021 (*Bottom*)**
682

Ensemble	Summer 2020				
	Accuracy - GM	Precision - GSD	Calibration	Information	CM Weight (Unnormalized)
Equal Weight	1.24	1.60	0.03	0.72	0.02
Performance – Predictive Weight	0.91	1.41	0.04	0.55	0.02
Performance – CM Weight*	1.12	1.57	0.54	0.81	0.43

Ensemble	Winter 2021				
	Accuracy - GM	Precision - GSD	Calibration	Information	CM Weight (Unnormalized)
Equal Weight	0.84	1.55	0.17	0.36	0.06
Performance – Predictive Weight	0.84	1.49	0.25	0.35	0.09
Performance - CM Weight*	0.85	1.45	0.44	0.43	0.19

683
684 * A hypothesis rejection significance level of 0.05 was used for this ensemble. More detail on why this was used is
685 available in *SI appendix*, **Notes 3 and 4**.
686
Abbreviations: CM = Classical Model; GM = geometric mean; GSD = geometric standard deviation

Table 2: Equal-weighted ensemble model performance for summer 2020 (*Top*) and winter 2021 (*Bottom*) stratified by different domains

Sub-Domain	Summer 2020				
	Accuracy - GM	Precision – GSD	Calibration	Information	CM Weight (Unnormalized)
High % non-Hispanic White (ID, ME; VT, WY)	1.30	1.86	0.60	0.86	0.52
High % non-Hispanic Black (LA, NY; GA, MD)	1.19	1.34	0.05	0.59	0.03
High Case Rate (ID, LA; VT, GA)	1.42	1.77	0.32	0.67	0.21
Low Case Rate (ME, NY; WY. MD)	1.09	1.39	0.14	0.78	0.11
One Week Ahead	0.95	1.41	0.27	0.53	0.14
Four Weeks Ahead	1.62	1.56	0.05	0.92	0.05

Sub-Domain	Winter 2021				
	Accuracy - GM	Precision – GSD	Calibration	Information	CM Weight (Unnormalized)
High % non-Hispanic White (ID, ME; VT, WY)	0.69	1.56	0.14	0.36	0.05
High % non-Hispanic Black (LA, NY; GA, MD)	1.02	1.42	0.73	0.37	0.27
High Case Rate (ID, LA; VT, GA)	0.81	1.76	0.66	0.41	0.27
Low Case Rate (ME, NY; WY. MD)	0.87	1.34	0.16	0.31	0.05
One Week Ahead	0.88	1.51	0.02	0.35	0.01
Four Weeks Ahead	0.80	1.62	0.06	0.37	0.02

Abbreviations: CM = Classical Model; GM = geometric mean; GSD = geometric standard deviation; GA = Georgia; ID = Idaho; LA = Louisiana; MD = Maryland; ME = Maine; NY = New York; VT = Vermont; WY = Wyoming
Note: Analogues of **Table 2** for the CM- and predictive-performance-weighted ensembles are also provided in *SI appendix*, **Table S7 and S8**, respectively.