

- 32 12. Laboratory of Medical Investigation in Pathogenesis and Directed Therapy in
33 Onco – Immuno – Hematology (LIM-31) HCFMUSP, University of São Paulo
34 Medical School, São Paulo, Brazil.
- 35 13. Epidemiologic Surveillance Center, Hospital das Clinicas – University of São
36 Paulo Medical School, São Paulo, Brazil.
- 37 14. Mathematical Sciences, University of Southampton, Southampton, United
38 Kingdom.
- 39 15. Department of Preventive Medicine, University of São Paulo Medical School
- 40 16. MRC Centre for Global Infectious Disease Analysis, J-IDEA, Imperial College
41 London, London, UK.
- 42 17. Oxford School of Global and Area Studies, University of Oxford, Oxford, United
43 Kingdom.

44

45 *These authors contributed equally.

46

47 #Corresponding authors:

48 1. Sabrina L. Li, School of Geography and the Environment, University of
49 Oxford, South Parks Rd, Oxford, OX1 3QY, United Kingdom. E-mail:
50 sabrina.li@ouce.ox.ac.uk

51

52 2. Rafael H.M. Pereira, Institute for Applied Economic Research (Ipea), SBS -
53 Quadra 1 - Bloco J - Ed. BNDES - CEP: 70076-900 - Brasília - DF – Brasil. E-
54 mail: rafael.pereira@ipea.gov.br

55

56

57

58 **Summary**

59

60 **Background:** Little evidence exists on the differential health effects of COVID-19 on
61 disadvantaged population groups. Here we characterise the differential risk of
62 hospitalisation and death in São Paulo state, Brazil and show how vulnerability to
63 COVID-19 is shaped by socioeconomic inequalities.

64 **Methods:** We conducted a cross-sectional study using hospitalised severe acute
65 respiratory infections (SARI) notified from March to August 2020, in the *Sistema de*
66 *Monitoramento Inteligente de São Paulo* (SIMI-SP) database. We examined the risk
67 of hospitalisation and death by race and socioeconomic status using multiple datasets
68 for individual-level and spatio-temporal analyses. We explained these inequalities
69 according to differences in daily mobility from mobile phone data, teleworking
70 behaviour, and comorbidities.

71 **Findings:** Throughout the study period, patients living in the 40% poorest areas were
72 more likely to die when compared to patients living in the 5% wealthiest areas (OR:
73 1.60, 95% CI: 1.48 – 1.74) and were more likely to be hospitalised between April and
74 July, 2020 (OR: 1.08, 95% CI: 1.04 – 1.12). Black and *Pardo* individuals were more
75 likely to be hospitalised when compared to White individuals (OR: 1.37, 95% CI: 1.32
76 – 1.41; OR: 1.23, 95% CI: 1.21 – 1.25, respectively), and were more likely to die (OR:
77 1.14, 95% CI: 1.07 – 1.21; 1.09, 95% CI: 1.05 – 1.13, respectively).

78 **Interpretation:** Low-income and Black and *Pardo* communities are more likely to die
79 with COVID-19. This is associated with differential access to healthcare, adherence
80 to social distancing, and the higher prevalence of comorbidities.

81 **Funding:**

82 This project was supported by a Medical Research Council-São Paulo Research
83 Foundation (FAPESP) CADDE partnership award (MR/S0195/1 and FAPESP
84 18/14389-0) (<http://caddecentre.org/>). This work received funding from the U.K.
85 Medical Research Council under a concordat with the U.K. Department for
86 International Development.

87

88 **Introduction**

89 The COVID-19 pandemic has amplified the effects of social inequalities on exposure
90 and death in low socioeconomic groups ¹, particularly in Brazil, where it has caused
91 significant mortality ². Few studies have addressed the uneven impact of COVID-19
92 by socioeconomic status and race ³⁻⁵, especially in low and middle-income countries,
93 in part because national surveillance systems seldom collect or report this information
94 ⁶. In Brazil, higher risk of COVID-19 death has been found for Black and *Pardo* (mixed
95 ethnicity) Brazilians ⁷. Nonetheless, there is still little information on how the differential
96 health outcomes of COVID-19 are shaped by broader social inequalities that
97 determine the capacity to adhere to social distancing and non-pharmaceutical
98 interventions (NPIs).

99 It is paramount to understand the potential social drivers of COVID-19 morbidity and
100 mortality, particularly in countries with high inequality such as Brazil ⁸. The first COVID-
101 19 cases in Brazil were detected in São Paulo ⁹, the most populous state and home
102 to diverse racial groups. In the Brazilian context of politically polarised public health
103 responses ¹⁰, São Paulo has been severely affected by COVID-19 ¹¹ and access to
104 testing has been limited for low-income populations ¹². By combining multiple high-
105 resolution data sources, we conducted a multi-scale analysis to investigate the risk of
106 hospitalisation and death from severe acute respiratory infections (SARI),
107 predominantly caused by COVID-19 ¹², by race and socioeconomic status in the state
108 of São Paulo. We examined potential drivers of these inequalities by evaluating local
109 levels of population adherence to social distancing, access to teleworking, and
110 prevalence of comorbidities

111 **Methods**

112 **Data Sources**

113 **Severe acute respiratory infections (SARI) and patient information**

114 Patient-level information on demographic characteristics, home address,
115 hospitalisation, and health outcomes were collected from the São Paulo State Health
116 Secretariat SARI hospitalisations database (SIMI-SP) ¹³. SARI can be caused by
117 SARS-CoV-2 and is defined by the Brazilian Ministry of Health as flu-like syndrome
118 plus one of the following: dyspnoea, persistent chest pain, or hypoxia. All SARI cases
119 and deaths are notified in the SIMI-SP database, regardless of hospitalisation.

120 We included all SARI related hospitalisations and deaths notified in São Paulo state
121 between March 15 and August 29, 2020. Given that recent data is incomplete due to
122 reporting delays ¹⁴ and to avoid biases, we limited our analysis to patients with
123 symptoms onset between these dates (epidemiological weeks 10 – 35) (figure 1B).
124 We also included SARI cases with unknown aetiology, as those are likely related to
125 COVID-19 but not lab-confirmed due to low rates of COVID-19 testing in Brazil ¹⁵ and
126 socioeconomic bias in testing ¹².

127 Zip code information was only available for cases reported in São Paulo state. Data
128 was geocoded using the patient's self-reported home address or postal code with
129 Galileo (www.img.com.br) and Google API. Information on the health facility where
130 each case was notified was linked to the National Registry of Health Facilities (CNES),
131 which includes information on the mode of health care provision (public and private).

132 The race of patients was partially self-declared and partially identified by a health
133 professional. Race was categorised as either "White", "Black", "Asian" (East or
134 Southeast Asian), "*Pardo*" (mixed ethnic ancestries with diverse skin colours)¹⁶ or
135 Indigenous. Race information was missing for 53 480 (23.9%) of retrieved SARI
136 cases, and were imputed using the racial distribution of the census tract of residence
137 (see Supplementary Appendix A for details). About 0.1% of the population in São

138 Paulo self-identified as indigenous ¹⁷ and since only 166 indigenous patients (0.07%)
139 were recorded, they were not considered in our analysis.

140 **Socioeconomic data**

141 We obtained data on municipality-level socioeconomic factors from the latest
142 population census (2010) compiled by the Brazilian Institute of Geography and
143 Statistics ([IBGE](https://www.ibge.gov.br/)). We selected household income per capita, population density, and
144 income inequality (Gini Index). We also determined the proportion of residents with a
145 primary education or lower, employment-to-population ratio, and the proportion of the
146 working population without a formal labour market contract or social security. The road
147 network distance from the centroid of each census tract to the nearest healthcare
148 facility was computed considering all facilities that hospitalised SARI patients via the
149 public healthcare system (SUS). Information on employment status and comorbidities
150 during the epidemic was retrieved from the National Household Sample Survey
151 (PNAD COVID-19), a national telephone survey conducted by IBGE with over 1 888
152 560 interviews between May and September 2020. Details are described in
153 Supplementary Appendix A.

154 **Daily mobility and non-pharmaceutical interventions**

155 To assess local-level adherence to social distancing, we used daily mobile phone data
156 provided by In-LoCo (<https://www.inloco.com.br/covid-19>)¹⁸ for the *Região*
157 *Metropolitana de São Paulo* (greater metropolitan area of São Paulo: RMSP). This
158 data was aggregated using a hexagonal grid based on the global H3 index at
159 resolution 8. Each cell has an edge of approximately 460 meters and an area of 0.74
160 km² (<https://h3geo.org/docs/core-library/restable>). For each H3 cell, the social
161 isolation index was measured as the number of people who did not leave their cell of
162 residence during the day, divided by the number of residents in that cell. Each mobile
163 phone was assigned to an H3 cell based on the owner's location of residence during
164 the evening and their previous travel history. The racial composition and income level
165 of each cell were determined using dasymetric interpolation (Supplementary Appendix
166 A).

167 We also used municipality and state-level data on NPIs from a continuous survey
168 conducted between May 13 and July 31, 2020¹⁹. The survey had 13 questions related
169 to the implementation and easing of social distancing measures, and responses were
170 obtained from 612 mayors in São Paulo (94.8% of the total).

171 **Data analysis**

172 **Probability of hospitalisation and death**

173 We conducted an individual-level analysis to estimate the probability of reporting a
174 SARI hospitalisation given a patient's race and average income level in their census
175 tract of residence. Census tracts were grouped by quantiles of income per capita into
176 six categories as presented in the results. Similarly, we determined the probability of
177 death from SARI given a patient's race, income, and administrative type of the health
178 facility where the patient was hospitalised (public or private). Both probabilities were
179 standardised by age and sex to account for demographic differences between groups.
180 They were calculated for every month between March and August 2020. Probabilities
181 for each age-sex group were estimated empirically using relative frequencies. Odds
182 ratios using White patients and the highest income level as reference groups were
183 computed. Confidence intervals were calculated using bootstrapping. Details are in
184 Supplementary Appendix A.

185 **Seroprevalence by socioeconomic status**

186 To assess the broader risk of SARS-CoV-2 infection beyond hospitalisation, we
187 analysed seroprevalence data collected as part of the national Covid-IgG study²⁰ from
188 blood donors aged 16 – 69 living in São Paulo city²⁰ (Supplementary Appendix A).
189 We calculated the proportion of individuals by education and race category with
190 detectable anti-SARS-CoV-2 antibodies between February and October 2020.

191 **Socioeconomic drivers and hospitalisation risk**

192 An ecological spatio-temporal regression analysis was conducted at the municipality
193 level for São Paulo state (n = 645 municipalities) to assess the monthly risk of

194 hospitalisation and its association with socioeconomic factors between March 1 and
195 August 29, 2020. To further understand the association between socioeconomic
196 conditions and COVID-19 risk, we conducted the same analysis at the census tract
197 level (n= 30 815) for the RMSP, where the majority of cases were concentrated. The
198 relative risk of hospitalisation was estimated using a hierarchical Bayesian model
199 composed of a generalised log-linear model with spatially structured and unstructured
200 random effects to account for spatial autocorrelation and time-varying random effect.
201 The spatial structure is characterised by population movement between municipalities
202 from March 1 to August 15, 2020 defined by In-LoCo mobile geo-location data
203 summarised elsewhere ¹⁸. A detailed description of the model and interpretation,
204 covariates, and diagnostics can be found in Supplementary Appendix A.

205 **Social adherence response to NPIs**

206 We used an event study design ²¹ to examine how different socioeconomic groups
207 changed their daily mobility levels in response to the introduction and relaxation of
208 NPIs in the RMSP. We compared changes in mobility patterns of the population living
209 in H3 cells with predominantly White versus predominantly Black or *Pardo* residents,
210 as well as of the population living in areas of the wealthiest and poorest income
211 quintiles. The daily social distancing index from hexagons were regressed on a set of
212 relative time dummies that indicated the number of days before and after the first NPI
213 introduction in São Paulo state. Hexagon-fixed effects controlled for unobserved time-
214 invariant determinants of social distancing, while day-fixed effects controlled for
215 temporal shocks common to all hexagons. We further included an additional time-
216 varying control variable, representing the number of days relative to the first confirmed
217 SARI case in each hexagon, and a dummy variable indicating the period of NPI
218 relaxation in each municipality. Sensitivity analyses were performed and discussed in
219 Supplementary Appendix A.

220 We employed a multinomial logistic regression to estimate the probability that
221 employed individuals would be working face-to-face, teleworking, or taking paid or
222 unpaid leave. Differences in the work status of individuals by race, education, and

223 occupation were calculated while controlling for age and sex. 35 groups of
224 employment occupations listed in the PNAD COVID-19 survey were aggregated to
225 ten SCO-08 1-digit occupational groups defined by the International Labour
226 Organisation. We further disaggregated health professionals and health technicians
227 (Supplementary Appendix A – Table A2).

228 **Comorbidities**


229 PNAD COVID-19 data was used to conduct a binomial logistic regression to estimate
230 the odds ratio of having at least one comorbidity, by race and education attainment
231 (pre-primary, primary, secondary, and tertiary), while controlling for age and sex for
232 São Paulo state. The comorbidities considered were chronic obstructive pulmonary
233 disease (COPD), diabetes, hypertension, or cardiovascular disease such as
234 myocardial infarction, angina or heart failure. Confidence intervals for the odds ratios
235 were calculated taking into account PNAD's complex sample design.

236

237 **Results**

238 Between March 1 and August 29, 2020, São Paulo state had the highest number of
239 SARI hospitalisations per 100 000 habitants compared to all states in Brazil (figure
240 1A). A time series illustrating the number of SARI hospitalisations is presented in
241 figure 1B. During this time, 232 540 patients were notified in the SIMI-SP database
242 (figure 1C), from which 230 794 (99.2%) had a confirmed COVID-19 diagnosis or were
243 diagnosed with SARI of unknown or missing aetiology. From these, 223 455 were
244 hospitalised (98.4%) or died without hospitalisation (1.6%). From the non-hospitalised
245 cases, we only selected deaths; 54 108 patients died, of which 52.5% were White,
246 20.2% were *Pardo*, 6.13% were Black, 1.96% were Asian, 0.052% were Indigenous
247 and 19.6% did not have race information. We geocoded 178 345 (79.8%) of all SARI
248 cases considered in our analyses with high accuracy at either street address, route,
249 or neighbourhood level without compromising personal privacy.


250 During the first month of the COVID-19 epidemic (March) in Brazil, hospitalised
251 patients were more likely to be White or Asian and come from census tracts with higher
252 income per capita (figure 2A and 2B). During this period, people living in low-income
253 areas, compared to high-income areas, were less likely (OR: 0.44, 95% CI: 0.42 –
254 0.46) to be hospitalised with SARI. This coincides with the early establishment of
255 COVID-19 and its first cases in higher-income travellers returning from overseas^{9,12}.



256
257 **Figure 1.** Number of SARI hospitalisations per 100 000 habitants by state in Brazil
258 between March 1 and August 29, 2020 (A). Number of SARI hospitalisations for the
259 state of São Paulo and the greater metropolitan area of São Paulo (RMSP) by week
260 of symptom onset (B). Flowchart of SIMI-SP data processing (Source:
261 <https://covid.saude.gov.br>) (C).
262

263 As the epidemic progressed from April onwards, patients were also more likely to be
264 from low-income census tracts (April to July, OR: 1.08, 95% CI: 1.04 – 1.12), except
265 for August, when patients were less likely to be from low-income census tracts (OR:
266 0.95, 95% CI: 0.91 – 1.00). Similarly, Black Brazilians and *Pardos* became more likely
267 to be hospitalised with SARI than Whites (OR: 1.37, 95% CI: 1.32 – 1.41; OR: 1.23,
268 95% CI: 1.21 – 1.25, respectively), while Asians became the least likely to be
269 hospitalised (OR: 0.90, 95% CI: 0.83 – 0.96). These results were further confirmed by

270 our seroprevalence findings, which showed that anti-SARS-CoV-2 antibodies were
 271 highest in Black blood donors and those with low educational attainment across all
 272 age groups (Supplementary Appendix B – figure S1).




273

274 **Figure 2.** Age-standardised odds ratio (OR) for SARI hospitalisation by race (A) and
 275 by income (B); Age-standardised OR of death among SARI cases by race (C) and
 276 income (D); and Age-standardised OR of death among SARI cases by hospital type
 277 for São Paulo state (E). *PPP: Purchasing power parity.*

278 Once hospitalised, Black and *Pardo* patients were more likely to die from SARI than
279 White patients (OR: 1.14, 95% CI: 1.07 – 1.21; 1.09, 95% CI: 1.05 – 1.13,
280 respectively) (figure 2C). This difference was more pronounced in March and was
281 reduced over time. We found that patients living in the poorest census tracts were
282 more likely to die from SARI compared to patients from the wealthiest tracts (OR: 1.60,
283 95% CI: 1.48 – 1.74) (figure 2D). Likewise, patients treated in public hospitals were
284 more likely to die than patients treated in private hospitals throughout the epidemic
285 (OR: 1.40, 95% CI: 1.34-1.46) (figure 2E). Racial differences in the probability of death
286 decreased when considering only patients hospitalised at public health facilities but
287 persisted among patients in private facilities (Supplementary Appendix B – figure S2).

288 To understand the geographical variation in SARI hospitalisation, we predicted and
289 mapped the relative risk of SARI hospitalisation for São Paulo state by month via a
290 model with a spatial structure defined by human movement fluxes derived from
291 anonymised mobile phone data (figure 3A) and covariates related to socioeconomic
292 status (figure 3B). This is shown in figure 3C. We found a lower risk of hospitalisation
293 in municipalities with high income per capita (fixed effect = -0.93, 95% CI: -1.17 – -
294 0.69) and high proportion of adult residents with a primary education or lower (-0.80,
295 95% CI: -1.0 – -0.60). Municipalities with fewer nearby public health facilities were
296 also found to have lower risk of hospitalisation (-0.24, 95% CI: -0.45 – -0.03). We also
297 found a higher risk of SARI hospitalisation in municipalities with higher population
298 density (0.24, 95% credible interval: 0.04 – 0.44).

299



300

301 **Figure 3:** Relative risk of SARI hospitalisation in the state of São Paulo at the
302 municipality level (A). Human movement between municipalities based on In-LoCo
303 mobile phone data retrieved from March to August 2020 (B). Fixed effects and 95%
304 credible intervals for socioeconomic covariates (C).


305

306 We found that over time, the risk of SARI hospitalisation increased particularly in
307 municipalities near and within the RMSP, where 70% of the SARI cases reported for
308 the state are concentrated (figure 4A). By mapping risk at the census tract level for
309 the RMSP, we detected increasing risk starting from São Paulo city (central region).
310 By June, almost all of the census tracts in and near the city centre were found to have
311 high relative risk, but this risk decreased by August.

312

313 Differential risk to SARI in the RMSP was also associated with daily mobility levels by
314 disadvantaged groups. Before the implementation of NPIs on March 13, mobility levels
315 were similar across all socioeconomic groups (figures 4B and 4C). However, 14 days
316 after the introduction of NPIs, isolation levels were 8.2% (95% CI: 7.2% – 9.2%) higher
317 in predominantly White areas compared to predominantly Black areas. Similarly, 27
318 days after the introduction of state-level NPIs, isolation levels were 11.2% (95% CI:
319 10.6% – 11.9%) higher in the wealthiest than in the poorest areas. Overall, we
320 detected a decreasing trend in isolation levels over time, and the magnitude of the
321 differences in social isolation levels between areas with predominantly White and
322 Black populations decreased to only 4.4% (95% CI: 3.3% – 5.5%) 151 days after the
323 introduction of the NPIs (figure 4D).

324



325


326 **Figure 4.** Relative risk of SARI hospitalisation for the RMSP (A). Seven-day moving
327 average of daily median levels in social distancing by race (B) and income (C).
328 Difference in daily social isolation by race (D) and income (E) after the introduction of
329 NPIs. In panels (B) and (C), solid lines show population-weighted median isolation
330 levels and shaded areas show population-weighted interquartile range (25% - 75%).
331 Dashed vertical lines indicate the dates of NPIs that enabled school closure (March
332 13 was the state NPI) and non-essential activities (March 18 and 22, municipal and
333 state NPIs, respectively).

334

335 Finally, we investigated the differential risk to SARI based on workers' position in the
336 labour market using data from the PNAD COVID-19 survey. After the introduction of
337 NPIs, workers employed in low-skilled jobs or essential services were more likely to
338 keep working face-to-face than workers in professional or managerial positions

339 (Supplementary Appendix B - figure S3). Workers with pre-primary education were
340 more likely to work in occupations that require in-person contact than workers with
341 tertiary education (Probability: 0.89, 95% CI: 0.87 – 0.90 compared to PR: 0.58, 95%
342 CI: 0.57 – 0.60) and less likely to work in occupations that allow teleworking (PR:
343 0.005, 95% CI: 0.004 – 0.007 versus PR: 0.36 95% CI: 0.35 – 0.37, respectively)
344 (figure 5A). When controlling for education and formal or informal employment, we
345 found no substantial difference between racial groups in the probability of working
346 face-to-face or teleworking (figure 5B).

347 We found that population groups at risk of death from SARI were also more likely to
348 have comorbidities. Compared to the population with tertiary education in São Paulo
349 state, individuals with primary education or lower are more likely to have one or more
350 comorbidities (OR: 1.36, 95% CI: 1.27 – 1.45) (figure 5C). Similarly, Black individuals
351 were also more likely to have one or more comorbidities than White individuals (OR:
352 1.29, 95% CI: 1.19 – 1.39) (figure 5D). Odds ratio estimates for each health condition
353 are summarised in Supplementary Appendix B – figure S4.



354

355 **Figure 5:** Probability of different working conditions by education attainment (A) and
 356 race (B). Odds ratio (OR = 1) of having multiple comorbidities: chronic obstructive
 357 pulmonary disease (COPD), diabetes, hypertension or cardiovascular disease such
 358 as infarction, angina, and heart failure by education attainment (C) and race (D).
 359 Horizontal lines show 95% confidence intervals. Source: PNAD COVID-19 /IBGE ¹⁷,
 360 July to September, 2020.

361

362 Discussion

363 Our study shows that disadvantaged hospitalised groups are disproportionately more
 364 likely to die from SARI. We find that the differential health outcomes can be explained
 365 by structural inequities linked to the incidence of comorbidities and to socioeconomic

366 status, which limit the ability of these groups to socially isolate and reduce their access
367 to optimal health services.

368

369 Social and racial inequalities shape the risk of SARI hospitalisation and death. After
370 the initial phase of international imports in Brazil ¹², Black or *Pardo* Brazilians residing
371 in low-income areas were more likely to be hospitalised and die with SARI compared
372 to White individuals, which aligns with recent findings ⁷. Our assessment of anti-SARS-
373 CoV-2 antibodies in blood donors categorised by demographic background further
374 confirms that Black Brazilians and those with lower socioeconomic status are
375 disproportionately exposed to COVID-19. While patients hospitalised in public health
376 facilities were more likely to die than those in private health facilities, racial inequalities
377 in death probability are attenuated when only considering patients within hospitals of
378 the same type, either public or private. Potential factors influencing this inequality
379 include higher comorbidities among poor Black patients and the lower access to
380 private care among low-income individuals that are disproportionately Black.

381

382 We found higher hospitalisation risk for those living in municipalities with low income
383 per capita and high population density compared to the rest of São Paulo state. These
384 populations mainly reside in the RMSP, which contains nearly half of the population
385 in São Paulo state and where bias in testing is evident in regions of lower
386 socioeconomic status ¹² (Supplementary Appendix B – figure S5). The risk of SARI
387 hospitalisation is particularly elevated in São Paulo city, where seroprevalence
388 estimates from blood donors show that anti-SARS-CoV-2 antibodies were highest in
389 older Black Brazilians and those with lower educational attainment.

390

391 We show that inequalities in the risk of SARI hospitalisation were partially explained
392 by differential mobility responses to social distancing guidelines, similar to the US ²².
393 In wealthier and predominantly White neighbourhoods, people were able to isolate
394 faster and sustain isolation for long periods of time. Among the working population,
395 low-income and Black workers were less likely to receive a furlough from work or
396 telework. Due to systemic inequalities in education and the labour market, these

397 groups are disproportionately employed in precarious job positions with no social
398 security and dependent on day-to-day income ²³, limiting their ability to reduce
399 mobility. Other important factors include the disadvantage of having multiple
400 comorbidities, which are more prevalent among Blacks and *Pardos* and those with
401 lower education.

402

403 Our study has limitations that may have underestimated the level of inequality. Firstly,
404 geocoding cases may have discarded patients from poor census tracts where
405 accuracy is limited ²⁴. Secondly, using data aggregated for various administrative
406 levels has inherent limitations due to ecological fallacy and the modifiable areal unit
407 problem ²⁵. Finally, the 2010 Brazilian population census and PNAD COVID-19 survey
408 may have limited the capture of socioeconomic changes in the last decade and
409 inclusion of extremely wealthy individuals ²⁶. Additionally, disadvantaged groups can
410 be underrepresented in health administrative records because of their lower access
411 to healthcare. Given that São Paulo is the wealthiest state and has the most robust
412 healthcare system in Brazil ²⁷, it is possible that the impact of inequalities is more
413 severe in other states.

414

415 Our findings on the difference in risk of SARI death reveal stark inequalities in access
416 to healthcare. Only 25% of Brazilians have access to private healthcare via health
417 insurance, reflecting how inequality in access to quality healthcare is largely driven by
418 income ²⁸. This leaves 75% of the population solely reliant on a chronically
419 underfunded public healthcare system, which highlights that disadvantaged
420 populations are more likely to be infected and deprived of care. Strengthening
421 healthcare access and its capacity will be critical for reducing health inequities during
422 this and forthcoming public health emergencies ²⁹.

423

424 Our findings on socioeconomic risk factors could help guide vaccine allocation in
425 diverse settings to achieve equitable access. Ensuring that disadvantaged groups,
426 especially those that have in-person occupations and live in crowded and deprived
427 areas receive vaccination will help prevent and slow down community transmission.

428 While race is not a risk factor in itself, it is critical to consider systemic inequalities that
429 lead Black and *Pardo* communities to be overrepresented among low socioeconomic
430 groups, to have higher rates of severe COVID-19 infection, and comorbidities that
431 exacerbate their risk of death. Therefore, including disadvantaged populations among
432 priority groups for vaccination could help reduce health inequities instead of
433 exacerbating them ³⁰.

434

435 Our study highlights the need for additional research to comprehend the effects of
436 social and health inequalities during pandemics. Firstly, an assessment of the
437 inequality in access to quality care within public and private health facilities and its risk
438 factors is needed to better understand mortality in different hospital settings. Secondly,
439 our study shows that population response to NPIs can vary significantly based on
440 social circumstances, suggesting that future studies should also consider
441 socioeconomic aspects when evaluating the effectiveness of NPIs. Thirdly, more data
442 is needed on whether social safety net programs that are guaranteeing income for
443 disadvantaged groups during the pandemic (e.g., Brazil's emergency cash payment),
444 may have enabled people to reduce their mobility. Nevertheless, our study has shown
445 the impact of social inequities on COVID-19 hospitalisation and death, thus informing
446 future research and policies related to the health impacts of COVID-19 in Latin
447 America.

448

449 **Contributors**

450 SLL and RHMP conceived the research questions and designed the study. RHMP,
451 CAPJ, LE, PJHA, PSP, CKB, AASS, WMS, RJB, LFB, AM, CAN, SCF, and NAS
452 collected the epidemiological, socioeconomical, mobility, NPIs, seroprevalence, and
453 comorbidities data. SLL, RHMP, CAPJ, AEZ, LE, PJHA, CKB, RJB, and VN conducted
454 exploratory and statistical analysis and interpreted the results. SLL, RHMP, and CAPJ
455 wrote the manuscript. All authors read and revised the final manuscript.

456

457 **Acknowledgement**

458 SLL is supported by the Oxford Martin School and the Canadian Social Sciences and
459 Humanities Doctoral Fellowship. CAPJ is supported by Coordenação de
460 Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) and Fundação
461 Faculdade de Medicina (FFM). AEZ is supported by The Oxford Martin Programme
462 on Pandemic Genomics. WMS is supported by the São Paulo Research Foundation,
463 Brazil (2017/13981-0 and 2019/24251-9). NRF is supported by a Wellcome Trust and
464 Royal Society Sir Henry Dale Fellowship (204311/Z/16/Z).

465

466

467 **References**

- 468 1 Ahmed F, Ahmed N, Pissarides C, Stiglitz J. Why inequality could spread COVID-
469 19. *Lancet Public Health* 2020; **5**: e240.
- 470 2 Walker PGT, Whittaker C, Watson OJ, *et al.* The impact of COVID-19 and strategies
471 for mitigation and suppression in low- and middle-income countries. *Science* 2020;
472 **369**: 413–22.
- 473 3 Drefahl S, Wallace M, Mussino E, *et al.* A population-based cohort study of socio-
474 demographic risk factors for COVID-19 deaths in Sweden. *Nat Commun* 2020; **11**:
475 5097.
- 476 4 UK Office for National Statistics. Coronavirus (COVID-19) related deaths by ethnic
477 group, England and Wales - Office for National Statistics. 2020
478 [https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriage](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020)
479 [s/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2marc](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020)
480 [h2020to10april2020](https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020) (accessed Nov 20, 2020).
- 481 5 Cummings MJ, Baldwin MR, Abrams D, *et al.* Epidemiology, clinical course, and
482 outcomes of critically ill adults with COVID-19 in New York City: a prospective
483 cohort study. *The Lancet* 2020; **395**: 1763–70.
- 484 6 Pan D, Sze S, Minhas JS, *et al.* The impact of ethnicity on clinical outcomes in
485 COVID-19: A systematic review. *EClinicalMedicine* 2020; **23**: 100404.
- 486 7 Baqui P, Bica I, Marra V, Ercole A, Schaar M van der. Ethnic and regional variations
487 in hospital mortality from COVID-19 in Brazil: a cross-sectional observational study.
488 *Lancet Glob Health* 2020; **8**: e1018–26.
- 489 8 Facundo A, Chancel, Lucas, Piketty, Thomas, Saez, Emmanuel, Zucman, Gabriel.
490 World Inequality Report 2018. 2017.
- 491 9 Jesus JG de, Sacchi C, Candido D da S, *et al.* Importation and early local
492 transmission of COVID-19 in Brazil, 2020. *Rev Inst Med Trop São Paulo* 2020; **62**:
493 e30.
- 494 10 Henriques CMP, Vasconcelos W, Henriques CMP, Vasconcelos W. Crises
495 dentro da crise: respostas, incertezas e desencontros no combate à pandemia da
496 Covid-19 no Brasil. *Estud Av* 2020; **34**: 25–44.
- 497 11 Rezende LFM, Thome B, Schweitzer MC, *et al.* Adults at high-risk of severe
498 coronavirus disease-2019 (Covid-19) in Brazil. *Rev Saúde Pública* 2020; **54**.
499 DOI:10.11606/s1518-8787.2020054002596.
- 500 12 de Souza WM, Buss LF, Candido D da S, *et al.* Epidemiological and clinical
501 characteristics of the COVID-19 epidemic in Brazil. *Nat Hum Behav* 2020; **4**: 856–
502 65.

- 503 13 Secretaria da saúde do estado de São Paulo. SIMI-SP: Pacientes internados por
504 Síndrome Respiratória Aguda Grave (SRAG).
505 <https://www.saopaulo.sp.gov.br/planosp/simi/> (accessed Oct 25, 2020).
- 506 14 Niquini RP, Lana RM, Pacheco AG, *et al.* SRAG por COVID-19 no Brasil:
507 descrição e comparação de características demográficas e comorbidades com
508 SRAG por influenza e com a população geral. *Cad Saúde Pública* 2020; **36**:
509 e00149420.
- 510 15 Hasell J, Mathieu E, Beltekian D, *et al.* A cross-country database of COVID-19
511 testing. *Sci Data* 2020; **7**: 345.
- 512 16 Telles E, Paschel T. Who Is Black, White, or Mixed Race? How Skin Color,
513 Status, and Nation Shape Racial Classification in Latin America. *Am J Sociol* 2014;
514 **120**: 864–907.
- 515 17 IBGE - Instituto Brasileiro de Geografia e Estatística (2020). Pesquisa Nacional
516 por Amostra de Domicílios Contínua (PNAD) COVID-19. Microdados. .
- 517 18 Peixoto PS, Marcondes D, Peixoto C, Oliva SM. Modeling future spread of
518 infections via mobile geolocation data and population dynamics. An application to
519 COVID-19 in Brazil. *PLOS ONE* 2020; **15**: e0235732.
- 520 19 De Souza Santos AA, Da Silva Cândido D, Marciel De Souza W, *et al.* SARS-
521 CoV-2 non-pharmaceutical interventions in Brazilian municipalities. 2020; :
522 1046406 bytes.
- 523 20 Buss LF, Prete CA, Abraham CMM, *et al.* Three-quarters attack rate of SARS-
524 CoV-2 in the Brazilian Amazon during a largely unmitigated epidemic. *Science*
525 2020; published online Dec 8. DOI:10.1126/science.abe9728.
- 526 21 Weill JA, Stigler M, Deschenes O, Springborn MR. Social distancing responses
527 to COVID-19 emergency declarations strongly differentiated by income. *Proc Natl*
528 *Acad Sci* 2020; **117**: 19658–60.
- 529 22 Chang S, Pierson E, Koh PW, *et al.* Mobility network models of COVID-19 explain
530 inequities and inform reopening. *Nature* 2020; : 1–8.
- 531 23 Lustig N, Pabon VM, Sanz F, Younger SD. The Impact of COVID-19 Lockdowns
532 and Expanded Social Assistance on Inequality, Poverty and Mobility in Argentina,
533 Brazil, Colombia and Mexico. ECINEQ, Society for the Study of Economic
534 Inequality, 2020 <https://ideas.repec.org/p/inq/inqwps/ecineq2020-558.html>
535 (accessed Nov 20, 2020).
- 536 24 Giest S, Samuels A. ‘For good measure’: data gaps in a big data world. *Policy*
537 *Sci* 2020; **53**: 559–69.

- 538 25 Duranton G, Overman HG. Testing for Localization Using Micro-Geographic
539 Data. *Rev Econ Stud* 2005; **72**: 1077–106.
- 540 26 Souza PHGF de. A distribuição de renda nas pesquisas domiciliares brasileiras:
541 harmonização e comparação entre Censos, PNADs e POFs. *Rev Bras Estud Popul*
542 2015; **32**: 165–88.
- 543 27 Paim J, Travassos C, Almeida C, Bahia L, Macinko J. The Brazilian health
544 system: history, advances, and challenges. *The Lancet* 2011; **377**: 1778–97.
- 545 28 Nogueira Avelar e Silva R, Russo G, Matijasevich A, Scheffer M. Covid-19 in
546 Brazil has exposed socio-economic inequalities and underfunding of its public
547 health system. *The BMJ*. 2020; published online June 19.
548 [https://blogs.bmj.com/bmj/2020/06/19/covid-19-in-brazil-has-exposed-deeply-](https://blogs.bmj.com/bmj/2020/06/19/covid-19-in-brazil-has-exposed-deeply-rooted-socio-economic-inequalities-and-chronic-underfunding-of-its-public-health-system/)
549 [rooted-socio-economic-inequalities-and-chronic-underfunding-of-its-public-health-](https://blogs.bmj.com/bmj/2020/06/19/covid-19-in-brazil-has-exposed-deeply-rooted-socio-economic-inequalities-and-chronic-underfunding-of-its-public-health-system/)
550 [system/](https://blogs.bmj.com/bmj/2020/06/19/covid-19-in-brazil-has-exposed-deeply-rooted-socio-economic-inequalities-and-chronic-underfunding-of-its-public-health-system/) (accessed Nov 20, 2020).
- 551 29 Castro MC, Carvalho LR de, Chin T, *et al*. Demand for hospitalization services
552 for COVID-19 patients in Brazil. *medRxiv* 2020; : 2020.03.30.20047662.
- 553 30 Emanuel EJ, Persad G, Kern A, *et al*. An ethical framework for global vaccine
554 allocation. *Science* 2020; **369**: 1309–12.
- 555