Title: A genomic epidemiology study of multidrug-resistant *Escherichia coli*, *Klebsiella*

pneumoniae and *Acinetobacter baumannii* in two intensive care units in Hanoi, Vietnam

Authors

Leah W. Roberts[1,2], Le Thi Hoi[3,4], Fahad A. Khokhar[2], Nguyen Thi Hoa[3], Tran Van Giang[3,4],

Cuong Bui[4], Tran Hai Ninh[3], Dao Xuan Co[5], Nguyen Gia Binh[5], Hoang Bao Long[4], Dang

Thi Huong[3], James E. Bryan[2], Archie Herrick[2], Theresa Feltwell[2], Behzad Nadjm[6,7], H.

Rogier van Doorn[7,8], Julian Parkhill[9], Nguyen Vu Trung[3,4], Nguyen Van Kinh[3], Zamin

Iqbal[1,*], M. Estée Török[2,10,*]

*Contributed equally

Affiliations

1. EMBL-EBI, Wellcome Genome Campus, Hinxton, Cambridge

2. Department of Medicine, University of Cambridge, Cambridge, United Kingdom

3. National Hospital for Tropical Diseases, Hanoi, Vietnam

4. Hanoi Medical University, Hanoi, Vietnam

5. Bach Mai Hospital, Hanoi Vietnam

6. Medical Research Council The Gambia at the London School of Hygiene & Tropical

Medicine, Fajara, The Gambia

7. Wellcome Trust Major Overseas Programme, Hanoi, Vietnam

8. Nuffield Department of Medicine, University of Oxford, Oxford, UK

9. Department of Veterinary Medicine, University of Cambridge, Cambridge, United

Kingdom

10. Departments of Infectious Diseases and Microbiology, Cambridge University

Hospitals NHS Foundation Trust, Cambridge, United Kingdom

Correspondence

M. Estée Török email: et317@cam.ac.uk

Zamin Iqbal, email: zi@ebi.ac.uk

Key Words

Antimicrobial resistance; whole genome sequencing; transmission; clusters; Escherichia coli;

Klebsiella pneumoniae; Acinetobacter baumannii; intensive care unit; Vietnam

Abstract (298 words)

Background

Vietnam has high rates of antimicrobial resistance (AMR) but limited genomic surveillance, impeding our ability to assess transmission dynamics. This study aimed to use whole genome sequencing (WGS) to examine the transmission of key AMR pathogens in two intensive care units in Hanoi, Vietnam.

Methods

A prospective surveillance study of all adults admitted to two intensive care units (ICUs) at the National Hospital for Tropical Diseases (NHTD) and Bach Mai Hospital (BMH) was conducted between June 2017 and January 2018. Clinical and environmental samples were cultured on selective media, characterised using MALDI TOF MS, and Illumina sequenced. Phylogenies based on the *de novo* assemblies (SPAdes) were constructed using Mafft (PARsnp), Gubbins and RAxML. Resistance genes were detected using Abricate against the NCBI database.

Findings

3,153 *Escherichia coli*, *Klebsiella pneumoniae* and *Acinetobacter baumannii* isolates from 369 patients were analysed. Phylogenetic analysis revealed predominant lineages within *A. baumannii* (global clone [GC]2, sequence types [ST]2, ST571) and *K. pneumoniae* (ST15, ST16, ST656, ST11, ST147) isolates. Colonisation was most common with *E. coli* (88.9%) followed by *K. pneumoniae* (62.4%). 91% of *E. coli* carried a *bla*CTX-M variant, while 81% of *K. pneumoniae* isolates carried *bla*NDM (54%) and/or *bla*KPC (45%). Transmission analysis using single nucleotide polymorphisms (SNPs) identified 167 clusters involving 251 (68%) patients, in some cases involving patients from both ICUs. There were no significant differences between the lineages or AMR genes recovered between the two ICUs.

Interpretation

This study represents the largest prospective surveillance study of key AMR pathogens in Vietnamese ICUs. Clusters of closely related isolates in patients across both ICUs suggests recent transmission prior to ICU admission in other healthcare settings or in the community.

Funding

This work was funded by the Medical Research Council Newton Fund, United Kingdom; the Ministry of Science and Technology, Vietnam (HNQT/SPĐP/04.16) and the Wellcome Trust, United Kingdom.

Research in context (269):

Evidence before this study:

Globally, antimicrobial resistance (AMR) is projected to cause 10 million deaths annually by 2050. While 90% of these deaths are expected to occur in African and Asian low- and middle-income countries (LMIC), attributing morbidity and mortality is difficult without the availability of comprehensive AMR data in these settings. Whilst efforts have been made to improve AMR surveillance in these settings, this is often hampered by limited infrastructure, training and financial resources.

Added value of this study:

This is the largest prospective surveillance study of three key AMR pathogens (*E. coli, K. pneumoniae* and *A. baumannii*) conducted in critical care settings in Vietnam. All patients were colonised or infected with one or more extended spectrum beta-lactamase (ESBL) producing and/or carbapenem-resistant organism. Colonisation with more than one organism was very common, with resistant *E. coli* predominantly isolated from stool. A small number of predominant lineages were identified for *K. pneumoniae* and *A. baumannii*, while the *E. coli* isolates were highly genetically diverse. A large number of genomic clusters were identified within the two ICUs, some of which spanned both ICUs. There were no significant differences between lineages or AMR genes between the two ICUs.

Implications of all the available evidence:

This study found high rates of colonisation and infection with three key AMR pathogens in adults admitted to two Vietnamese ICUs. Whilst transmission was common within ICUs the finding of similar lineages and AMR genes in both ICUs suggests that dissemination of AMR occurs prior to ICU admission, from either referral sites or in community settings prior to hospital admission. Strategies to tackle AMR in Vietnam will need to account for this by extending surveillance beyond ICU to hospital and community settings.

Introduction

Low- and middle-income countries (LMICs) have reported widespread antimicrobial resistance (AMR) in healthcare, community and agricultural settings. In South-East Asia, dense human populations, intensive animal farming, unrestricted access to antibiotics and limited laboratory infrastructure have all contributed to the rapid expansion of AMR (*1, 2*).

Much of this burden comes as a result of excessive use of antimicrobials in human and animal populations. In Vietnam, antimicrobial usage has been estimated to be two times higher in humans, and 1.5 times higher in animals, as compared to the European Union (*3*). Despite legal restrictions in Vietnam, antibiotics are often dispensed without prescriptions in the community (*4*). Broad-spectrum antibiotics are also commonly administered in healthcare settings to mitigate the effects of limited capacity for microbiological testing and infection control (*4, 5*). Detection of both resistant bacteria and antimicrobials have been recorded in the environment (*6, 7*), hospital waste (*8*) and food sources (*9, 10*).

Extensive AMR has led to increased pressure on hospitals and is particularly problematic in critical care settings. A study conducted in Thailand between 2008 and 2012 found that almost 80% of nosocomial infections were caused by resistant bacteria, accounting for roughly 100,000 AMR cases and 50,000 deaths annually (*11*). *Klebsiella pneumoniae* is considered a dire threat because of high rates of AMR and virulence (*12*) and widespread resistance to last-line treatments, such as carbapenems and polymyxins (*13*). *Acinetobacter baumannii* is a leading cause of ventilator-associated pneumonia (VAP) in critically ill patients, and is commonly treated with colistin due to widespread carbapenem resistance (*14, 15*).

Whilst AMR surveillance based on phenotypic antimicrobial susceptibility testing in Vietnam has improved in recent years, the infrastructure required for systematic genomic surveillance remains to be established. This is particularly important to determine circulating lineages and elucidate potential transmission events, as other methods do not provide the same level of resolution (*16*). Over the past decade, various studies have demonstrated the utility of WGS in characterising AMR, transmission routes, and dominant lineages (*17-19*). LMICs however remain relatively understudied, with few studies conducted in Vietnamese hospitals (*20-23*). With increasing globalisation, understanding the dynamics of circulating lineages and

evolving AMR in these regions is necessary to address both local and global efforts to detect,

monitor and manage resistant bacterial infections.

In order to address this knowledge gap, we conducted a prospective genomic surveillance

study of key AMR pathogens in two hospitals in Vietnam. We targeted intensive care unit

(ICU) patients as we hypothesised that these would be most likely to have been treated with

antibiotics and to harbour AMR pathogens. Furthermore, we focussed our analysis on the

three most commonly isolated species (*Escherichia coli, K. pneumoniae*, and *A. baumannii*)

that were extended-spectrum beta-lactamase (ESBL) producers and/or carbapenem-resistant.

An additional in-depth analysis focussed primarily on a subset of the *K. pneumoniae* isolates

was also performed in a separate project (Pham et al., personal communication).

Methods

Study design, setting and participants

This prospective observational cohort study was conducted in two hospitals, the National

Hospital for Tropical Diseases (NHTD) and Bach Mai Hospital (BMH) in Hanoi, Vietnam,

between June 2017 to January 2018. All patients (aged 18 years or older) admitted to the

adult ICUs of the two hospitals were eligible for inclusion in the study. NHTD is a specialist

hospital for infectious and tropical diseases with a 22-bedded ICU which receives up to 400

patients per year. BMH is a large tertiary referral hospital, with a 45-bedded ICU that

receives up to 1,200 patients per year. Both hospitals are located in the same area of Hanoi

but operate independently of each other and do not share laboratory facilities, equipment or

staff. Patients are not commonly transferred between the two hospitals.

Study procedures

Screening specimens were collected from ICU patients on admission, on discharge and

weekly during their ICU stay. Specimens included stool/rectal swabs, urine, skin/wound

swabs and sputum/tracheal aspirates. Environmental samples were collected using flocked

swabs (from door handles, bed rails, medical equipment and patient tables) on a monthly

basis. Clinical data related to the ICU admission were collected from the medical records and

entered into a case record form and then into an electronic database. Laboratory data were

collected and recorded in an electronic database.

Laboratory methods and sequencing

All patient and environmental specimens were cultured on selective media (CHROMagar™
ESBL, CHROMagar™ mSuperCARBA™, CHROMagar™ VRE, CHROMagar, France).
Single colony picks of target organisms (*Escherichia coli*, *Acinetobacter baumannii* and
Klebsiella pneumoniae) were selected and identified using MALDI-TOF MS (Bruker
Diagnostics, Bremen, Germany) and stored at -80 °C. Stored isolates were shipped in two
batches to the University of Cambridge, United Kingdom, where they were sub-cultured, re-
identified using MALDI-TOF MS, and underwent antimicrobial susceptibility testing (Vitek-
2, BioMérieux, Marcy L'Étoile, France). Isolate DNA was extracted using QIACube and the
QIAamp 96 DNA QIACube HT kit (Qiagen, Hilden, Germany) prior to shipping to the
Wellcome Sanger Institute for sequencing. DNA was sequenced in two batches on an
Illumina HiSeq X10 machine (Illumina Inc., San Diego (CA), USA).
181
182   *Read quality control*
183   Raw Illumina reads were checked for quality using fastQC (v0.11.8) (*24*) and MultiQC (v
184   1.0.dev0) (*25*). Raw Illumina reads were also checked for contamination using Kraken2
185   (v2.0.7-beta) (*26*) and Bracken (v2.5) (*27*).
186
187   *Assembly*
188   Illumina reads were *de novo* assembled using SPAdes (v3.13.1) (*28*) and checked for quality
189   using Quast (v5.0.2) (*29*) and CheckM (v1.0.18) (*30*). Further filtering and quality control
190   methods for all assemblies is provided in the Supplementary Methods.
191
Phylogenetic construction
Prior to phylogenetic construction, reads were mapped to filtered assemblies and sites that
had <90% consensus compared to the reference allele were masked. Masked assemblies were
then aligned using PARsnp (v1.2) under default settings (*31*). Core multi-alignments
produced using PARsnp were filtered for recombination using Gubbins (v.2.3.5) (*32*).
Phylogenies were then constructed using RAxML (GTR-GAMMA model) (v8.2.12) (*33*) as
implemented through Gubbins. *E. coli* phylogroups were determined using ClermonTyping
(*34*). Trees were visualised using iTol v5.6.2 (*35*). Details on global reference selection are
provided in the Supplementary Methods.
201
Antibiotic resistance gene detection

Resistance genes and plasmid replicons were detected from draft assemblies using Abricate

(v1.0.1) (*36*) and the NCBI database (for resistance genes) (*37*)**.** Genes were considered

present if there was 90% coverage at 90% nucleotide identity (*38*).

Multi-locus sequence typing (MLST)

Sequence types were determined using mlst (v2.19.0) (https://github.com/tseemann/mlst) and

the associated species scheme (specifically the Pasteur scheme for *A. baumannii*) (*39-41*).

Transmission cluster analysis

Transmission clusters were constructed using Transcluster (using the makeSNPClusters

method, which ignores time of sampling and uses a pure SNP-distance cut-off) (*42*) using

single nucleotide polymorphisms (SNPs) determined after recombination filtering using

Gubbins. Transmission cut-offs were evaluated based on intra- and inter-patient SNP

diversity within each species phylogeny (Supplementary Figure 1).

217
Results
Samples included in the study

Between June 2017 to January 2018, a total of 3,367 isolates were cultured, comprising

Escherichia coli (n=765), *Klebsiella pneumoniae* (n=1,372) and *Acinetobacter baumannii*

(n=1,230). Thirty-one isolates were excluded from the analysis because of poor assembly

quality. A further 150 isolates were excluded because of suspected inter-species

contamination, and 33 isolates were excluded because of suspected intra-species (strain-level)

contamination (Supplementary Figure 2). Thus 3,153 isolates (93.6%), comprising 2,901

isolates from 369 patients and 252 environmental isolates, passed quality filtering and were

included in the final analyses.

Clinical data

Of the 3,153 isolates, 1,042 (33%) were collected from BMH, while 2,111 (67%) were

collected from NHTD (Table 1). Both hospitals recruited a similar number of patients and the

average age was 53-55 years (BMH median age 55, NHTD median age 57.5). The average

length of stay (LOS) in BMH was 7 days (median 6 days), and 21 days (median 16 days) at

NHTD (Supplementary Figure 3). Patient outcomes are summarised in Table 1.

240 **Table 1: Summary of isolates and patients included in the study**

Variable	Bach Mai Hospital	National Hospital of Tropical Diseases
Isolates:		
Total	1042	2111
Clinical	993 (95%)	1898 (90%)
Environmental	49 (5%)	213 (10%)
Patients:		
Total	182	187
Male	104 (57%)	114 (61%)
Female	71 (39%)	50 (27%)
Gender not recorded	7 (4%)	23 (12%)
Age:		
Male	53 years (range 16-85)	55.2 years (range 5-92)
Female	53.4 years (range 19-91)	55.7 years (range 10-90)
Age not recorded	n=5 (3%)	n=19 (10%)
Length of stay:		
	7.5 days (range 0 – 35)	21 days (range 1 – 75)
Stay not recorded	n=4 (2.2%)	n=7 (3.7%)
Outcome at discharge from ICU		
Stable, discharged home	10 (6%)	38 (20%)
Improved, transferred to another ward	117 (64%)	83 (44%)
Deteriorated, transferred to another ward	3 (2%)	7 (4%)
Discharged home to die	44 (24%)	43 (23%)
Died in hospital	4 (2%)	9 (5%)
Not recorded	4 (2%)	7 (4%)

All of the 369 patients were colonised or infected with one or more of the three species: *E. coli*, *A. baumannii* and *K. pneumoniae*: 146 patients (40%, 55 at BMH and 91 at NHTD) were colonised or infected with all three species; 133 patients (36%, 66 at BMH and 67 at NHTD) were colonised or infected with two of the three species; and 90 patients (24%, 61 at MNH and 29 at NHTD) only had one species detected.

Both *E. coli* and *K. pneumoniae* were isolated primarily from stool / rectal swabs (627/721 [87.0%] and 822/1316 [62.5%], respectively). *K. pneumoniae* was also isolated from other sites including sputum (325/1316 [24.7%]), urine samples (63/1316 [4.8%]) and skin swabs (17/1316 [1.3%]). In contrast, *A. baumannii* isolates were mostly isolated from sputum (621/1116 [55.6%]), followed by stool / rectal swabs (247/1116 [22.1%]), urine (49/1116 [4.4%]) and skin swabs (36/1116 [3.2%]). *A. baumannii* also accounted for the highest number of environmental isolates (161/1116, [14.4%]), compared to 6.5% (85/1316) for *K. pneumoniae* and 2.2% (16/721) for *E. coli*.

Whole genome sequencing reveals predominant circulating lineages

Phylogenetic trees for each species were constructed to explore lineage diversity within the dataset. The *E. coli* isolates were found to be genomically diverse, with isolates spread over eight phylogroups and 80 sequence types (STs) (Figure 1). The most prevalent ST was ST648 (phylogroup A; 11.8%), followed by ST410 (phylogroup C; 9.7%), ST617 (phylogroup A; 9.2%), ST131 (phylogroup B2; 7.9%) and ST1193 (phylogroup B2; 7.4%). Overall, 33 of the 80 STs only had one representative isolate in this dataset.

In contrast, the *K. pneumoniae* and *A. baumannii* isolates appeared to be centred around specific dominant lineages. More than 80% of the *K. pneumoniae* isolates were from one of five STs, including ST15 (n=34%), ST16 (n=20%), ST656 (n=12%), ST11 (n=11%) and ST147 (n=7%). The majority of *A. baumannii* were global clone (GC)2 (n=832, 74.6%) (*43*) and mainly belonged to ST2 (n=48%) and ST571 (n=24%) (based on the Pasteur scheme). There did not appear to be any relationship between STs and hospitals, with all of the major ST lineages detected in both ICUs.

Figure 1: Whole genome phylogenies for [A] *E. coli,* **[B]** *A. baumannii***, and [C]** *K. pneumoniae:*
recombination-filtered core-SNP trees with mid-point root. Tree metadata includes (from left to right column
beside trees): MLST, source and hospital. Outermost purple bars indicate environmental isolates. Branches
corresponding to *E. coli* phylogroups are coloured accordingly. Main STs are highlighted in the image using the
pale-yellow boxes.

In order to gain broader insight into the lineages, we selected globally representative strains
to contextualise our dataset. Addition of these global representatives into the *E. coli*
phylogeny showed that most isolates belonged to a globally diverse set of STs that were not
unique to Vietnam, but found across parts of North America, Europe and Asia
(Supplementary Figure 4). Similarly, several of the major *K. pneumoniae* lineages were
represented globally, particularly ST147, ST11 (mainly from China and the USA) and ST15
(mainly Asian countries) (Supplementary Figure 5). However, it was also clear that local
expansion was prominent, particularly among the ST656, ST16 and ST15 lineages. For *A.*
baumannii, we focused primarily on GC2 isolates (Supplementary Figure 6). There was very
little representation of global strains similar to lineages within our dataset, and those that
were available consisted mainly of strains from other parts of Asia.

Closer inspection of the global representatives found several strains in each species that were
closely related (<5 core SNPs) to isolates in our dataset (Supplementary Table 1). Most of
these global representatives were also isolated in Asian countries. The exception was *E. coli*,
where two closely related global representatives were from the United Kingdom and
Australia (ST1193 and ST131, respectively). Agricultural isolates were also linked to these
lineages, as the other two closely related *E. coli* representatives were originally isolated from
poultry (biosample SAMEA104188722) and a farm worker in Vietnam (biosample
SAMEA5277968).

High prevalence of antibiotic resistance genes among majority of isolates
Almost all isolates carried acquired resistance genes belonging to at least 3 antibiotic classes,
with 90% of *E. coli*, 97% of *K. pneumoniae* and 41% *A. baumannii* carrying genes across 5
antibiotic classes (Figure 2). There were no discernible differences based on sample source or
hospital, with the exception of *E. coli* detected in pus/skin swabs (n=6) which appeared on
average to carry resistance to more antibiotic classes. *K. pneumoniae* isolates tended to fall
into one of three "peaks" (Figure 2). This was due to lineage-specific carriage of acquired
resistance genes, where ST15 isolates tended to carry resistance to more classes, compared to
ST16 which often carried the least. The other three main lineages (ST656, ST11, ST147) fell
between these two peaks. The exception was in the environmental samples, where only two
peaks can be seen. This is likely due to very few ST16 isolates detected in the environment.

Figure 2: Summary of isolates and the number of antibiotic resistance classes separated by species, hospital and sample type.

Resistance to antibiotics classes varied across the *E. coli* phylogeny, reflective of the diversity of strains within the dataset (Supplementary Figure 7). *bla*CTX-M genes were found in most *E. coli*, with *bla*CTX-M-15 (36%), *bla*CTX-M-27 (30%) and *bla*CTX-M-55 (17%) the most prevalent (Table 2). *bla*KPC-2 (13%) and *bla*NDM-[1,4,5,7] (24%) were present sporadically across the phylogroups, suggesting independent acquisitions events. Only 4% (n=28) of isolates carried *mcr* genes conferring resistance to colistin. Again, these seemed to be independent acquisitions, with the exception of an ST206 cluster (phylogroup A; n=11) involving three patients from NHTD.

Conversely, MDR gene presence across the *K. pneumoniae* isolates appeared consistent with the main lineages, suggesting clonal expansion rather than diverse sampling of the species

(Supplementary figure 8). Similar to the *E. coli*, incidence of *bla*CTX-M-15 (37.5%) was

high, but less so overall compared to blaKPC-2 (45%) and *bla*NDM (54%) (NDM-4 [27.9%],

NDM-1 [24.7%] and NDM-5 [1.8%]) (Table 2).

Acquired AMR genes were overall less prevalent among the *A. baumannii* isolates. Similar to

the *K. pneumoniae*, resistance to specific classes tended to be a feature of each distinct

lineage, suggesting clonal expansion (Supplementary Figure 9). The carbapenemase gene

*bla*OXA-23 was present in 83% of the dataset, with *bla*OXA-58 and *bla*OXA-72 present at

much lower frequencies (5% and 0.2% respectively) (Table 2). The aminoglycoside

resistance gene *armA* was also highly prevalent (76%).

Overall, 133 AMR genes were detected in BM and 154 were detected in NHTD. 49 genes

were unique to either hospital (35 in NHTD, 14 in BM), but were only detected at a

prevalence of less than 0.1%, suggesting sporadic cases. The remaining 129 genes were the

same across both hospitals. The genes with the highest prevalence (at least 1%) were found to

be almost identical in both hospitals, with the exception of *bla*NDM-4 (0.98%), *dfrA12*

(0.92%), *rmtB1* (0.86%), *qnrB6* (0.74%) and *bla*OXA-181 (0.56%) which were below 1%

prevalence in NHTD.

In order to determine if certain time points throughout the study had different gene burdens

(potentially indicative of mobile genetic element [MGE]-mediated transmission), we plotted

gene presence versus date for genes equivalent to 1% prevalence in either hospital

(Supplementary Figures 10). Overall, we found that both ICUs had a consistently high

burden, making it difficult to distinguish significant gene fluctuations over time. In NHTD,

we observed three genes that seemed to peak between November to December 2017.

Examination of our dataset for isolates with these three genes (*bla*NDM-4, *bla*OXA-181 and

rmtB1) revealed a subset of ST16 *K. pneumoniae* that carried all three genes as well as a

single ST11 isolate from BM. We plotted the presence of these isolates over time, which

mirrored the rise in prevalence in NHTD over the November to December period

(Supplementary Figure 11). No *E. coli* nor *A. baumannii* isolates in this dataset carried all

three genes.

370 **Table 2: Summary of resistance genes found in the three species**

Resistance gene class	*Escherichia coli*	*Acinetobacter baumannii*	*Klebsiella pneumoniae*
Tetracycline	563 (78.1%)	701 (62.8%)	753 (57.2%)
Sulphonamide	649 (90%)	746 (66.8%)	969 (73.6%)
Fluoroquinolone	161 (22.3%)	19 (1.7%)	1187[3] (90.2%)
Colistin	28 (3.9%)	0 (0%)	10 (0.8%)
Fosfomycin	48 (6.7%)	5 (0.4%)	1316[3] (100%)
MLS	579 (80%)	816 (73.1%)	623 (47.3%)
Trimethoprim	621 (86.1%)	41 (3.7%)	982 (74.6%)
Phenicols	274 (38%)	176 (15.8%)	675 (51.3%)
Bleomycin	179 (24.8%)	36 (3.2%)	717 (54.5%)
β-lactamase	718 (99.6%)	1018 (91.2%)	1286 (97.7%)
Class C:			
EC	721[1]	2	0
ACT	0	1	0
CMY	209	1	2
DHA	26	2	24
Class A:			
LAP	17	0	142
CARB	0	63	0
PER	0	68	0
TEM	347	697	686
SHV	7	9	1292[3]
VEB	0	12	3
CTX	613	4	681
KPC	94	3	593
Class D:			
OXA	251	996[2]	611
Class B:			
IMP	0	6	1
NDM	173	35	716
Rifamycin	114 (15.8%)	63 (5.6%)	810 (61.6%)
Aminoglycoside	680 (94.3%)	1115 (99.9%)	1290 (98%)
Streptothricin	11 (1.53%)	8 (0.7%)	0 (0%)

371 [1] blaEC intrinsic in *E. coli*

372 [2] blaADC and blaOXA intrinsic in *A. baumannii* (except OXA-[1,10,23,58,72])

 [3] fosA (fosfomycin), oqxAB (fluoroquinolone) and blaSHV intrinsic in *K. pneumoniae*

ESBL resistance genes are highlighted in grey and carbapenem resistance genes in yellow

Most patients carried several AMR strains

Over half of the patients in this study had multiple isolates of the same species during their

stay (*E. coli*: 51%, *K. pneumoniae*: 57%, *A. baumannii*: 58%). Of these patients, 60-70% had

different sequence types (ST) (*E. coli*: 67%, *K. pneumoniae*: 68%, *A. baumannii*: 64%)

(Figure 3). For *E. coli* and *K. pneumoniae*, the majority of patients only had isolates detected

in stool (80% and 54%, respectively). Conversely, most patients with *A. baumannii* were

detected only in sputum (38%), or sputum and stool (32%).

Figure 3: Overview of strain diversity, recurrence and source among study patients: "Patients" refers to

the total number of patients in this study that had at least one isolate of that species. Within each species, we

evaluated whether patients had (i) only a single isolate for that species, or (ii) multiple isolates. If only a single

isolate, we determined whether it was collected on admission to the ICU or after. For multiple isolates, we

determined if the patient's isolates were the same ST (recurrent) or a different ST. We finally looked at how

many patients had isolates from one site (urine, swab, stool or sputum) or mixed sites (any combination of
sites).
393

394   *Evidence of extensive transmission between ICU patients*
395   Temporal observation of the isolates found no obvious association of any time point with any
396   ST to suggest an outbreak of a specific lineage. In order to investigate potential clusters
397   within the ICUs at a higher resolution, we examined plausible short-term transmission events
398   using single nucleotide polymorphisms (SNPs).
399

To identify closely related strains that could indicate recent transmission, we evaluated
clusters based on SNP distances across the core genome of each species for this dataset.
Given the short sampling period, none of the three major species were likely to acquire more
than 1 SNP while in the hospital. As such, we looked at samples with genomic evidence of
most recent transmission; zero SNP clusters. Clusters were defined when they involved >1
patient. Clusters involving a single patient and environmental samples were not included.
406

Most clusters were detected in *K. pneumoniae* and *A. baumannii* isolates, with 71 and 74
clusters representing 38% and 52% of total isolates for that species, respectively (Figure 4).
K. pneumoniae had some of the largest clusters, ranging in size from 2 to 79 isolates, while *A.*
baumannii clusters were smaller, between 2 to 33 isolates. Only 22 clusters were detected in
E. coli and were generally small (median 3 isolates, range 2 to 9 isolates), representing only
13% of the *E. coli* dataset.
413

414   For all three species, the majority of clusters were detected between patients within a single
415   ICU. Evaluating admission and discharge dates further confirmed patient overlap in these
416   clusters (Figure 4, Supplementary Figures 12 to 17).
417

Figure 4: Summary of 0 SNP clusters in all species: Clusters were defined as (i) multiple patients: samples were derived from at least two different patients, or (ii) same patient: isolates were derived from the same patient, or only a single patient and the environment. Epidemiological evidence to support clusters was defined as (i) confirmed patient overlap: all patient ICU stays overlap with another in the same cluster, (ii) some patient overlap: at least 2 patient ICU stays overlap, and (iii) zero patient overlap between all patients in cluster. ENV isolate in clusters: clusters were counted if an environmental isolate was found in that cluster. Colonisation vs. infection: clusters were counted if they (a) had only isolates from stool (i.e. colonisation) or (b) had isolates from urine, swabs and/or sputum with or without isolates from stool (i.e. infection).

A. baumannii clusters were most often associated with environmental isolates (24/74 clusters, Figure 4). The largest *A. baumannii* cluster within the same hospital ICU involved 33 isolates from eight patients and eight environmental samples (Supplementary Figure 13; ST451, cluster number 13). Admissions for patients in this cluster overlapped with detection of the same strain in the environment, which was also detected in the hospital environment several months later. Only two *E. coli* clusters contained related environmental isolates. *K. pneumoniae* environmental isolates were more often found in within-hospital (ICU) clusters (n=11) compared to between-hospital clusters (n=4).

To broadly evaluate infection risk for each outbreak cluster, we determine whether clusters contained (i) stool sample / rectal isolates only, indicating colonisation or (ii) isolates from urine, skin swab or sputum samples, which could represent infection or colonisation at

multiple sites. The majority of *A. baumannii* clusters contained isolates from non-stool /

rectal swab samples (Figure 4). Conversely, colonisation-only clusters were common for *E.*

coli; the largest *E. coli* cluster involving both hospital ICUs contained only 5 isolates from

four patients, which were all isolated from stool (Supplementary Figure 14; ST617, cluster

number 37). *K. pneumoniae* clusters were a mix, with both colonisation-only clusters and

infection clusters.

In addition to suspected within-ICU transmission, we also detected a number of clusters

involving patients from both hospital ICUs (Figure 4). The most pronounced example of this

was a large ST15 *K. pneumoniae* cluster involving 79 isolates from 38 patients and 6

environmental samples (Supplementary Figure 18). Most were collected between July to

September, with some late occurrences in October and November. All patients from NHTD

between July to September had overlapping timelines, consistent with spread within the ICU

(Supplementary figure 16; ST15, cluster number 15). Only one patient had no evidence of

overlap (ND162) but did cluster with environmental isolates from the same timeframe,

indicating a possible environmental source. Patients from BM appeared to have both patient

overlap and consecutive acquisitions without patient overlap as time progressed. This

suggests that patient transmission and also transmission via other routes (e.g. inadequate

cleaning before the next patient, transmission via healthcare workers) may have been

important factors in the spread of this strain.

The identification of closely related isolates between independently operating ICUs

suggested that there may have been a common source located outside the ICU e.g admission

to the same location prior to admission to ICU. To determine if certain lineages were more

associated with acquisition within the ICU, we assessed diversity on arrival (i.e. the patients

first sample) versus diversity within the ICU (all other samples). Based on ST alone, we

found a slight increase in diversity in the ICU versus on arrival (Supplementary Figure 19).

However, the unique STs recovered in either setting only represented a small portion of the

isolates overall. All of the main lineages for each species were found on both admission and

within the ICU (Supplementary Figure 19 and 20).

Figure 5: Scatterpie showing the number of clusters in patients across all species: y-axis represents patients

from BM or NHTD involved in at least one 0 SNP cluster. X-axis represents length of stay for that patient; one

pie is plotted per patient at the duration of their stay. Each circle represents 0 SNP clusters in a single patient.

The size of the clusters corresponds to the number of clusters, while the colour relates to the species. The left

plot shows all 0 SNP clusters, while the right plot shows clusters condensed at 5 SNPs.

Identification of multiple transmission clusters per patients

Overall, there were 251 patients (representing 68% of the cohort) involved in 167 clusters

across the three species collected over the course of this study. 112 patients were only

involved in a single cluster during their time in the ICU (Figure 5). However, the remaining

139 patients were involved in at least two clusters, with one patient involved in 12 clusters

detected at 0 SNPs. For patients with at least two clusters, 20 had clusters from all three

species, 94 had clusters from two species and 25 had only one species. Overall, we saw a

general trend towards more clusters in a single patient as they spent more time in the ICU

ward.

To determine if our clusters were potentially derived from a single original cluster predating

their time in the ICU, we looked at SNP distances between clusters of the same ST

(Supplementary Figure 21). At a threshold of five SNPs, several of the prominent STs within

each species formed large clusters, including ST804 in *A. baumannii* and ST16 in *K.*

pneumoniae. At this threshold, we found 29 clusters in the *A. baumannii* dataset (originally

74), 23 clusters in the *K. pneumoniae* (originally 71) and 19 clusters in *E. coli* (originally 22).

By readjusting our analysis per patient using these clusters, we found that 123 patients had

only a single cluster during their stay (Figure 5). 128 patients had 2 clusters, with the

maximum number of clusters in a single patient being 7 (n=3 patients).

Discussion

Here we present a large prospective surveillance study of three key AMR pathogens from

two hospital ICUs in Vietnam. We used WGS to capture a high-resolution snapshot of the

dominant circulating lineages over a six-month period. In this study we focused on *E. coli, K.*

pneumoniae, and *A. baumannii*, as these were the most commonly isolated species from both

ICUs. These three species have also been reported as highly prevalent in other Vietnamese

studies (*44*), and are amongst the most clinically significant Gram-negative bacteria, having

been designated as "critical" priority pathogens for research and development of new

antibiotics by the World Health Organisation (*45*).

Phylogenetic analysis of all three species suggested local dominance of specific lineages for

K. pneumoniae and *A. baumannii*. Comparison to global representatives found few closely

linked strains, with an overall preference for local clustering. This was particularly evident

for *K. pneumoniae* ST15, ST656, ST11 and ST16 and many of the GC2 isolates for *A.*

baumannii. One notable exception was from a study conducted in southern Vietnam (*46*),

where a number of GC2 *A. baumannii* strains were found to be closely related to ST571

isolates from this study. Similar studies in other Vietnamese hospital settings have also

identified *K. pneumoniae* isolates corresponding to ST15 (*23*), ST16 (*47*) and ST11 (*48*) with

similar antimicrobial resistance profiles carrying *bla*KPC and *bla*NDM. This suggests that

these lineages may not be restricted to referral hospitals in Hanoi but may potentially occur

throughout Vietnam. In contrast, analysis of the study *E. coli* isolates and comparison with

global references did not identify evidence for locally dominant lineages, instead showing

large dispersal of global strains throughout the phylogeny.

Despite limiting our study samples to ESBL-producing and/or carbapenem-resistant isolates

belonging to three species, we identified a large number of isolates with an average of 16.55

and 9.68 isolates/day from patients in BMH and NHTD, respectively. This is an exceedingly

high number compared to other countries, such as the United Kingdom (UK), where a similar

study only found 199 ESBL-producing Enterobacteriaceae over the course of one year (0.5

isolates/day) from three hospital sites between 2008 to 2009 (*49*). A point-prevalence survey

conducted in a UK hospital in 2017 also identified no positive carbapenemase-producing

Enterobacteriaceae (CPE) from 540 samples (*50*).

Nearly all of the isolates presented in this study were characterised as multi-drug resistant

(MDR) on the basis of resistance to three or more antimicrobial drug classes (*51*). *K.*

pneumoniae was found to be particularly resistant, with a number of isolates carrying genes

to more than nine antibiotic classes. The only drug to which we did not observe extensive

resistance was *mcr* gene-mediated colistin resistance, which has been reported in animal

farming and agriculture in Vietnam (*52-55*). It is possible that the *mcr* gene has not yet

become disseminated to more urban areas of Vietnam, despite the use of colistin in clinical

settings. A limited number of isolates displayed mutational changes related to colistin

resistance, such as interruption of the *mgrB* gene in *K. pneumoniae* (*56*). We did not further

investigate the role of mutational colistin resistance as we did not have the capacity to

corroborate genotypic resistance with phenotypic measurements in this study.

Colonisation was found to be a large reservoir for AMR, with the majority of *E. coli* and

more than half of the *K. pneumoniae* isolated from stool samples, as has been documented

previously in Vietnamese hospitals (*57*). High community usage of antibiotics in Vietnam is
likely to promote colonisation with AMR bacteria, as prior treatment with antibiotics is
known to lead to colonisation (*58*). Colonisation itself has been identified as a risk factor for
subsequent infection (*59-61*), and has been previously documented in ICU patients in
southern Vietnam (*62*). Colonisation with AMR *E. coli* has also been identified as a risk for
transferral of resistance to other colonising pathogens, such as *Shigella* (*63*).

Infection control is critically important for reducing the risk of hospital-acquired infections
(HAIs) and mortality in ICUs. Here we found evidence for numerous recent transmission
events involving multiple patients, where *A. baumannii* and *K. pneumoniae* were more often
found in transmission clusters. *A. baumannii* is particularly problematic to control in hospital
environments owing to its ability to resist desiccation and cleaning (*64*) and to survive for
long periods of time on surfaces (*65, 66*). *K. pneumoniae* is also commonly responsible for
outbreaks in healthcare settings globally (*67-69*), and requires immediate and appropriate
intervention due to its propensity to be highly resistant and virulent (*70*). Detection and
prevention of AMR transmission in LMICs is difficult for a number of reasons. These include
limited capacity for microbiological testing, overcrowding of ICUs, inability to isolate ICU
patients, and inadequate staff training or knowledge of infection control procedures (*71, 72*).
The high level of AMR in this study would have made it difficult to discern specific clusters
using phenotypic methods alone. WGS enables accurate investigation of transmission events
but its use is limited by lack of infrastructure, expertise and high cost, particularly in
resource-constrained settings.

In addition to evidence for recent transmission between patients on the same ICU, we also
identified clusters involving patients from both hospital ICUs. This result was unexpected, as
there was no direct transfer of patients between the two ICUs. The most likely explanation
for these clusters is a source outside of the ICUs, such as other wards or other hospitals which
may have referred patients to ICU. Another possibility is that AMR strains may have been
acquired in the community, reflecting high rates antibiotic usage in the community and AMR
detection in livestock and food. Based on the similarity between lineages and AMR across
both ICUs, we suggest that transmission, particularly of the predominant *A. baumannii* and *K.*
pneumoniae lineages, is likely already circulating outside of the ICUs, where it is then further
propagated.

We acknowledge some limitations to our study. First, contamination of some of the isolates
prior to WGS necessitated additional filtering steps for the majority of samples. Secondly, we
did not extensively explore plasmid profiles amongst the samples because of the limitations
of short read sequence data. Thirdly, although serial samples were collected and cultured, we
selected single colony picks for sequencing. It is therefore possible that some diversity may
have been lost at different timepoints during the study sampling period. Finally, this study
focused on patient and environmental samples only. We were therefore unable to investigate
potential transmission events involving hospital staff and/or visitors.
593
Nevertheless, we present the largest prospective surveillance study of multidrug-resistant *E.*
coli, *A. baumannii* and *K. pneumoniae* in Vietnamese critical care patients to date. The
extensive transmission and AMR detected within and between ICU wards suggests dominant
circulating lineages of *A. baumannii* and *K. pneumoniae* existing both within hospitals, and
potentially in community settings in Vietnam. Further work should be conducted to expand
genomic surveillance in hospital and community settings to determine the levels of AMR and
prominent lineages in order to inform AMR control strategies in Vietnam.
601
Data sharing
Genome sequence data have been deposited in the European Nucleotide Archive (ENA)
under the Bioproject PRJEB29424. A list of the sample accession numbers is available in
Supplementary Dataset 1. Isolate genome assemblies (heterogenous sites masked and
unmasked)   are   available   on   Figshare   under   the   following   DOI:
10.6084/m9.figshare.13303253,
10.6084/m9.figshare.13302728.
609
Ethical statement
The study protocol was approved by the Scientific and Ethical Committees of the National
Hospital for Tropical Diseases and Bach Mai Hospital and by the University of Cambridge
Human Biology and Research Ethics Committee (reference: HBREC 2017.09). Written
informed consent was obtained from the patient or from their relative prior to enrolment in
the study.
616
617   *Declaration of interests*
618   All authors have no conflicts of interest to declare.

Funding statement

This work was funded by the Medical Research Council Newton Fund, the Vietnamese Ministry of Science and Technology (HNQT/SPĐP/04.16), and the Wellcome Trust. LWR is supported by an EMBL-EBI biomedical postdoctoral research fellowship. MET was supported by a Clinician Scientist Fellowship (funded by the Academy of Medical Sciences and the Health Foundation) and the NIHR Cambridge Biomedical Research Centre. The funding source had no role in study design; in the collection, analysis, and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Author contributions

Conceptualisation: LWR, ZI, MET

Data collection: FK, NTHM, LTH, NGB, DXC, LTH, NTH, TG, CB, HNT, BN, RvD, NVT

Sample processing: NTH, FK, JB, JH, TF

Methodology: LWR, ZI

Formal Analysis: LWR, ZI

Writing (original draft): LWR

Writing (review/editing): ZI, MET

Supervision: NVT, RvD, ZI, MET

Project administration: LTH, NVT, RvD, MET

Funding acquisition: MET, JP and NVK

Acknowledgements

The authors would like to thank the patients for participating in this study and the clinical and laboratory staff of the National Hospital for Tropical Diseases and Bach Mai Hospital for their assistance with this study. We also acknowledge the sequencing team at the Wellcome Sanger Institute for their assistance with sequencing the samples including in the study.

References

1. R. M. Zellweger *et al.*, A current perspective on antimicrobial resistance in Southeast Asia. *J Antimicrob Chemother* **72**, 2963-2972 (2017).
2. B. A. Walther *et al.*, Biodiversity and health: Lessons and recommendations from an interdisciplinary conference to advise Southeast Asian research, society and policy. *Infect Genet Evol* **40**, 29-46 (2016).

3.   J. J. Carrique-Mas, M. Choisy, N. Van Cuong, G. Thwaites, S. Baker, An estimation
of total antimicrobial usage in humans and animals in Vietnam. *Antimicrob Resist*
Infect Control **9**, 16 (2020).

4.   K. V. Nguyen *et al.*, Antibiotic use and resistance in emerging economies: a situation
analysis for Viet Nam. *BMC Public Health* **13**, 1158 (2013).

5.   M. E. de Kraker, A. J. Stewardson, S. Harbarth, Will 10 Million People Die a Year
due to Antimicrobial Resistance by 2050? *PLoS Med* **13**, e1002184 (2016).

6.   N. H. Tran *et al.*, Occurrence and risk assessment of multiple classes of antibiotics in
urban canals and lakes in Hanoi, Vietnam. *Sci Total Environ* **692**, 157-174 (2019).

7.   S. Yamasaki, T. D. Le, M. Q. Vien, C. Van Dang, Y. Yamamoto, Prevalence of
extended-spectrum beta-lactamase-producing Escherichia coli and residual
antimicrobials in the environment in Vietnam. *Anim Health Res Rev* **18**, 128-135
(2017).

8.   H. A. Duong *et al.*, Occurrence, fate and antibiotic resistance of fluoroquinolone
antibacterials in hospital wastewaters in Hanoi, Vietnam. *Chemosphere* **72**, 968-973
(2008).

9.   N. T. Nhung *et al.*, Antimicrobial residues and resistance against critically important
antimicrobials in non-typhoidal Salmonella from meat sold at wet markets and
supermarkets in Vietnam. *Int J Food Microbiol* **266**, 301-309 (2018).

10.   S. P. T. S. S. A. K. Anal, Addressing the antibiotic resistance and improving the food
safety in food supply chain (farm-to-fork) in Southeast Asia. *Food Control* **108**,
(2020).

11.   T. Phodha, A. Riewpaiboon, K. Malathum, P. C. Coyte, Annual relative increased in
inpatient mortality from antimicrobial resistant nosocomial infections in Thailand.
Epidemiol Infect **147**, e133 (2019).

12.   K. L. Wyres *et al.*, Genomic surveillance for hypervirulence and multi-drug resistance
in invasive Klebsiella pneumoniae from South and Southeast Asia. *Genome Med* **12**,
11 (2020).

13.   M. D. Malchione, L. M. Torres, D. M. Hartley, M. Koch, J. L. Goodman,
Carbapenem and colistin resistance in Enterobacteriaceae in Southeast Asia: Review
and mapping of emerging and overlapping challenges. *Int J Antimicrob Agents* **54**,
381-399 (2019).

14.   V. Le Minh *et al.*, In vitro activity of colistin in antimicrobial combination against
carbapenem-resistant Acinetobacter baumannii isolated from patients with ventilator-
associated pneumonia in Vietnam. *J Med Microbiol* **64**, 1162-1169 (2015).

15.   D. J. Biedenbach *et al.*, Antimicrobial-resistant Pseudomonas aeruginosa and
Acinetobacter baumannii From Patients With Hospital-acquired or Ventilator-
associated Pneumonia in Vietnam. *Clin Ther* **38**, 2098-2105 (2016).

16.   S. Quainoo *et al.*, Whole-Genome Sequencing of Bacterial Pathogens: the Future of
Nosocomial Outbreak Analysis. *Clin Microbiol Rev* **30**, 1015-1063 (2017).

17.   M. S. Toleman *et al.*, Prospective genomic surveillance of methicillin-resistant
Staphylococcus aureus (MRSA) associated with bloodstream infection, England, 1
October 2012 to 30 September 2013. *Euro Surveill* **24**, (2019).

18.   H. T. T. Phan *et al.*, Illumina short-read and MinION long-read WGS to characterize
the molecular epidemiology of an NDM-1 Serratia marcescens outbreak in Romania.
J Antimicrob Chemother **73**, 672-679 (2018).

19.   E. M. Harrison *et al.*, Transmission of methicillin-resistant Staphylococcus aureus in
long-term care facilities and their related healthcare networks. *Genome Med* **8**, 102
(2016).

20. A. Santona *et al.*, Emergence of unusual vanA/vanB2 genotype in a highly mutated
vanB2-vancomycin-resistant hospital-associated E. faecium background in Vietnam.
Int J Antimicrob Agents **52**, 586-592 (2018).
21. H. Chung The *et al.*, A high-resolution genomic analysis of multidrug-resistant
hospital outbreaks of Klebsiella pneumoniae. *EMBO Mol Med* **7**, 227-239 (2015).
22. B. Berglund *et al.*, Colistin- and carbapenem-resistant Klebsiella pneumoniae
carrying mcr-1 and blaOXA-48 isolated at a paediatric hospital in Vietnam. *J*
Antimicrob Chemother **73**, 1100-1102 (2018).
23. B. Berglund *et al.*, Molecular and phenotypic characterization of clinical isolates
belonging to a KPC-2-producing strain of ST15 Klebsiella pneumoniae from a
Vietnamese pediatric hospital. *Antimicrob Resist Infect Control* **8**, 156 (2019).
24. . vol. 2020.
25. P. Ewels, M. Magnusson, S. Lundin, M. Kaller, MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047-3048
(2016).
26. D. E. Wood, J. Lu, B. Langmead, Improved metagenomic analysis with Kraken 2.
Genome Biol **20**, 257 (2019).
27. J. L. F. P. B. P. T. S. L. Salzberg, Bracken: estimating species abundance in
metagenomics data. *PeerJ Computer Science* **3:e104**, (2017).
28. A. Bankevich *et al.*, SPAdes: a new genome assembly algorithm and its applications
to single-cell sequencing. *J Comput Biol* **19**, 455-477 (2012).
29. A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072-1075 (2013).
30. D. H. Parks, M. Imelfort, C. T. Skennerton, P. Hugenholtz, G. W. Tyson, CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043-1055 (2015).
31. T. J. Treangen, B. D. Ondov, S. Koren, A. M. Phillippy, The Harvest suite for rapid
core-genome alignment and visualization of thousands of intraspecific microbial
genomes. *Genome Biol* **15**, 524 (2014).
32. N. J. Croucher *et al.*, Rapid phylogenetic analysis of large samples of recombinant
bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* **43**, e15 (2015).
33. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis
of large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
34. J. Beghain, A. Bridier-Nahmias, H. Le Nagard, E. Denamur, O. Clermont,
ClermonTyping: an easy-to-use and accurate in silico method for Escherichia genus
strain phylotyping. *Microb Genom* **4**, (2018).
35. I. Letunic, P. Bork, Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Res* **47**, W256-W259 (2019).
36. T. Seemann, Abricate, Github https://github.com/tseemann/abricate.
37. M. Feldgarden *et al.*, Validating the AMRFinder Tool and Resistance Gene Database
by Using Antimicrobial Resistance Genotype-Phenotype Correlations in a Collection
of Isolates. *Antimicrob Agents Chemother* **63**, (2019).
38. M. Inouye *et al.*, SRST2: Rapid genomic surveillance for public health and hospital
microbiology labs. *Genome Med* **6**, 90 (2014).
39. L. Diancourt, V. Passet, A. Nemec, L. Dijkshoorn, S. Brisse, The population structure
of Acinetobacter baumannii: expanding multiresistant clones from an ancestral
susceptible genetic pool. *PLoS One* **5**, e10034 (2010).
40. L. Diancourt, V. Passet, J. Verhoef, P. A. Grimont, S. Brisse, Multilocus sequence
typing of Klebsiella pneumoniae nosocomial isolates. *J Clin Microbiol* **43**, 4178-4182
(2005).

41. K. A. Jolley, J. E. Bray, M. C. J. Maiden, Open-access bacterial population genomics:
BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open*
Res **3**, 124 (2018).
42. J. Stimson *et al.*, Beyond the SNP Threshold: Identifying Outbreak Clusters Using
Inferred Transmissions. *Mol Biol Evol* **36**, 587-603 (2019).
43. J. F. Turton *et al.*, Identification of Acinetobacter baumannii by detection of the
blaOXA-51-like carbapenemase gene intrinsic to this species. *J Clin Microbiol* **44**,
2974-2976 (2006).
44. T. V. D. Vu *et al.*, Antimicrobial susceptibility testing and antibiotic consumption
results from 16 hospitals in Viet Nam: The VINARES project 2012-2013. *J Glob*
Antimicrob Resist **18**, 269-278 (2019).
45. W. H. Organisation. (2017), vol. 2020.
46. T. Tada *et al.*, Dissemination of clonal complex 2 Acinetobacter baumannii strains co-
producing carbapenemases and 16S rRNA methylase ArmA in Vietnam. *BMC Infect*
Dis **15**, 433 (2015).
47. T. Tada *et al.*, Dissemination of Carbapenem-resistant Klebsiella pneumoniae clinical
isolates with various combinations of Carbapenemases (KPC-2, NDM-1, NDM-4, and
OXA-48) and 16S rRNA Methylases (RmtB and RmtC) in Vietnam. *BMC Infect Dis*
17, 467 (2017).
48. J. Chang *et al.*, Emergence of NDM-4-producing Klebsiella pneumoniae in a Korean
hospital due to a patient hospitalized in Vietnam and case review. *J Infect Chemother*
25, 909-912 (2019).
49. D. A. Enoch *et al.*, Epidemiology of extended-spectrum beta-lactamase-producing
Enterobacteriaceae in a UK district hospital; an observational study. *J Hosp Infect* **81**,
270-277 (2012).
50. H. J. Wilson *et al.*, Point-prevalence survey of carbapenemase-producing
Enterobacteriaceae and vancomycin-resistant enterococci in adult inpatients in a
university teaching hospital in the UK. *J Hosp Infect* **100**, 35-39 (2018).
51. A. P. Magiorakos *et al.*, Multidrug-resistant, extensively drug-resistant and pandrug-
resistant bacteria: an international expert proposal for interim standard definitions for
acquired resistance. *Clin Microbiol Infect* **18**, 268-281 (2012).
52. Y. Yamamoto *et al.*, Wide dissemination of colistin-resistant Escherichia coli with the
mobile resistance gene mcr in healthy residents in Vietnam. *J Antimicrob Chemother*
74, 523-524 (2019).
53. T. Yamaguchi *et al.*, The presence of colistin resistance gene mcr-1 and -3 in ESBL
producing Escherichia coli isolated from food in Ho Chi Minh City, Vietnam. *FEMS*
Microbiol Lett **365**, (2018).
54. N. V. Trung *et al.*, Zoonotic Transmission of mcr-1 Colistin Resistance Gene from
Small-Scale Poultry Farms, Vietnam. *Emerg Infect Dis* **23**, 529-532 (2017).
55. S. Malhotra-Kumar *et al.*, Colistin-resistant Escherichia coli harbouring mcr-1
isolated from food animals in Hanoi, Vietnam. *Lancet Infect Dis* **16**, 286-287 (2016).
56. L. Poirel *et al.*, The mgrB gene as a key target for acquired resistance to colistin in
Klebsiella pneumoniae. *J Antimicrob Chemother* **70**, 75-80 (2015).
57. D. M. Tran *et al.*, High prevalence of colonisation with carbapenem-resistant
Enterobacteriaceae among patients admitted to Vietnamese hospitals: Risk factors and
burden of disease. *J Infect* **79**, 115-122 (2019).
58. J. Ruiz *et al.*, Influence of antibiotic pressure on multi-drug resistant Klebsiella
pneumoniae colonisation in critically ill patients. *Antimicrob Resist Infect Control* **8**,
38 (2019).

59.   C. M. Manaia, Assessing the Risk of Antibiotic Resistance Transmission from the
Environment to Humans: Non-Direct Proportionality between Abundance and Risk.
Trends Microbiol **25**, 173-181 (2017).
60.   M. E. Flokas, S. A. Karageorgos, M. Detsis, M. Alevizakos, E. Mylonakis,
Vancomycin-resistant enterococci colonisation, risk factors and risk for infection
among hospitalised paediatric patients: a systematic review and meta-analysis. *Int J*
Antimicrob Agents **49**, 565-572 (2017).
61.   J. Asir *et al.*, Simultaneous gut colonisation and infection by ESBL-producing
Escherichia coli in hospitalised patients. *Australas Med J* **8**, 200-207 (2015).
62.   D. B. Thuy, Colonisation and Infection with Antibiotic-Resistant Organisms in
Patients in the Adult Intensive Care Unit, Hospital for Tropical Diseases, Ho Chi
Minh City, Vietnam. *PhD thesis, University of Oxford*, (2018).
63.   P. Thanh Duy *et al.*, Commensal Escherichia coli are a reservoir for the transfer of
XDR plasmids into epidemic fluoroquinolone-resistant Shigella sonnei. *Nat Microbiol*
5, 256-264 (2020).
64.   A. Y. Peleg, H. Seifert, D. L. Paterson, Acinetobacter baumannii: emergence of a
successful pathogen. *Clin Microbiol Rev* **21**, 538-582 (2008).
65.   P. J. van den Broek *et al.*, Epidemiology of multiple Acinetobacter outbreaks in The
Netherlands during the period 1999-2001. *Clin Microbiol Infect* **12**, 837-843 (2006).
66.   A. Jawad, H. Seifert, A. M. Snelling, J. Heritage, P. M. Hawkey, Survival of
Acinetobacter baumannii on dry surfaces: comparison of outbreak and sporadic
isolates. *J Clin Microbiol* **36**, 1938-1941 (1998).
67.   E. S. Snitkin *et al.*, Tracking a hospital outbreak of carbapenem-resistant Klebsiella
pneumoniae with whole-genome sequencing. *Sci Transl Med* **4**, 148ra116 (2012).
68.   D. Marchaim *et al.*, Outbreak of colistin-resistant, carbapenem-resistant Klebsiella
pneumoniae in metropolitan Detroit, Michigan. *Antimicrob Agents Chemother* **55**,
593-599 (2011).
69.   D. Gu *et al.*, A fatal outbreak of ST11 carbapenem-resistant hypervirulent Klebsiella
pneumoniae in a Chinese hospital: a molecular epidemiological study. *Lancet Infect*
Dis **18**, 37-46 (2018).
70.   K. E. Holt *et al.*, Genomic analysis of diversity, population structure, virulence, and
antimicrobial resistance in Klebsiella pneumoniae, an urgent threat to public health.
Proc Natl Acad Sci U S A **112**, E3574-3581 (2015).
71.   M. Vukoja *et al.*, A survey on critical care resources and practices in low- and
middle-income countries. *Glob Heart* **9**, 337-342 e331-335 (2014).
72.   M. J. Schultz *et al.*, Current challenges in the management of sepsis in ICUs in
resource-poor settings and suggestions for the future. *Intensive Care Med* **43**, 612-624
(2017).
839