

1 **Concurrent validity of an Estimator of Weekly Alcohol** 2 **Consumption (EWAC) based on the Extended AUDIT**

3

4 **Authors**

5 Dutey-Magni, PF,^{1,4,*} Brown, J,² Holmes, J,³ Sinclair, JMA⁴

6

7 1 Institute of Health Informatics, University College London, London, UK

8 2 Institute of Epidemiology and Health Care, University College London, London, UK

9 3 School of Health and Related Research, University of Sheffield, Sheffield, UK

10 4 Faculty of Medicine, University of Southampton, London, UK

11

12 *** Corresponding author:**

13 Peter F. Dutey-Magni

14 Institute of Health Informatics

15 University College London

16 222 Euston Road

17 London

18 NW1 2DA

19 UK

20 Telephone: (+44) 020 3108 6424

21 Email: p.dutey-magni@ucl.ac.uk

22

23 **Running head**

24 Estimator of Weekly Alcohol Consumption EWAC

25 **Word count**

26 3496 words

27 **Protocol registration**

28 Dutey-Magni, PF, Sinclair, JMA, Brown, J. 2018. 'Concurrent Validity of an Estimator
29 of Weekly Alcohol Consumption (EWAC) Based on the Extended AUDIT.' OSF.

30 November 12. doi:[10.17605/OSF.IO/7WE4M](https://doi.org/10.17605/OSF.IO/7WE4M).

31 **Abstract**

32 **Background and Aims:** The 3-question Alcohol Use Disorders Identification Test (AUDIT-
33 C) is commonly employed in healthcare to screen for levels of alcohol consumption. AUDIT-
34 C scores (0–12) have no direct interpretation and do not provide information on alcohol
35 intake, an important variable for behaviour change. The study aimed to (a) develop a
36 continuous metric from the Extended AUDIT-C, offering equivalent accuracy, and providing a
37 direct estimator of weekly alcohol consumption (EWAC); (b) evaluate the EWAC's bias and
38 error using the Graduated-Frequency (GF) questionnaire as a reference standard of alcohol
39 consumption.

40 **Design:** Cross-sectional diagnostic study based on a nationally-representative survey.

41 **Settings:** Community-dwelling households in England.

42 **Participants:** 22,404 household residents aged ≥ 16 years reporting drinking alcohol at least
43 occasionally.

44 **Measurements:** Computer-assisted personal interviews consisting of (a) Extended AUDIT
45 and (b) GF. The primary outcomes were: mean deviation < 1 UK unit (metric of bias); root
46 mean squared deviation < 2 UK units (metric of total error) between EWAC and GF. The
47 secondary outcome was an equivalent receiver operating characteristic area under the curve
48 for predicting alcohol consumption in excess of 14 and 35 UK units compared to AUDIT-C
49 and AUDIT.

50 **Findings:** EWAC had a positive bias of 0.2 UK units [95% confidence interval: 0.04, 0.3]
51 compared to GF. Deviations were skewed: while the mean error was ± 11 UK units/week
52 [9.8, 12.0], in half of participants the deviation between EWAC and GF was between 0 and
53 ± 2.1 UK units/week. EWAC predicted consumption in excess of 14 UK units/week with a
54 significantly greater area under the curve (0.921 [0.917, 0.925]) than AUDIT-C (0.871 [0.866,
55 0.876]) or the full AUDIT (0.854 [0.849, 0.860]).

56 **Conclusions:** The EWAC <https://ewac.netlify.app> is designed to estimate weekly alcohol
57 consumption using answers to the Extended AUDIT-C questionnaire. Using the detailed GF
58 as a reference standard, the EWAC met the targeted bias tolerance. Its accuracy was
59 superior to that of both AUDIT-C and the full AUDIT in relation to consumption thresholds,
60 making it a reliable complement to the Extended AUDIT-C for health promotion
61 interventions.

62 **Keywords**

63 alcohol consumption; self report; alcohol use disorder; screening programs, diagnostic;
64 preventive health services

65

66 1 Introduction

67 Alcohol consumption is responsible for 5% of disability-adjusted life years [1]. This burden
68 extends far beyond the health burden of *alcohol use disorders*, as defined in the
69 International Statistical Classification of Diseases (ICD-10 F10.1/F10.2 [2]) or the Diagnostic
70 and Statistical Manual of Mental Disorders [3]. Clinical guidelines aiming to prevent [4], treat
71 and reduce [5] harm from alcohol consumption recommend systematic screening for alcohol
72 consumption using a validated diagnostic tool. However, conceptual differences (exemplified
73 by the diagnostic classifications above) remain in how best to diagnose, measure, and
74 communicate harm [6].

75 A global standard has emerged in the 10-item Alcohol Use Disorders Identification Test
76 (AUDIT) [7]. Its first three questions, known as AUDIT-C and focussing on consumption,
77 have equivalent predictive capability [8]. AUDIT-C is easy to use for patients and clinicians
78 alike, making it an attractive choice for alcohol interventions in healthcare and other settings.
79 Yet, AUDIT-C exhibits two limitations:

- 80 1. **Ceiling effect:** AUDIT-C's maximum response options for alcohol consumption
81 frequency and quantity are heavily right-censored (Table 1). This creates a ceiling
82 effect making the AUDIT-C poorly responsive to change in individuals with a high
83 baseline score, even if they reduce consumption by up to thirty percent (e.g. frequency
84 of drinking down from 7 to 5 days or quantity down from 16 to 11 drinks per day).
- 85 2. **AUDIT score intelligibility:** when presented alone, the AUDIT-C's ordinal score
86 (range: 0–12) is not easily interpretable to patients and clinicians alike, who must be
87 refer to cut-offs informed by diagnostic accuracy studies [9] and set with great variation
88 internationally [10]. The fact that the AUDIT-C score is not easy to interpret poses
89 challenges in the delivery of brief interventions. Healthcare professionals lack
90 confidence in discussing alcohol consumption [11–13], and the abstract AUDIT-C score
91 requires training on how to deliver feedback [11]. Similar challenges are faced when

92 communicating AUDIT-C results in self-administered interventions such as web apps
93 [\[14\]](#).

94 The 'Extended AUDIT-C' addresses the first of these limitations thanks to a greater range of
95 response options on quantity and frequency (Table 1). It has been used in UK research as
96 part of two trials [\[15,16\]](#) and one continuous household survey [\[17\]](#) to measure
97 characteristics of consumption that could not have been measured with the right-censored
98 AUDIT-C.

99 The present study proposes to address the second of these limitations, by developing and
100 validating a fast Estimator of Weekly Alcohol Consumption (EWAC) computed from the
101 Extended AUDIT-C. The EWAC is designed to offer equivalent diagnostic capabilities in
102 relation to alcohol use disorders (being based on the AUDIT-C), while providing a continuous
103 and more directly interpretable metric of alcohol consumption. This is intended facilitate the
104 delivery of brief interventions, since measuring alcohol consumption is a crucial part of
105 behaviour change techniques (self-monitoring, feedback on behaviour, social comparison)
106 commonly employed in self- [\[18\]](#) and clinician-administered [\[19\]](#) interventions.

107 This structure of the paper is as follows. The paper reports the development of the EWAC,
108 which is the product of quantity (AUDIT-1) by frequency (AUDIT-2) with adjustment for
109 occasional intense drinking (AUDIT-3). Coefficients are estimated in a hierarchical Bayesian
110 response model using Extended AUDIT-C and Graduated-Frequency (GF) data from a large
111 English household survey, the Alcohol Toolkit Study [\[17\]](#). The same data are then used to
112 compare EWAC and GF, derive metrics of bias and error, and detect potential subgroup
113 variations in accuracy. Finally, the paper tests the equivalence of receiver operating
114 characteristics of EWAC compared with AUDIT-C and the full AUDIT in predicting alcohol
115 consumption exceeding 112g/week (14 UK units) and 280g/week (35 UK units).

116 **2 Methods**

117 **2.1 Design**

118 Neither the AUDIT-C nor the Extended AUDIT-C provide a direct measure of alcohol
119 consumption. Instead, we employ methods developed for quantity-frequency-variability
120 instruments [20]. For every individual i , the EWAC is computed as the product of F_i and Q_i
121 (AUDIT questions 1 and 2 respectively) adjusted with the frequency of intense drinking V_i
122 (AUDIT-3):

$$EWAC_i = F_i Q_i + V_i b$$

123 where b denotes the average units of alcohol consumed in an intense drinking day.

124 Coefficients F, Q, V and b are unknown. The present study considers two sets of candidate
125 coefficients:

- 126 • AUDIT response item interval midpoint (e.g. 2.5 for '2 to 3 times per week')
- 127 • coefficients estimated empirically from a sample of individuals with measurements of
128 Extended AUDIT-C and GF, using a hierarchical Bayesian response model with the
129 estimating equation $GF_i = F_i Q_i + V_i b + e_i$, where e denotes independently normally
130 distributed errors. We set parabola-shaped informative priors on coefficients F, Q, V .
131 Details on model fitting, convergence evaluation and prior tuning is reported in
132 supplementary information S1.

133 **2.2 Participants**

134 Data originate from baseline measures in waves 110–133 (November 2015–October 2017)
135 of the Alcohol Toolkit Study (ATS), a computer-assisted personal interview of residents of
136 private English households aged ≥ 16 years [17]. Participants were included in the analysis if
137 they completed both the Extended AUDIT and the GF questionnaires. Out of 40,832
138 participants, 14,408 (35%) reported 'never' consuming alcohol in AUDIT question 1 and
139 were not asked any further AUDIT or GF questions. They were thus excluded.

140 **2.3 Measures**

141 Index measures underpinning the EWAC were the three questions making up the Extended
142 AUDIT-C (supplementary information S2), in which participants described their drinking
143 *during the last 6 months*.

144 The reference standard used is the ATS GF schedule (supplementary information S3), in
145 which participants described how many times they consumed given quantities of alcohol
146 *during the last 4 weeks* [21]. A longstanding obstacle in alcohol research and treatment lies
147 in the absence of undisputed 'gold standard' or biomarker for objectively determining alcohol
148 consumption. Instead, a number of instruments measure self-reported consumption with
149 varying validity and reliability over different durations. Comprehensive reviews [22–27]
150 indicate that yesterday recall and prospective diaries tend to record higher (and more
151 accurate) alcohol consumption by minimising recall bias, followed by GF measures. The GF
152 schedule's main advantage lies in measuring occasional heavy consumption, which can
153 constitute an important proportion of total consumption. Although widely employed in
154 population surveys, the GF schedule is uncommon in clinical practice.

155 We examined the EWAC's coverage of per-capita alcohol retail sales for England in 2014
156 [28], which capture all alcohol produced/processed in or imported to England for sale or
157 consumption.

158 We also cross-examined the EWAC's distribution against other reference estimates obtained
159 from 8,610 household residents aged ≥ 16 years participating in the 2011 Health Survey for
160 England (HSE) [29]. On that particular year, the HSE included (a) a computer-assisted
161 interviewer-led beverage-specific quantity-frequency questionnaire; (b) a yesterday recall
162 question [27]; and (c) a prospective 7-day diary [30].

163 **2.4 Missing data**

164 We excluded 4,020/26,424 participants (15.2%) who did not have a valid GF alcohol
165 consumption record (Table S7). In the remaining 22,404 valid observations, GF data were

166 assumed to be missing at random conditionally on the Extended AUDIT-C responses. In
167 subgroup analyses, a further 530 observations (2.4%) assumed to be missing at random
168 were excluded.

169 **2.5 Analyses**

170 The protocol was pre-registered [31]. Results are reported in UK alcohol units (8g or 10mL of
171 pure alcohol). Analyses were conducted in R [32–34]. Scripts are available online [35].

172 **2.5.1 Overall bias and error**

173 The agreement between the EWAC and the GF was quantified as follows:

174 • bias was estimated by the **mean deviation** to the reference standard
175 $MD = n^{-1} \sum_{i=1}^n (EWAC - GF)$. We tested the hypothesis that the MD does not exceed 1
176 UK unit using a two-sided *t*-test.

177 • precision was estimated by the **root mean squared deviation**
178 $RMSD = \sqrt{n^{-1} \sum_{i=1}^n (EWAC - GF)^2}$, a measure of total error capturing both bias and
179 random deviation from the reference standard. For example, an RMSD of 2 signifies that
180 the EWAC is on average with ± 2 UK units of the reference standard. We tested the
181 hypothesis that the RMSD does not exceed 2 UK units using a one-sided χ^2
182 homogeneity test.

183 Two sets of candidate coefficients were considered (section 2.1). We only report findings for
184 the candidate set producing the lowest bias and error.

185 **2.5.2 Subgroup bias and error**

186 Multivariate regression models tested whether the EWAC's validity varies across population
187 subgroups:

188 • the simple deviation ($EWAC - GF$) was regressed in a linear model to test subgroup
189 differences in MD

190 • the squared deviation $(EWAC - GF)^2$ was regressed in a log-transformed linear model to
191 test subgroup differences in the geometric mean squared deviation. Model coefficients
192 were then back-transformed (square root of the exponential) into relative RMSD
193 estimates; these are interpreted as the ratio of the subgroup RMSD to the reference
194 RMSD, a ratio >1 indicating worse precision than in the reference category.

195 Both models (supplementary Tables S5–S6) included the following predictors: sex by age
196 group; ethnic group; highest educational qualification; religion; smoking status. Additional
197 models (supplementary Table S6) were fitted solely in respondents with an AUDIT-C score
198 ≥ 5 or an AUDIT score ≥ 8 , for whom additional characteristics were recorded during
199 interview: favourite drink (beer; wine; spirits alone; mixed spirits; cider; other); and whether
200 the respondent had attempted to restrict alcohol intake in the last 12 months.

201 **2.5.3 Receiver Operating Characteristics**

202 We tested the EWAC's superiority to the traditional AUDIT and AUDIT-C scores in predicting
203 consumption exceeding 14 or 35 UK units/week. These correspond to UK thresholds for
204 characterising alcohol use as 'increasing risk' (predicted by an AUDIT-C score of 5–7), and
205 'higher risk' (AUDIT-C ≥ 8) which is above 35 units for women and 50 units for men [36]. We
206 tested the hypothesis that the EWAC has an identical receiver operating characteristic full
207 area under the curve (AUC) to the AUDIT-C and the full AUDIT scores using nonparametric
208 paired AUC tests [37]. AUDIT-C and AUDIT scores were calculated from the Extended
209 AUDIT by capping the contribution of each question to 4.

210 **2.5.4 Aggregate concurrent validity**

211 We compared the empirical cumulative distributions of (1) the EWAC computed in the ATS;
212 (2) the GF estimator in the ATS; (3) the beverage-specific estimator in the 2011 HSE; (4) the
213 prospective diary estimator in the 2011 HSE in adults aged ≥ 18 years. We report the
214 proportions of on-trade and off-trade alcohol sales [28] accounted for by each method.
215 Poststratification survey weights adjusted for nonresponse bias in sources (1-3), and self-
216 selection into prospective diary data collection in source (4).

217 **3 Results**

218 **3.1 Overall bias and accuracy**

219 EWAC coefficients estimated empirically (supplementary information S4) had smaller bias
220 and error and were used for the remainder of the analysis. With those, the EWAC's
221 Pearson's correlation with GF was estimated at $r = 0.71$ [0.71, 0.72] (Kendall's rank
222 correlation $\tau = 0.63$).

223 The mean deviation (MD) was 0.2 alcohol units/week [95% CI: 0.04, 0.3]. This bias is
224 smaller than the preregistered ± 1 -unit bias tolerance ($p = 1.000$).

225 The root mean squared deviation (RMSD) at 10.9 units/week [95% CI: 9.8, 12.0] was
226 significantly greater than the pre-registered 2-unit total error tolerance ($p < 0.001$), suggesting
227 that the EWAC falls on average 11 units away from the GF reference standard.

228 The RMSD masked a dispersed and skewed distribution of error (Table 2). In 50% of
229 participants, the EWAC fell within ± 2 UK units of the GF weekly consumption estimate.
230 Thus, an interval defined as the EWAC ± 2 units (e.g. '10 to 14 units') would contain the
231 reference standard for half of individuals, while an interval ranging ± 3 units (e.g. '9 to 15
232 units') would contain the reference standard for 60% of individuals.

233 Plots of EWAC against GF (Figure 1) indicate a slight positive bias for consumptions up to
234 10–14 units/week, and a slight negative bias beyond. The EWAC only starts losing
235 granularity above 70 units/week (99th percentile of its distribution), where it provides just 6
236 possible values (82; 83; 92; 93; 100; 125 units/week; see Figure 1(b)).

237 **3.2 Subgroup accuracy**

238 MDs and RMSDs were regressed against respondent characteristics to identify potential
239 subgroup differences in bias or precision from the reference category (White British females
240 aged 25–34 years without educational qualifications, who never smoked). Model predictors
241 accounted for <2% of the variance, indicating stability in MD and RMSD across subgroups
242 (Table S5).

243 Figure 2 shows subgroups with a predicted MD exceeding ± 1 unit/week and a p -value <0.05 .
244 Respondents of Black/Other/White Other ethnic groups had an overestimated EWAC: their
245 MDs exceeded the reference MD by 2.0 units [0.9, 3.1]; 2.7 units [0.8, 4.5]; and 1.1 units
246 [0.5, 1.7] respectively. MDs of men aged 55–64 years, or ≥ 75 years exceeded the reference
247 MD by 1.3 units [0.2, 2.4] and 1.5 units [0.3, 2.8] respectively. There was no evidence of an
248 effect of favourite drink or attempts to reduce alcohol intake in increasing-risk drinkers (Table
249 S6).

250 Figure 3 shows subgroups with an RMSD at least 25% smaller/greater than the reference.
251 RMSD was 58% [50, 67%] greater in current smokers and 34% [14, 56%] greater in
252 respondents who stopped smoking in the past year. It was also 44% [29, 60%] and 34% [19,
253 50%] greater respectively in men and respondents aged 16–24 years. In contrast, error was
254 35 to 70% smaller in Black/Other/White Other ethnic groups.

255 Figure 4 presents the same metrics for increasing-/higher-risk drinkers (AUDIT-C ≥ 5 or
256 AUDIT ≥ 8). Educational qualifications significantly improved the agreement between EWAC
257 and the reference standard in this group (Table S7). School and degree-level qualifications
258 reduced RMSD by 24% [12, 37%] and 37% [23, 52%] respectively, suggesting that
259 respondents may better evaluate and recall alcohol beverage content. Conversely, the
260 RMSD of respondents who attempted to reduce alcohol use was 23% [16, 30%] larger than
261 the reference RMSD.

262 **3.3 Receiver Operating Characteristics**

263 We examined the EWAC's ability to predict consumption exceeding 14 or 35 UK units/week.
264 The full areas under the receiver operating characteristic curves (AUC, Figure S7) are
265 presented along sensitivity and specificity at the best thresholds in Tables 3–4.

266 **14 UK units/week (increasing-risk):** EWAC increases the AUC by 5 percentage points
267 compared to the AUDIT-C score ($p<0.001$); and 7 percentage points compared to the full
268 AUDIT score ($p<0.001$). The cut-off maximising the sum of specificity and sensitivity on the

269 EWAC is 10 units/week. The sensitivity at this threshold is identical to AUDIT-C, but
270 specificity gains 13 percentage points. Using the nominal cut-off of 14 units/week on the
271 EWAC raises specificity to 0.922, at the cost of a reduction in sensitivity to 0.705 (Table 3).

272 **35 units/week (higher-risk):** EWAC provides small increases in AUC compared with the
273 AUDIT-C score ($p<0.001$) and the full AUDIT score ($p<0.001$). The best cut-off for detecting
274 consumption in excess of 35 units/week using the EWAC is 17 units/week (Table 4).

275 **3.4 Empirical distribution**

276 Table 5 estimates adult residents' total alcohol consumption in England using four different
277 sources, and compares them with alcohol retail sales. The HSE exhibits the highest
278 estimates and coverage of alcohol sales. EWAC amounts to 71% of the total consumption
279 estimated by the HSE's prospective diary, and 48% of retail sales.

280 Figure 5 suggests that the EWAC, like the ATS GF, estimates a greater prevalence of lower-
281 risk (≤ 14 units/week) and increasing-risk alcohol use than HSE.

282 **4 Discussion**

283 **4.1 Main findings**

284 We developed a continuous metric of estimated weekly alcohol consumption using the 3-
285 question Extended AUDIT-C. In a representative household survey, this metric estimated
286 usual alcohol consumption with a mean precision of ± 11 units/week and a median precision
287 of ± 2 units when compared with the GF reference standard.

288 **4.2 Strengths and limitations**

289 This paper is the first to (a) develop an EWAC using a well-accepted and validated alcohol
290 screening tool such as the AUDIT; and (b) quantify its bias and precision with respect to a
291 continuous measure of alcohol consumption. One study [38] previously reported mean
292 consumption by AUDIT-C score, but without quantifying bias or precision of such a measure.
293 Other studies have evaluated the AUDIT-C's accuracy in estimating alcohol consumption,
294 but only in relation to predicting consumption in excess of predefined thresholds [9]. Such

295 studies achieved AUCs ranging 0.83–0.96. Our study found the EWAC to be superior to both
296 the AUDIT-C and the full AUDIT in predicting GF exceeding 14 and 35 units/week. At the 14-
297 unit threshold, the specificity gain from 0.684 (AUDIT-C) to 0.876 places the EWAC with the
298 other best-performing diagnostic tools.

299 Our study provides strong confidence in the internal and external validity of findings in
300 England on account of the large sample size and extensive range of subgroup analyses
301 reported. Bias was mostly consistent across subgroups examined (age/sex, education,
302 smoking status, religion), with one exception. EWAC overestimated alcohol consumption by
303 2-3 UK units/week in Black/Other ethnic groups; though overall precision was better in
304 Black/Asian/Other ethnic groups than in White British respondents. Variation in the
305 sensitivity of AUDIT-C across ethnic groups has previously been noted in the US [39].

306 Repurposing a well-known tool such as the AUDIT-C has several advantages. It is already
307 translated in many languages and adapted to the varying standard drink sizes adopted
308 internationally [7]. The Extended AUDIT scores can be converted into traditional AUDIT
309 scores by capping items to 4, thereby offering a point of comparison with existing evidence.
310 The AUDIT's properties are also well understood in diverse contexts and modes of
311 administration, based on the last 30 years of international research. For instance, a previous
312 study which found the AUDIT-C to be responsive to changes of 70g/week [40] can suggest
313 that the EWAC's own responsiveness to change should be equivalent, if not greater than the
314 AUDIT-C's, given the Extended AUDIT-C's additional response items.

315 Despite this, the EWAC's design does not escape limitations shared by all methods of
316 screening or categorising alcohol use disorders. The conceptualisation of alcohol use
317 disorders is related to, but does not exclusively depend upon the amount of alcohol
318 consumed. Since Jellinek's description of 'the disease concept of alcoholism' [41] there have
319 been numerous attempts to categorise the range of phenotypes characterising alcohol use
320 disorders in the absence of any biomarker to 'verify' the presence of a particular pathology.
321 The EWAC, by limiting itself to an estimation of alcohol consumption is transparent across a

322 wide range of alcohol use disorders but does not measure the other factors underpinning
323 this complex and heterogeneous condition [6,42].

324 We note other study limitations. Our findings may not apply to other countries, or to
325 subpopulations with an atypical alcohol consumption (e.g. patients seeking care for severe
326 alcohol dependence, alcohol-related liver disease). Like the HSE, the ATS does not cover
327 populations excluded from most sampling frames (e.g. communal/carceral institutions,
328 homeless/no fixed abode populations).

329 We also note that the reference standard, as all self-reported measures, is imperfect. This
330 may introduce bias into estimates of precision: by definition, the reference standard's own
331 error will inflate the RMSD. In other words, it is likely that a proportion of the RMSD is
332 attributable to error in the GF measures rather than the EWAC.

333 Previous research offers reassurance that the EWAC's error is comparable to that of other
334 (even the more time-consuming) questionnaires. One telephone interview study [43]
335 measured a Pearson's correlation $r = 0.86$ and 0.87 in drinking frequencies and volumes
336 between a 12-month GF interview and a subsequent 28-day prospective diary. A postal
337 study [44] measured a correlation $r = 0.89$ between a 30-day GF and a prospective diary
338 (identical reference periods). Heeb *et al.* [22] measured a Kendall's rank correlation $\tau = 0.41$
339 between a 7-day GF and a 7-day prospective diary. For reference, our study measured a
340 correlation coefficient $r = 0.71$ and $\tau = 0.63$ between a 6-month EWAC and a 4-week GF
341 and suggests that EWAC's agreement with GF is similar to that of diaries.

342 **4.3 Potential applications**

343 Being equivalent or superior to the AUDIT in speed, accuracy, and international
344 standardisation, the EWAC appears suitable for use in clinical practice to support brief
345 interventions and to feed back a reliable 4-unit wide interval estimate of alcohol consumption
346 (eg: '9–13 units/week' or '70–100 g/week'). One such calculator is available online:

347 <https://ewac.netlify.app>

348 EWAC is already compatible with medical records information models developed in the
349 Systematised Nomenclature of Medicine Clinical Terms (SNOMED CT, Alcohol intake
350 (observable entity) [45]) and by the English Royal College of Physicians [46]. Such
351 information can have secondary uses as a variable in other disease risk scores, or to
352 prospectively recording of long-term alcohol exposure, an important risk factor for a range of
353 medical conditions.

354 Compared to an AUDIT-C score, alcohol consumption is a relatable scale, with health
355 education/promotion qualities and potential to support interventions focusing on recognising
356 the alcohol content and volume of drinks. Knowledge of alcohol beverage content is
357 generally poor [47], and many countries have not adopted a measure of standard drinks [48].
358 Two thirds of English drinkers could assess the standard drink equivalent of one alcohol unit
359 [49].

360 Assessment of alcohol consumption is not well embedded in clinical practice [50]. The
361 EWAC calculator fills a gap in resources empowering individuals to understand, monitor and
362 control their alcohol consumption with—or without—the involvement of healthcare
363 professionals. The EWAC appears particularly suitable for digital interventions.

364 Nutt, Rehm *et al.* [51,52] argued that alcohol-related harm is best prevented if individuals
365 know their consumption level, and health professionals in all settings can engage patients
366 effectively to manage risks with evidence-based interventions, in a similar way to other risk
367 factors for disease, for example blood pressure or cholesterol. The EWAC's dimensional
368 rather than categorical format may facilitate this while avoiding the stigma sometimes
369 associated with clinical categorisations of alcohol use disorders.

370 **Declarations**

371 **Acknowledgements**

372 This research was supported by the Medical Research Council [grant reference
373 MR/P016960/1]. The ATS data collection was funded primarily by the National Institute for
374 Health Research (NIHR) School for Public Health Research [grant reference
375 SPHR□SWP□ALC□WP5] and Public Health Research Programme [grant reference
376 15/63/01]. The EWAC online calculator development was funded by the Wessex Academic
377 Health Science Network. The views expressed are those of the author(s) and not necessarily
378 those of the NHS, the NIHR or the Department of Health.

379 **Ethics**

380 This study was approved by the University of Southampton's Faculty of Medicine Ethics
381 Committee (ERGO 44682).

382 **Consent for publication**

383 Not applicable.

384 **Competing interests**

385 JB has received unrestricted research funding to study smoking cessation from companies
386 who manufacture smoking cessation medications. PD, JH and JMAS declare no competing
387 interests.


388 References

- 389 [1] Shield K, Manthey J, Rylett M, Probst C, Wettlaufer A, Parry CDH, et al. National, regional, and global
390 burdens of disease from 2000 to 2016 attributable to alcohol use: a comparative risk assessment study.
391 *The Lancet Public Health* 2020;5:e51–61. doi:[10.1016/S2468-2667\(19\)30231-2](https://doi.org/10.1016/S2468-2667(19)30231-2).
- 392 [2] World Health Organisation. ICD-10: International statistical classification of diseases and related health
393 problems: Tenth revision. 2004.
- 394 [3] American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. Washington, DC:
395 American Psychiatric Association; 2013. doi:[10.1176/appi.books.9780890425596](https://doi.org/10.1176/appi.books.9780890425596).
- 396 [4] Health NI of, Excellence C. Alcohol-use disorders: prevention. Public health guideline [PH24]. 2010.
- 397 [5] National Institute of Health and Care Excellence. Alcohol-use disorders: diagnosis, assessment and
398 management of harmful drinking and alcohol dependence. Clinical guideline [CG115]. 2011.
- 399 [6] Saunders JB, Degenhardt L, Reed GM, Poznyak V. Alcohol Use Disorders in ICD-11: Past, Present, and
400 Future. *Alcoholism: Clinical and Experimental Research* 2019;43:1617–31. doi:[10.1111/acer.14128](https://doi.org/10.1111/acer.14128).
- 401 [7] Babor TF, Higgins-Biddle JC, Saunders JB, Monteiro MG. *The Alcohol Use Disorders Identification Test.*
402 *Guidelines for Use in Primary Care*. Second Edition. Geneva: World Health Organisation, Department of
403 Mental Health; Substance Dependence; 2001.
- 404 [8] Bush K. The AUDIT Alcohol Consumption Questions (AUDIT-C): An Effective Brief Screening Test for
405 Problem Drinking. *Archives of Internal Medicine* 1998;158:1789. doi:[10.1001/archinte.158.16.1789](https://doi.org/10.1001/archinte.158.16.1789).
- 406 [9] Meneses-Gaya C de, Zuardi AW, Loureiro SR, Crippa JAS. Alcohol Use Disorders Identification Test (AUDIT):
407 An updated systematic review of psychometric properties. *Psychology & Neuroscience* 2009;2:83–97.
408 doi:[10.3922/j.psns.2009.1.12](https://doi.org/10.3922/j.psns.2009.1.12).
- 409 [10] Nadkarni A, Garber A, Costa S, Wood S, Kumar S, MacKinnon N, et al. Auditing the AUDIT: A systematic
410 review of cut-off scores for the Alcohol Use Disorders Identification Test (AUDIT) in low- and middle-
411 income countries. *Drug and Alcohol Dependence* 2019;202:123–33.
412 doi:[10.1016/j.drugalcdep.2019.04.031](https://doi.org/10.1016/j.drugalcdep.2019.04.031).
- 413 [11] Johnson M, Jackson R, Guillaume L, Meier P, Goyder E. Barriers and facilitators to implementing screening
414 and brief intervention for alcohol misuse: a systematic review of qualitative evidence. *Journal of Public*
415 *Health (Oxford, England)* 2011;33:412–21. doi:[10.1093/pubmed/fdq095](https://doi.org/10.1093/pubmed/fdq095).
- 416 [12] Hutchings D, Cassidy P, Dallolio E, Pearson P, Heather N, Kaner E. Implementing screening and brief
417 alcohol interventions in primary care: Views from both sides of the consultation. *Primary Health Care*
418 *Research and Development* 2006;7:221–9. doi:[10.1191/1463423606pc292oa](https://doi.org/10.1191/1463423606pc292oa).
- 419 [13] McCormick KA, Cochran NE, Back AL, Merrill JO, Williams EC, Bradley KA. How Primary Care Providers
420 Talk to Patients About Alcohol. *Journal of General Internal Medicine* 2006;060721075157048.
421 doi:[10.1111/j.1525-1497.2006.00490.x](https://doi.org/10.1111/j.1525-1497.2006.00490.x).
- 422 [14] Beyer F, Lynch E, Kaner E. Brief Interventions in Primary Care: an Evidence Overview of Practitioner and
423 Digital Intervention Programmes. *Current Addiction Reports* 2018;5:265–73. doi:[10.1007/s40429-018-](https://doi.org/10.1007/s40429-018-0198-7)
424 [0198-7](https://doi.org/10.1007/s40429-018-0198-7).
- 425 [15] Kaner E, Bland M, Cassidy P, Coulton S, Dale V, Deluca P, et al. Effectiveness of screening and brief
426 alcohol intervention in primary care (SIPS trial): pragmatic cluster randomised controlled trial. *BMJ*
427 2013;346:e8501–1. doi:[10.1136/bmj.e8501](https://doi.org/10.1136/bmj.e8501).
- 428 [16] Crane D, Garnett C, Michie S, West R, Brown J. A smartphone app to reduce excessive alcohol
429 consumption: Identifying the effectiveness of intervention components in a factorial randomised control
430 trial. *Scientific Reports* 2018;8:4384. doi:[10.1038/s41598-018-22420-8](https://doi.org/10.1038/s41598-018-22420-8).
- 431 [17] Beard E, Brown J, West R, Acton C, Brennan A, Drummond C, et al. Protocol for a national monthly survey
432 of alcohol use in England with 6-month follow-up: 'The Alcohol Toolkit Study' Health behavior, health
433 promotion and society. *BMC Public Health* 2015;15. doi:[10.1186/s12889-015-1542-7](https://doi.org/10.1186/s12889-015-1542-7).
- 434 [18] Kaner EF, Beyer FR, Garnett C, Crane D, Brown J, Muirhead C, et al. Personalised digital interventions for
435 reducing hazardous and harmful alcohol consumption in community-dwelling populations. *Cochrane*
436 *Database of Systematic Reviews* 2017. doi:[10.1002/14651858.CD011479.pub2](https://doi.org/10.1002/14651858.CD011479.pub2).
- 437 [19] Michie S, Whittington C, Hamoudi Z, Zarnani F, Tober G, West R. Identification of behaviour change
438 techniques to reduce excessive alcohol consumption. *Addiction* 2012;107:1431–40. doi:[10.1111/j.1360-](https://doi.org/10.1111/j.1360-0443.2012.03845.x)
439 [0443.2012.03845.x](https://doi.org/10.1111/j.1360-0443.2012.03845.x).
- 440 [20] Lemmens P, Tan ES, Knibbe R a. Measuring quantity and frequency of drinking in a general population
441 survey: a comparison of five indices. *Journal of Studies on Alcohol* 1992;53:476–86.
442 doi:[10.15288/jsa.1992.53.476](https://doi.org/10.15288/jsa.1992.53.476).
- 443 [21] Greenfield TK. Ways of measuring drinking patterns and the difference they make: experience with
444 graduated frequencies. *Journal of Substance Abuse* 2000;12:33–49. doi:[10.1016/S0899-3289\(00\)00039-](https://doi.org/10.1016/S0899-3289(00)00039-0)
445 [0](https://doi.org/10.1016/S0899-3289(00)00039-0).
- 446 [22] Heeb J-L, Gmel G. Measuring alcohol consumption: A comparison of graduated frequency, quantity
447 frequency, and weekly recall diary methods in a general population survey. *Addictive Behaviors*
448 2005;30:403–13. doi:[10.1016/j.addbeh.2004.04.022](https://doi.org/10.1016/j.addbeh.2004.04.022).
- 449 [23] Livingston M, Callinan S. Underreporting in Alcohol Surveys: Whose Drinking Is Underestimated? *Journal of*
450 *Studies on Alcohol and Drugs* 2015;76:158–64. doi:[10.15288/jsad.2015.76.158](https://doi.org/10.15288/jsad.2015.76.158).
- 451 [24] Rehm J. Measuring Quantity, Frequency, and Volume of Drinking. *Alcoholism: Clinical and Experimental*
452 *Research* 1998;22:4s–14s. doi:[10.1111/j.1530-0277.1998.tb04368.x](https://doi.org/10.1111/j.1530-0277.1998.tb04368.x).

- 453 [25] Stockwell T, Donath S, Cooper-Stanbury M, Chikritzhs T, Catalano P, Mateo C. Under-reporting of alcohol
454 consumption in household surveys: A comparison of quantity-frequency, graduated-frequency and recent
455 recall. *Addiction* 2004;99:1024–33. doi:[10.1111/j.1360-0443.2004.00815.x](https://doi.org/10.1111/j.1360-0443.2004.00815.x).
- 456 [26] Stockwell T, Zhao J, Macdonald S. Who under-reports their alcohol consumption in telephone surveys and
457 by how much? An application of the 'yesterday method' in a national Canadian substance use survey.
458 *Addiction* (Abingdon, England) 2014;109:1657–66. doi:[10.1111/add.12609](https://doi.org/10.1111/add.12609).
- 459 [27] Stockwell T, Zhao J, Greenfield T, Li J, Livingston M, Meng Y. Estimating under- and over-reporting of
460 drinking in national surveys of alcohol consumption: identification of consistent biases across four
461 English-speaking countries. *Addiction* 2016;111:1203–13. doi:[10.1111/add.13373](https://doi.org/10.1111/add.13373).
- 462 [28] Public Health England. Alcohol sales in England in 2014: Analysis to assess suitability for inclusion as an
463 indicator in the Local Alcohol Profiles for England. 2017.
- 464 [29] NatCen Social Research, Royal Free and University College Medical School. Health Survey for England,
465 2011 [computer file]. Colchester, Essex: UK Data Archive [distributor], April 2013. SN: 7260. Colchester,
466 Essex: 2013. doi:[10.5255/UKDA-SN-7260-1](https://doi.org/10.5255/UKDA-SN-7260-1).
- 467 [30] Boniface S, Kneale J, Shelton N. Drinking pattern is more strongly associated with under-reporting of alcohol
468 consumption than socio-demographic factors: evidence from a mixed-methods study. *BMC Public Health*
469 2014;14:1297. doi:[10.1186/1471-2458-14-1297](https://doi.org/10.1186/1471-2458-14-1297).
- 470 [31] Dutey-Magni P, Sinclair J, Brown J. Concurrent validity of an Estimator of Weekly Alcohol Consumption
471 (EWAC) based on the Extended AUDIT 2018. doi:[10.17605/OSF.IO/7WE4M](https://doi.org/10.17605/OSF.IO/7WE4M).
- 472 [32] R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for
473 Statistical Computing; 2017.
- 474 [33] Wickham H. Tidyverse: Easily install and load the 'tidyverse' 2017.
- 475 [34] Stan Development Team. RStan: The R interface to Stan 2018.
- 476 [35] Dutey-Magni PF. Concurrent validity of an Estimator of Weekly Alcohol Consumption (EWAC) based on the
477 Extended AUDIT: Computer scripts 2020. doi:[10.5281/zenodo.4315024](https://doi.org/10.5281/zenodo.4315024).
- 478 [36] Lavoie D. Alcohol identification and brief advice in England: A major plank in alcohol harm reduction policy.
479 *Drug and Alcohol Review* 2010;29:608–11. doi:[10.1111/j.1465-3362.2010.00224.x](https://doi.org/10.1111/j.1465-3362.2010.00224.x).
- 480 [37] DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the Areas under Two or More Correlated Receiver
481 Operating Characteristic Curves: A Nonparametric Approach. *Biometrics* 1988;44:837.
482 doi:[10.2307/2531595](https://doi.org/10.2307/2531595).
- 483 [38] Rubinsky AD, Dawson DA, Williams EC, Kivlahan DR, Bradley KA. AUDIT-C Scores as a Scaled Marker of
484 Mean Daily Drinking, Alcohol Use Disorder Severity, and Probability of Alcohol Dependence in a U.S.
485 General Population Sample of Drinkers. *Alcoholism: Clinical and Experimental Research* 2013;37:1380–
486 90. doi:[10.1111/acer.12092](https://doi.org/10.1111/acer.12092).
- 487 [39] Frank D, DeBenedetti AF, Volk RJ, Williams EC, Kivlahan DR, Bradley KA. Effectiveness of the AUDIT-C as
488 a Screening Test for Alcohol Misuse in Three Race/Ethnic Groups. *Journal of General Internal Medicine*
489 2008;23:781–7. doi:[10.1007/s11606-008-0594-0](https://doi.org/10.1007/s11606-008-0594-0).
- 490 [40] Bradley KA, McDonnell MB, Bush K, Kivlahan DR, Diehr P, Fihn SD. The AUDIT Alcohol Consumption
491 Questions: Reliability, Validity, and Responsiveness to Change in Older Male Primary Care Patients.
492 *Alcoholism: Clinical and Experimental Research* 1998;22:1842–9. doi:[10.1111/j.1530-0277.1998.tb03991.x](https://doi.org/10.1111/j.1530-0277.1998.tb03991.x).
- 493 [41] Jellinek EM. The disease concept of alcoholism. New Haven: Hillhouse Press; 1960. doi:[10.1037/14090-000](https://doi.org/10.1037/14090-000).
- 494 [42] Leggio L, Kenna GA, Fenton M, Bonenfant E, Swift RM. Typologies of Alcohol Dependence. From Jellinek to
495 Genetics and Beyond. *Neuropsychology Review* 2009;19:115–29. doi:[10.1007/s11065-008-9080-z](https://doi.org/10.1007/s11065-008-9080-z).
- 496 [43] Greenfield TK, Kerr WC, Bond J, Ye Y, Stockwell T. Improving Graduated Frequencies Alcohol Measures for
497 Monitoring Consumption Patterns: Results from an Australian National Survey and a US Diary Validity
498 Study. *Contemporary Drug Problems* 2009;36:705–33. doi:[10.1177/009145090903600320](https://doi.org/10.1177/009145090903600320).
- 499 [44] Hilton ME. A comparison of a prospective diary and two summary recall techniques for recording alcohol
500 consumption. *British Journal of Addiction* 1989;84:1085–92. doi:[10.1111/j.1360-0443.1989.tb00792.x](https://doi.org/10.1111/j.1360-0443.1989.tb00792.x).
- 501 [45] UK Health and Social Care Information Centre. UK SNOMED CT Drug Extension, RF2: Full, Snapshot &
502 Delta 2018.
- 503 [46] Haroon S, Wooldridge D, Hoogewerf J, Mittal A, Bhala N, O'Donnell A, et al. Information standards for
504 recording alcohol use in electronic health records: Project report 2018.
- 505 [47] Mongan D, Long J. Standard drink measures throughout Europe; peoples' understanding of standard drinks.
506 RARHA: Joint Action on Reducing Alcohol Related Harm; 2015.
- 507 [48] Kalinowski A, Humphreys K. Governmental standard drink definitions and low-risk alcohol consumption
508 guidelines in 37 countries. *Addiction* 2016;111:1293–8. doi:[10.1111/add.13341](https://doi.org/10.1111/add.13341).
- 509 [49] UK Office for National Statistics. Drinking: adults' behaviour and knowledge in 2008. Opinions (Omnibus)
510 Survey Report No. 39. 2009.
- 511 [50] Brown J, West R, Angus C, Beard E, Brennan A, Drummond C, et al. Comparison of brief interventions in
512 primary care on smoking and excessive alcohol consumption: A population survey in England. *British*
513 *Journal of General Practice* 2016;66:e1–9. doi:[10.3399/bjgp16X683149](https://doi.org/10.3399/bjgp16X683149).
- 514 [51] Nutt DJ, Rehm J. Doing it by numbers: A simple approach to reducing the harms of alcohol. *Journal of*
515 *Psychopharmacology* 2014;28:3–7. doi:[10.1177/0269881113512038](https://doi.org/10.1177/0269881113512038).
- 516 [52] Rehm J, Anderson P, Manthey J, Shield KD, Struzzo P, Wojnar M, et al. Alcohol Use Disorders in Primary
517 Health Care: What Do We Know and Where Do We Go? *Alcohol and Alcoholism* 2016;51:422–7.
518 doi:[10.1093/alcalc/agg127](https://doi.org/10.1093/alcalc/agg127).
- 519

520

521 **Figures and tables**




522

523 **Figure 1.** Plots of EWAC against GF in (a) low/increasing risk ATS respondents

524 (n=21,338) and (b) all ATS respondents (n=22,373)


525



526

527 **Figure 2.** Forest plot of modelled MD for selected subgroups


It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



528

529 **Figure 3.** Forest plot of RMSD ratio (selected subgroups to reference category)


530



531

532 **Figure 4.** Forest plot of RMSD ratio (selected subgroups to reference category) in

533 respondents with a hazardous/harmful alcohol use (AUDIT-C>=5 or AUDIT>=8)



534

535 **Figure 5.** Empirical cumulative distribution function of weekly alcohol consumption in
536 England according to four alcohol schedules in residents aged 18 years and over

537

538 **Table 1.** Comparison of AUDIT-C and Extended AUDIT-C (questions 1 and 2)

Response items							
AUDIT-1: 'How often do you have a drink containing alcohol?'							
AUDIT-C	Never	Monthly or less	2 - 4 times per month	2 - 3 times per week	4+ times per week	-	
Extended AUDIT-C	Never	Monthly or less	2 - 4 times per month	2 - 3 times per week	4-5 times per week	6+ times per week	
AUDIT-2: 'How many units of alcohol do you drink on a typical day when you are drinking?'							
AUDIT-C	0 - 2	3 - 4	5 - 6	7 - 9	10+	-	
Extended AUDIT-C	0 - 2	3 - 4	5 - 6	7 - 9	10-12	13-15	16+

539

540

541 **Table 2.** Percentiles of the absolute deviation between EWAC and GF schedule (n =
542 22,404)

Absolute deviation percentiles	10%	20%	30%	40%	50%	60%	70%	80%	90%	95%	99%
	0.4	0.7	1.0	1.5	2.1	3.0	4.2	6.2	10.6	17.0	38.7

543

544 **Table 3.** Receiver operating characteristics of AUDIT-C score and EWAC for consumption
545 ≥ 14 UK units or 112g/week (n = 22,404)

Index test	AUC	95% CI	Best threshold	Sensitivity	Specificity
AUDIT-C score	0.871	[0.866, 0.876]	4.5	0.882	0.684
Full AUDIT score	0.854	[0.849, 0.860]	5.5	0.791	0.751
EWAC	0.921	[0.917, 0.925]	10	0.876	0.816

546 *Note:* The best threshold refers the cut-off value that maximises the sum of sensitivity and
547 specificity.
548

549

550 **Table 4.** Receiver operating characteristics of AUDIT-C score and EWAC for consumption
551 ≥ 35 UK units or 280g/week (n = 22,404)

Index test	AUC	95% CI	Best threshold	Sensitivity	Specificity
AUDIT-C score	0.913	[0.904, 0.921]	6.5	0.862	0.809
Full AUDIT score	0.901	[0.893, 0.909]	6.5	0.903	0.741
EWAC	0.936	[0.929, 0.943]	16.8	0.865	0.866

552 *Note:* The best threshold refers the cut-off value that maximises the sum of sensitivity and
553 specificity.
554

555

556 **Table 5.** Summary statistics on alcohol consumption in England in residents aged 18
557 years and over (excluding abstainers)

Study	Mean (UK units/week)	Median (UK units/week)	Variance	N	% of alcohol sold
HSE beverage-specific QF	14.0	7.3	474.6	6,545	72.6
HSE prospective diary	13.0	8.0	264.7	4,640	67.6
ATS GF	8.5	5.1	242.0	22,136	44.0
ATS EWAC	9.3	5.2	148.9	25,882	48.0
Retail sales	19.3	–	–	–	–

558