The COVID University Challenge: a Hazard Analysis of Critical Control Points

Assessment of the Return of Students to Higher Education Establishments

Kelly L. Edmunds[1], Laura Bowater[2], Julii Brainard[2], Jean-Charles de Coriolis[1], Iain Lake[3], Rimsha R. Malik[2], Lorraine Newark[4], Neil Ward[5], Kay Yeoman[1], Paul R. Hunter[2]

[1]Centre for Ecology, Evolution and Conservation, School of Biological Sciences, University of East Anglia, Norwich, NR4 7TJ, UK.

[2]Norwich Medical School, University of East Anglia, Norwich, NR4 7TJ, UK.

[3]School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK.

[4]Learning and Teaching Service, University of East Anglia, Norwich, NR4 7TJ, UK.

[5]Vice Chancellor's Office, University of East Anglia, Norwich, NR4 7TJ, UK.

Address for Corresponding author: Professor Paul Hunter, Norwich Medical School, University of East Anglia, Norwich, NR4 7TJ, UK. Tel: +44 (0)1603 456161

email: paul.hunter@uea.ac.uk

Acknowledgements: Professors Hunter and Lake as well as Dr. Brainard to the National Institute for Health Research Health Protection Research Unit (NIHR HPRU) in Emergency

Preparedness and Response at King's College London in partnership with Public Health England (PHE) and the University of East Anglia. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, UEA, the Department of Health or Public Health England. The authors are grateful to Jim Hunter of the University of East Anglia Safety Services staff for his valuable input and thank all those who have provided comment at any stage of the HACCP assessment or on this manuscript. As this study did not directly involve human subjects or data about individuals, ethical approval was not required.

ABSTRACT

The COVID-19 pandemic disrupted economies and societies throughout the world in 2020. Education was especially affected, with schools and universities widely closed for long periods. People under 25 years have the lowest risk of severe disease but their activities can be key to persistent ongoing community transmission. A challenge arose for how to provide education, including university level, without the activities of students increasing wider community SARS-CoV-2 infections.

We used a Hazard Analysis of Critical Control Points (HACCP) framework to assess the risks associated with university student activity and recommend how to mitigate these risks. This tool appealed because it relies on multi-agency collaboration and interdisciplinary expertise and yet is low cost, allowing rapid generation of evidence-based recommendations.

We identified key critical control points associated with university student' activities, lifestyle and interaction patterns both on-and-off campus. Unacceptable contact thresholds and the most up-to-date guidance were used to identify levels of risk for potential SARS-CoV-2 transmission, as well as recommendations based on existing research and emerging evidence for strategies that can reduce the risks of transmission. Employing the preventative measures we suggest can reduce the risks of SARS-CoV-2 transmission among and from university students. Reduction of infectious disease transmission in this demographic will reduce overall community transmission, lower demands on health services and reduce risk of harm to clinically vulnerable individuals while allowing vital education activity to continue. HACCP assessment proved a flexible tool for risk analysis in a specific setting in response to an emerging infectious disease threat.

Keywords: COVID-19, SARS-CoV-2, HACCP, rapid response, university, education

1. INTRODUCTION

Emerging infectious diseases present substantial threats to global health[1]. The current outbreak of SARS-CoV-2 that causes COVID-19 disease caught countries unaware and presented unprecedented challenges. To date the pandemic is linked to over 57.8 million reported cases and 1.3 million deaths (as at 22/11/20; WHO, 2020a). The long-term socio-economic harms of the pandemic are forecast with alarming numbers across multiple sectors including employment, global GDP, national debt and public service provision (World Bank, 2020). It is people under 25 years that are least in danger from the virus clinically (Russell et al., 2020), but this group arguably will shoulder the greatest long-term burden that result from for the current efforts to control the virus in-order to reduce societal harms. These dis-benefits/disadvantages include reduced education, reduced employment opportunities, living longer with degraded quality public services that in turn includes education that is likely to be highly resource-restricted in the near future (Maher, Hoang, & Hindery, 2020).

Early analyses suggests that closures of educational facilities had helped to reduce the outbreak in individual countries (Davies et al., 2020; Hunter et al., 2020; Li et al., 2020), through a variety of mechanisms. Although it remains unclear exactly how likely young children are to be infected by or transmit SARS-CoV-2, educational establishments contain adults that are interacting, collaborating and mixing. Also, young children are typically escorted to schools by adults who encounter other adults en route. Secondary and tertiary students often travel to school using public transport interacting highly with both each other and other adults en route and on site.

The prospect of shutting down educational facilities for long durations during the pandemic was not viable because the resultant unacceptable harm to both individuals and the wider economy was highly likely. For the age range 18-24 there was an additional consideration: a University College London survey found that loneliness during early lockdown was common in the age demographic (Leavey, Eastaugh, & Kane, 2020), an age when counts of social contacts normally peak rather than contract (Mossong et al., 2008). Full time home-study could increase loneliness and was often physically hampered by practical problems: many young adults live in rented accommodation without adequate quiet and dedicated space at home to study effectively (Leavey, Eastaugh, & Kane, 2020). Good internet access from home, or access to a PC/laptop, could also not be assured for all even in high income countries (Auxier & Anderson, 2020; Wakefield 2020). For individuals, a decision to suspend education/training and enter the labour market temporarily was not appealing as unemployment rates rose sharply (CTV News, 2020; Ergocun, 2020; Heath, 2020). For those students starting the new academic year in September 2020, approximately a third would have been enrolling in their first year of university study, which would bring with it additional unknowns.

A societal problem with the prospect of suspending university education for an uncertain but long duration was the medium-long term effects on professional training. Most high income countries recruit each year large numbers of newly trained graduates into a wide range of key worker professions, including nursing, medicine, allied health professions, social care, engineering, science and teaching. Suspending university education would increase existing shortages of essential degree-qualified workers across multiple sectors (Garcia & Weiss, 2019; Gilroy, 2019). Replacement recruitment of professionals from other parts of the world was

unlikely given long training times required and pandemic disruption of education everywhere. It was also apparent that some universities faced severe financial crisis if students did not return to conventional tertiary education provision environment (Drayton & Waltmann, 2020); contraction of the tertiary education sector is not desirable as pandemic conditions demonstrated the value of degree trained professionals for response and infectious disease control.

Ensuring the continuation of tertiary education during the pandemic was imperative. Some degree courses can be effectively delivered in a tele-learning (online) format, but many courses still depend on practical work in laboratories, using university equipment as well as placement opportunities in external facilities that deliver in-person training. This reliance on specific facilities and types of training is especially true for vocational courses such as medicine, nursing, midwifery, physiotherapy is a challenge faced by our own institution which normally graduates hundreds of qualified health professionals each year. We were also aware that many non-health care students in our community, studying entirely online university degrees, lived on or near to campus and might still rely on other types of community locations (field work) or university facilities (such as the library or laboratories).

Therefore, the aim of this research was to identify the critical control points among the usual and necessary activities of university students – points where the risks of transmission were greatest but could also be mitigated and reduced –to make recommendations for mitigating these risks. The approach for this research involved the use of the Hazard Analysis of Critical Control Points (HACCP) framework; a low-cost tool for rapid assessment, that is simple for stakeholders to

engage in, encourages multi-agency collaboration and interdisciplinary expertise whilst generating evidence-based recommendations (Edmunds et al., 2013). HACCP assessments originated in risk management of food production systems, but have been increasingly used to assess risks related to an emerging health threats and infectious diseases (Edmunds et al., 2013; Edmunds et al., 2016).

We applied the HACCP approach to the activities of university students and propose this method as a way to generate recommendations when dealing with a rapidly evolving infectious disease outbreak. We identify behaviours and practices linked to typical student activities which are likely to present risks for the transmission, both direct and indirect, of SARS-CoV-2. HACCP assessments are potentially powerful methods that can be used as rapid response tools in specific settings during emerging infectious disease outbreaks, as a precursor to more time-consuming and costly quantitative data collection, biomedical testing and clinical studies in specific settings.

2. METHODS

To complete our assessment of the risks of SARS-CoV-2 transmission during the return of students to UK university settings during the COVID-19 pandemic, we followed the first three HACCP principles (as used in previous studies into EIDs; Edmunds et al., 2013; Edmunds et al., 2016). The first stage of the process involved a series of virtual workshops and meetings amongst the HACCP team; all of whom are based at the higher education setting of the University of East Anglia (UEA) in Norwich, UK; a campus-based university with a student population of 17,429 in 2019/20 (of which 8,067 (46.3%) were new starters) (UEA 2020). The HACCP team includes experts in the fields of microbiology, infectious diseases, EIDs, public

health, environmental health, science communication, student safety and student administrative services. The first workshop and subsequent follow-up discussions led to the creation of early versions of six flowcharts which consider different scenarios for student interactions on campus. These were i) on-campus teaching; ii) off-campus teaching; iii) living on campus; iv) living off-campus; v) non-teaching on-campus activities and vi) social activities. Following further consultation, the flowcharts were then critically analysed by the team and finalised, simplified versions produced. A hazard was considered to be a process that could take place within a higher education university setting that would provide the opportunity, at an unacceptable level of risk, for the transmission of SARS-CoV-2 to another person. Taking into account the likelihood of a situation occurring and the likelihood of SARS-CoV-2 transmission being able to occur within that situation, the hazards were then grouped into high, medium and low-risk categories. We generated recommendations for high-risk critical control points. The focus on high-risk CCPs is because our key objective was to make a set of recommendations that would do most to reduce transmission of SARS CoV-2 without being too onerous and likely to be followed. In concurrent circumstances (late 2020), an eradication strategy was not feasible in the UK. However, effective suppression of wide transmission to strongly limit harms from COVID-19 was stated in government policy and supported by most institutions and public experts (BBC, 2020; UK Government, 2020a). Therefore, in this HACCP exercise, we focused our recommendations on only situations which generate the majority of transmissions ("high risk"), but this HACCP exercise did not address situations where likelihood of transmission was low or medium. By focusing on high risk contexts, we sought to provide recommendations that would have maximum effect in keeping both the university and wider community prevalence of the infections at relatively low and thus manageable levels.

Following validation of the flowchart, the same team of experts focused on all of the high risk hazards and determined appropriate Critical Control Points (CCPs). A CCP is a point at which there is the opportunity to control, prevent or eliminate the risks for pathogen transmission. Critical limits were then set for each of the CCPs identified. These critical limits are thresholds used as preventative measures to control the hazards within the system. The outputs of each of these three principles were cross-referenced with the existing literature on SARS-CoV-2 epidemiology, prevention and control.

3. RESULTS

3.1 Hazard Analysis

Our HACCP assessment of the risks of SARS-CoV-2 transmission for students in UK university settings during the COVID-19 pandemic identified high-risk potentially hazardous practices linked to social interactions, teaching sessions and day-to-day on-campus activities (Table 1). Primary concerns were shared use of social spaces such as sports changing rooms, prayer facilities, study facilities including teaching rooms and rooms for the examination of patients for those students on medical and health science courses. When considering the sharing of student residences (both on and off campus), all those ordinarily living within the student residence were considered to come from the same household. Thus, social distancing and other protective measures such as the use of face coverings during interactions between them was not deemed necessary, unless one or more members of the household were presenting with symptoms of COVID-19 or had received a positive test result in the previous 14 days. All other activities and practices linked to students at higher education establishments were deemed to present a medium or low level of risk for the transmission of SARS CoV-2.

3.2 Critical Control Points and Critical Limits

In total, 23 high-risk CCPs were identified by this HACCP assessment. Each CCP is a point at which there is an opportunity to adopt measures to reduce the risks of transmission of SARS CoV-2 during the return of students to UK university settings. Following extensive consultation and cross-referencing with the existing literature, critical limits were then proposed for each high-risk CCP. In many cases, multiple recommendations were made for each CCP (Table 1).

Several of the critical limits suggested include fundamental infection control practices for coronaviruses, namely the correct and proper use of face coverings, employing appropriate hand hygiene measures and adopting appropriate social distancing measures, such as concurrently recommended physical distancing in the UK at a minimum of 1m+.

4 DISCUSSION

The HACCP framework allows for a rapid identification of the risks associated to a specific setting and evolving disease outbreak, in this case SARS-CoV-2 transmission to, from and between university students. The COVID-19 pandemic has highly disrupted education at all levels. It seems clear that shutting educational facilities can reduce community prevalence of SARS-CoV-2 but the harms long-term closure would cause to individuals and wider society are unacceptable. Therefore, strategies need to be found that enable most education delivery to continue, including forms of training that require physical use of education facilities.

No degree of pandemic preparedness can ever be fully comprehensive. This is because the nature of microbial contagions is that they tend to have unique features specific to the

pathogenicity of the disease. The case of SARs-CoV-2 infection is distinctive because it is mostly harmless to a key demographic (young people) while COVID-19. However, there are many aspects of effective COVID-19 control that require tremendous disruptions to the lives of young people. Our HACCP assessment used a structured methodology to evaluate opportunities for mitigating risks of SARS-CoV-2 transmission associated with usual activities of university students, in an effort to facilitate their continued and essential education and training.

It seems clear that a proportion of SARS-CoV-2 transmission is from infected persons who lack symptoms: either pre-symptomatic and probably completely asymptomatic infections (Gandhi, Yokoe, & Havlir, 2020). The diversity of symptoms experienced is also large: COVID-19 is not simply an influenza-like illness with fever and cough as diagnostic symptoms (Sahoo, Jhunjhunwala, & Jolly, 2020). This situation presents unique challenges in terms of identifying cases and preventing them from passing the virus on. In peak prevalence periods, infection control relies on assuming that oneself and all other persons are potentially infected and therefore the only way to limit spread is through dramatically decreased contact with all other persons – even those with no contact or recent symptom history (Menon, 2020).

Our results feature recurring mitigation measures due to understanding of common disease transmission pathways and dynamics. Physical distancing from others (at least 2 metres) is believed to be especially effective because the predominant transmission pathway is via saliva droplets and possibly aerosols (WHO, 2020b). There is mixed quality evidence about the utility of wearing masks or face coverings to prevent COVID-19 in individuals (Brainard et al., 2020). In our own systematic review we concluded that wearing masks had some benefit for reducing transmission both when an infected person wore the mask and when a currently uninfected

person wears one. However, we did not consider that mask wearing provided a guarantee of protection and was certainly not an alternative to other social distancing measures. The overall benefits of requiring populations to wear face coverings on wider community prevalence and subsequent mortality (Lyu & Wehby, 2020) is still not absolutely clear (Hunter et al., 2020).

High ventilation rates and outdoor settings are believed to be linked to lower risk of transmission (Bhagat et al., 2020). Good hand hygiene is basic infection control practice, as are measures to reduce the chances of fomite transmission (by sanitising any potentially shared objects).   How to follow these principles in avoiding infection depends somewhat on setting and activity, such as meetings, religious worship, returns from family visits, queuing, interactions with household members, etc. We acknowledge in Table 1 that guidance and understanding of best infection control practices regarding COVID-19 are still emerging and recommend that students consult local guidance as it is published. Each transmission opportunity can be minimised or even eliminated through careful implementation of the critical limits outlined in Table 1.

At our own university, the response to COVID-19 has had three main components, covering communications, social distancing and testing and monitoring. Regular communications were sent to all students about COVID-19 safety measures and the University's expectations around responsible behaviour. These were reinforced by enhanced security and monitoring of students on-campus, in accommodation and in other social settings, to discourage and prevent social gatherings that span households. Hand sanitiser and associated signage provided in teaching rooms and at key points in communal areas helped maintain a visible reminder of COVID-19 risk. To help ensure adequate distancing, one-way systems for navigating corridors within buildings and the narrower pathways outside on campus were introduced. In addition, a

precautionary approach to 2m distancing in teaching rooms was pursued, based on shoulder-to-shoulder measurement and assuming some potential movement of torsos while sitting at desks (teaching room capacities were between 10 and 20% of normal).

At the start of autumn term, the University introduced an asymptomatic testing scheme in partnership with the Earlham Institute, a genomics research laboratory also located on the Norwich Research Park (Berger Gillam et al., 2020). The 'Norwich Testing Initiative' had capacity to analyse 500 PCR tests per day. All members of the university community (students and staff) who regularly visited campus or lived there were encouraged to get tested. Positive case students and their households were isolated for 10 days with non-symptomatic, non-positive housemates isolating for 14 days, with regular welfare support for those living off campus. There was daily monitoring of case numbers and patterns by University Safety Services and members of the University's Executive Team, through a daily Major Operations Group meeting, to review trends and patterns and adjust the University's response as necessary.

After students arrived back at UEA at the start of autumn term, the number of actively positive student cases being managed peaked at just 69 after 3 weeks, but fell to below 25 by week 6. There were just 2 cases that tested positive in the previous 10 days, by 9[th] December. Of 360 cases in total detected over the first 11-weeks of monitoring, just three were members of staff involved in teaching. This suggests that the transmission reduction measures taken to prevent spread in teaching spaces were successful.

5 CONCLUSIONS

The COVID pandemic has delivered many epidemiological lessons and revealed numerous shortcomings in both local and international responses to emerging infectious diseases (EID). In the early stages of responding to an EID outbreak, the use of a response tool such as the HACCP framework allows for the rapid identification of risks and hazardous behaviours. Better risk reduction strategies can be generated after implementation of in-depth epidemiological studies, but such studies can take months or years to complete and produce results and recommendations. In contrast, the HACCP framework can be implemented fairly quickly using diverse informants and stakeholders. The long term consequences of the emergence of COVID-19 are proving difficult to predict, but we can still do much to limit pandemic harms. There is considerable merit in rapid responses methods to minimise harms linked to future disease outbreaks.

References

Auxier, B. & M. Anderson, (2020). As schools close due to the coronavirus, some U.S. students face a digital 'homework gap'. *Pew Research Center,* 16 March 2020, available at https://www.pewresearch.org/fact-tank/2020/03/16/as-schools-close-due-to-the-coronavirus-some-u-s-students-face-a-digital-homework-gap/

BBC. (2020). Coronavirus: WHO head calls herd immunity approach 'immoral'. *BBC*, 13

October 2020, available at https://www.bbc.co.uk/news/world-54518286

Bhagat, R. K., Wykes, M. D., Dalziel, S. B., & Linden, P. F. (2020). Effects of ventilation on the

indoor spread of COVID-19. *Journal of Fluid Mechanics*, 903, F1, doi:10.1017/jfm.2020.720.

Brainard J.S., Jones, N., Lake, I., Hooper, L and Hunter, P. (2020). Community use of facemasks

and similar barriers to prevent respiratory illness such as COVID-19: A rapid scoping

review. *Eurosurveillance,* In Press.

Centers for Disease Control and Prevention. (2020a). Keeping the workplace safe, available at:

https://www.cdc.gov/coronavirus/2019-ncov/downloads/workplace-school-and-home-

guidance.pdf

Centers for Disease Control and Prevention. (2020b). Handwashing: Clean Hands Save Lives.

When and How To Wash Your Hands, 24 November 2020, available at:

https://www.cdc.gov/handwashing/when-how-handwashing.html

Centers for Disease Control and Prevention. (2020c). Cleaning and Disinfection for Households, 10 July 2020, available at: https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cleaning-disinfection.html

CTV News, (2020). Canada's jobless rate soars to 13 per cent in April. *CTV News*, 08 May 2020, available at https://www.ctvnews.ca/business/canada-s-jobless-rate-soars-to-13-per-cent-in-april-1.4930397

Davies, N. G., Kucharski, A. J., Eggo, R. M., Gimma, A., & Edmunds, W. J., on behalf of the Centre for the Mathematical Modelling of Infectious Diseases COVID-19 working group. (2020). Effects of non-pharmaceutical interventions on COVID-19 cases, deaths, and demand for hospital services in the UK: a modelling study. *The Lancet Public Health,* 5(7), E375-E385, doi.org/10.1016/S2468-S2667(20)30133-X.

Drayton, E & B. Waltmann. (2020). Will universities need a bailout to survive the COVID-19 crisis? *Institute for Fiscal Studies*, 06 July 2020, available at https://www.ifs.org.uk/publications/14919

Edmunds, K. L., Hunter, P.R., Few, R., Bell, D.J. (2013). Hazard Analysis of Critical Control Points Assessment as a Tool to Respond to Emerging Infectious Disease Outbreaks. *PLoS ONE* 8(8): e72279. doi:10.1371/journal.pone.0072279.

Edmunds K.L., Abd Elrahman, S., Bell, D.J., Brainard, J., Dervisevic, S., Fedha, T.P., Few, R., Howard, G., Lake, I., Maes, P., Matofari, J., Minnigh, H., Mohamedani, A. A., Montgomery, M., Morter, S., Muchiri, E., Mudau, L.S., Mutua, B.M., Ndambuki, J.M., Pond, K., Sobsey, M.D., van der Es, M., Zeitoun, M & Hunter, P. (2016). Dealing with Ebola-infected waste: a Hazard Analysis of Critical Control Points for reducing the risks to public health. *Bulletin of the World Health Organization*, 94:424-432. doi: http://dx.doi.org/10.2471/BLT.15.163931

Ergocun, G. (2020). EU unemployment in August rises to 7.4%. *Anadolu Agency*, 01 October 2020, available at https://www.aa.com.tr/en/economy/eu-unemployment-in-august-rises-to-74-/1992138

European Centre for Disease Prevention and Control. (2020). Contact tracing: Public health management of person, including healthcare workers, having had contact with COVID-19 cases in the European Union. ECDC, Stockholm, available at: https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-public-health-management-contact-novel-coronavirus-cases-EU.pdf

Gandhi, M., Yokoe, D. S. & Havlir, D. V. (2020). Asymptomatic transmission, the Achilles' heel of current strategies to control COVID-19. *New England Journal of Medicine,* doi.10.1056/NEJMe2009758

Garcia, E & E. Weiss. (2019). The teacher shortage is real, large and growing, and worse than we thought. *Economic Policy Institute*, 26 March 2019, available at https://www.epi.org/publication/the-teacher-shortage-is-real-large-and-growing-and-worse-than-we-thought-the-first-report-in-the-perfect-storm-in-the-teacher-labor-market-series/

Gillam, T. B., Cole, J., Gharbi, K., Angiolini, E., Barker, T., Bickerton, P., Brabbs, T., Chin, J., Coen, E., Cossey, S., Davey, R., Davidson, R., Durrant, A., Edwards, D., Hall, N., Henderson, S., Hitchcock, M., Irish, N., Lipscombe, J., Jones, G., Parr, G., Rushworth, S., Shearer, N., Smith, R and Steel, N. (2020). Norwich COVID-19 testing initiative pilot: evaluating the feasibility of asymptomatic testing on a university campus. *Journal of Public Health* (Oxf), published online 30 Oct 2020, doi: 10.1093/pubmed/fdaa194

Gilroy, R. (2019). NHS workforce being 'hollowed out' by registered nurse shortages. *Nursing Times*, 28 November 2019, available at https://www.nursingtimes.net/news/workforce/nhs-workforce-being-hollowed-out-by-registered-nurse-shortages-28-11-2019/

Global Heat Health Information Network, (2020). Do air conditioning and ventilation systems increase the risk of virus transmission? If so, how can this be managed? Available at: http://www.ghhin.org/heat-and-covid-19/ac-and-ventilation

Heath, M. (2020). Australian Unemployment Soars on Record Back-to-Back Job Losses. *Bloomberg*, 18 June 2020, available at https://www.bloomberg.com/news/articles/2020-06-18/australian-unemployment-jumps-to-7-1-in-may-currency-declines

Hunter, P. R., Colon-Gonzalez, F., Brainard, J. S & Rushton, S. (2020). Impact of non-pharmaceutical interventions against COVID-19 in Europe: a quasi-experimental study. *MedRxiv,* preprint doi.org/10/1101/2020.05.01.20088260

Leavey, C., Eastaugh, A. & Kane, M. (2020). Generation COVID-19. Building the case to protect young people's future health. *The Health Foundation*, 30 August 2020, available at https://www.health.org.uk/publications/long-reads/generation-covid-19

Li, Y., Campbell, H., Kulkarni, D., Harpur, A., Nundy, M., Wang, X., Nair, H. for the Usher Network for COVID-19 Evidence Reviews (UNCOVER) group. (2020). The temporal association of introducing and lifting non-pharmaceutical interventions with the time-varying

reproduction number (R) of SARS-CoV-2: a modelling study across 131 countries. *The Lancet Infectious Diseases*, doi.org/10.1016/S1473-3099(20)30785-4

Lyu, W & Wehby, G. L. (2020) Community Use Of Face Masks And COVID-19: Evidence From A Natural Experiment Of State Mandates In The US: Study examines impact on COVID-19 growth rates associated with state government mandates requiring face mask use in public. *Health Affairs*, 39(8):1419-25.

Macias, A. E., de la Torre, A., Moreni-Espinosa, S., Leal, P. E., Bourlon, M. T. & Ruiz-Palacios G. M. (2010) Controlling the novel A (H1N1) influenza virus: don't touch your face! *The Journal of Hospital Infection,* 73(3), 280-281, doi.org/10.1016/j.jhin.2009.06.017.

Maher, C.S., Hoang, T. & Hindery, A. (2020). Fiscal Responses to COVID□19: Evidence from Local Governments and Nonprofits. *Public Administration Review*, 80(4), 644-650, doi.org/10.1111/puar.13238

Menon, P. (2020). 'Act like you have COVID-19': PM Arden says as New Zealand heads into lockdown, *Reuters World News*, 25 March 2020, available at https://uk.reuters.com/article/uk-health-coronavirus-newzealand-emergen/act-like-you-have-covid-19-pm-ardern-says-as-new-zealand-heads-into-lockdown-idUKKBN21C065

Morawska, L. & D. K. Milton. (2020). It is time to address airborne transmission of Coronavirus

Disease 2019 (COVID-19). *Clinical Infectious Diseases*, 71(9), 2311-2313,

doi.org/10.1093/cid/ciaa939

Mossong, J., Hens, N., Jit, M., Beutels, P., Auranen, K., Mikolajczyk, R., Massari, M., Salmaso,

S., Tomba, G. S., Wallinga, J., Heijne, J., Sadkowska-Todys, M., Rosinska, M. & Edmunds, W.

J. (2008). Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS

Med*. 5(3):e74, doi:10.1371/journal.pmed.0050074

Nielsen, B. F., & Sneppen, K. (2020). Superspreaders provide essential clues for mitigation of

COVID-19. *MedRxiv*, preprint doi.org/10/1101/2020/09/15/20195008

Qian, H., Miao, T., Liu, L., Zheng, X., Luo, D., & Li, Y. (2020). Indoor transmission of SARS-

CoV-2. *MedRxiv,* pre-print doi.org/10.1101/2020.04.04.20053058

Russell, T. W., Hellewell, J., Jarvis, C. I., Van Zandvoort, K., Abbott, S., Ratnayake, R.,

CMMID COVID-19 working group, Flasche, S., Eggo, R. M., Edmunds, W. J. & Kucharski A.

J. (2020). Estimating the infection and case fatality ratio for coronavirus disease (COVID-19)

using age-adjusted data from the outbreak on the Diamond Princess cruise ship, February 2020. *Eurosurveillance*, *25*(12), doi.org.10.2807/1560-7917.ES.2020.25.12.2000256.

Sahoo, S., Jhunjhunwala, S. & Jolly, M. K. (2020). The Good, The Bad and The Ugly: A Mathematical Model Investigates the Differing Outcomes Among CoVID-19 Patients. *Journal of the Indian Institute for Science*, doi.org/10/1007/s41745-020-00205-1

UK Government. (2020a). A joint statement on coronavirus (COVID-19) from the UK Government, the Northern Ireland Executive, the Scottish Government, and the Welsh Government. *Cabinet Office*, UK Government, 25 September 2020, available at https://www.gov.uk/government/publications/joint-statement-on-covid-19/joint-statement-on-coronavirus-covid-19

UK Government. (2020b). Celebrating religious festivals during coronavirus (COVID-19). *Ministry of Housing, Communities & Local Government*, UK Government. 10 December 2020, available at https://www.gov.uk/government/publications/covid-19-guidance-for-the-safe-use-of-places-of-worship-during-the-pandemic-from-4-july/celebrating-religious-festivals-during-coronavirus-covid-19

UK Government. (2020c). Coronavirus (COVID-19): safer travel guidance for passengers. *Department for Transport*, UK Government, 9 December 2020, available at https://www.gov.uk/guidance/coronavirus-covid-19-safer-travel-guidance-for-passengers

UK Government. (2020d). Stay at home: guidance for households with possible or confirmed coronavirus (COVID-19) infection. Coronavirus (COVID-19): safer travel guidance for passengers. *Public Health England*, UK Government, 5 November 2020, available at https://www.gov.uk/government/publications/covid-19-stay-at-home-guidance/stay-at-home-guidance-for-households-with-possible-coronavirus-covid-19-infection

UK Government. (2020e). Local restriction tiers: what you need to know. *Department of Health and Social Care*, UK Government, 30 November 2020, available at https://www.gov.uk/guidance/local-restriction-tiers-what-you-need-to-know

UK Government. (2020f). Keeping workers and clients safe during COVID-19 in close contact services. *HM* Government. 5 November 2020, available at https://assets.publishing.service.gov.uk/media/5ef2889986650c12970e9b57/working-safely-during-covid-19-close-contact-041120.pdf

University of East Anglia. (2020). UEA Student Facts and Figures 2019/20. Available at: https://www.uea.ac.uk/documents/37663/2822180/Facts+and+Figures+Jan+2020.pdf/b97dfeec-5306-4490-8e07-1c77a533897b?t=1595321923736, last updated 20/01/20.

van Doremalen, N., Bushmaker, T., Morris, D. H., Holbrook, M. G., Gamble, A., Williamson, B. N., Tamin, A., Harcourt, J. L., Thornburg, N. J., Gerber, S. I., Lloyd-Smith, J. O., de Wit, E., Munster, V. J. (2020). Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of Medicine*, 382:1564-1567, doi.org/10.1056/NEJMc2004973

Wakefield, J. (2020). Coronavirus: Coping with lockdown on poor broadband. *BBC News*, 02 May 2020, available at https://www.bbc.co.uk/news/technology-52414280

Wikipedia. (2020) Arm span. Available at: https://en.wikipedia.org/wiki/Arm_span

World Bank. 2020. Global Economic Prospects, June 2020. Washington, DC: World Bank. © World Bank. https://openknowledge.worldbank.org/handle/10986/33748, License: CC BY 3.0 IGO.

World Health Organisation. (2020a). *COVID-19 Weekly Epidemiological Update*. Updated 22

November 2020, available at https://www.who.int/publications/m/item/weekly-epidemiological-

update---24-november-2020

World Health Organisation. (2020b). Transmission of SARS-CoV-2: implications for infection

prevention precautions. *Scientific Brief, World Health Organisation,* 09 July 2020, available at

https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-

for-infection-prevention-precautions

World Health Organisation. (2020c). Coronavirus disease (COVID-19): Ventilation and air

conditioning in health facilities. *World Health Organisation,* 29 July 2020, available at

https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-ventilation-and-air-

conditioning-in-health-facilities

World Health Organisation. (2020d). Advice on the use of masks in the context of COVID-19.

World Health Organisation, 5 June 2020, WHO reference number: WHO/2019-

nCov/IPC_Masks/2020.4, available at https://apps.who.int/iris/rest/bitstreams/1279750/retrieve