1   The long-term success of mandatory vaccination laws at

2   implementing world's first vaccination campaign in rural Finland

3   Susanna Ukonaho[1], Virpi Lummaa[1]*, Michael Briga[1]*

[1] Department of Biology, University of Turku, Finland

*co-senior authors

Short title: Vaccination law effectiveness

Abstract:

Infectious diseases are a major public health concern and socio-economic burden at all ages,

but children are by far the most vulnerable to infections. In high income countries, infectious

diseases are on the rise, a phenomenon in part attributed to the recent surge of vaccination

hesitancy. To combat vaccination hesitancy, several countries recently made vaccinating

children mandatory, but the effectiveness of such vaccination laws in increasing the vaccination

coverage remains debated and the long-term consequences are unknown. Here we quantified

the consequences of vaccination laws on the vaccination coverage monitoring for a period of

63 years world's first vaccination campaign against the highly lethal childhood infection

smallpox in rural Finland. We found that annual vaccination campaigns were focussed on

children up to 1 year old, but that their vaccination coverage was low and declined with time

until the start of the vaccination law, which stopped the declining trend and was associated

with an abrupt coverage increase of 20 % to cover >80 % of all children. The spatio-temporal

variance in vaccination coverage decreased with time but this was not directly associated with

the vaccination law. Our results indicate that vaccination laws had a long-term beneficial effect

at increasing the vaccination coverage and will help public health practitioners to make

informed decisions on how to act against vaccine hesitancy and optimize the impact of

vaccination programmes.

1. Introduction

Infectious diseases are a major public health concern and socio-economic burden at all ages (1), but children are by far the most vulnerable to infections (2). In many low and middle-income countries, pertussis and measles remain among the top ten of all-cause mortality for children below age five. In high income countries, these infections are currently on the rise (3,4), despite the ready availability of vaccines.

One major factor driving the increasing burden of childhood infections in high-income countries is vaccine hesitancy, i.e. partial or delayed acceptance or refusal of vaccination despite the availability of vaccines (5). This vaccine hesitancy can stem either from passive neglect or active refusal and can arise for example from physical (e.g. poor accessibility and long distances) or psychological reasons (e.g. lack of communication, health literacy and cultural appropriateness) (5). Vaccine hesitancy is a key concern for public health institutions, because vaccination is our main tool to reduce and ultimately reach the local elimination of childhood infections, a target of many national public health institutions and of the World Health Organization (WHO) (6). For the childhood infection measles and pertussis, reaching elimination requires a vaccination coverage of 95 and 98% of the population respectively (7,8), and such levels of vaccination coverage that cannot be achieved with the current levels of hesitancy in high-income countries (9).

One strategy to combat vaccination hesitancy is to adopt vaccination laws, i.e. make vaccination compulsory. Within the last five years, several European countries have adopted vaccination laws, for example in Italy since 2017 children are required to be vaccinated against ten childhood infections to be allowed to public schools and similar laws have been adopted in France (2018) and Germany (2020). However, to what extent vaccination laws are effective at increasing vaccination coverage is debated (10–15). On the one hand, vaccination laws can increase vaccination coverage, but on the other hand vaccination laws can trigger a protest response and increase vaccination refusal. So far, preliminary studies and short-term monitoring indicate that recent vaccination laws have been successful at generating a short-term increase in vaccination coverage in Italy and France (16,17) and in California, where a recent vaccination policy eliminated nonmedical exemptions from school entry requirements (18). However, to what extent this is a general result and whether the increase lasts on the long-term remain unknown.

In this study, we quantified the vaccination coverage and the long-term impact of vaccination laws on the vaccination coverage and its spatio-temporal dynamics during world's first vaccination program against smallpox in 19th century Finland. In brief, smallpox was one of

the most lethal diseases in human history, affecting especially European and Native American populations (19–21). For example, in our study population in 18th and 19th century Finland, smallpox was a major cause of death in children, killing up to 15% of Finland's population around 1800 (22,23). The smallpox vaccination campaign started in Finland in 1802 and qualitative historical records indicate it was met with high hesitancy (24,25). Despite continuing opposition to vaccination, the Finnish government adopted a vaccination decree in 1883, enforced from 1885 onwards, which required mandatory vaccinations for all children under the age of two as well as for the anyone in the population who had not already been vaccinated (26).

For the purpose of this study, we digitized a series of unique historical vaccination records (63 years, 1800 pages and 46,000 individuals) that were maintained by the local church and the district doctor from the first vaccination campaign against smallpox in eight rural parishes of Finland (Fig. 1A, Fig. S1). Records contained, per parish and year, the name, parent's occupation, date and age of people who were called for vaccination, the date of the follow-up examination, and the presence or absence from vaccination and whether the vaccination was a success based on the reaction from the vaccine (effective, mild or no reaction) (Fig. S1). To estimate vaccination coverage per age class, we combined these individual-based vaccination records with church-maintained records on births and deaths from a newly available high-resolution long-term database provided by the Genealogical Society of Finland.

Here we used these records for two specific aims. First, we described the seasonal and annual dynamics of the smallpox vaccination campaigns in rural Finland and which age classes were targeted. Little is known on how the smallpox vaccination campaign was implemented in the 19th century and to what extent it was successful (20), especially in rural areas where vaccination hesitancy remains high even today in many parts of the world (27). Second, we tested two specific hypotheses, namely: did the vaccination law cause (i) a long-term increase in vaccination coverage and (ii) a long-term decrease in variance in vaccination coverage. Increasing vaccination coverage and stabilizing its spatio-temporal variance are important because high variance can create pockets of susceptibles from which epidemics can re-occur (28). The high vaccine hesitancy in the 19th century together with the rare long-term monitoring of vaccination coverage after the vaccination law provide a unique opportunity to quantify the long-term impact of vaccination laws in a high-opposition setting which is not yet possible in countries that recently implemented vaccination laws.

3. Results

3.1. Seasonality and age distribution of the vaccination campaign

Here we present data covering eight rural parishes in South-West Finland (Fig. 1A), with

26,328 records (57%) during the 46 years of the pre-mandatory era from 1837 to 1882 (average

572 records per year) and 19,935 records (43%) during the 17 years of mandatory era from

1883 until 1899 (average 1173 records per year; Table 1).

During both eras, vaccination campaigns started in May or June, occurred predominantly in

summer and often ended in August or September or occasionally as late as October or

November (Fig. 1B). This seasonal pattern was confirmed by wavelet analysis, where we

observed a statistically significant annual vaccination periodicity ($p<0.05$), both on average

over the whole observation period ($p<0.05$, Fig. S2A) and in the time dependent analyses

($p<0.05$, Fig. S2B).

We then investigated the age distribution of the people vaccinated during the campaigns. There

was a clear bias towards children under age one, with in the pre-mandatory era on average 69%

of the records being under age one, 21% between one and three years, 5.3% between three and

five years and 4.5% five years old or older ($\chi^2= 1143$, $p<0.0001$, Fig. 2A). In the mandatory

era, the bias towards one-year olds was 10% higher at 80% of the records and this increase was

statistically significant ($\chi^2= 39.73$, $p=2.9e-10$, Fig. 2A). We did not observe a lower limit for

the age at first vaccination (Fig. 2B): 872 of the 34,496 records (2.5%) of the children under

the age of one were below the age of 3 months. Thus, there was a bias to vaccinate before age

one which increased with time and with no apparent lower age limit.

3.2 Spatio-temporal dynamics of vaccination coverage

The number of vaccinations increased with time (Fig. 2A, Table S1A). Threshold models

showed that this increase accelerated after the break point in 1871 (1871<4AICCI<1871; Fig.

S4A) and this result was supported by gamm models (Fig. S5A&B). This break point coincided

shortly after the drop in births during the famine years 1867-68. Because vaccination dynamics

can be driven mostly by the dynamics of births (Fig. 2A), we studied birth-independent

spatiotemporal dynamics of vaccinations by using vaccination coverage, i.e. the number of

individuals vaccinated relative to the number of individuals in the age class and focused the

analyses below on children until age one year.

We tested the hypothesis that the vaccination law increased the vaccination coverage. In the

pre-mandatory   era,   the   vaccination   coverage   was   on   average   68%   (SD=27.8,

18.7<[95CI]<147) and in the mandatory era this increased with 20% to 88% (SD=21.5,

57.4<[95CI]<143; Fig. 3). Analysing these time series with threshold models showed an abrupt

increase in vaccination coverage in 1882 (1882<4AICCI<1882; ΔAICc relative to a linear

increase=-30.4; Table S1B; Fig. 4, Fig S4B). Before the threshold the vaccination coverage

decreased with time, while after the threshold we observed an increase, where the 95%CI

coefficient before the threshold is statistically significant but after the threshold overlaps with

zero and hence is not significant (respectively: β_1=-4.27, -8.35<[95CI]<-0.20, p=0.04; β_2=9.22,

-6.89<[95CI]<25.33; Table S1B). Gamm models supported the conclusions of threshold

models with the gam derivative indicating the steepest increase in coverage in 1882 (Fig.

S5C&D). Hence, the implementation of the vaccination law was associated with (i) a stop in

declining coverage and (ii) an abrupt 20% increase in coverage which (iii) persistent on the

long-term.

We then tested whether the variance in vaccination coverage changed over time and Breusch-

Pagan test indicated a decrease in variance (ΔAICc=-15.49). We tried to decompose whether

this decrease in variance is due to a spatial (between parishes) and/or a temporal (between

years) component. Vaccination coverage varied substantially and consistently between

parishes: parish identity captured 9.0% of the variance in the pre-mandatory era which

increased to 46% in the mandatory era (model in Table S1B) with a repeatability of respectively

0.083 (0.00<[95CI]<0.21) and 0.32 (0.069<[95CI]<0.55). The parish-level SD and CV in

vaccination coverage decreased over time and showed an abrupt change around 1873 (SD:

1869<4AICCI<1876, CV: 1872<4AICCI<1874; Table S1C & S1D; Fig. 3B, Fig. S4C&D) and

gamm models confirmed these conclusions (Fig. S5E-H). The interannual SD decreased as

well with time, from 12.94 in the pre-mandatory era to 6.48 in the mandatory era (Breusch-

Pagan test p=0.46). Hence the variance in vaccination coverage stabilized between parishes

and over time, but the parish-level stabilization started approximately ten years before the

introduction of the vaccination law.

4. Discussion

Commonly preventable childhood infections are on the rise in many contemporary high-income countries following increases in vaccination hesitancy, which has led to vaccination laws against infections such as measles and pertussis e.g. in Europe and Australia. However, we know almost nothing on the long-term effects of vaccination laws on vaccination coverage and its dynamics. In this study, we show that the world's first vaccination campaigns in rural Finland was mainly executed during the summer, often in the middle of farming season and most of the vaccinated individuals were young children under the age of one year. We show an abrupt increase in vaccination coverage increase which coincided with the start of a vaccination law in Finland and persisted for at least 15 years. However, we also found differences in vaccination coverage between parishes, and although spatio-temporal variance in vaccination coverage decreased with time, this variance persisted and its sudden decrease ten years before the vaccination law appeared not to be associated with the vaccination law. Here we discuss three implications of our results the impact of vaccination campaigns and vaccination policy.

First, one notable characteristic of the campaign was that children seemed to be vaccinated without lower age limit. Smallpox epidemics were still prevalent during the 19[th] century Finland and as vaccinators travelled to rural areas only once a year, early vaccination may at first hand seem the optimal solution. However, we now know that early vaccination can be ineffective, for example through maternal blunting, where maternal antibodies of vaccinated mothers interfere with infant immune response and lower the infant's responsiveness to vaccines (46), hence for example in contemporary societies a recommended age at first vaccination of at least three months for several childhood infections and one year for smallpox (47). This ineffective early vaccination likely contributed to later smallpox outbreaks. In addition, early vaccination can also increase the risk of complications, especially central nervous system disease, including postvaccinial encephalopathy (PVE) (47,48), which could cause an increase in vaccine hesitancy, although our results show no evidence of such response.

Second, low vaccination coverage in rural areas is a challenge that contemporary vaccination campaigns try to solve, especially in low and middle income countries (49–51). Increasing their vaccination coverage is important because pockets of low vaccination coverage can be a source for outbreaks towards other areas (28). Our finding of repeatable parish-specific vaccination coverage show that vaccination coverage is a geographically local characteristic. The reasons behind the poor and fluctuating coverage in rural communities are complex (5,49,50), but at least two possibilities stand out. Firstly, remote parishes were often more difficult to access for vaccinators and possibly less well managed than more urban parishes, a characteristic also

found in contemporary societies (52). Secondly, people living in remote rural parishes might

be less motivated, more hesitant or even completely opposed to vaccination (53,54). Although

we observed a decline in the spatio-temporal variation in vaccination coverage, indicating

improvements in the management of the vaccination campaign, the reasons behind this decline

remain to be identified.

Finally, recent estimates on the growth of the anti-vaccine movement show an alarming ability

to reach and recruit vaccine-hesitant individuals (55). Finland's smallpox vaccination

campaign 200 years ago was also initially likely met with high hesitancy (26). The vaccination

decree introduced in 1883 and enforced after 1885 required mandatory vaccinations for all

children under the age of two as well as for the anyone in the population who had not already

been vaccinated. Our results show an abrupt increase in the vaccination coverage at the start of

the vaccination law. Similarly, short-term studies in contemporary societies observed

immediate increases in vaccination coverage after the introduction of the vaccination laws

(16,56) and our study suggested that the observed increases will persist on a longer term. In

our study the increase in vaccination coverage associated with the start of the vaccination law

could be due to a law-abiding decrease in vaccine hesitancy or alternatively due to a sudden

improvement in vaccine access or delivery. In the latter scenario, we would also expect sudden

improvements in vaccine access or delivery to also cause abrupt changes in the variance in

vaccination coverage, a result which we do not find. This indicates that the vaccination law

increased people's incentive to get vaccinated and is in contrast with the stabilisation

(decreasing SD and CV) in vaccination coverage that we observed a decade before the

vaccination law, which probably reflects an improvement in Finland's access to vaccines.

The reasons behind vaccine hesitancy remain complex and hence other measures than

vaccination laws, such as training medical professionals about vaccines, actively

recommending vaccines and good communication with the lay public are also needed to

increase the vaccination coverage (15,57). Vaccination laws should be implemented in

association with clear and consistent communication and these can help reach infection-

specific thresholds for herd immunity and subsequent elimination or eradication (28,58). Our

study indicates that vaccination laws will be an effective long-term tool in the public health

battle to increase vaccination coverage and towards the elimination of childhood infections,

even in socio-demographic contexts (rural areas, pre-health care periods) where vaccines were

met with high reluctance.

2. Materials and methods

2.1 Study population and data collection

Vaccination records were manually photographed and digitised from extensive church records held in Finnish national and provincial archives. These data include details of the villagers and their lives, that were by law required to be collected by local clergymen from 1749 onwards. The clergymen recorded the name, address, parents' occupation, age, and the level of success of the vaccination (as indicated by reported reaction to the vaccine: effective, mild or no reaction) for each person receiving the vaccination at a given time, as well as previously vaccinated, absent or refused individuals (Fig. S1). We obtained data from eight rural parishes (Honkajoki, Ikaalinen, Jämijärvi, Karvia, Kustavi, Parkano, Rymättylä, Tyrvää) from 1837 until 1899, but we lacked vaccination records from 1847-1851, 1854 and 1856-1858 for all parishes excluding Ikaalinen and Tyrvää, which only lacked campaign season data for these years (Fig. S3). These records include information on a total of 46,232 vaccinations, 26,328 records from pre-mandatory and 19,935 records from mandatory era (Table 1).

In order to estimate the number of vaccinated children per age group, we gathered birth and death records for each parish from the national and provincial archives, the Genealogical Society of Finland and Finland's Family History Association digitized archives. These records consisted of yearly births and deaths per age cohort (e.g. 0-1, 1-3, 3-5), which we used to calculate the number of individuals per age cohort per parish. We estimated vaccination coverage based on the church records and total cohort sizes by calculating the yearly proportion of vaccinated per cohort and parish from 1837 to 1899. Every five to ten years we had review tables of age-specific cohort sizes per parish from the historical records which confirmed our annually estimated cohort sizes.

2.2 Statistical analyses

2.2.1. Seasonality and age distribution of the vaccination campaign

The first aim of the study is to describe the occurrence and seasonal dynamics of the vaccination campaigns. To detect whether vaccination campaigns were seasonally recurring we conducted wavelet analysis. In brief, wavelet analysis decomposes time series data using functions (wavelets) simultaneously as a function of both time (year of observation) and period, with a period of 12 months indicating a seasonally recurring pattern (29). We fitted a Morlet wavelet with the function 'analyze.wavelet' of the package 'WaveletComp' (30). We present the monthly analyses by plotting a period from six months to two years, where 12 months indicated a yearly seasonal signal. To circumvent any local peaks, we smoothened and detrended the

results following standard protocol (30). We determined statistical significance by comparing
the observed periodicity from the data with that of 1000 'white noise' simulated datasets with
[95%] significance level.

We then describe the age structure of the vaccination campaigns. For these analyses, we used
a matrix with the number of vaccinated per age group per year and divided age into four groups
(0-1, 1-3, 3-5, 5+). We tested for the bias towards children under one year old using non-
parametric permutation tests with the function 'independence_test' of the package 'coin' (31)
based on 10,000 permutations.

2.2.2. Spatio-temporal dynamics of vaccination coverage

To quantify the dynamics in vaccination coverage over time and across parishes and whether
the vaccination law affected these dynamics, we used generalized linear mixed models (glms)
and generalized additive mixed models (gamms). We ran four statistical models with as the
dependent variable either: (a) the number of vaccinations, (b) the vaccination coverage, (c) the
parish-level annual SD in vaccination coverage and (d) the parish-level annual CV in
vaccination coverage (for explanation see below, Table S1A-D). In all these models the
predictor variable was year standardized to the mean of zero and an SD of one.

We performed all statistical analyses with the software R (version: 3.6.1, R Core Team, 2020).
We fitted glms with the functions 'glm', 'lme' and 'gls' respectively of the packages 'stats'(R
Core Team 2020) and 'nlme' (33). These gave consistent results and we here present the results
using 'glm' for the number of vaccinated (Poisson distributed counts), 'lme' for vaccination
coverage (Gaussian distributed) and 'gls' for parish-level SD and CV (Gaussian distributed
with temporal autocorrelation, see below). To quantify the statistical 'significance' of predictor
variables, we compared the model fits on the data with the second order Akaike Information
Criterion (AICc; (34,35) using the function 'AICc' of the package 'MuMIn' (36). In brief,
model selection ranks the models based on their AICc value, where better fitting models are
indicated by lower AICc, models within 4 ΔAICc are considered plausible, and increasingly
implausible up to 14 ΔAICc after which they are implausible (34,35).

To account for the fact that people are clustered into parishes, we included parish identity as a
random intercept in the model (b), respectively the vaccination coverage analyses, which
improved the model fit (ΔAICc=-29 in Table S1B). To have an idea of the importance of parish
level differences, we reported the proportion of variance in the model explained by the random
intercept and estimated the repeatability of parish-level differences with the function 'rpt' of
the package 'rptR' (37). The repeatability is an intra-class correlation coefficient that captures

the between-parish variance (by parish identity as a random intercept) relative to the total

variance and ranges from zero (no parish-level differences) to one (all variance in the dependent

variable is explained by parish-level differences). We estimated the 95% confidence intervals

(95CI) around the repeatability based on 1000 bootstraps.

For model (a) number of vaccinations, we used Poisson distribution and for all other models

(b-d) we used a Gaussian distribution. All model residuals were correctly distributed and

fulfilled all other assumptions as checked with the function 'simulateResiduals' of the package

'DHARMa' (38). In the model (b), there was however heteroscedasticity as identified with the

functions (i) 'testQuantiles' from the package 'DHARMa' (38), and (ii) the Breusch-Pagan test

performing a linear model using as dependent variable the squared value of the conditional

residuals of the model (b) and standardized year as predictor variable (39). In model (b), we

corrected for heteroscedasticity by including the variance power ('varPower') in the function

weights (40,41). We then tried to identify two sources of changes in heteroscedasticity by

decomposing vaccination coverage into a parish-level component and an annual component.

For the parish-level component, we used as the dependent variable the annual parish-level SD

and CV in vaccination coverage (respectively models (c) and (d), Table S1 C & D). To identify

whether there was an annual component in heteroscedasticity we used the Breusch-Pagan test

as described above on the annual mean in vaccination coverage (excluding parish-level

variance) and using pre-mandatory and mandatory era as a two-level categorical predictor.

We controlled for temporal autocorrelation following (42) by including standardized year as

an auto-regressive factor of order 1 (corAR1). In models of number of vaccinated (i) and

vaccination coverage (ii), this worsened the model fit (ϕ=0.57±0.25, ΔAICc=+155 for model

b in Table S1B), indicating no temporal autocorrelation, a conclusion also supported with the

function 'acf' (ACF<0.1) of the package 'stats' (R Core Team 2020). For these models, we

hence presented the results without temporal autocorrelation, but analyses including temporal

autocorrelation gave consistent results (not shown). Models (c) and (d), respectively parish-

level SD and CV in vaccination coverage were autocorrelated and thus we included the corAR1

autocorrelation structure in these models.

To investigate whether there were abrupt changes in vaccination coverage or its spatio-

temporal variance which coincided with the start of the vaccination law, we fitted threshold

models following Douhard et al. (43) and Briga et al (44). In brief, for each model (a-d), we

tested a series of threshold models, with one year interval between each threshold and identified

the best fitting threshold model based on the model's AICc value. To identify whether threshold

models were appropriate at all, we compared the model fit of the best fitting threshold model

with that of a linear model without threshold (Table S1 A-D). We identified the confidence interval around the threshold year by including all models within 4 ΔAICc of the best fitting threshold model (43,44), abbreviated as 4AICCI (Fig. S4 A, B, C & D). To confirm the results of threshold models, we performed all analyses also using generalized additive mixed models (gamms), with the function 'gamm' of the package 'mgcv', and to identify the year of change (inflection points) we used their derivatives with the function 'fderiv' of the package 'gratia' (45). All gamm analyses confirmed the conclusions obtained from the threshold models (Fig. S5 A-H).

Acknowledgements

Many thanks to Kimmo Pokkinen, Sara Itkonen and Tuija Koivisto for help with historical data, to the National Archives of Finland for allowing us to digitize and use the vaccination records and to Finland's Family History Association and the Genealogical Society of Finland for access to the birth and death data. We are grateful for the funding from the Doctoral Programme in Biology, Geography and Geology, University of Turku (SU), the Academy of Finland (292368, VL), ERC (CoG 648766, VL) and the Ella & Georg Ehrnrooth Foundation (MB).

References

1. Morens DM, Folkers GK & Fauci AS (2004). The challenge of emerging and re-emerging infectious diseases. Nature 430(July):242–249. https://doi.org/10.1038/nature02759

2. Naghavi M, Abajobir A, Abbafati C, Abbas K, Abd-Allah F, Abera S-F & Aboyans V (2017). Global, regional and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet 390(10100):1151–1210. https://doi.org/10.1016/S0140-6736(17)32152-9

3. Dabbagh A, Patel MK, Dumolard L, Gacic-Dobo M, Mulders MN, Okwo-Bele JM, Kretsinger K, Papania MJ, Rota PA, & Goodson JL (2017). Progress toward regional measles elimination- worldwide, 2000-2016. Morbidity and Mortality Weekly Report. 66(42):1148–1153. https://doi.org/10.15585/mmwr.mm6642a6

4. Cherry JD (2012). Epidemic Pertussis in 2012 — The Resurgence of a Vaccine-Preventable Disease. *New England Journal of Medicine* 367(9):785–787. https://doi.org/10.1056/NEJMp1209051

5. World Health Organisation (2014). *Report of the SAGE working group on vaccine hesitancy.*

6. World Health Organisation (2010). *Global eradication of measles: report by the Secretariat.*

7. Guerra FM, Bolotin S, Lim G, Heffernan J, Deeks SL, Li Y & Crowcroft NS (2017). The basic reproduction number (R0) of measles: a systematic review. *The Lancet Infectious Diseases* 17(12):e420-e428. https://doi.org/10.1016/S1473-3099(17)30307-9

8. Kretzschmar M, Teunis PFM & Pebody RG (2010). Incidence and reproduction numbers of pertussis: Estimates from Serological and Social Contact Data in Five European Countries. *PLoS Medicine* 7(6): e1000291. https://doi.org/10.1371/journal.pmed.1000291

9. Wellcome Trust (2019). *Chapter 5: Attitudes to vaccines in Wellcome Global Monitor 2018.*

10. Akmatov MK, Rübsamen N, Deyneko IV, Karch A & Mikolajczyk RT (2018). Poor knowledge of vaccination recommendations and negative attitudes towards vaccinations are independently associated with poor vaccination uptake among adults – Findings of a population-based panel study in Lower Saxony, Germany. *Vaccine* 36(18):2417–2426.

374   https://doi.org/10.1016/j.vaccine.2018.03.050

11. Maltezou HC, Theodoridou K, Ledda C, Rapisarda V & Theodoridou M (2019). Vaccination of healthcare workers: is mandatory vaccination needed? *Vaccines* 18(1):5–13. https://doi.org/10.1080/14760584.2019.1552141

12. Gualano MR, Olivero E, Voglino G, Corezzi M, Rossello P, Vicentini C, Bert F & Siliquini R (2019). Knowledge, attitudes and beliefs towards compulsory vaccination: a systematic review. *Human Vaccines and Immunotherapeutics* 15(4):918–931. https://doi.org/10.1080/21645515.2018.1564437

13. Hadjipanayis A, Dornbusch HJ, Grossman Z, Theophilou L & Brierley J (2020). Mandatory vaccination: a joint statement of the Ethics and Vaccination working groups of the European Academy of Paediatrics. *European Journal of Pediatrics* 179(4):683–687. https://doi.org/10.1007/s00431-019-03523-4

14. Vaz OM, Ellingson MK, Weiss P, Jenness SM, Bardají A, Bednarczyk RA & Omer SB (2020). Mandatory vaccination in Europe. *Pediatrics*. 145(2):e20190620. https://doi.org/10.1542/peds.2019-0620

15. Musa S, Skrijelj V, Kulo A, Habersaat KB, Smjecanin M, Primorac E, Becirovic D & Jackson C (2020). Identifying barriers and drivers to vaccination: A qualitative interview study with health workers in the Federation of Bosnia and Herzegovina. *Vaccine* 38(8):1906–1914. https://doi.org/10.1016/j.vaccine.2020.01.025

16. Gori D, Costantino C, Odone A, Ricci B, Ialonardi M, Signorelli C, Vitale F & Fantini MP (2020). The impact of mandatory vaccination law in Italy on mmr coverage rates in two of the largest italian regions (Emilia-romagna and sicily): An effective strategy to contrast vaccine hesitancy. *Vaccines* 8(1):57. https://doi.org/10.3390/vaccines8010057

17. Lévy-Bruhl D, Fonteneau L, Vaux S, Barret A-S, Antona D, Bonmarin I, Che D, Quelet S & Coignard B (2019). Assessment of the impact of the extension of vaccination mandates on vaccine coverage after 1 year, France, 2019. *Euro Surveillance : Bulletin Europeen Sur Les Maladies Transmissibles = European Communicable Disease Bulletin* 24(26):1900301. https://doi.org/10.2807/1560-7917.ES.2019.24.26.1900301

18. Nyathi S, Karpel HC, Sainani KL, Maldonado Y, Hotez PJ, Bendavid E & Lo NC (2019). The 2016 California policy to eliminate nonmedical vaccine exemptions and changes in vaccine coverage: An empirical policy analysis. *PLoS Medicine* 16(12): e1002994. https://doi.org/10.1371/journal.pmed.1002994

19.   Duncan SR, Scott S & Duncan CJ (1993). The Dynamics of Smallpox Epidemics in
Britain, 1550-1800. *Demography* 30(3):405–423. https://doi.org/10.2307/2061648

20.   van Panhuis WG, Grefenstette J, Jung SY, Chok NS, Cross A, Eng H, Lee BY,
Zadorozhny V, Brown S, Cummings D & Burke DS (2013). Contagious Diseases in the
United States from 1888 to the Present. *New England Journal of Medicine*
369(22):2152–2158. https://doi.org/10.1056/NEJMms1215400

21.   Krylova O & Earn DJD (2019). Patterns of smallpox mortality in London, England, over
three centuries. BioRXiV. https://doi.org/10.1101/771220

22.   Mielke JH, Jorde LB, Trapp PG, Anderton DL, Pitkanen K & Eriksson AW (1984).
Historical Epidemiology of Smallpox in Åland, Finland: 1751-1890. *Demography*
21(3):271–295.

23.   Briga M, Ketola T & Lummaa V. The immediate effectiveness of the world's first
vaccination programme against childhood infections in a developing society. *In review.*

24.   Björkstén J. *Vaccinationens historia i Finland II*. Helsingfors; 1908.

25.   Björkstén J. *Vaccinationens historia i Finland, I*. Helsingfors; 1902.

26.   Pitkänen K (1990). Myrkkyä, sanoi Tohtori Lybeck. Rokotustaistelu 1900-luvun
alkuvuosikymmenillä. In *Arki ja murros. Tutkielmia keisariajan lopun Suomesta* (pp.
127–139). SHS, Helsinki.

27.   Ijarotimi IT, Fatiregun AA, Adebiyi OA, Ilesanmi OS & Ajumobi O (2018). Urban–
rural differences in immunisation status and associated demographic factors among
children 12-59 months in a southwestern state, Nigeria. *PLoS One* 13(11):e0206086.
https://doi.org/10.1371/journal.pone.0206086

28.   Keeling MJ & Rohani P (2011). *Modeling infectious diseases in humans and animals*.
Princeton, 278 USA: Princeton University Press.

29.   Cazelles B, Chavez M, Constantin de Magny G, Guégan J-F & Hales S (2007). Time-
dependent spectral analysis of epidemiological time-series with wavelets. *Journal of the
Royal Society, Interface* 4(15):625–636. https://doi.org/10.1098/rsif.2007.0212

30.   Roesch A & Schmidbauer H (2018). *WaveletComp: Computational Wavelet Analysis*.

31.   Hothorn T, Bretz F & Westfall P (2008). Simultaneous inference in general parametric
models. *Biometrical Journal* 50(3):346–363. https://doi.org/10.1002/bimj.200810425

32.   R Core Team, R (2020). *A language and environment for statistical computing.* R

Foundation for Statistical Computing, Vienna, Austria. https://www.r-project.org/

33.   Pinheiro JC, Bates DM, DehRoy S & Sarkar D (2020). nlme: Linear and Nonlinear
Mixed Effects Models. *R Core Team.* https://cran.r-project.org/package=nlme

34.   Burnham KP, Anderson DR & Huyvaert KP (2011). AIC model selection and
multimodel inference in behavioral ecology: Some background, observations, and
comparisons. *Behavioral Ecology and Sociobiology* 65(1):23–35.
https://doi.org/10.1007/s00265-010-1029-6

35.   Burnham KP & Anderson DR (2002). *Model selection and multimodel inference : a*
practical information-theoretic approach. Springer-Verlag New York.
https://doi.org/10.1007/b97636

36.   Barton K (2020). *MuMIn: Multi-model inference.* https://cran.r-
project.org/package=MuMIn

37.   Nakagawa S & Schielzeth H (2010). Repeatability for Gaussian and non-Gaussian data:
A practical guide for biologists. *Biological Reviews* 85(4):935–956.
https://doi.org/10.1111/j.1469-185X.2010.00141.x

38.   Hartig F (2020). *DHARMa: residual diagnostics for hierarchical (multi-level/mixed)*
regression models. https://cran.r-project.org/package=DHARMa

39.   Breusch TS & Pagan AR (1979). A Simple Test for Heteroscedasticity and Random
Coefficient Variation. *Econometrica* 47(5):1287–1294.
https://doi.org/10.2307/1911963

40.   Zuur AF, Ieno EN, Walker N, Saveliev AA & Smith GM (2009). *Mixed effects models*
and extensions in ecology with R. Springer-Verlag New York.
https://doi.org/10.1007/978-0-387-87458-6

41.   Pinheiro JC & Bates DM (2000). *Mixed-Effect Models in S and S-plus.* Springer-Verlag
New York. https://doi.org/10.1007/b98882

42.   Woods S (2017). *Generalized Additive Models. An Introduction with R* (2nd ed.).
Chapman & Hall, Boca Raton, FI. https://doi.org/10.1201/9781315370279

43.   Douhard F, Gaillard J-M, Pellerin M, Jacob L & Lemaître J-F (2017). The cost of
growing large: costs of post-weaning growth on body mass senescence in a wild
mammal. *Oikos* 126(9):1329–1338. https://doi.org/10.1111/oik.04421

44.   Briga M, Jimeno B & Verhulst S (2019). Coupling lifespan and aging? The age at onset
of body mass decline associates positively with sex-specific lifespan but negatively with

environment-specific   lifespan.   *Experimental   Gerontology*   119:111–119.
https://doi.org/10.1016/j.exger.2019.01.030

45.   Simpson GL (2019). *gratia: Graceful 'ggplot'-based graphics and other functions for*
GAMs fitted using "mgcv". https://cran.r-project.org/package=gratia

46.   Zimmermann P, Perrett KP, Messina NL, Donath S, Ritz N, van der Klis FRM & Curtis
N (2019). The Effect of Maternal Immunisation During Pregnancy on Infant Vaccine
Responses. *EClinicalMedicine* 13:21–30. https://doi.org/10.1016/j.eclinm.2019.06.010

47.   Kempe C (1960). Studies Smallpox and Complications of Smallpox Vaccination.
Pediatrics 26:176–189.

48.   Cono J, Casey C, Bell D & Centers for Disease Control and Prevention (2003). Smallpox
Vaccination   and   Adverse   Reactions.   Guidance   for   Clinicians.   *MMWR.*
Recommendations   and   reports: Morbidity   and   mortality   weekly   report.
Recommendations and reports 52(RR-4):1–28.

49.   Odusanya OO, Alufohai EF, Meurice FP & Ahonkhai VI (2008). Determinants of
vaccination   coverage   in   rural   Nigeria.   *BMC   Public   Health*   8(1):381.
https://doi.org/10.1186/1471-2458-8-381

50.   Singh PK (2013). Trends in Child Immunization across Geographical Regions in India:
Focus   on   Urban-Rural   and   Gender   Differentials.   *PLoS   One*   8(9):e73102.
https://doi.org/10.1371/journal.pone.0073102

51.   Shikuku DN, Muganda M, Amunga SO, Obwanda EO, Muga A, Matete T & Kisia P
(2019). Door - To - door immunization strategy for improving access and utilization of
immunization Services in Hard-to-Reach Areas: A case of Migori County, Kenya. *BMC*
Public Health 19(1):1–11. https://doi.org/10.1186/s12889-019-7415-8

52.   Gram L, Soremekun S, ten Asbroek A, Manu A, O'Leary M, Hill Z, Danso S, Amenga-
Etego S, Owusu-Agyei S & Kirkwood BR (2014). Socio-economic determinants and
inequities in coverage and timeliness of early childhood immunisation in rural Ghana.
Tropical   Medicine   &   International   Health   19(7):802–811.
https://doi.org/10.1111/tmi.12324

53.   Wagner AL, Huang Z, Ren J, Laffoon M, Ji M, Pinckney LC, Sun X, Prosser LA,
Boulton ML & Zikmund-Fisher BJ (2020). Vaccine Hesitancy and Concerns About
Vaccine Safety and Effectiveness in Shanghai, China. *American Journal of Preventive*
Medicine S0749-3797(20)30402-5. https://doi.org/10.1016/j.amepre.2020.09.003

54.   Thomas TL, Caldera M, & Maurer J (2019). A short report: parents HPV vaccine
502   knowledge in rural South Florida. *Human vaccines & immunotherapeutics*, 15(7-
503   8):1666–1671. https://doi.org/10.1080/21645515.2019.1600986

504   55.   Johnson NF, Velásquez N, Restrepo NJ, Leahy R, Gabriel N, Oud S El, Zheng M,
505   Manrique P & Wuchty S (2020). The online competition between pro- and anti-
506   vaccination views. *Nature* 582:230–233. https://doi.org/10.1038/s41586-020-2281-1

507   56.   Partouche H, Gilberg S, Renard V & Saint-Lary O (2019). Mandatory vaccination of
508   infants in France: Is that the way forward? *European Journal of General Practice*
509   25(1):49–54. https://doi.org/10.1080/13814788.2018.1561849

510   57.   Larson HJ, Jarrett C, Eckersberger E, Smith DMD & Paterson P (2014). Understanding
511   vaccine hesitancy around vaccines and vaccination from a global perspective: A
512   systematic review of published literature, 2007-2012. *Vaccine* 32(19):2150–2159.
513   https://doi.org/10.1016/j.vaccine.2014.01.081

514   58.   Anderson RM & May R (1991). *Infcetious diseases of humans: dynamics and control.*
515   Oxford University Press

Figures and tables

Table 1. Overview of the data, where we describe year, population and sample size, average number of vaccinated per year per parish, average timing and vaccination coverage per age class and interannual and interparish standard deviation (SD) and coefficient of variation (CV) of vaccination coverage for both pre-mandatory and mandatory eras.

	Pre-mandatory era	Mandatory era
Years	1837-1882	1883-1899
Average population size per parish	2600	3700
N vaccinated	26328	19935
Average n vaccinated per year per parish	108	148
Average [95CI] of month vaccinated	7.40 ± 0.018	6.72 ± 0.011
Average % of vaccinated that are under 1 year old	68.1	87.5
Average % of vaccinated between 2 and 5 years	16.3	10.3
Interannual and -parish SD of vaccination coverage	27.8	21.5
Interannual and -parish CV of vaccination coverage	40.9	24.6

Figure 1. Geographical and temporal distributions of the data, showing (A) map of the eight rural parishes in south-west Finland with a map of Northern Europe on the left and a zoomed-in map of south-west Finland on the right, and (B) the predominant seasonal timing of annual vaccination campaigns occurring in summer.

Figure 2. Smallpox vaccinations were biased towards the youngest age groups (A) with no apparent minimum age
(B). The number of vaccinated per age group (A) and age distribution for children under the age of one year (B).
The dashed black line in (A) indicates the number of annual births in the eight study parishes. The red line in (B)
indicates contemporary recommendation for age at first vaccination, which is 3 months in Finland for most
vaccines. For smallpox, the recommended age at first vaccination is one year (47).

Figure 3. The vaccination coverage increased after the introduction of vaccination law in 1883 (A) together with
the decrease in the variance between parishes (B).

Figure 4. The temporal dynamics in vaccination coverage of children until age one year are best captured by a
threshold model with an increase in vaccination coverage in 1882. Before the threshold, the vaccination coverage
decreases with time, while after the threshold there is an increase but the 95% CI of both coefficients overlap with
zero and hence are not statistically significant. The predicted values are plotted as a light grey line against the
black observed datapoints and the grey bands represent the [95%] confidence intervals.

Supplementary material:

The long-term success of mandatory vaccination laws at implementing world's first vaccination campaign in rural Finland

Susanna Ukonaho[1], Virpi Lummaa[1]*, Michael Briga[1]*

[1] Department of Biology, University of Turku, Finland

*co-senior authors

Figure S1. Illustration of Finnish historical vaccination records (Community: Ikaalinen, year: 1876, page 1). Columns represent (from left to right): I, ii, iii: Location, village and house; iv: Name and occupation of the parent (mother or father); v, vi: Names and birthdate of person called for vaccination; vii, viii: date of vaccination; ix, x: date of follow-up examination; xi-xiii: success of vaccination (successful, partial or no reaction); xiv-xv: absent from vaccination or follow-up examination. People who disagreed to vaccination are symbolised by '/' or 'x' and do not have a date (not on this page); xvi: remarks and notes.

Figure S2. Wavelet periodogram (A) and wavelet power spectrum (B) for season at vaccination show a consistent
average around one year from 1837 until 1899. A) A smaller peak can be observed at 6 months. Red dots represent
significance at the 5% level and blue dots 10% level. The axis describe period in months on the y-axis and wavelet
power levels on the x-axis. B) Legend describes wavelet power levels, where red indicates strong periodicity and
green to blue signals weak periodicity. The blue gaps around 1850-1860 represent a gap in the data from all
parishes at that time. The black line highlights the one-year mark in both plots.

Figure S3. The number of vaccinated individuals increased towards the end of the 19[th] century as monitored from
1837 to 1899.

Table S1. The model selection table indicated a quadratic increase in the number of vaccinated (count),
vaccination coverage and coefficient of variation over time, but was not a significant predictor in standard
deviation models. The table describes model type, model intercept, threshold intercept (Intercept2), scaled year
coefficient, scaled threshold year coefficient (year2), period (before vs. after the law), model AICc, delta AICc,
Akaike weight and threshold year for each dependent variable. The vaccination coverage model in B includes
parish as a random variable. Best fitting models are indicated in italic.

	Model output				Model selection			
Model	Intercept	Intercept2	year	year2	AICc	ΔAICc	weight	threshold
A) Vaccination count								
Linear	6.51		0.43		4892.08	995.02	0.00	
No year-related change	6.60				12781.81	8884.75	0.00	
Threshold	*6.22*	*0.60*	*0.14*	*0.041*	*3897.06*	*0.00*	*1.00*	*1871*
B) Vaccination coverage								
Linear	74.90		5.21		3469.68	30.26	0.00	
No year-related change	75.20				3481.65	42.23	0.00	
Threshold	*66.62*	*15.41*	*-4.28*	*9.27*	*3439.42*	*0.00*	*1.00*	*1882*
C) Parish-level standard deviation in vaccination coverage								
Linear	23.29		-1.70		475.92	12.22	0.00	
No year-related change	23.26				477.60	13.90	0.00	
Threshold	*29.98*	*-8.73*	*5.56*	*-7.05*	*463.70*	*0.00*	*1.00*	*1873*
D) Parish-level coefficient of variation in vaccination coverage								
Linear	32.49		-3.74		520.38	30.13	0.00	
No year-related change	32.43				524.63	25.88	0.00	
Threshold	*48.09*	*-15.70*	*13.57*	*-21.17*	*494.50*	*0.00*	*1.00*	*1873*

581   Figure S4. The number of vaccinated showed statistically significant threshold in 1871 (A, Table S1A), while the

582   vaccination coverage showed a statistically significant threshold in 1882 (B, Table S1B). Both parish-level SD

583   and CV models showed a statistically significant threshold in 1873 (C&D, Table S1C&D).

585 Figure S5. Gamm models show the steepest increase for number of vaccinated in 1871 (A) and vaccination

586 coverage in 1882 (C), and for both parish-specific SD and CV a decrease starting from 1873 onwards (E&G).

587 Shown here are the predicted values (A, C, E, G) and the derivatives of the GAMM models (B, D, F, H) for

588 vaccination count (eq. model Table S1A; A&B), vaccination coverage (eq. model Table S1B; C&D) and parish-

589 specific SD (eq. model table S1C; E&F) and parish-specific CV (eq. model Table S1D; G&H). Vaccination

590 coverage models included parish identity as a random factor. All models used autocorrelation structure AR1 for

591 year and estimated 95% confidence intervals (grey band) around the predicted slopes (blue line).