The Impact of COVID-19 Lockdowns on the Behavior of Italian Citizens and Particulate Matter 10 and 2.5 Emissions in Lombardy

Alessandro Rovetta[1,2]

[1] Mensana srls, Research and Disclosure Division, Brescia, Italy
[2] Redeev srl, Technologic and Scientific Research, Napoli, Italy

e-mail: rovetta.mresearch@gmail.com, ORCID: 0000-0002-4634-279X

Keywords: Consumption, COVID-19, Italy, Lockdown, PM10, PM2.5, Sustainability

Abstract

Italy was one of the first nations in the world to have been heavily affected by the first wave of COVID-19. To date, it is among the first countries for both total cases and deaths. Numerous containment measures were taken from February to December 2020 to mitigate the pandemic, including total lockdowns across the entire country. The purpose of this research is to evaluate how these restrictions influenced Italian citizens' consumption habits (like online shopping, smart working and distance learning) and particulate matter (PM) 10 and 2.5 concentrations in the Lombardy region. In particular, this survey is aimed at assessing the environmental sustainability of the behaviors adopted in response to the limitations of personal freedoms on a regional and national level and quantifying the effect of the latter on particulate concentrations in Lombardy, the region most damaged by both COVID-19 and air pollution. Many tools and online platforms have been used to collect data, such as Google Trends, web portals for sharing statistical and demographic information like the National Institute of Statistics (ISTAT) and AdminStat Italia, and surveys conducted by the Department of Civil Protection, other scientific studies, and the main national newspapers. Technical data on particulate matter was collected from the website of the Regional Agency for the Protection of the Environment (ARPA). To highlight any significant change, the average daily concentrations of PM10 and 2.5 during 2020 in all the provinces of Lombardy were compared with those of the previous year. The comparison between the mean values was made through the t-test. Two values were considered as statistically confident when $t < 1.9$. However, since the real significance of these thresholds is not easily determined, we kept some margins of suspicious confidence. Finally, we looked for possible causal correlations between changes in citizens' behavior and specific key events related to COVID-19. To do this, Pearson and Spearman correlations were used. The P-value threshold was indicatively set at 0.05. Microsoft Excel 2020 and Google Sheets software were used for data analysis. In conclusion, this paper showed a substantial ineffectiveness of total and partial lockdowns in reducing PM10 and PM2.5 concentrations in Lombardy. Furthermore, it estimates that COVID-19 has

strongly changed the consumption habits of Italian citizens, forcing them to use home delivery services more actively which, at present, poses a potential additional threat to the environment. Among the positive aspects of this change there was the learning of the most modern technologies for distance teaching and smart working, a point from which to start for the construction of an energetically and environmentally sustainable policy.

1. Introduction

The COVID-19 disease has raged around the world, radically changing habits and feelings of global citizens of all ages and status and causing deep economic and health crises in all stricken countries (World Health Organization, 2020[1], Figueroa et al., 2020, Francisco et al., 2020). Among these, Italy is one of the most affected: to date (December 5, 2020), it counts almost 1.7 million official cases and 59,000 official deceased, making it the sixth nation in the world for number of deaths from COVID-19 (Worldometers, 2020)[2]. However, during the first pandemic wave, many media, websites, and environmental protection bodies highlighted copious beneficial effects of the lockdown on pollution and air quality (FASDA, 2020)[3]. In this regard, this research focuses mainly on two aspects: i) the concentrations of particulate matter pollutants with a diameter ranging from 2.5 μm to 10 μm (PM10) and less than 2.5 μm (PM2.5) in the Lombardy region, and ii) the increase in the national demand for home food delivery services, food shopping delivery services, general shopping delivery services, smart working, and distant learning. In particular, the purpose of this paper is to quantify the variation in the concentration of PM10 and PM2.5 as well as evaluate the sustainability of the behavior adopted by citizens in response to restrictions on personal freedom, containment measures, and lockdowns imposed by the government to face the widespread of SARS-CoV-2. In this regard, Lombardy was chosen as it is the region most affected by both COVID-19 and particulate matter, and there is a deep epidemiological link between the latter and SARS-CoV-2 spread (Rovetta and Castaldo, 2020). This makes it very suitable for conducting a reliable investigation and generalizing the results through statistical inference. On the contrary, since the danger, once it arrived in the country, was perceived similarly in all regions (Rovetta and Castaldo, 2020), the survey on the behavioral response of citizens has been extended nationally. To do this, Google Trends was used. Google Trends is an online tool, provided by Google™, that allows users to search for any query and visualize and quantify the trend over time of the web interest towards it in any region of the world. This infoveillance tool is increasingly exploited by the scientific community for the investigation of users' behavior and creating predictive models (Jimenez et al., 2020).

---

[1] https://www.who.int/news/item/13-10-2020-impact-of-covid-19-on-people's-livelihoods-their-health-and-our-food-systems
[2] https://www.worldometers.info/coronavirus/
[3] https://www.fasda.it/effetti-quarantena-ambiente/

2. Methods

2.1 Data collection

We used Google Trends[4] to explore Italian citizens' web interest in home food delivery services, food shopping services, shopping services, and online job and learning platforms, during and after the first COVID-19 restrictions and containment measures. Google Trends returns normalized values, called "relative search volumes (RSV)", ranging from 0 (very low relative interest) to 100 (very high relative interest). All the search parameters and queries are shown in Table 1. PM10 and 2.5 daily averages data were collected from all the measuring stations of all the provinces of Lombardy. To do that, the Regional Agency for Environmental Protection (ARPA) official website[5] was used. Data on COVID-19 cases and deaths was collected from the Department of Civil Protection official website[6]. Meteorological data was collected from the website "Il Meteo.it"[7].

Topic	Query term	Category	Sampling frequency
Home food delivery services	just eat	All	1/week
	uber eats		
	deliveroo		
	glovo		
	foodracers		
Food shopping delivery services	esselunga online	All	1/week
	spesa online (online food shopping)		
	carrefour online		
	conad online		
	iperal online		
Shopping delivery services	amazon	All	1/week
	ebay		
Smart working and distance learning	DAD + Teams + Zoom + Meet + Skype	All	1/week

Table 1. Google Trends queries and search parameters.

2.2 Statistical Analysis

We quantified the impact of COVID-19 restrictions and containment measures on the web queries RSVs and average concentrations of PM10 and 2.5 through the

---

percentage increases $\Delta\% = (x_1 - x_0)/x_0 \cdot 100$ and their related t-test $t = (x_1 - x_0)/\sigma$. All statistical errors σ were calculated with the standard propagation formula (Taylor, 1997). Each mean value $\underline{x}$ was reported together with its own standard error of the mean (SEM) according to the following modality: $\underline{x} \pm SEM$ (Taylor, 1997). Data normality was verified both with the z-tests $k/\sigma_k < 2$ and $s/\sigma_s < 2$ for kurtosis k and skewness s and graphically (Tabachnick and Fidell, 1996). For doubtful cases $(2 < z < 1.5)$, we also used the Shapiro-Wilk test. When the number of measures n was greater than 30, we exploited the central limit theorem to validate the t-test when dealing with not normally distributed data (Kwak and Kim, 2017).

2.2.1 Google Trends

We compared the weekly web query trends of 2020 with those of 2019 and 2018, searching for significant discrepancies and focusing on the first lockdown and pre-lockdown period (from 20 February to 18 May 2020). When 2018 and 2019 data series were periodic or stationary, the trends of 2020 queries were compared to those of the previous two years. When 2018 and 2019 data series were aperiodic and non-stationary, anomalous peaks of RSV were highlighted[8]. Two mean values were considered confident only when $t < 1.5$ and not confident only when $t \geq 1.9$; in the intermediate $1.5 \leq t < 1.9$ range, non-confidence was interpreted as suspicious. Pearson and Spearman correlations were searched between RSVs and COVID-19 total cases and deaths. The possible causal nature of the latter was tested by researching the same correlations between the 2019 RSVs and the COVID-19 total cases and deaths during 2020. In order to highlight any hidden correlation, Pearson and Spearman coefficients were used together (Rovetta, 2020). The chosen P-value significance threshold was $\alpha = .05$; however, all correlations with $.05 < P < .1$ were classified as suspicious (Amrhein et al., 2017).

2.2.2 PM10 and 2.5

We compared the average concentrations of PM10 and PM2.5 during 2020 with those of 2019, searching for significant discrepancies and focusing on the first lockdown (period 1: from March 9 to May 18, 2020) and during 2020 (period 2: January 1 to November 19, 2020). Two mean values were considered confident only when $t < 1.5$ and not confident only when $t \geq 1.9$; in the intermediate $1.5 \leq t < 1.9$ range, non-confidence was interpreted as suspicious. Meteorological data, such as minimum, maximum, and average temperatures, % relative humidity, and maximum wind speed, were analyzed according to the same criteria and in the same periods. To provide an idea of the coverage of territory of the measuring stations and the demographic importance of all provinces, the number of inhabitants and surface area of each province were also reported in the results section. These data were taken from the official websites of National Institute of Statistics (ISTAT)[9] and AdminStat[10].

---

[8] Anomalous peaks are defined as upward trends possibly related to lockdowns or other key events as these were not present in the previous two years trends.
[9] https://www.istat.it/

3. Results

Substantial weather changes were identified in Lombardy between 2019 and 2020. In particular, significant or suspected decreases in the maximum wind speed occured in 100% of cases during period 1 ($\Delta_{av}\% = -18.0 \pm 3.3$) and in 66.7% of cases during period 2 ($\Delta_{av}\% = -9.2 \pm 1.1$). In table 1, the t-test results between the meteorological quantities of 2019 and 2020 in 9 of the 12 provinces of Lombardy are shown.

2020 vs 2019		Period 1 (1 March - 18 May)					Period 2 (1 January - 19 November)				
		AT	T min	T max	HR	W	AT	T min	T max	HR	Ws
Bergamo	t-test	2.3	2.3	1.8	-1.3	-2.2	-0.2	-0.1	-0.3	0.5	-1.3
Brescia	t-test	1.2	0.2	1.6	0	-1.7	-0.7	-0.8	-0.7	-0.8	-2.1
Como	t-test	1.7	1.8	1.5	1.6	-2.8	-0.2	0.2	-0.1	1.9	-2.6
Cremona	t-test	2.1	1.8	1.8	-2.3	-1.7	0.1	0.4	-0.1	-2.8	-1.4
Lecco	t-test	2.3	2.3	1.8	-1.3	-2.2	-0.2	-0.1	-0.3	0.5	-1.3
Mantova	t-test	2.3	1	2.1	-2.7	-2.1	-0.3	-0.4	0.1	0.4	-2.1
Milano	t-test	1.9	1.3	2	-1.4	-2.3	0	0.2	0	1	-2.6
Pavia	t-test	1.9	1.3	2	-1.4	-2.3	0	0.2	0	1	-2.6
Varese	t-test	1.7	1.8	1.5	1.6	-2.8	-0.2	0.2	-0.1	1.9	-2.6

Table 1. Weather differences between 2019 and 2020. AT = average temperature, T min = minimum temperature, T max = maximum temperature, HR = relative humidity, Ws = wind speed.

Therefore, the results on the concentrations of particulate matter must be interpreted in the light of the above.

3.1 Effect of COVID-19 restrictions on particulate matter concentrations

3.1.0 Lombardy region
During period 1 (i.e. the comparison between the periods 9 March - 18 May 2020 and 2019), significant or suspected increases in the mean concentrations of PM10 and 2.5 were observed in 57% of cases (53/93); significant or suspected reductions occurred only in 3.2% of cases (3/93). In the remaining 39.8% of cases, we found no significant or suspected changes (37/93). During period 2 (i.e. the comparison between the period 1 January - 19 November, 2020 and 2019), significant or suspected increases in the mean concentrations of PM10 and 2.5 were observed in

---

[10] https://ugeo.urbistat.com/AdminStat/it

20.4% of cases (19/93); significant or suspected reductions occurred only in 6.4% of cases (6/93). In the remaining 73.2% of cases, no significant or suspected changes were found (68/93).

3.1.1 Province of Bergamo

The province of Bergamo has about 1.108 million inhabitants and extends for 2,755 km². Here there are 13 measuring stations: 8 for PM10 and 5 for PM2.5. During period 1, 3 significant changes were observed for the average concentrations of PM10 ($\Delta_1\% = +20.0 \pm 11.0$, $t_1 = 2.1$; $\Delta_2\% = +28.8 \pm 11.7$, $t_2 = 2.7$; $\Delta_3\% = 19.0 \pm 10.4$, $t_3 = 2.1$) and 2 for the average concentrations of PM2.5 ($\Delta_4\% = +36.5 \pm 12.7$, $t_4 = 3.5$; $\Delta_5 = +23.6 \pm 10.8$, $t_5 = 2.5$). 2 suspected cases for PM10 and 1 for PM2.5 were also observed ($\Delta_6\% = +16.4 \pm 9.7$, $t_6 = 1.8$; $\Delta_7\% = +14.8 \pm 8.6$, $t_7 = 1.7$; $\Delta_8\% = +15.2 \pm 11.1$, $t_8 = 1.5$). Therefore, the first lockdown caused no significant changes in PM concentrations in 38.5% of cases and significant or suspected increases in 61.5% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 92.3% of cases; only 1 significant change were observed for PM10 ($\Delta_9\% = -13.3 \pm 5.3$, $t_9 = 2.4$).

3.1.2 Province of Brescia

The province of Brescia has about 1.264 million inhabitants and extends for 4,786 km². Here there are 9 measuring stations: 6 for PM10 and 3 for PM2.5. Only 1 significant change in PM10 average concentration was observed during period 1 ($\Delta\% = +23.1 \pm 10.5$, $t = 2.4$). During period 2, 1 significant change in average PM2.5 concentration ($\Delta_1\% = -10.5 \pm 5.5$, $t_1 = 2.3$) and 1 suspected change in average PM10 concentration ($\Delta_2\% = -7.5 \pm 5.7$, $t_2 = 1.5$) were observed. Therefore, the first lockdown caused no significant changes in PM concentrations in 88.9% of cases and significant increases in 11.1% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 77.8% of cases and significant or suspected decreases in 22.2% of cases.

3.1.3 Province of Como

The province of Como has about 600,000 inhabitants and extends for 1,279 km². Here there are 4 measuring stations: 3 for PM10 and 1 for PM2.5. During period 1, 2 significant changes in average PM10 concentrations were observed ($\Delta_1\% = +25.0 \pm 7.8$, $t_1 = 3.0$; $\Delta_2\% = +22.4 \pm 7.0$, $t_2 = 2.7$); one suspected change in average PM2.5 concentrations was also found ($\Delta_3\% = +11.8 \pm 6.6$, $t_3 = 1.7$). No significant nor suspected changes in PM10 or 2.5 concentrations were observed during period 2. Therefore, the first lockdown caused no significant changes in PM concentrations in 25.0% of cases and significant or suspected increases in PM concentrations in 75.0% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations.

3.1.4 Province of Cremona

The province of Cremona has about 360,000 inhabitants and extends for 1,770 km². Here there are 10 measuring stations: 6 for PM10 and 4 for PM2.5. During period 1, 3 significant changes were observed for average concentrations of PM10 ($\Delta_1\% = +33.7 \pm 12.2$, $t_1 = 3.3$; $\Delta_2\% = +17.1 \pm 7.6$, $t_2 = 2.0$; $\Delta_3\% = 16.2 \pm 7.8$, $t_3 = 1.9$) and 1 for average concentrations of PM2.5 ($\Delta_4 = +28.6 \pm 7.3$, $t_4 = 3.4$). 2 suspected cases for PM10 and 1 for PM2.5 were also observed ($\Delta_5\% = +12.7 \pm 8.0$, $t_5 = 1.8$; $\Delta_6\% = +14.2 \pm 8.3$, $t_6 = 1.6$; $\Delta_7\% = +12.6 \pm 7.7$, $t_7 = 1.5$). During period 2, only 1 significant change in average PM2.5 concentrations has been observed ($\Delta_1\% = +12.9 \pm 5.6$, $t_1 = 2.2$). Therefore, the first lockdown caused no significant changes in PM concentrations in 30.0% of cases and significant or suspected increases in PM concentrations in 70.0% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 90.0% of cases and significant increases in 10.0% of cases.

3.1.5 Province of Lecco

The province of Lecco has about 340,000 inhabitants and extends for 806 km². Here there are 8 measuring stations: 5 for PM10 and 3 for PM2.5. During period 1, 3 significant changes were observed for average concentrations of PM10 ($\Delta_1\% = +46.3 \pm 6.5$, $t_1 = 3.9$; $\Delta_2\% = +27.9 \pm 7.6$, $t_2 = 2.6$; $\Delta_3\% = 34.6 \pm 6.3$, $t_3 = 3.4$) and 2 for average concentrations of PM2.5 ($\Delta_4\% = +30.4 \pm 7.8$, $t_4 = 2.6$; $\Delta_5\% = +29.1 \pm 8.0$, $t_5 = 2.5$). During period 2, 2 significant changes in average PM10 concentrations ($\Delta_1\% = +11.0 \pm 4.9$, $t_1 = 1.9$; $\Delta_2\% = -18.2 \pm 7.1$, $t_2 = 3.5$) and 1 for PM2.5 ($\Delta_2\% = -13.1 \pm 6.6$, $t_2 = 2.5$) were observed. 1 suspected change was also found ($\Delta_3\% = +9.5 \pm 5.2$, $t_3 = 1.6$). Therefore, the first lockdown caused no significant changes in PM concentrations in 37.5% of cases and significant or suspected increases in PM concentrations in 62.5% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 50.0% of cases, significant decreases in 25.0% of cases, and significant or suspected increases in the remaining 25.0%.

3.1.6 Province of Lodi

The province of Lodi has about 230,000 inhabitants and extends for 783 km². Here there are 9 measuring stations: 7 for PM10 and 2 for PM2.5. During period 1, significant changes were observed in all measuring stations of PM10 ($\Delta_1\% = +21.9 \pm 8.2$, $t_1 = 1.9$; $\Delta_2\% = +29.4 \pm 8.6$, $t_2 = 2.3$; $\Delta_3\% = 25.9 \pm 7.4$, $t_3 = 2.5$; $\Delta_4\% = +26.0 \pm 8.2$, $t_4 = 2.2$; $\Delta_5\% = +23.6 \pm 8.5$, $t_5 = 2.0$; $\Delta_6\% = 27.5 \pm 7.7$, $t_6 = 2.4$; $\Delta_7 = +25.6 \pm 7.5$, $t_7 = 2.4$) and PM2.5 ($\Delta_8\% = +27.1 \pm 9.8$, $t_8 = 1.9$; $\Delta_9\% = +19.4 \pm 7.8$, $t_9 = 1.9$). During period 2, 1 significant and 1 suspected changes in average concentrations of PM10 ($\Delta_1\% = +12.4 \pm 4.8$, $t_1 = 2.2$; $\Delta_2\% = +8.3 \pm 4.6$, $t_2 = 1.6$) and 1 suspected change in average concentrations of PM2.5 ($\Delta_1\% = +10.9 \pm 6.2$, $t_1 = 1.5$) were observed. Therefore, the first lockdown caused significant increases in PM concentrations in 100.0% of cases. Considering

period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 66.7% of cases and significant or suspected increases in the remaining 33.3%.

3.1.7 Province of Mantova

The province of Mantova has about 413,000 inhabitants and extends for 2,341 km². Here there are 13 measuring stations: 9 for PM10 and 4 for PM2.5. During period 1, 5 significant and 1 suspected changes were observed for average concentrations of PM10 $(\Delta_1\% = +28.4 \pm 7.8, t_1 = 2.5; \Delta_2\% = +25.2 \pm 8.1, t_2 = 2.1; \Delta_3\% = 24.8 \pm 7.9, t_3 = 2.2; \Delta_4\% = +36.3 \pm 8.1, t_4 = 2.8; \Delta_5\% = +30.0 \pm 8.3, t_5 = 2.3; \Delta_6\% = +22.6 \pm 9.1, t_6 = 1.8)$. During period 2, 1 significant and 2 suspected changes in PM10 concentrations were recorded $(\Delta_1\% = +13.9 \pm 4.6, t_1 = 2.5; \Delta_2\% = +8.2 \pm 4.6, t_2 = 1.6, \Delta_3\% = 10.1 \pm 4.8, t_3 = 1.8)$. Therefore, the first lockdown caused no significant changes in PM concentrations in 53.8% of cases and significant or suspected increases in the remaining 46.2%. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 84.6% of cases and significant or suspected increases in the remaining 15.4%.

3.1.8 Province of Milano

The province of Milano has about 3.26 million inhabitants and extends for 1,575 km². Here there are 13 measuring stations: 10 for PM10 and 3 for PM2.5. During period 1, 5 significant changes were observed for average concentrations of PM10 $(\Delta_1\% = +24.3 \pm 11.2, t_1 = 2.4; \Delta_2\% = +26.4 \pm 11.4, t_2 = 2.5; \Delta_3\% = 27.9 \pm 12.4, t_3 = 2.7; \Delta_4\% = -34.2 \pm 11.4, t_4 = 3.7; \Delta_5\% = -21.8 \pm 8.4, t_5 = 2.2)$ and 1 for average concentrations of PM2.5 $(\Delta_6\% = +24.8 \pm 13.5, t_6 = 2.1)$. During period 2, 4 significant changes were observed for average concentrations of PM10 $(\Delta_1\% = +10.8 \pm 6.1, t_1 = 1.9; \Delta_2\% = +11.3 \pm 5.9, t_2 = 2.0; \Delta_3\% = -20.1 \pm 3.8, t_3 = 4.9; \Delta_4\% = +15.3 \pm 5.9, t_4 = 2.8)$ and 1 for average concentrations of PM2.5 $(\Delta_5\% = +18.0 \pm 7.6, t_5 = 2.5)$; 1 suspected change in average PM10 concentrations were also recorded $(\Delta_6\% = +8.3 \pm 5.7, t_6 = 1.5)$. Therefore, the first lockdown caused no significant changes in PM concentrations in 53.8% of cases, significant increases in 30.8% of cases, and significant decreases in 15.4% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 66.7% of cases and significant or suspected increases in the remaining 33.3% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 53.8% of cases, significant or suspected increases in 38.5% of cases, and significant decreases in 7.7% of cases.

3.1.9 Province of Monza e Brianza

The province of Monza e Brianza has about 872,000 inhabitants and extends for 406 km². Here there are 4 measuring stations: 3 for PM10 and 1 for PM2.5. During

period 1, 3 significant changes for average concentrations of PM10 ($\Delta_1\% = +15.7 \pm 7.3$, $t_1 = 1.9$; $\Delta_2\% = +28.0 \pm 7.2$, $t_2 = 3.0$; $\Delta_3\% = 20.5 \pm 6.8$, $t_3 = 2.8$) and 1 for average concentrations of PM2.5 ($\Delta_4\% = +33.0 \pm 7.2$, $t_4 = 3.3$) were observed. During period 2, 1 significant change in average concentrations of PM10 ($\Delta_1\% = +12.9 \pm 5.4$, $t_1 = 2.2$) and 1 suspected change in average concentration of PM2.5 ($\Delta_2\% = +10.0 \pm 5.2$, $t_2 = 1.8$) were recorded. Therefore, the first lockdown caused significant increases in PM concentrations in 100.0% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 50.0% of cases and significant or suspected increases in the remaining 50.0%.

3.1.10 Province of Pavia
The province of Pavia has about 548,000 inhabitants and extends for 2,969 km². Here there are 11 measuring stations: 6 for PM10 and 5 for PM2.5. During period 1, 3 significant changes in average concentrations of PM10 ($\Delta_1\% = +24.3 \pm 7.4$, $t_1 = 2.3$; $\Delta_2\% = +24.3 \pm 7.4$, $t_2 = 2.3$; $\Delta_3\% = 20.4 \pm 7.7$, $t_3 = 2.0$) and 3 in average concentrations of PM2.5 ($\Delta_4\% = 32.1 \pm 6.4$, $t_4 = 3.2$; $\Delta_5\% = 39.7 \pm 7.7$, $t_5 = 3.0$; ($\Delta_6\% = 35.8 \pm 6.5$, $t_6 = 3.4$); 2 suspected changes in average concentrations of PM10 and 1 suspected changes in average concentrations of PM2.5 were also found ($\Delta_7\% = +14.7 \pm 7.4$, $t_7 = 1.6$; $\Delta_8\% = -13.0 \pm 9.4$, $t_8 = 1.7$; $\Delta_9\% = 20.5 \pm 10.3$, $t_9 = 1.5$). During period 2, 1 significant change in average concentrations of PM2.5 ($\Delta_1\% = +18.6 \pm 5.3$, $t_1 = 2.6$) and 3 suspected change in average concentrations of PM10 ($\Delta_2\% = -11.0 \pm 7.4$, $t_2 = 1.8$; $\Delta_3\% = +11.3 \pm 6.4$, $t_3 = 1.5$; $\Delta_4\% = +12.2 \pm 6.6$, $t_4 = 1.5$). Therefore, the first lockdown caused significant or suspected increases in PM concentrations in 72.7% of cases, suspected decreases in 9.1% of cases, and no significant or suspected changes in 18.2% of cases. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 63.6% of cases, significant or suspected increases in 27.3% of cases, and suspected decreases in the remaining 9.1%.

3.1.11 Province of Sondrio
The province of Sondrio has about 182,000 inhabitants and extends for 3,196 km². Here there are 6 measuring stations: 4 for PM10 and 2 for PM2.5. During period 1, 3 significant changes for average concentrations of PM10 ($\Delta_1\% = +21.6 \pm 6.4$, $t_1 = 2.5$; $\Delta_2\% = +17.7 \pm 5.5$, $t_2 = 2.5$; $\Delta_3\% = 27.0 \pm 6.4$, $t_3 = 3.0$) and 1 for average concentration of PM2.5 ($\Delta_4\% = +20.5 \pm 5.9$, $t_4 = 2.7$); 1 suspected change has occurred for average concentrations of PM10 ($\Delta_5\% = +14.0 \pm 6.8$, $t_5 = 1.7$). During period 2, no significant or suspected changes were recorded. Therefore, the first lockdown caused significant increases in PM concentrations in 83.3% of cases and no significant changes in the remaining 16.7%. Considering period 2, no significant or suspected changes were observed.

3.1.12 Province of Varese

The province of Sondrio has about 890,000 inhabitants and extends for 1,198 km². Here there are 6 measuring stations: 4 for PM10 and 2 for PM2.5. During period 1, 1 significant change in average concentrations of PM10 ($\Delta_1\% = +21.6 \pm 7.6$, $t_1 = 2.1$) and 1 in average concentrations of PM2.5 ($\Delta_2\% = +27.9 \pm 7.9$, $t_2 = 2.4$) were observed. During period 2, only 1 suspected change in average PM2.5 concentrations was found ($\Delta_1\% = +11.9 \pm 6.4$, $t_1 = 1.7$). Therefore, the first lockdown caused no significant changes in PM concentrations in 66.7% of cases and significant increases in the remaining 33.3%. Considering period 2, all COVID-19 restrictions caused no significant changes in PM concentrations in 83.3% of cases and significant or suspected increases in the remaining 50.0%.

3.2 Consumer behavior

3.2.1 Home food delivery services
The web interest of Italian citizens towards home food delivery services followed a periodic trend during 2018 and 2019 (Figure 1). During the first lockdown and pre-lockdown (20 February - 18 May, 2020), the queries "just eat, uber eats, deliveroo, foodracers and glovo" recorded a significant percentage increase with respect the same period in 2019 ($\Delta_1\% = +30.6 \pm 6.5$, $t_1 = 4.8$; $\Delta_2\% = +80.7 \pm 58.0$, $t_2 = 2.9$; $\Delta_3\% = +38.7 \pm 25.1$, $t_3 = 5.8$; $\Delta_4\% = +269.2 \pm 60.5$, $t_4 = 5.2$; $\Delta_5\% = +173.0 \pm 44.8$, $t_5 = 8.3$). It must be pointed out that three of these queries underwent a significant increase between 2018 and 2019 (uber eats, deliveroo, and glovo); however, the highest RSV peak detected in spring 2020, during the first lockdown, was not present in 2018 and 2019 series (Figure 1). Furthermore, such trends have been heavily influenced also in the months following the first wave of COVID-19 (Figure 1). Significant correlations were observed between RSVs and COVID-19 total cases and total deaths ($r_1 = .78$, $P < .0001$; $r_2 = .72$, $P = .0003$); nevertheless, such correlations were also identified between the same web queries during 2019 and COVID-19 total cases and total deaths.

Figure 1. Italian citizens' web interest in home food delivery services from November 2017 to November 2020.

3.2.2 Online food shopping services and online supermarkets

The web interest of Italian citizens towards food shopping services and online supermarkets underwent a significant percentage increase during the first lockdown compared to the trend during 2018 and 2019 (Figure 2). The queries "esselunga online, spesa online (online food shopping) + supermercato online (online supermarket) + supermercati online (online supermarkets), carrefour online, conad online, iperal online" recorded the following percentage increases:

.

Moreover, the interest in these queries remained relatively high during the rest of 2020 compared to those of the previous two years, growing again with the arrival of the second period of restrictions (Figure 2). Significant hidden correlations were observed between RSVs and COVID-19 total cases and total deaths

; nevertheless, such correlations were also identified between the same web queries during 2019 and COVID-19 total cases and total deaths.

Figure 2. Italian citizens' web interest in food shopping delivery services from November 2017 to November 2020.

3.2.3 Online shopping

The web interest of Italian citizens towards home food delivery services followed a periodic trend during 2018 and 2019 (Figure 3). During the first lockdown and pre-lockdown (20 February - 18 May, 2020), the query "amazon" recorded a significant percentage increase with respect the same period in 2019

. This increase continued, to varying degrees, throughout the spring and early summer (Figure 3). The query "ebay" also underwent an anomalous rise during the first lockdown period, reaching the annual RSV peak despite what happened in 2018 and 2019 (Figure 3). Only low and non significant correlations were identified between such queries and COVID-19 total cases and deaths   .

3.2.4 Smart working and distance learning

The web interest of Italian citizens towards smart working and distance learning platforms such as DAD, Teams, Zoom, Meet, and Skype, underwent a drastic surge during and after the first lockdown (during period 1 and   during period 2). Among all regions, Calabria is the one that has shown the highest interest in these technologies with an

. The other regions had an   in the interval   i.e. they showed a more similar interest   .

Figure 3. Italian citizens' web interest in shopping delivery services from November 2017 to November 2020.

4. Discussion

Several studies have been conducted on the impact of the lockdown on Lombardy's air quality (Cameletti, 2020, Piccoli et al., 2020, Putaud et al., 2020, and Collivignarelli et al., 2020). However, not all results are fully concordant: in fact, while Putaud et al. and claim that they have not observed any significant change in PM10 concentrations in Milan, Collivignarelli et al. argue that the lockdown had a positive effect on the latter as well as on PM2.5 and other air pollutants. In this context, Cameletti highlights how a reduction in particulate matter emissions does not necessarily imply a reduction in its concentrations, underlining the complexity in the analysis of these types of phenomena. Such evidence could help explain the discrepancy found between previous authors and between the various reports from environmental protection agencies and universities. But not only: these differences could also be explained by the different approaches adopted in the aforementioned studies, since some have used predictive models inherent exclusively to 2020 while others have compared the concentrations of pollution in the current year with those of previous years. This paper is among the latter, looking for the differences in average concentrations of PM10 and PM2.5 between specific periods during 2020 and 2019. Piccoli et al. showed that the months of February and March 2019 and 2020 presented similar temperatures in Milan, making them suitable for a comparative analysis. This supports the assumption of the absence of unusual weather regimes. However, a widespread reduction in wind speed was highlighted in this paper which certainly contributed to the increase in concentrations of particulate matter across the Lombardy region. Moreover, non-uniform and significant variations

in temperature and relative humidity may have further biased the estimate of particulate matter emissions in an unpredictable manner. Nonetheless, the most important result of this paper is that even a total lockdown strategy is absolutely insufficient to reduce the concentrations of PM10 and PM2.5 in a sustainable way. This again calls into question the role of motor vehicles in the production of this type of air pollution. Abu-Allaban et al. showed that the fractional contributions of motor vehicles to ambient PM in urban areas in the USA were in the ranges 20-76% and 35-92% for PM2.5 and PM10, respectively. The Italian National Research Council (CNR)[11] measured a reduction in traffic of 48-60% in Italian cities which was obviously not followed by an equal reduction in particulates, confirming the great variance of the above ranges. Therefore, this research suggests that industries, private and household activities, and ambiental factors, could contribute very significantly to air pollution in Lombardy. Alongside that, the demand for online delivery and shopping services has dramatically increased both during and after lockdowns. Such e-commerce services, including Amazon, when providing very fast delivery times, can have a very negative environmental impact: in fact, there is not only the pollution due to the emissions of the transport vehicles but also the plastic packaging with which the products, shipped by the companies, are typically packaged[12,13] (CNN, 2019, SCSC, 2019). In this regard, Amazon says it is planning a sustainable delivery strategy[14] (Amazon, 2020); never before is it important for the company to keep its word. Besides that, it is also essential that states and organizations all over the world impose, in their administrative policies, more stringent and effective criteria for monitoring environmental impact. Anyway, since the growing web interest has been matched by a real increase in orders placed by Italian users (AGI, 2020)[15], Google Trends has proved to be an effective infoveillance tool for this type of scientific survey. Finally, smart working and distance learning could be very valid alternatives in terms of environmental impact as they do not involve travelling or excessive use of materials such as textbooks, notepads, notebooks, chalks or markers and others. However, this leads to an increase in household consumption of electricity and, during the winter, of heat; nonetheless these are also present in work structures, private homes in Italy often do not have systems of the same energy class (ABC Costruzioni, 2018). Moreover, there are many challenges to be faced in order to take this path (The Planet Mark, 2016, Hodder, 2020, Sarti et al., 2020) and the debate between parents, teachers and psychologists about online lessons is still heated (Daniel, 2020, Mukhtar et al., 2020, Abbasi et al., 2020, Abuhammad, 2020). What is certain is that the population has had to forcibly learn to use some of the most modern technologies and techniques of file sharing, databases, and online communication software and tools; thus, in the future, it will be easier and less impacting on an emotional-psychological level to

---

[11] https://www.cnr.it/it/comunicato-stampa/9702/l-inquinamento-in-italia-durante-il-lockdown

[12] https://edition.cnn.com/2019/07/15/business/fast-shipping-environmental-impact/index.html

[13] https://sustainability.aboutamazon.com/environment/packaging-and-products

[14] https://sustainability.aboutamazon.com/environment/packaging-and-products

[15] https://www.agi.it/economia/news/2020-04-22/fase-2-coronavirus-amazon-spedizioni-8411103/

insert smart working and online learning methods in everyday life, looking for the right compromise between the psychosocial health safeguard and environmental protection. Finally, the correlations found between COVID-19 cases and deaths, and web searches related to home delivery services in 2020 and 2019, could prove how the virus has spread easier in regions with more frequent commercial relationships.

4.1 Strengths and limitations
As far as the author knows, this is the first study that compares PM10 and PM2.5 concentrations of 2019 and 2020 in Lombardy including data from all the measuring stations of all its provinces also quantifying the meteorological differences of the latter. Furthermore, this study demonstrates a substantial reliability of Google Trends in estimating the demand for e-commerce and home delivery services of Italian web users as well as the ineffectiveness of lockdowns in the reduction of PM10 and PM2.5 concentrations in the most active region of Italy. However, some limitations must be considered: the search for statistical correlations cannot provide details on any possible causal nature of the phenomena investigated. The web interest of Google users is not certain to reflect the general interest of all Italian citizens. In some provinces, the number of measuring stations could be insufficient to represent in detail the overall trend of PM10 and PM2.5 concentrations. Finally, some weather differences may have biased the comparison between the particulate emissions of 2019 and 2020.

5. Conclusions

This study shows that even a forced total lockdown strategy, capable of reducing traffic in Italian cities by 48-60%, was absolutely insufficient to reduce the concentrations of PM10 and PM2.5 in the Lombardy region. Environmental factors, industries, and household activities, can therefore have non-marginal or even preponderant roles in the production of this type of pollution. At the same time, COVID-19 has shown that citizens and the government are not yet ready to manage policies that support smart working or distance learning. Furthermore, the increase in demand for home delivery services and online shopping during and after the lockdowns must draw further attention from government authorities on the environmental issue. In conclusion, we can state that COVID-19 has drastically changed many of the habits of Italian citizens. Among the positive aspects of this change there was the learning of new technological methods and tools for online collaboration and remote communication via the Internet. This can be a starting point for the integration of policies aimed at preserving both the mental and physical health of citizens and the environment on which the latter depends.

References

1. World Health Organization (WHO, 2020). Impact of COVID-19 on people's livelihoods, their health and our food systems. Accessed: December 5, 2020. https://www.who.int/news/item/13-10-2020-impact-of-covid-19-on-people's-livelihoods-their-health-and-our-food-systems

2. Figueroa, C.A., and Aguilera, A. (2020). The Need for a Mental Health Technology Revolution in the COVID-19 Pandemic. Front. Psychiatry 11:523. doi: 10.3389/fpsyt.2020.00523

3. Francisco, R., Pedro, M., Delvecchio, E., Espada, J.P., Morales, A., Mazzeschi, C., and Orgilés, M. (2020). Psychological Symptoms and Behavioral Changes in Children and Adolescents During the Early Phase of COVID-19 Quarantine in Three European Countries. Front. Psychiatry 11:570164. doi: 10.3389/fpsyt.2020.570164

4. Worldometer (2020). COVID-19 coronavirus pandemic. Accessed: December 5, 2020.
   https://www.worldometers.info/coronavirus/

5. FASDA Fund. Gli effetti della quarantena sull'ambiente (Italian website). Accessed: December 5, 2020.
   https://www.fasda.it/effetti-quarantena-ambiente/

6. Rovetta, A., and Castaldo, L. (2020). Relationships between Demographic, Geographic, and Environmental Statistics and the Spread of Novel Coronavirus Disease (COVID-19) in Italy. Cureus 12(11): e11397. doi:10.7759/cureus.11397.

7. Rovetta, A., and Castaldo, L. (2020). The Impact of COVID-19 on Italian Web Users: A Quantitative Analysis of Regional Hygiene Interest and Emotional Response. Cureus 12(9): e10719. doi:10.7759/cureus.10719

8. Jimenez, A., Estevez-Reboredo, R.M., Santed, M.A., and Ramos, V. COVID-19 symptom Google search surges, precede local incidence surges: evidence from Spain. J Med Internet Res. 2020 Oct 26. doi: 10.2196/23518

9. Google Trends. Accessed: December 4, 2020.
   https://trends.google.com/trends/

10. Regional Agency for Environmental Protection (ARPA). Accessed: from November 20 to November 27, 2020.
    https://www.arpalombardia.it/Pages/Aria/Richiesta-Dati.aspx

11. Il meteo (Italian website). Accessed: November 20, 2020.

https://www.ilmeteo.it/portale/archivio-meteo/Lombardia

12. Department of Civil Protection. Accessed: December 1, 2020.
https://github.com/pcm-dpc/COVID-19

13. National Institute of Statistics (ISTAT). Accessed. December 2, 2020.
https://www.istat.it/

14. AdminStat. Accessed: December 2, 2020.
https://ugeo.urbistat.com/AdminStat/it

15. Taylor, J.R. (1997). An Introduction to Error Analysis: The Study of Uncertainties in Physical Measurements. Univ Science Books. ISBN-13: 978-0935702422

16. Tabachnick, B.G., and Fidell, L.S. (1996). Using multivariate statistics (3rd ed.). New York: HarperCollins.

17. Kwak, S.G., and Kim J.H. (2017). Central limit theorem: the cornerstone of modern statistics. Korean Journal of Anesthesiology 2017;70(2):144-156. doi: 10.4097/kjae.2017.70.2.144

18. Rovetta, A. (2020). Raiders of the Lost Correlation: A Guide on Using Pearson and Spearman Coefficients to Detect Hidden Correlations in Medical Sciences. Cureus 12(11): e11794. doi:10.7759/cureus.11794

19. Amrhein, V., Korner-Nievergelt, F., Roth, T (2017). The earth is flat (p > 0.05): significance thresholds and the crisis of unreplicable research. PeerJ. 2017 Jul 7;5:e3544. doi: 10.7717/peerj.3544. eCollection 2017

20. Cameletti, M (2020). The Effect of Corona Virus Lockdown on Air Pollution: Evidence from the City of Brescia in Lombardia Region (Italy). Atmosp. Environ. 2020, 239, 117794. doi: 10.1016/j.atmosenv.2020.117794

21. Piccoli, A., Agresti, V., Balzarini, A., Bedogni, M., et al. (2020). Modeling the E□ect of COVID-19 Lockdown on Mobility and NO2 Concentration in the Lombardy Region. Atmosphere 2020, 11(12), 1319. doi: 10.3390/atmos11121319

22. Putaud, J.-P., Pozzoli, L., Pisoni, E., Martins Dos Santos, et al. (2020). Impacts of the COVID-19 lockdown on air pollution at regional and urban background sites in northern Italy, Atmos. Chem. Phys. Discuss. doi: 10.5194/acp-2020-755, in review, 2020.

23. Collivignarelli, M.C., Abbà, A., Bertanza, G., Pedrazzani, R., et al (2020). Lockdown for CoViD-2019 in Milan: What are the effects on air quality? Sci. Total Environ. 2020, 732, 139280. doI: 10.1016/j.scitotenv.2020.139280

24. Abu-Allaban, M., Gillies, J.A., Gertler, A.W., Clayton, R., et al. (2006). Motor vehicle contributions to ambient PM10 and PM2.5 at selected urban areas in the USA. Environ Monit Assess 2007 Sep;132(1-3):155-63. doi: 10.1007/s10661-006-9511-3.

25. Consiglio Nazionale delle Ricerche (CNR, 2020). L'inquinamento in Italia durante il lockdown (Italian website). Accessed: December 10, 2020. https://www.cnr.it/it/comunicato-stampa/9702/l-inquinamento-in-italia-durante-il-lockdown

26. CNN Business (2020). America's addiction to absurdly fast shipping has a hidden cost. Accessed: December 11, 2020. https://edition.cnn.com/2019/07/15/business/fast-shipping-environmental-impact/index.html

27. Supply Chain Solutions Center (SCSS, 2019). Packaging waste 101: the problem. Accessed: December 11, 2020. https://supplychain.edf.org/resources/sustainability-101-packaging-waste-the-problem/

28. Agenzia Giornalistica Italiana (AGI, 2020). Come lavora Amazon durante l'emergenza coronavirus (Italian website). Accessed: December 11, 2020. https://www.agi.it/economia/news/2020-04-22/fase-2-coronavirus-amazon-spedizioni-8411103/

29. ABC Costruzioni srl (2018). Classe energetica: qual è la situazione delle case italiane (Italian website). https://abccostruzioni.it/classe-energetica-qual-e-la-situazione-delle-case-italiane/

30. The Planet Mark, Eden Project (2016). Smartworking and sustainability. Accessed: December 12, 2020. https://www.theplanetmark.com/smartworking-and-sustainability/

31. Hodder, A. (2020). New Technology, Work and Employment in the era of COVID19: reflecting on legacies of research. New Technol Work Employ. 2020 Aug 3 : 10.1111/ntwe.12173. doi: 10.1111/ntwe.12173

32. Sarti, D., De Salvatore, M., Gazzola, S., Pantaleoni, C., et al. (2020). So far so close: an insight into smart working and telehealth reorganization of a Language and Learning Disorders Service in Milan during COVID-19

pandemic. Neurol Sci. 2020 May 29 : 1–4. Neurol Sci. 2020 May 29 : 1–4. doi: 10.1007/s10072-020-04481-8

33. Daniel, J. (2020). Education and the COVID-19 pandemic. Prospects (Paris). 2020 Apr 20 : 1–6. doi: 10.1007/s11125-020-09464-3

34. Mukhtar, K., Javed, K., Arooj, M., Sethi, A. (2020). Advantages, Limitations and Recommendations for online learning during COVID-19 pandemic era. Pak J Med Sci. 2020 May; 36(COVID19-S4): S27–S31. doi: 10.12669/pjms.36.COVID19-S4.2785

35. Abbasi, S., Ayoob, T., Malik, A., Memon, S.I. (2020). Perceptions of students regarding E-learning during Covid-19 at a private medical college.

36. Abuhammad, S. (2020). Barriers to distance learning during the COVID-19 outbreak: A qualitative review from parents' perspective. Heliyon. 2020 Nov; 6(11): e05482. doi: 10.1016/j.heliyon.2020.e05482