Vitamins D_2 and D_3 have overlapping but different effects on human gene expression revealed through analysis of blood transcriptomes in a randomised double-blind placebo-controlled food-fortification trial

Louise R. Durrant[,1, 2], Giselda Bucca[a,b, 1,3], Andrew Hesketh[b, 1], Carla Möller-Levet[a], Laura Tripkovic[a], Huihai Wu[a], Kathryn H. Hart[a], Ruan M. Elliott[a], John C. Mathers[c], Susan A. Lanham-New[a] & Colin P. Smith[a,b, 3*]

[a] Department of Nutritional Sciences, School of Biosciences and Medicine, Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK
[b] School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, UK
[c] Human Nutrition Research Centre, Population Health Sciences Institute, Newcastle University, UK

[1] These authors contributed equally
[2] Current address: Yakult UK Ltd., Anteros, Odyssey Business Park, West End Road, South Ruislip, HA4 6QQ; formerly known as Louise Wilson.
[3] Current address: School of Pharmacy and Biomolecular Sciences, University of Brighton, Brighton, UK
*For correspondence: Prof Colin P. Smith, School of Pharmacy and Biomolecular Sciences, University of Brighton, Huxley Building, Moulsecoomb, Brighton, BN2 4GJ, UK; Tel. +44 (0)1273 642048. Email: genomics@cpsmith.net

Key words: Vitamin D_3, vitamin D_2, 25-hydroxyvitamin D, human transcriptome, immunity, ethnic differences, COVID-19

Abstract

For the first time, we report the influence of vitamin D_2 and vitamin D_3 on genome-wide gene expression in whole blood from healthy women representing two ethnic groups, white European and South Asian. In this randomised placebo-controlled trial, participants were given daily physiological doses (15 µg) of either vitamin D_2 or D_3 for 12 weeks and changes in the transcriptome were compared relative to the transcriptome at baseline. While there was some overlap in the repertoire of

differentially expressed genes after supplementation with each vitamin D source, most changes were specific to either vitamin D_3 or vitamin D_2, suggesting that each form of the vitamin may have different effects on human physiology. Notably, following vitamin D_3 supplementation, the majority of changes in gene expression reflected a down-regulation in the activity of genes, many encoding components of the innate and adaptive immune systems. These are consistent with the emerging concept that vitamin D orchestrates a shift in the immune system towards a more tolerogenic status. The profound differences in gene expression after supplementation with vitamin D_2 compared with vitamin D_3 warrant a more intensive investigation of the biological effects of the two forms of vitamin D on human physiology.

Author Contributions

CPS and SL-N conceived of the study. CPS, GB and SL-N supervised the project. LD, GB and LT carried out the experiments, AH, GB, CPS, LD, CM-L and HW analyzed and interpreted the transcriptome data, CPS, AH and GB wrote the first draft of the manuscript, AH generated the figures and all authors contributed to the final manuscript.

Significance statements

This study suggests that the influence of vitamins D_2 and D_3 on human physiology may not be the same, as deduced from differences in gene expression within whole blood.

South Asian participants were found to respond differently to vitamin D supplementation at the transcriptome level from white Europeans.

The differentially expressed immune pathways identified in this study are consistent with vitamin D orchestrating a more tolerogenic immune status and this could be relevant in the context of the severity of immune response to viral infections such as COVID-19.

Introduction

Vitamin D is a pro-hormone that is critical for good health. While vitamin D's function in maintaining the musculo-skeletal system is best understood, it fulfils myriad roles in human physiology, including maintaining a healthy immune system (Prietl et al., 2013, Martens et al., 2020). Vitamin D deficiency (defined as serum concentrations of 25-hydroxyvitamin D (25(OH)D) below 25 nmol/L) or inadequacy (defined as serum level of 25(OH)D below 50 nmol/L) is considered a global pandemic and a public health issue of great importance in the human population, especially in older people, individuals not exposed to sufficient sunlight and, importantly, ethnic minority groups with dark skin tone (Kohlmeier, 2020, Lanham-New et al., 2020). Vitamin D deficiency is considered to have many detrimental effects on human health which include increased risk of osteoporotic and stress fractures, increased risk of developing cardiovascular diseases and some cancers, and poor modulation of the immune system (Holick, 2007, Bouillon, 2018). Low vitamin D status is associated with higher mortality, including death from cancer (Zhang et al., 2019).

Vitamin D_3 (cholecalciferol) is produced in the skin by the action of ultraviolet B radiation from the sun. Following two successive hydroxylation steps, in the liver and kidney, respectively, the active form of the vitamin, 1,25-dihydroxyvitamin D_3 (1,25(OH)D_3) binds to the intracellular vitamin D receptor (VDR) and, in complex with the retinoid X receptor (RXR), the heterodimer regulates expression of hundreds of genes through the vitamin D response element (VDRE) (Long et al., 2015). Vitamin D can also be obtained from foods or as a supplement. Both the plant/fungus-derived vitamin D_2 (ergocalciferol) and the animal-derived D_3 forms are available as nutritional supplements. Both can be hydroxylated into their active forms and bind to the VDR with similar affinity, but there are differences in their catabolism and in their binding affinity to vitamin D binding protein (DBP), the major vitamin D transport protein in blood. Vitamin D_2 binds DBP with lower affinity and is catabolised faster (Hollis, 1984, Houghton and Vieth, 2006).

The functional equivalence, or otherwise, of vitamins D_2 and D_3 for human health has been a subject of much debate in recent years, with some authors suggesting that the two compounds have equal efficacy while others provide evidence that vitamin

D_3 increases circulating serum 25(OH)D concentration more efficiently than vitamin D_2 (Tripkovic et al., 2012, Wilson et al., 2017). Current guidance given to the general public by the US National Institutes of Health (NIH), the UK Department of Health and Social Care, and other government bodies around the world lack acknowledgement that the two forms of vitamin D are not considered equivalent or equally effective when used at nutritional (physiological) doses, while at high doses vitamin D_2 is less potent. The NIH state that most evidence indicates that vitamin D_3 increases serum 25(OH)D levels to a greater extent and maintains these higher levels longer than vitamin D_2, even though both forms are well absorbed in the gut (ODS, 2020).

Although vitamin D is best known for its role in maintaining bone health and calcium homeostasis, it also exerts a broad range of extra-skeletal effects on cellular physiology and on the immune system (Pilz et al., 2019, Schwalfenberg, 2011, Christakos et al., 2016, Rosen et al., 2012) and multiple studies have linked poor vitamin D status with the pathogenesis of immune mediated inflammatory diseases (Damoiseaux and Smolders, 2018). Vitamin D is essential for human health and it is recommended that individuals maintain serum concentrations of at least 25 nmol/L of 25(OH)D (the accepted biomarker of systemic vitamin D status), throughout the year and throughout the life course (SACN, 2016). However, it can be difficult to sustain such levels of serum 25(OH)D year-round from diet and sunlight alone, particularly amongst 'at-risk' groups, such as those with limited sun exposure or those with darker skin tone living in northern latitudes (above 40ºN).

In acute studies comparing the efficacy of vitamins D_2 and D_3 in raising serum 25(OH)D concentration, vitamin D_2 is less effective than D_3 when given as single bolus (Tripkovic et al., 2012). However, the findings from studies following the daily administration of D_2 or D_3 over longer time-periods are more equivocal, with clinical trials showing higher efficacy of D_3 (Zittermann et al., 2014, Tripkovic et al., 2017) or equal efficacy (Holick et al., 2008). In a recent meta-analysis, supplementation with vitamins D_2 and D_3 were found equally effective in raising 25(OH)D concentrations in infants up to 1 year of age (Zittermann et al., 2020). It should be noted that supplementation with one form of vitamin D can reduce circulating concentrations of the other form. For example, our recent comparative trial of D_2 versus D_3 revealed

that serum $25(OH)D_3$ concentration was reduced over the 12-week intervention in participants given vitamin D_2, compared with the average 12-week concentration in participants given placebo ((Tripkovic et al., 2017), see Fig. 1C, for example). The implications of such reciprocal depletion remain to be explored.

There is limited understanding of the effects of vitamin D supplementation on gene expression *in vivo* in humans because of the diversity of experimental designs used. This includes: (i) diverse sampling intervals, ranging from hours to years following vitamin D supplementation (Berlanga-Taylor et al., 2018, Neme et al., 2019); (ii) use of substantially different doses ranging from physiological (moderate) to supra-physiological doses; and (iii) relatively small numbers of participants were recruited so that most studies were considerably underpowered (Neme et al., 2019, Hossein-nezhad et al., 2013). Currently, there is no robust evidence, from *in vivo* human genome-wide expression analysis, about which specific cellular pathways are influenced by vitamin D supplementation. Moreover, the influence of vitamin D_2, as distinct from vitamin D_3, on gene expression in humans has not yet been evaluated, even though vitamin D_2 is used widely as a supplement and food fortificant.

We have addressed this gap in knowledge by investigating gene expression in a relatively large cohort of healthy white European and South Asian women who participated in a randomised double-blind placebo-controlled trial that compared the relative efficacy of vitamins D_2 and D_3 in raising serum $25(OH)D$ concentration. The study concluded that vitamin D_3 was superior to vitamin D_2 in raising serum $25(OH)D$ concentration (Tripkovic et al., 2017). As an integral part of the original study design, we also investigated gene expression in the participants over the 12-week trial period. This allowed us to examine the effects of vitamin D supplementation on the transcriptome and to determine whether these effects might differ depending on supplementation with vitamin D_2 compared with vitamin D_3. We therefore sampled blood at the beginning, middle and end of the 12-week intervention and quantified changes in the whole blood transcriptome. Surprisingly, we observed that the two vitamins exerted overlapping but different effects on the human blood transcriptome. Our findings support the hypothesis that the biological effects of vitamin D_2 and D_3 may differ in humans and suggest that a more comprehensive analysis of the biological effects of the two forms of vitamin D on human physiology is warranted. In

this context, it is important to note that a recent meta-analysis of vitamin D supplementation trials found that reduced cancer mortality was seen only with vitamin D_3 supplementation, not with vitamin D_2 supplementation, and indicated that all-cause mortality was significantly lower in trials with vitamin D_3 supplementation than in trials with vitamin D_2 (Zhang et al., 2019).

Results

The study volunteers were South Asian (SA) and white European (WE) women based in the United Kingdom, aged between 20 and 64 years (n=335; (Tripkovic et al., 2017)) and participation was for 12 weeks over the winter months (October to March, in Surrey, UK; latitude, 51°14' N). Serum measurements, including concentrations of total 25(OH)D, 25(OH)D_2 and 25(OH)D_3, were determined from fasting blood samples taken at the start (baseline defined as Visit 1 (V1)) and at weeks 6 (V2) and 12 (V3). To determine whether changes in serum 25(OH)D concentration as a result of vitamin D supplementation are associated with physiological changes at the level of gene expression, we investigated the whole blood transcriptome from a representative subset of the study participants (n=98) using total RNA isolated from V1 and V3 blood samples and Agilent Human Whole Genome 8× 60K v2 DNA microarrays (Fig. 1). Participants for transcriptome analysis were selected to provide comparable numbers between the placebo and the two vitamin D treatment groups, covering the full range of serum responses to supplementation within the D_2 and D_3 treatment groups, as judged from the measured changes in serum 25(OH)D_2 or 25(OH)D_3 concentration between V1 and V3 (**Supplementary Data File 1**).

Metadata for the transcriptome analysis cohort

Following microarray data quality control, transcriptome data were available for both the V1 and V3 samples of 97 study participants (Fig. 1b); the RNA from one participant failed quality control and was excluded. Similar to data reported previously for the entire study cohort (Tripkovic et al., 2017), the sub-set of participants in the present study showed increased concentrations of 25(OH)D_2 and 25(OH)D_3 only after supplementation with vitamin D_2 and vitamin D_3, respectively,

and higher total 25(OH)D in both treatment groups (Fig. 1c). In the vitamin D_3 treatment groups the mean 25(OH)D_3 serum levels rose by 59% (WE) and 116% (SA) over the 12-week intervention. Conversely, mean serum 25(OH)D_3 concentrations fell across the 12 weeks of the study in the placebo groups who did not receive additional vitamin D; 25(OH)D_3 baseline vitamin D levels in the SA ethnic group tended to be lower than for the WE group and dropped, respectively, by 23% and 29% relative to baseline V1 (Fig. 1c; **Supplementary Data File 1**). It is notable that, in the vitamin D_2 treatment group, serum 25(OH)D_3 concentration decreased to a greater extent over the 12-week period, by 52% (SA) and 53% (WE); the implications of this are considered further in the Discussion. Serum 25(OH)D_2 was low in the absence of specific supplementation, typically less than 5 nmol/L (152/162 samples analysed). Serum calcium concentration, appropriately adjusted for serum albumin concentration, was maintained within a normal clinical range across all sample groups, while the concentration of parathyroid hormone (PTH) was stable between V1 and V3 only within the SA placebo group, the WE vitamin D_2 and WE vitamin D_3 intervention groups. PTH concentrations showed an increase at V3 compared with V1 in the WE placebo group, but a decrease in both the vitamin D_2 and D_3 intervention groups in the SA ethnic cohort.

Effects of supplementation with either vitamin D_2 or D_3 on global gene expression

Filtering of the normalised microarray data to select for probes showing signals at least 10% above background in at least 41 (~20%) of the arrays yielded transcript abundance data from 20,662 probes corresponding to 17,588 different genomic features (12,436 of which were annotated with an ENTREZ gene identifier). Using the filtered data, we identified significant differences in the transcriptional responses occurring across the 12-week V1 to V3 period of the study *between* the vitamin D_2, vitamin D_3 and placebo treatment groups for each ethnic cohort and, in a separate analysis, to determine significant changes *within* each group between the V3 and V1 sampling points (Fig. 2; **Supplementary Data File 2**). The former was tested in a 'difference-in-difference' analysis according to the generalised null hypothesis [treatment1.V3 – treatment1.V1] - [treatment2.V3 – treatment2.V1] = 0 (Fig. 2a, rows 7-12), and the latter using treatment.V3 – treatment.V1 = 0 (Fig. 2a, rows 1-6).

No significant changes in probe signals (adj.P.Val<=0.05) were identified in the 'difference-in-difference' analysis for the data for the WE ethnic group, but five were found in the corresponding analysis of the SA cohort (Fig. 2a). Furthermore, difference-in-difference analysis of the entire study data, ignoring ethnicity, did not yield statistically significant changes between the D_2 or D_3 treatment groups and the placebo control (data not shown). We note that another 'difference-in-difference' study on the influence of vitamin D on gene expression (the BEST-D trial) failed to identify any significant changes in gene expression following long-term vitamin D treatment in a Caucasian cohort (Berlanga-Taylor et al., 2018). The observed $\log_2$ fold-change ($\log_2$FC) in signal abundances between V3 and V1 in the study typically fall well within +/- 1, and in such a context it is likely that the sample sizes used within the study restrict our ability to reliably identify individual gene expression responses using this highly conservative approach. We consider that inter-individual differences in human populations are likely to hinder detection of small statistically significant gene expression changes *across* treatment groups from microarray-derived data; however, statistically significant differences are detected when evaluating gene expression changes *within* an individual across time (as described below where we examine paired observations with two time points per subject). The five significant changes that were identified as occurring in response to treatment of the SA ethnic group with vitamin D_3, relative to the placebo, are summarised in **Supplementary Fig. 1**. These are driven primarily by the marked changes in the placebo group between the V1 and V3 sampling points, and may be associated with the sustained low serum 25(OH)D_3, or high PTH, concentrations observed in this group of subjects. One of these differentially expressed genes encodes the cAMP response element binding protein CREB1 (Fig. 2b) which is part of the cAMP-PKA-CREB signaling pathway in bone cells which contributes to the regulation of skeletal metabolism in response to PTH concentrations (Zhang et al., 2011).

In contrast to the difference-in-difference analysis, the direct *within*-group comparisons of transcript abundance measurements at V3 versus V1 per subject identified large numbers of significant changes in gene expression, most notably in the WE D_2, WE D_3 and WE placebo groups (Fig. 2a and c, **Supplementary Data File 2**). These significant gene expression changes in the WE ethnic cohort arising in

the vitamin D treatment groups are considered in more detail below. While there is some overlap in the groups of differentially expressed genes, only 13% of down-regulated differentially expressed genes (102 of 774) were common between the two vitamin D treatment groups, while 28% (216 of 774) and 59% (456 of 774) were uniquely down-regulated by D_2 and D_3, respectively (excluding those additionally down-regulated in the placebo group over the equivalent 12-week intervention)(Fig 2c). As previously noted above, the serum $25(OH)D_3$ concentration fell over the 12-week intervention period in the vitamin D_2-treated group; thus, changes in gene expression observed in this group could *a priori* either be due to the influence of vitamin D_2 itself, or could be attributable to the depletion of the endogenous $25(OH)D_3$ reserves. It was therefore important to have included comparative transcriptomic analysis of the placebo group in this study to distinguish gene expression changes resulting from vitamin D_2 *per se* from those changes arising due to the depletion of $25(OH)D_3$ levels in this vitamin D_2 treatment group.

It is notable that expression of a large number of genes was altered between the first and last visits (V1 to V3) in the non-treated placebo group. While some of these changes will be attributable to the effects of the vitamin D depletion observed in this group over the course of the study, there is significant over-representation of genes known to exhibit seasonal differences in gene expression (Dopico et al., 2015) (**Supplementary Fig. 2**). There is however no unique association of seasonally expressed genes with the data for the WE placebo group; a similar significant enrichment is also observed for the significant changes in gene expression identified in the WE D_2 and WE D_3 treatment groups (**Supplementary Fig. 2**) and therefore it is unlikely that this apparent seasonal effect exclusively reflects vitamin D-specific effects. There was no overlap between genes significantly down-regulated exclusively in the placebo group and those significantly up-regulated exclusively in the vitamin D treatment groups (or vice versa). Consistent with the 12-week period of the study, seasonal changes in gene expression are therefore a background feature of all the data collected, and is therefore a factor that needs to be considered when undertaking such time course studies.

In the group of probe signals with significantly reduced abundance in the V3 samples compared with V1 in the data for the vitamin D_2 and D_3 supplemented participants,

but not in the placebo participants, Probe A_33_P3275751, detecting *SEC14L1* gene transcription showed the largest decrease in response to both forms of vitamin D (Fig. 2d). This probe was originally designed to detect 'transcript variant 7' and hybridizes to a location at the 3' end of the *SEC14L1* locus, detecting several splice variants. High expression of *SEC14L1* is significantly associated with lymphovascular invasion in breast cancer patients, where transcript abundance correlates positively with higher grade lymph node metastasis, and poor prognostic outcome (Sonbul et al., 2018). Its overexpression is also frequent in prostate cancer where *SEC14L1* has been identified as a potential biomarker of aggressive progression of the disease (Agell et al., 2012, Burdelski et al., 2015).

Genes significantly down-regulated by both D_2 and D_3, but not placebo, are enriched for functions associated with immune responses

Given the relatively small sample sizes used in this study it is considered more appropriate to examine enrichment of cellular pathways among differentially expressed genes rather than focusing on changes in individual genes. Functional enrichment analysis of the genes represented by the probes specifically repressed by vitamins D_2 and D_3 (blue data points in Fig. 2d) suggests that both supplements have suppressive effects on the immune response in the WE ethnic group (Fig. 3 and **Supplementary Data File 3**). Indeed, a significant (p = 1.98E-06) protein-protein interaction network derived from the *Homo sapiens* medium confidence interactions curated in the STRING database (Szklarczyk et al., 2017b) is centred on a highly connected group of proteins associated with the innate immune response, neutrophil degranulation and leukocyte activation (Fig. 3a; all protein groups are detailed in **Supplementary Data File 3**). This includes the histone acetyltransferase, EP300, known to function as a transcriptional coactivator with VDR, the vitamin D receptor protein (Fetahu et al., 2014). Collectively, these findings are consistent with the emerging view that vitamin D exposure leads to a shift from a pro-inflammatory to a more tolerogenic immune status (Prietl et al., 2013, Martens et al., 2020).

Genes significantly up-regulated by both D_2 and D_3, but not placebo, include components of histone H4 and the spliceosome

A similar analysis of the probes specifically induced by D_2 and D_3 (red points in Fig. 2d) produced a significant (p = 4.99E-09) protein-protein interaction network enriched primarily for mitochondrial and ribosomal proteins, but also involving two subunits of histone H4 and SNRPD2, a core component of the SMN-Sm complex that mediates spliceosomal snRNP assembly (Fig. 3b).

Comparative functional enrichment analysis supports roles for vitamins D_2 and D_3 in the suppression of immunity, and in chromatin modification and spliceosome function

As a complementary approach to the functional enrichment analysis of specific subsets of genes whose expression is altered by supplementation with vitamin D_2 and D_3 but not by placebo treatment, a comparative functional enrichment analysis of all significant changes in each treatment group was performed (see Methods). This aimed to identify functional categories that are more extensively affected by vitamin D_2, or by vitamin D_3, or by both vitamins D_2 and D_3, than by the placebo, and took into consideration the changes occurring in the placebo treatment group across the 12 weeks of the study (Figs. 4 and 5; **Supplementary Data File 4**). Consistent with the earlier analysis, functional categories associated with immunity and immune response pathways are prominent among those genes repressed by vitamin D supplementation (Fig. 4), while mitochondrial, ribosomal and spliceosomal functions are prominent in the induced genes (Fig. 5). Although the two vitamin treatments share many common categories identified from this analysis, the results also highlight some differences between the respective effects of D_2 or D_3 supplementation. For example, 'histone exchange' is significant only in the vitamin D_2 up-regulated genes, and 'chromatin modifying enzymes' are significant only in the D_3 down-regulated genes.

Overall, the observed differences in gene expression from the blood transcriptome presented in this study suggest that the physiological effects of vitamin D_3 and D_2 may be dissimilar.

Weighted gene correlation network analysis (WGCNA) identifies modules of co-expressed genes that significantly correlate with serum markers of vitamin D supplementation in the WE and SA ethnic groups

WGCNA quantifies both the correlations between individual pairs of genes or probes across a data set, and also the extent to which these probes share the same neighbours (Langfelder and Horvath, 2008). The WGCNA process creates a dendrogram that clusters similarly abundant probes into discrete branches, and subsequent cutting of the dendrogram yields separate co-expression modules, representing putatively co-regulated sets of genes. The first principal component of the expression matrix of each module defines the expression profile of the eigengene for the module, and this can then be correlated with experimental metadata. By allowing phenotypic traits to be associated with relatively small numbers (tens) of module eigengenes, instead of thousands of individual variables (*i.e.* gene probes), WGCNA both alleviates the multiple testing problem associated with standard differential expression analysis and also directly relates experimental traits to gene expression data in an unsupervised way that is agnostic of the experimental design.

WGCNA was used to construct separate signed co-expression networks for the WE and SA ethnic groups as described in the Methods, and Pearson correlations between expression of the module eigengenes and serum $25(OH)D_2$, $25(OH)D_3$, total $25(OH)D$ and parathyroid hormone (PTH) concentrations were calculated (Fig. 6 and **Supplementary data files 6 - 8**). These results therefore ignore whether the data originates from the vitamins D_2, D_3 or placebo treatment groups and focus solely on the relationship between the serum metabolite concentrations and gene expression. A significant negative correlation was observed between $25(OH)D_2$ concentration and expression modules in the WE co-expression network that are enriched for immune-associated functions (midnight blue and pink modules in Fig. 6a). This is consistent with the results from the different analytical approaches (presented in Figs 3 and 4). Similarly, the only module exhibiting a significant positive correlation with $25(OH)D_2$ (and total $25(OH)D$ concentration), the red expression module, is enriched for GO categories associated with the ribosome and

mRNA processing (**Supplementary Data File 6**). No significant correlations with serum 25(OH)D$_3$ concentrations were detected in the WE cohort.

Interestingly, stronger and more numerous correlations were observed in the co-expression network for the SA cohort (Fig. 6b). In agreement with the WE network, an expression module significantly associated with the ribosome (black module in Fig. 6b) showed a general positive correlation with total 25(OH)D concentration whereas a module enriched for histone binding (turquoise) had a significant negative correlation with 25(OH)D concentration. Strikingly however, modules enriched for immune response functions (midnightblue, orange, purple, yellow in Fig. 6b) were consistently positively correlated with total 25(OH)D and usually 25(OH)D$_3$, but not 25(OH)D$_2$ concentrations in the SA network, and negatively correlated with PTH concentrations. This raises the possibility that vitamin D supplementation may exert different effects on the immune system depending on ethnicity of the individual, may indicate that PTH status has an influence on the outcome and may reflect physiological differences resulting from the low baseline vitamin D status in SA women. In this context we observed that PTH concentrations at the V1 sampling point in the SA cohort were elevated compared with those in the WE cohort (Fig. 1c).

Discussion

Vitamin D intake down-regulates expression of many genes including those encoding multiple pathways involved in innate and adaptive immunity

In the present study we shed light on the physiological effects of the two forms of vitamin D commonly used as food-fortificants or vitamin supplements. Using data from the same intervention study, previously we reported that vitamin D$_3$ supplementation is more effective than supplementation with vitamin D$_2$ in raising serum 25(OH)D concentration (Tripkovic et al., 2017). However, other studies have reported that the two forms of vitamin D are equally effective in raising serum 25(OH)D concentration (Holick et al., 2008, Zittermann et al., 2020). In the present study we examined gene expression changes *in vivo* following supplementation with either vitamin D$_2$ or D$_3$ (or placebo) in healthy women from the South East of the UK. Here we extend the results reported by Tripkovic et al. (2017) by conducting whole

genome transcriptome analysis in 97 of the original cohort of 335 women from two ethnic backgrounds, South Asian (SA) and white European (WE). We characterised gene expression changes in the human blood transcriptome in response to vitamin D_2 or D_3 supplementation over a 12-week treatment intervention over the winter months; the randomised groups were provided with physiological (15 µg) daily doses of vitamin D_2 or vitamin D_3 fortified foods, or identical foods without added vitamin D (placebo). We found extensive changes in gene expression in all three treatment groups, some of which were unique to the vitamin D_2-treated or D_3-treated groups, with the majority exhibiting downregulation of transcription over the 12-week intervention period. Our success in detecting statistically significant differential gene expression in this study stems partly from the longitudinal design where it was possible to evaluate statistical differences in transcript levels within each group (from baseline to 12-week measurement). This approach partly circumvents the problems with inter-individual differences in human studies which make it difficult to detect robust changes between different groups of individuals. Furthermore, the statistical analysis approach examined pairwise differences in gene expression from the V1 and V3 samples, which helps to reduce the impact of inter-individual differences on statistical significance.

This *in vivo* human transcriptome study has demonstrated long-term effects on gene expression. The majority of differentially expressed genes identified in this study were suppressed by vitamin D supplementation, and many of these encoded pathways were involved in immunity. Many of these observed gene expression changes are consistent with vitamin D exerting a modulating effect on the immune system, leading towards a more tolerogenic state, a concept reviewed by (Prietl et al., 2013). The findings from this study are based on an analysis of 97 individuals divided equally between the three treatment groups. Our ability to detect differences in the effects of vitamin D_2 and vitamin D_3 may have been negatively impacted by the inclusion of two different ethnic groups among the 97 participants, since the transcriptome results from the two ethnic groups are clearly different. It will be very important to independently replicate this study using a much larger cohort in order to verify, or otherwise, the key findings from this study. From power calculations it is considered that for a whole human microarray-based gene expression study of this nature, and with gene expression changes of the magnitude we observe, at least

400 participants of each ethnic group should be recruited for each treatment, giving a total cohort size of 2,400. The biological interpretation of our findings in the present study should be considered as preliminary in this context, requiring independent verification.

It is difficult to compare the findings of the present study with other reported *in vivo* studies because of the considerable differences in experimental design, including the use of supra-physiological vitamin D doses (up to 2,000 μg single bolus doses), different population types and small sample sizes which make statistical analysis not feasible (e.g. (Hossein-nezhad et al., 2013, Neme et al., 2019)). Furthermore, our study is unique in that it compared gene expression in participants given either of the two commonly used forms of vitamin D, vitamin D_2 and vitamin D_3. Intriguing, this revealed that there were considerable differences in the gene expression profiles following each nutritional treatment. For example, excluding genes that were also differentially expressed over the 12-week intervention in the placebo group, only 13% of down-regulated differentially expressed genes were common between the two treatment groups while 28% and 59% were uniquely down-regulated by vitamins D_2 and D_3, respectively (Figure 2c).

It is questionable whether it is useful to extrapolate the results from single high dose bolus and *in vitro* studies such as those described by (Koivisto et al., 2020) with this sustained dietary intake *in vivo* study. In the former study 15 'vitamin D-responsive' genes were identified as important mediators of innate and adaptive immunity. Five of these 15 genes are also identified as differentially expressed in our present study (*CD93*, *CEBPB*, *THBD*, *THEMIS2* and *TREM1*). However, in the present study these five genes were *down-regul*ated by vitamin D intake (**Supplementary Data File 2**) whereas they are reported to be up-regulated in the studies reported by (Koivisto et al., 2020). These contrasting results highlight the difficulties with comparing data from different experimental designs. As argued previously, because of the relatively small sample sizes investigated and the small effect sizes, this study has focussed on identification of statistically significant pathway enrichment following sustained vitamin D supplementation, rather than a gene-focussed analysis. Nevertheless, we consider it useful to consider the respective roles of these five specific gene products

since their *down*-regulation by vitamin D would make sense from a biological perspective. In our study the gene encoding thymocyte selection-associated family member 2 (*THEMIS2*) is down-regulated after vitamin D_2 supplementation, the CD93-encoding gene is down-regulated following D_2 and D_3 supplementation, while the genes encoding thrombomodulin (*THBD*) and the CCAAT enhancer-binding protein beta (*CEBPB*) are down-regulated following D_3 supplementation. *THBD* is a key vitamin D target in peripheral blood mononuclear cells (PBMCs) with the function of reducing blood clots while at the same time preventing the pro-inflammatory consequences of NF-kB signalling. The *THBD* and *CD93* genes are next to each other, share the same super-enhancer, and their transcription start site regions respond to $1,25(OH)_2 D_3$. *CEBPB* encodes a transcription factor important in the regulation of genes involved in immune and inflammatory responses. *TREM1* encodes triggering receptor expressed on myeloid cells which stimulates neutrophil and monocyte-mediated inflammatory responses. It triggers release of pro-inflammatory chemokines and cytokines, as well as increased surface expression of cell activation markers. TREM-1 amplifies inflammatory responses that are triggered by bacterial and fungal infections and is a crucial mediator of septic shock (Bouchon et al., 2001).

Another cellular function that is affected by vitamin D_2 and D_3 supplementation is chromatin modification and remodelling. The *EP300* gene is down-regulated by both vitamins, but unaffected in the placebo group. *EP300* encodes a histone acetylase that regulates transcription by chromatin remodelling, in particular histone acetylation of H3 at lysine-27, representing an epigenetic modification which activates transcription (Hatzi et al., 2013). Importantly, EP300 is also a coactivator of VDR.

Functional categories of genes enriched among the upregulated genes, following supplementation with either vitamin D_2 or $D_3,$ include translation, mitochondrial and spliceosome function (Fig. 3b, **Supplementary Data File 4**); statistically enriched biological cellular component terms in these functional categories are ribosomal proteins, components of the mitochondrial respiratory chain, two subunits of the histone H4 and snRNP Sm protein components of the spliceosome assembly. It is established that, in addition to influencing transcription, vitamin D can also influence post-transcriptional events by recruiting co-regulators. In this context it is relevant

that components of the spliceosome such as snRNP Sm proteins that mediate both transcriptional control and splicing decisions, leading to alternatively spliced transcripts (Zhou et al., 2015), were upregulated by vitamin D supplementation in this study. These statistically significant gene expression changes were detected as temporal differences *within* each treatment group (differences at baseline (V1) versus 12 weeks after the intervention (V3) *within* single individuals). Differences in gene expression *across* treatment groups were only found in the South Asian group (Fig. 2a, b). The SA treatment group that received supplementation with vitamin D_3, revealed three genes significantly upregulated and two down-regulated after the 12-week intervention relative to the placebo group (Fig. 2; **Supplementary Fig. 1**). We note that the gene encoding cAMP-responsive element binding protein 1 (*CREB1*) is one of the three upregulated genes. It encodes a transcription factor that binds as a homodimer to the cAMP-responsive element. CREB1 protein is phosphorylated by several protein kinases, and induces transcription of genes in response to hormonal stimulation of the cAMP pathway. The protein kinase A (PKA) pathways are activated by the parathyroid hormone PTH in response to low serum calcium levels to maintain serum calcium homeostasis, primarily by promoting vitamin D 1α-hydroxylation in the kidney. Both PTH and 1,25 $(OH)_2$ vitamin D have similar effects in promoting the maturation of osteocytes and opposing the differentiation of osteoblasts into osteocytes (St John et al., 2015, St John et al., 2014). It is relevant to note in this context that all treatment groups in the SA cohort showed elevated serum PTH at the baseline sampling point relative to the WE cohort.

Vitamins D_2 and D_3 do not influence expression of the same genes

Our results indicate that the cellular responses to the two forms of vitamin D supplementation have some commonalities but also show some clear differences (catalogued in **Supplementary Data File 4**). For example, some biological processes such as histone modification and covalent chromatin modification are downregulated following vitamin D_3 supplementation only, while spliceosomal function are upregulated by vitamin D_2 only. In light of the finding that vitamins D_2 and D_3 influence expression of different genes in the human blood transcriptome, recently we undertook a parallel *in vitro* study with a model rat cell line (Mengozzi et al., 2020). We examined the respective influences of the two physiologically active

forms of vitamin D, 1,25(OH)D$_3$ and 1,25(OH)D$_2$ on differentiation and global gene expression in differentiating rat CG4 oligodendrocyte precursor cells and revealed considerable differences in the influence of the two types of vitamin D on gene expression at 24 h and after 72 h following onset of differentiation. We demonstrated that 1,25(OH)D$_3$ and 1,25(OH)D$_2$ respectively influenced expression of 1,272 and 574 genes at 24 h following addition of the vitamin to the culture, where many of the changes in expression were specific to one or the other form of the vitamin (Mengozzi et al., 2020). This study provides evidence of the different direct effects of the two active vitamin D metabolites on gene expression *in vitro* in cultured cells and provides some evidence that the changes we observed in the *in vivo* study may reflect the influence of the physiologically active forms of the vitamin, 1,25(OH)D$_3$ and 1,25(OH)D$_2$.

Influence of vitamin D on expression of genes encoding immune pathways

In common to both the D$_2$ and D$_3$ treatment groups, but not the placebo group, we found that many different pathways of the immune system are differentially expressed (largely down-regulated) by vitamin D, consistent with the emerging theme that one physiological role of vitamin D is to suppress, or balance, the activity of the immune system (Prietl et al., 2013, Martens et al., 2020)(Fig 3a, Fig. 4; Supplementary data files 3, 4 and 5). The immunomodulatory effects of vitamin D on both innate and adaptive immunity are well documented (Liu et al., 2006, Hayes and Nashold, 2018, Teles et al., 2013). Vitamin D supplementation has the potential to ameliorate the symptoms of autoimmune diseases such as multiple sclerosis, as supported by several lines of evidence: the role in promoting differentiation of oligodendrocyte progenitor cells and therefore remyelination of CNS neurons (Gomez-Pinedo et al., 2020); suppression of the differentiation of potentially pathogenic T-cells and enhancement of regulatory T-cell differentiation (Wasnik et al., 2020). However, a recent Cochrane review found little benefit of vitamin D in this context (Jagannath et al., 2018).

We have also found some differences in response to vitamin D in relation to ethnicity (with the caveat that the sample size of the SA group was smaller than the WE group). Ethnic differences in response are suggested from the weighted gene

correlation network analysis (WGCNA). A direct correlation between the stimulation of the immune response with an increase in serum 25(OH)D$_3$ concentration (and a corresponding negative correlation with the serum concentration of PTH) was evident in the SA group only. This is the opposite of that observed in the WE group where the effect of vitamin D supplementation was to suppress immune pathways.

Conclusions

The transcriptome results from this study differ from those of previous *in vivo* studies which used supra-physiological doses of vitamin D over a short time-frame, or used *in vitro* cultured cells. Indeed, where there is overlap in the identity of the differentially expressed genes identified in the respective studies vitamin D appears to exert the opposite effect relative to our longer-term repeated physiological dose study. Short-term responses to large boluses are relatively simple to do, and less prone to noise, but their effect on gene expression does not seem to be consistent with that observed with longer term repeated physiological dose studies like the present study and therefore may not represent the real-world effects of repeated dietary intake of vitamin D. Thus, despite the challenges and limitations of doing longer term *in vivo* human studies, that are complicated by seasonal variation in gene expression, inter-individual human variation, subtle physiological changes in gene expression, more 'noise' through intake of other vitamin D sources during the treatment intervention, our work suggests it may be important to conduct more such longer term studies since the results from the shorter term interventions may not be representative of the influence of vitamin D on normal physiological gene expression. One potential criticism of whole blood transcriptome studies is that the results may not represent the gene expression changes that occur in other tissues as a result of vitamin D supplementation. However, we have shown, in comparable whole blood transcriptome studies of human sleep physiology, that the changes in the human blood transcriptome correlate with the changes that are observed in other tissues and organs in comparable mammalian animal studies, giving us some confidence in the physiological relevance of the changes identified in this vitamin D transcriptome study (Archer et al., 2014, Moller-Levet et al., 2013).

The transcriptome results differ considerably between the two different ethnic groups. While some of the differences may be attributable to differences in sample sizes studied between the two groups it is also possible that some of the differences represent genuine differences in the respective physiological responses of the two ethnic groups. Alternatively, the differences may reflect the starting physiological status of the SA participant group, who were found to have considerably lower baseline vitamin D concentrations (and higher PTH concentrations) than the WE group. The importance of clarifying these different responses is given particular impetus by the emerging evidence for the interplay between ethnicity, skin tone, vitamin D status and susceptibility to viral infection, COVID-19 being of particular current relevance here (Lanham-New et al., 2020, Kohlmeier, 2020, Kmietowicz, 2020, Weir et al., 2020). The transcriptomic data presented in this study might provide a useful context for further studies aimed at understanding the role of vitamin D in influencing the immune response to SARS-CoV-2 infection, particularly in relation to severe COVID-19.

In summary our results indicate that whilst the two forms of vitamin D (D_2 and D_3) exert some overlapping roles in human physiology, each form may elicit different molecular responses, findings which warrant a more detailed consideration of the system-wide effects of these two forms vitamin D. Clearly, this type of study requires replication, using a much larger independent cohort, with balanced representation from different ethnic groups. This is perhaps of particular importance to at-risk ethnic groups, including black and South Asian populations who reside in northern latitudes. The studies would need to be time course-based (temporal) to track gene expression changes *within* individuals from baseline activity to defined sampling times and would need in-built control to account for seasonal gene expression changes. They would need to be designed specifically to answer whether different ethnicity or different vitamin D baseline levels give rise to different responses to vitamin D supplementation.

Since some pathways appear to be regulated specifically by vitamin D_3, future studies should investigate whether vitamin D_2 supplementation might counteract some of the benefits of vitamin D_3 on human health. This possibility is prompted by the findings from this cohort that the circulating concentration of $25(OH)D_3$ within

vitamin D_2-treated participants was significantly lower after the 12-week intervention than in the placebo group who received no vitamin D supplements – suggesting that the former might be depleted by the latter. The results from this study suggest that guidelines on food fortification and supplementation with specific forms of vitamin D may need revisiting.

Materials and Methods

Blood Transcriptome analysis

The recruitment of individuals as part of the BBSRC D2-D3 Study was described previously (Tripkovic et al., 2017). This study received ethical approval from the South-East Coast (Surrey) National Health Service Research Ethics Committee (11/LO/0708) and the University of Surrey Ethics Committee (EC/2011/97/FHMS). All of the participants gave written informed consent in agreement with the Helsinki Declaration before commencing study activities. Briefly, 335 women of both South Asian (SA) and white European (WE) descent were randomised to one of three intervention groups for 12-weeks and provided with daily doses of vitamin D within fortified foods: placebo, 15 µg/d vitamin D_2 or 15µg/d vitamin D_3. Of the 335 participants, 97 were selected for transcriptome analysis (32 were WE, 32 were A and 33 were selected at random from both ethnic groups to form a balanced placebo group). High responders (>50% increase in 25OHD concentrations) and low responders (<50% increase in 25OHD concentrations) were included within both vitamin D_2 and D_3 supplementation groups to maximize the chance of observing differences in gene expression among subjects. The selected time points for transcriptome analysis were V1 representing the baseline and V3, 12 weeks after treatment commenced.

Whole peripheral blood (2.5 ml) was collected using PAXgene Blood RNA tubes (Becton Dickinson). PAXgene Blood RNA Tubes were inverted ten times immediately after drawing blood, stored upright at 15-25°C for 24 hours, followed by a -20°C freezer for 24 hours and then into a -80°C freezer for long-term storage.

Transcriptomic analysis was conducted essentially as described in previous studies (Archer et al., 2014, Moller-Levet et al., 2013). Total RNA was isolated using the

PAXgene Blood RNA Kit (Qiagen) following the manufacturers recommendations. cRNA was synthesised and fluorescently labelled with Cy3-CTP from 200 ng of total RNA using the Low Input QuickAmp Labelling Kit, One Color (Agilent Technology). Labelled cRNA was hybridised on a Sure Print G3 Human Gene Expression 8 x 60K v2 microarray slide (Agilent Technologies). Standard manufacturer's instructions for one colour gene-expression analysis were followed for labelling, hybridsation and washing steps. Extracted RNA was quantified using NanoDrop ND2000 spectrophotometer (Thermo Scientific). RNA quality and integrity was evaluated using either the Bioanalyzer 2100 or the TapeStation 4200 (Agilent Technologies). Only RNA samples with an RNA Integrity Number (RIN) of >7.0 were subjected to DNA microarray analysis. Microarrays were hybridised at 65°C for 17 hours in an Agilent Hybridization Oven on a rotisserie at 10 rpm. The washed microarrays were scanned using an Agilent Microarray Scanner with a resolution of 2 μm.

Transcriptome data processing and differential expression analysis.
Raw scanned microarray images were processed using Agilent Feature Extraction software (v11.5.1.1) with the Agilent 039494_D_F_20140326_human_8x60K_v2 grid, and then imported into R for normalization and analysis using the LIMMA package (Ritchie et al., 2015). Microarray data were background-corrected using the 'normexp' method (with an offset of 50) and quantile normalised, producing expression values in the log base 2 scale. The processed data were then filtered to remove probes exhibiting low signals across the arrays, retaining non-control probes that are at least 10% brighter than negative control probe signals on at least 41 arrays (~20% of the arrays in the analysis). Data from identical replicate probes were then averaged to produce expression values at the unique probe level. Initial data exploration identified one sample (participant 0017, V3 time point) with array data that was a notable outlier from the group and therefore both the V1 and V3 microarrays for this subject were excluded from all subsequent analysis, reprocessing the data as above in their absence before proceeding.

Tests for differential expression were performed using LIMMA, applying appropriate linear model designs to identify: (i) significant differences in the transcriptional responses occurring across the 12-week V1 to V3 period of the study between the treatment and placebo groups for each ethnic cohort (example contrasts tested, in

the format ethnicity_treatment_time: (WE_D2_V3 - WE_D2_V1) - (WE_P_V3-
WE_P_V1) = 0 , and (WE_D3_V3 - WE_D3_V1) - (WE_P_V3 - WE_P_V1) = 0); and
ii) to determine significant changes for each ethnic cohort within each treatment
group between the V3 and V1 sampling points, blocking on subject identity (example
contrasts tested: WE_D2_V3 - WE_D2_V1 = 0, WE_D3_V3 - WE_D3_V1 = 0).
Blocking on subject identity was used to control for inter-subject variability.
Significance p-values were corrected for multiplicity using the Benjamini and
Hochberg method, obtaining adjusted p-values (adj.P.Val). Normalised data for all
participants (V3 – V1) was assessed by principal component analysis to screen for
any batch effects (**Supplementary Figure 3**).

Functional enrichment and network analysis

Functional enrichment analysis of lists of genes of interest possessing valid ENTREZ
gene identifiers was performed using the R package clusterProfiler (Yu et al., 2012)
The software produces adjusted p-values (p.adjust) using the Benjamini and
Hochberg correction method. Construction and analysis of protein-protein interaction
networks from sets of genes was undertaken in Cytoscape (Shannon et al., 2003)
using the STRING plugin (Szklarczyk et al., 2017a). Cytoscape was also used to
construct and visualise commonality in the functional categories identified as being
significantly enriched in the genes responding to the experimental treatments. To
visualise categories identified from the D_2 or D_3 treatment groups but not from the
placebo (e.g. Figs. 4 and 5) significantly enriched categories (p.adjust < 0.01) from
all groups were imported such that treatment group nodes (D_2, D_3 and placebo (P))
are linked to functional category nodes by edges assigned the corresponding
p.adjust values. All nodes with an edge connection to the placebo treatment node
were then removed and the resulting networks further filtered to retain only those
nodes with at least one edge connection with p.adjust<=0.001.

Weighted gene co-expression network analysis was performed in R using the
WGCNA (Langfelder and Horvath, 2008) and CoExpNets (Botia, 2019) packages. A
normalised expression data matrix generated by filtering to retain probes with signals
more markedly above background (30% brighter than negative control probe signals
on at least 41 arrays) was used as input, consisting of 12,169 unique probes. Signed
scale-free networks were constructed separately for the data for the SA and WE

ethnic groups using 50 iterations of the k-means clustering option in the CoExpNets 'getDownstreamNetwork' function to refine the clustering process (**Supplementary Data File 6**). Pearson correlations between cluster module eigengenes and metadata for serum 25(OH)D$_2$, 25(OH)D$_3$, total 25(OH)D and PTH concentrations were calculated using pairwise complete observations.

Data availability: All microarray data are available in the ArrayExpress database (http://www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-8600.

Acknowledgements

This work was funded by BBSRC (UK) DRINC award BB/I006192/1 to SLN and CPS, by a University of Brighton Innovation Seed Fund Award (to CPS). The authors would like to particularly thank Michael Chowen CBE DL and Maureen Chowen for their generous philanthropic donation (to CPS).

Declaration of interests

All authors confirm that they have no conflicts of interest.

Figure legends

Figure 1. (a) Selection of 98 subject participants from the Tripkovic et al. (2017) study (Tripkovic et al., 2017) for transcriptomic analysis of their V1 (baseline) and V3 (12-week) samples in the present study; (b) Ethnicity and treatment group membership for the 97 subjects for which transcriptomic data was obtained (data for one South Asian subject from the placebo group did not pass quality control); (c) Metadata on serum concentrations of $25(OH)D_2$, $25(OH)D_3$, total $25(OH)D$, PTH and calcium (albumin-adjusted) for the 97 subjects (see **Supplementary Data File 1** for details). Participants were selected to provide comparable numbers between the placebo and the two vitamin D treatment groups, covering the full range of serum responses to supplementation within the D_2 and D_3 treatment groups, as judged from the measured changes in serum $25(OH)D_2$ or $25(OH)D_3$ concentrations between V1 and V3.

Figure 2. (a) Summary of limma differential expression test results identifying significant changes (adj.P.Val<=0.05) in transcript probe abundance within and between the experimental groups (see **Supplementary Data File 2** for full details). Both the number of unique microarray probes, and the number of unique genome loci they represent, are indicated. WE = "white European"; SA = "South Asian"; D2 = vitamin D_2 treatment group; D3 = vitamin D_3 treatment group; P = placebo group. (b) The CREB1 gene probe signal is significantly different (adj.P.Val=0.021) between the vitamin D_3-treated group and the placebo group over the course of the V1 to V3 study period in the South Asian cohort (i.e. [SA D3 V3 v V1] v [SA P V3 v V1] comparison from Fig. 2a). (c) Venn diagrams showing the numbers of probes in the white European (WE) cohort that are specifically significantly (adj.P.Val<=0.05) down- or up-regulated in V3 compared to V1 in each treatment group. (d) Comparison of the $\log_2$ fold-changes (FC) in abundance occurring from V1 to V3 in the WE cohort in the D_2 and D_3 treatment groups. Probes significantly up-regulated in both D_2 and D_3 groups but not the placebo are coloured red, while those similarly down-regulated are shown in blue (see **Supplementary Data File 3**). A probe detecting *SEC14L1* shows the largest decreases in abundance in those down-regulated by D_2 and D_3 but not placebo.

Figure 3. Protein-protein interaction networks for gene products corresponding to the probes (a) significantly down-regulated or (b) significantly up-regulated in both the D_2 and D_3 treatment groups of the WE cohort, but not the placebo group. Details given in **Supplementary Data File 3**. The networks were generated using the STRING database of *Homo sapiens* medium confidence (0.4) interactions, and only connected nodes are shown. Networks for both (a) and (b) are significantly enriched for interactions compared to randomised sets, yielding p-values of 1.98×10^{-6} and 4.99×10^{-9}, respectively.

Figure 4. Networks illustrating the functional categories significantly enriched in the gene products represented by the probes significantly down-regulated (adj.P.Val $\leq$ 0.05) in the D_2 and D_3 treatment groups of the WE cohort, but not in the placebo group. Gene products represented by the significantly down-regulated probes in the comparisons WE D2 V3 v V1, WE D3 V3 v V1 and WE P V3 v V1 from Fig. 2(a), and possessing ENTREZ identifiers, were subjected separately to functional enrichment analysis using compareCluster (Yu et al., 2012). The details for each group are given in **Supplementary Data File 4**. Significantly enriched categories (p.adjust $\leq$ 0.01) from all groups were imported into Cytoscape (Shannon et al., 2003) to visualise categories identified from the D_2 or D_3 treatment groups but not by the placebo (see Methods for details). The thickness of lines (edges) drawn to connect the nodes shown increases with increasing significance (p.adjust-values) and only categories where at least one edge is $\leq 1 \times 10^{-3}$ are shown. The complete networks for each differentially expressed group of genes are shown in **Supplementary Data File 5.**

Figure 5. Networks illustrating the functional categories significantly enriched in the gene products represented by the probes significantly up-regulated (adj.P.Val $\leq$ 0.05) in the D_2 and D_3 treatment groups of the WE cohort, but not in the placebo group. Gene products represented by the significantly up-regulated probes in the comparisons WE D2 V3 v V1, WE D3 V3 v V1 and WE P V3 v V1 from Fig. 2(a), and possessing ENTREZ identifiers, were subjected separately to functional enrichment analysis using compareCluster as detailed in the legend to Fig. 4. The complete networks for each differentially expressed group of genes are shown in **Supplementary Data File 5.**

Figure 6. Gene co-expression networks for the WE (a) and SA (b) ethnic groups. For each group of subjects, signed scale-free networks were constructed from the expression data using WGCNA to cluster probes with similar expression characteristics across all samples into discrete modules (colours on the y-axis: colours in (a) and (b) are independent). Module membership for the modules of genes identified in the WGCNA analysis of the data from the South Asian and white European cohorts are detailed in **Supplementary Data File 6**. The colour scale to the right of each panel represents the Pearson correlation (from -1 to +1) between the expression profile for each module's eigengene and the respective serum $25(OH)D_2$, $25(OH)D_3$, total $25(OH)D$ and PTH concentrations (x-axis). The Pearson correlation coefficients are also provided in each box, followed in brackets by an adjusted p-value, testing for the significance of each correlation. Statistically significant correlations (adjusted p-value ≤ 0.05) are circled in red. A headline significant GO functional enrichment category (p.adjust ≤ 0.01) for each significant module is shown to the right; GO test results for each module are summarised in **Supplementary data files 7 and 8**).

Supplementary Figures and Supplementary Data Files

Supplementary Figure 1.

Summary of the probes identified as being significantly different in the 'difference-in-difference' analysis of the SA cohort (comparison [SA D3 V3 v V1] v [SA P V3 v V1] in Fig. 2a). (A) Differential expression test results; (B) Probe signal abundance ($\log_2$) distribution across the experimental treatments in the SA cohort.

Supplementary Figure 2.

Seasonal gene expression as a feature of the 12-week study (P = placebo, D2 = vitamin D_2, D3 = vitamin D_3). (a) Summary of the timing of the V1 samples taken in the study (for each individual, V3 samples were taken 12 weeks after the V1 sample); (b) Significant over-representation of the seasonal genes identified by Dopico et al (2015) *Nat Commun*, **6**, 7000 (BABYDIET dataset) in the genes identified as significantly changing in the white European (WE) placebo, D_2 and D_3 treatment groups.

Supplementary Figure 3.

Principal component analysis of the transcriptome data for the 97 subjects passing quality control. Data for samples from visits V1 and V3 are processed as the difference between V3 and V1 (V3 minus V1) for each subject. The distribution is relatively random.

Supplementary Data Files

Supplementary Data File 1

Metadata on serum concentrations of $25(OH)D_2$, $25(OH)D_3$, total $25(OH)D$, PTH and calcium (albumin-adjusted) for the 97 subjects subjected to transcriptomic analysis (selected from the cohort described by Tripkovic et al. 2017).

Supplementary Data File 2 [Note: data file is 9.1 MB.]

Significant differences in the transcriptional responses occurring across the 12-week V1 to V3 period of the study *between* the vitamin D_2, vitamin D_3 and placebo treatment groups for each ethnic cohort, and significant changes *within* each group between the V3 and V1 sampling points (see Fig. 2).

Supplementary Data File 3

Functional analysis of the genes represented by the probes specifically repressed by D_2 and D_3 in the white European cohort (blue data points in Fig. 2d) or induced by D_2 and D_3 in the white European cohort (red data points in Fig. 2d).

Supplementary Data File 4

Comparative functional enrichment analysis of all significant changes in each treatment group in the white European cohort (Fig. 2c; and see Figs 4 and 5).

Supplementary Data File 5

Networks illustrating all the functional categories significantly enriched (p.adjust < 0.01) in analysis of the gene products represented by the probes significantly up- or down-regulated (adj.P.Val < 0.05) in the white European cohort treatment groups (from data presented in **Supplementary Data File 4**).

Supplementary Data File 6

Module membership for the modules of genes identified in the WGCNA analysis of the data from the South Asian and white European cohorts, as presented in **Fig. 6**.

Supplementary Data File 7

Gene ontology functional enrichment analysis results for the modules of genes identified in the WGCNA analysis of the data from the white European cohort.

Supplementary Data File 8

Gene ontology functional enrichment analysis results for the modules of genes identified in the WGCNA analysis of the data from the South Asian cohort.

References

AGELL, L., HERNANDEZ, S., NONELL, L., LORENZO, M., PUIGDECANET, E., DE MUGA, S., JUANPERE, N., BERMUDO, R., FERNANDEZ, P. L., LORENTE, J. A., SERRANO, S. & LLORETA, J. 2012. A 12-gene expression signature is associated with aggressive histological in prostate cancer: SEC14L1 and TCEB1 genes are potential markers of progression. *Am J Pathol,* 181**,** 1585-94.

ARCHER, S. N., LAING, E. E., MOLLER-LEVET, C. S., VAN DER VEEN, D. R., BUCCA, G., LAZAR, A. S., SANTHI, N., SLAK, A., KABILJO, R., VON SCHANTZ, M., SMITH, C. P. & DIJK, D. J. 2014. Mistimed sleep disrupts circadian regulation of the human transcriptome. *Proc Natl Acad Sci U S A,* 111**,** E682-91.

BERLANGA-TAYLOR, A. J., PLANT, K., DAHL, A., LAU, E., HILL, M., SIMS, D., HEGER, A., EMBERSON, J., ARMITAGE, J., CLARKE, R. & KNIGHT, J. C. 2018. Genomic Response to Vitamin D Supplementation in the Setting of a Randomized, Placebo-Controlled Trial. *EBioMedicine,* 31**,** 133-142.

BOTIA, J. A. 2019. CoExpNets: Co-expression network management based on WGCNA + k-means. R package version 0.1.0. 0.1.0 ed.

BOUCHON, A., FACCHETTI, F., WEIGAND, M. A. & COLONNA, M. 2001. TREM-1 amplifies inflammation and is a crucial mediator of septic shock. *Nature,* 410**,** 1103-7.

BOUILLON, R. 2018. Extra-Skeletal Effects of Vitamin D. *Front Horm Res,* 50**,** 72-88.

BURDELSKI, C., BARREAU, Y., SIMON, R., HUBE-MAGG, C., MINNER, S., KOOP, C., GRAEFEN, M., HEINZER, H., SAUTER, G., WITTMER, C., STEURER, S., ADAM, M., HULAND, H., SCHLOMM, T., TSOURLAKIS, M. C. & QUAAS, A. 2015. Saccharomyces cerevisiae-like 1 overexpression is frequent in prostate cancer and has markedly different effects in Ets-related gene fusion-positive and fusion-negative cancers. *Hum Pathol,* 46**,** 514-23.

CHRISTAKOS, S., DHAWAN, P., VERSTUYF, A., VERLINDEN, L. & CARMELIET, G. 2016. Vitamin D: Metabolism, Molecular Mechanism of Action, and Pleiotropic Effects. *Physiol Rev,* 96**,** 365-408.

DAMOISEAUX, J. & SMOLDERS, J. 2018. The Engagement Between Vitamin D and the Immune System: Is Consolidation by a Marriage to Be Expected? *EBioMedicine,* 31**,** 9-10.

DOPICO, X. C., EVANGELOU, M., FERREIRA, R. C., GUO, H., PEKALSKI, M. L., SMYTH, D. J., COOPER, N., BURREN, O. S., FULFORD, A. J., HENNIG, B. J., PRENTICE, A. M., ZIEGLER, A. G., BONIFACIO, E., WALLACE, C. & TODD, J. A. 2015. Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat Commun,* 6**,** 7000.

FETAHU, I. S., HOBAUS, J. & KALLAY, E. 2014. Vitamin D and the epigenome. *Front Physiol,* 5**,** 164.

GOMEZ-PINEDO, U., CUEVAS, J. A., BENITO-MARTIN, M. S., MORENO-JIMENEZ, L., ESTEBAN-GARCIA, N., TORRE-FUENTES, L., MATIAS-GUIU, J. A., PYTEL, V., MONTERO, P. & MATIAS-GUIU, J. 2020. Vitamin D increases remyelination by promoting oligodendrocyte lineage differentiation. *Brain Behav,* 10**,** e01498.

HATZI, K., JIANG, Y., HUANG, C., GARRETT-BAKELMAN, F., GEARHART, M. D., GIANNOPOULOU, E. G., ZUMBO, P., KIROUAC, K., BHASKARA, S., POLO, J. M., KORMAKSSON, M., MACKERELL, A. D., JR., XUE, F., MASON, C. E., HIEBERT, S. W., PRIVE, G. G., CERCHIETTI, L., BARDWELL, V. J., ELEMENTO, O. & MELNICK, A. 2013. A hybrid mechanism of action for BCL6 in B cells defined by formation of functionally distinct complexes at enhancers and promoters. *Cell Rep,* 4**,** 578-88.

HAYES, C. E. & NASHOLD, F. E. 2018. Vitamin D and Multiple Sclerosis. *In:* FELDMAN, D. (ed.). Elsevier Inc.

HOLICK, M. F. 2007. Vitamin D deficiency. *N Engl J Med,* 357**,** 266-81.

HOLICK, M. F., BIANCUZZO, R. M., CHEN, T. C., KLEIN, E. K., YOUNG, A., BIBULD, D., REITZ, R., SALAMEH, W., AMERI, A. & TANNENBAUM, A. D. 2008. Vitamin D2 is as effective as vitamin

D3 in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab,* 93**,** 677-81.

HOLLIS, B. W. 1984. Comparison of equilibrium and disequilibrium assay conditions for ergocalciferol, cholecalciferol and their major metabolites. *J Steroid Biochem,* 21**,** 81-6.

HOSSEIN-NEZHAD, A., SPIRA, A. & HOLICK, M. F. 2013. Influence of vitamin D status and vitamin D3 supplementation on genome wide expression of white blood cells: a randomized double-blind clinical trial. *PLoS One,* 8**,** e58725.

HOUGHTON, L. A. & VIETH, R. 2006. The case against ergocalciferol (vitamin D2) as a vitamin supplement. *Am J Clin Nutr,* 84**,** 694-7.

JAGANNATH, V. A., FILIPPINI, G., DI PIETRANTONJ, C., ASOKAN, G. V., ROBAK, E. W., WHAMOND, L. & ROBINSON, S. A. 2018. Vitamin D for the management of multiple sclerosis. *Cochrane Database Syst Rev,* 9**,** CD008422.

KMIETOWICZ, Z. 2020. Sixty seconds on . . . vitamin D. *BMJ,* 371**,** m3872.

KOHLMEIER, M. 2020. Avoidance of vitamin D deficiency to slow the COVID-19 pandemic. *BMJ Nutrition, Prevention & Health,* 3.

KOIVISTO, O., HANEL, A. & CARLBERG, C. 2020. Key Vitamin D Target Genes with Functions in the Immune System. *Nutrients,* 12.

LANGFELDER, P. & HORVATH, S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics,* 9**,** 559.

LANHAM-NEW, S. A., WEBB, A. R., CASHMAN, K. D. & AL, E. 2020. Vitamin D and SARS-CoV-2 virus/COVID-19 disease. *BMJ Nutrition, Prevention & Health* 3.

LIU, P. T., STENGER, S., LI, H., WENZEL, L., TAN, B. H., KRUTZIK, S. R., OCHOA, M. T., SCHAUBER, J., WU, K., MEINKEN, C., KAMEN, D. L., WAGNER, M., BALS, R., STEINMEYER, A., ZUGEL, U., GALLO, R. L., EISENBERG, D., HEWISON, M., HOLLIS, B. W., ADAMS, J. S., BLOOM, B. R. & MODLIN, R. L. 2006. Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science,* 311**,** 1770-3.

LONG, M. D., SUCHESTON-CAMPBELL, L. E. & CAMPBELL, M. J. 2015. Vitamin D receptor and RXR in the post-genomic era. *J Cell Physiol,* 230**,** 758-66.

MARTENS, P. J., GYSEMANS, C., VERSTUYF, A. & MATHIEU, A. C. 2020. Vitamin D's Effect on Immune Function. *Nutrients,* 12.

MENGOZZI, M., HESKETH, A., BUCCA, G., GHEZZI, P. & SMITH, C. P. 2020. Vitamins D3 and D2 have marked but different global effects on gene expression in a rat oligodendrocyte precursor cell line. *Mol Med,* 26**,** 32.

MOLLER-LEVET, C. S., ARCHER, S. N., BUCCA, G., LAING, E. E., SLAK, A., KABILJO, R., LO, J. C., SANTHI, N., VON SCHANTZ, M., SMITH, C. P. & DIJK, D. J. 2013. Effects of insufficient sleep on circadian rhythmicity and expression amplitude of the human blood transcriptome. *Proc Natl Acad Sci U S A,* 110**,** E1132-41.

NEME, A., SEUTER, S., MALINEN, M., NURMI, T., TUOMAINEN, T. P., VIRTANEN, J. K. & CARLBERG, C. 2019. In vivo transcriptome changes of human white blood cells in response to vitamin D. *J Steroid Biochem Mol Biol,* 188**,** 71-76.

ODS, N. 2020. *Vitamin D* [Online]. National Institutes of Health. Available: https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional/ [Accessed].

PILZ, S., ZITTERMANN, A., TRUMMER, C., THEILER-SCHWETZ, V., LERCHBAUM, E., KEPPEL, M. H., GRUBLER, M. R., MARZ, W. & PANDIS, M. 2019. Vitamin D testing and treatment: a narrative review of current evidence. *Endocr Connect,* 8**,** R27-R43.

PRIETL, B., TREIBER, G., PIEBER, T. R. & AMREIN, K. 2013. Vitamin D and immune function. *Nutrients,* 5**,** 2502-21.

RITCHIE, M. E., PHIPSON, B., WU, D., HU, Y., LAW, C. W., SHI, W. & SMYTH, G. K. 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res,* 43**,** e47.

ROSEN, C. J., ADAMS, J. S., BIKLE, D. D., BLACK, D. M., DEMAY, M. B., MANSON, J. E., MURAD, M. H.
& KOVACS, C. S. 2012. The nonskeletal effects of vitamin D: an Endocrine Society scientific
statement. *Endocr Rev,* 33**,** 456-92.
SACN. 2016. SACN (Scientific Advisory Committee on Nutrition) Vitamin D and Health Report. .
SCHWALFENBERG, G. K. 2011. A review of the critical role of vitamin D in the functioning of the
immune system and the clinical implications of vitamin D deficiency. *Mol Nutr Food Res,* 55**,**
96-108.
SHANNON, P., MARKIEL, A., OZIER, O., BALIGA, N. S., WANG, J. T., RAMAGE, D., AMIN, N.,
SCHWIKOWSKI, B. & IDEKER, T. 2003. Cytoscape: a software environment for integrated
models of biomolecular interaction networks. *Genome Res,* 13**,** 2498-504.
SONBUL, S. N., ALESKANDARANY, M. A., KUROZUMI, S., JOSEPH, C., TOSS, M. S., DIEZ-RODRIGUEZ,
M., NOLAN, C. C., MUKHERJEE, A., MARTIN, S., CALDAS, C., ELLIS, I. O., GREEN, A. R. &
RAKHA, E. A. 2018. Saccharomyces cerevisiae-like 1 (SEC14L1) is a prognostic factor in breast
cancer associated with lymphovascular invasion. *Mod Pathol,* 31**,** 1675-1682.
ST JOHN, H. C., BISHOP, K. A., MEYER, M. B., BENKUSKY, N. A., LENG, N., KENDZIORSKI, C.,
BONEWALD, L. F. & PIKE, J. W. 2014. The osteoblast to osteocyte transition: epigenetic
changes and response to the vitamin D3 hormone. *Mol Endocrinol,* 28**,** 1150-65.
ST JOHN, H. C., MEYER, M. B., BENKUSKY, N. A., CARLSON, A. H., PRIDEAUX, M., BONEWALD, L. F. &
PIKE, J. W. 2015. The parathyroid hormone-regulated transcriptome in osteocytes: parallel
actions with 1,25-dihydroxyvitamin D3 to oppose gene expression changes during
differentiation and to promote mature cell function. *Bone,* 72**,** 81-91.
SZKLARCZYK, D., MORRIS, J. H., COOK, H., KUHN, M., WYDER, S., SIMONOVIC, M., SANTOS, A.,
DONCHEVA, N. T., ROTH, A., BORK, P., JENSEN, L. J. & VON MERING, C. 2017a. The STRING
database in 2017: quality-controlled protein-protein association networks, made broadly
accessible. *Nucleic Acids Res,* 45**,** D362-D368.
SZKLARCZYK, D., MORRIS, J. H., COOK, H., KUHN, M., WYDER, S., SIMONOVIC, M., SANTOS, A.,
DONCHEVA, N. T., ROTH, A., BORK, P., JENSEN, L. J. & VON MERING, C. 2017b. The STRING
database in 2017: quality-controlled protein–protein association networks, made broadly
accessible. *Nucleic Acids Research,* 45**,** D362-D368.
TELES, R. M., GRAEBER, T. G., KRUTZIK, S. R., MONTOYA, D., SCHENK, M., LEE, D. J., KOMISOPOULOU,
E., KELLY-SCUMPIA, K., CHUN, R., IYER, S. S., SARNO, E. N., REA, T. H., HEWISON, M., ADAMS,
J. S., POPPER, S. J., RELMAN, D. A., STENGER, S., BLOOM, B. R., CHENG, G. & MODLIN, R. L.
2013. Type I interferon suppresses type II interferon-triggered human anti-mycobacterial
responses. *Science,* 339**,** 1448-53.
TRIPKOVIC, L., LAMBERT, H., HART, K., SMITH, C. P., BUCCA, G., PENSON, S., CHOPE, G., HYPPONEN,
E., BERRY, J., VIETH, R. & LANHAM-NEW, S. 2012. Comparison of vitamin D2 and vitamin D3
supplementation in raising serum 25-hydroxyvitamin D status: a systematic review and
meta-analysis. *Am J Clin Nutr,* 95**,** 1357-64.
TRIPKOVIC, L., WILSON, L. R., HART, K., JOHNSEN, S., DE LUSIGNAN, S., SMITH, C. P., BUCCA, G.,
PENSON, S., CHOPE, G., ELLIOTT, R., HYPPONEN, E., BERRY, J. L. & LANHAM-NEW, S. A. 2017.
Daily supplementation with 15 mug vitamin D2 compared with vitamin D3 to increase
wintertime 25-hydroxyvitamin D status in healthy South Asian and white European women:
a 12-wk randomized, placebo-controlled food-fortification trial. *Am J Clin Nutr,* 106**,** 481-490.
WASNIK, S., SHARMA, I., BAYLINK, D. J. & TANG, X. 2020. Vitamin D as a Potential Therapy for
Multiple Sclerosis: Where Are We? *Int J Mol Sci,* 21.
WEIR, E. K., THENAPPAN, T., BHARGAVA, M. & CHEN, Y. 2020. Does vitamin D deficiency increase the
severity of COVID-19? *Clin Med (Lond),* 20**,** e107-e108.
WILSON, L. R., TRIPKOVIC, L., HART, K. H. & LANHAM-NEW, S. A. 2017. Vitamin D deficiency as a
public health issue: using vitamin D2 or vitamin D3 in future fortification strategies. *Proc*
Nutr Soc, 76**,** 392-399.

YU, G., WANG, L. G., HAN, Y. & HE, Q. Y. 2012. clusterProfiler: an R package for comparing biological
themes among gene clusters. *Omics,* 16**,** 284-7.
ZHANG, R., EDWARDS, J. R., KO, S. Y., DONG, S., LIU, H., OYAJOBI, B. O., PAPASIAN, C., DENG, H. W. &
ZHAO, M. 2011. Transcriptional regulation of BMP2 expression by the PTH-CREB signaling
pathway in osteoblasts. *PLoS One,* 6**,** e20780.
ZHANG, Y., FANF, F., TANG, J., JIA, L., FENG, Y., XU, P. & FARAMAND, A. 2019. Association between
vitamin D supplementation and mortality: systematic review and meta-analysis. *British*
Medical Journal, 366**,** l4673.
ZHOU, R., CHUN, R. F., LISSE, T. S., GARCIA, A. J., XU, J., ADAMS, J. S. & HEWISON, M. 2015. Vitamin D
and alternative splicing of RNA. *J Steroid Biochem Mol Biol,* 148**,** 310-7.
ZITTERMANN, A., ERNST, J. B., GUMMERT, J. F. & BORGERMANN, J. 2014. Vitamin D
supplementation, body weight and human serum 25-hydroxyvitamin D response: a
systematic review. *Eur J Nutr,* 53**,** 367-74.
ZITTERMANN, A., PILZ, S. & BERTHOLD, H. K. 2020. Serum 25-hydroxyvitamin D response to vitamin
D supplementation in infants: a systematic review and meta-analysis of clinical intervention
trials. *Eur J Nutr,* 59**,** 359-369.

a)

Tripkovic *et al.* participants, n=335
WE n=245 , SA n=90

Selected 32 subjects from each of the D2
and D3 treatment groups, and 34 from the
placebo

For D2 and D3, a range of serum [25OHD]
responses to supplementation were chosen

Placebo subjects were chosen at random

Subjects for transcriptome analysis, n=98
WE n=67 , SA n=31

b)

Group	White Euro (n)	South Asian (n)	Total (n)
D2	21	11	32
D3	21	11	32
Placebo	25	8	33

c)

Fig. 1.

Fig. 2.

a) Significantly down-regulated by D_2 and D_3, but not placebo (blue points in Fig. 2d)

b) Significantly up-regulated by D_2 and D_3, but not placebo (red points in Fig. 2d)

Significant functional enrichment (FDR<0.05)

- 🟥 Immune system process
- 🟦 Innate immune system
- ⬛ Transcription factor complex
- 🟩 Neutrophil degranulation
- 🟪 Leukocyte activation

Significant functional enrichment (FDR<0.05)

- 🟪 Mitochondrion
- 🟩 Ribosomal protein
- 🟨 snRNP Sm proteins
- ⬜ Histone H4

Fig. 3.

a) Significant GO BP functional categories associated
    with down-regulated probes in the D2 or D3 treatment
    groups, but not the placebo

b) Significant Reactome PA categories associated with
    down-regulated probes in the D2 or D3 treatment
    groups, but not the placebo

Fig. 4.

a) Significant GO CC functional categories associated with up-regulated probes in the D2 or D3 treatment groups, but not the placebo

b) Significant GO BP functional categories associated with up-regulated probes in the D2 or D3 treatment groups, but not the placebo

Fig. 5.

a) Co-expression modules from WE subject data
    (n= 67, 134 microarrays)

b) Co-expression modules from SA subject data
    (n=30, 60 microarrays)

Fig. 6.