Systematic review reveals multiple sexually antagonistic polymorphisms affecting human disease and complex traits

Jon Alexander Harper[1]*, Tim Janicke[2,3] and Edward H. Morrow[4]

* Corresponding author

[1]Evolution, Behaviour and Environment Group, School of Life Sciences, John Maynard Smith Building, University of Sussex, Brighton, BN1 9QG, UK.

[2]Centre d'Écologie Fonctionnelle et Évolutive, UMR 5175, CNRS, Université de Montpellier, École Pratique des Hautes Études, Montpellier, France.

[3]Applied Zoology, Technical University Dresden, Zellescher Weg 20b, 01062 Dresden, Germany.

[4]Department of Environmental and Life Sciences, Karlstad University, 651 88, Karlstad, Sweden.

Jon Alexander Harper ORCID: 0000-0002-1765-9990

Tim Janicke ORCID: 0000-0002-1453-6813

Edward H Morrow ORCID: 0000-0002-1853-7469

Short title: Systematic review of sexually antagonistic genes in human disease

Abstract

Sex differences in human disease risk and incidence are widely documented but their origins are poorly understood. An evolutionary model for differential disease risk between the sexes posits that alleles that are a risk factor (deleterious) in one sex may actually be protective (beneficial) in the opposite sex, and as such these sexually antagonistic (SA) alleles may be maintained in the population at frequencies higher than expected under purifying selection against unconditionally deleterious alleles. Sexual antagonism at genetic loci is thought to arise due to fundamental differences in how Darwinian fitness is maximized in the two sexes. Evidence for the occurrence of SA alleles comes from a number of non-human vertebrate and invertebrate models, which indicate they are common, although only very few specific SA loci have been identified. While there are many examples of loci with sex-limited effects in humans, there are no known examples of specific genetic variants that experience opposing patterns of selection between the sexes in humans. We hypothesized that this may be due to a mismatch in terminology between scientific disciplines rather than a genuine lack of examples. We therefore undertook a two-stage review of the literature for evidence of genetic polymorphisms in human disease and complex traits that exert their effects in a SA pattern, extracting effect sizes and allele frequencies where possible. In the first stage, we used search terms from evolutionary biology that aimed to identify examples of genes specifically referred to as having SA effects or were implicated in intralocus sexual conflict. In the second, we used terminology that was designed to capture as many examples as possible from the biomedical literature (e.g. sex-dependent, sex-opposite, men/women, boys/girls), which may have been missed in the first stage due to differences in terminology between the disciplines. The two searches gave very different results. The first stage returned 34 papers in total, none of which included evidence from humans of genetic loci with opposite effects on the two sexes. The second stage returned 881 papers, which after screening resulted in 70 papers, 32 were deemed eligible, and described 51 different genetic loci or haplotypes with sex-opposite effects on complex traits or disease. Of these, 22 loci increase disease risk or severity in one sex but are protective in the other. These loci may therefore be considered SA loci if the effects on disease risk translate into marginal effects on fitness through fecundity selection or mortality, although in no case were they referred to in terms of sexual

antagonism. As predicted, alleles with net positive effects across the sexes occurred at a

higher frequency than those with net negative effects. Our review reveals that significant

long-term barriers to communication are occurring between scientific fields as a result of

discipline-specific terminology. Less specific search terms revealed evidence that SA variants

are maintained at relatively high frequencies in humans, apparently because while they

contribute to the propensity for one sex to develop a range of diseases and disorders, they

protect the other.

495 words

Introduction

In species with separate sexes an evolutionary conflict at the level of individual genetic loci can occur, where alleles that are favoured by selection in one sex are selected against in the other (i.e intralocus sexual conflict, Parker, 1979). This sexually antagonistic (SA) form of selection is thought to be driven by differences in how the two sexes maximize their fitness and concomitant unequal variances in reproductive success. Research into SA selection has expanded recently with the recognition that it feeds into several important evolutionary processes. It is primarily thought to drive the evolution of sexual dimorphism and trait diversification (Lande, 1980; Pennell et al., 2016; Rice, 1984), including sex-biased gene expression (Ellegren and Parsch, 2007). As a potential driver of balancing selection, it has also been implicated in the maintenance of genetic variation (Connallon and Clark, 2013; Grieshop and Arnqvist, 2018) that would otherwise be eroded by directional selection. Most recently, it has been suggested that genetic variation at SA loci could contribute to the occurrence of a number of common human diseases (Morrow, 2015; Morrow and Connallon, 2013), which also show considerable variation between the sexes in terms of their prevalence, severity and age of onset (Ober et al., 2008; Rigby and Kulathinal, 2015).

Despite the interest in sexual antagonism as an evolutionary force, determining the identity of SA loci remains a major challenge (Ruzicka et al., 2020). An early empirical milestone in the field of sexual conflict was achieved by quantitative genetic studies demonstrating that genomes in a laboratory adapted population of *Drosophila melanogaster* harbour significant amounts of SA standing genetic variation (Chippindale et al., 2001; Rice, 1992). A number of other systems have shown similar results, albeit using different methods, including invertebrates and vertebrates from both lab and wild populations (Bonduriansky and Chenoweth, 2009; Foerster et al., 2007; Mills et al., 2011). With the advent of advanced genomic tools, two model systems report specific examples of SA genetic loci: in Atlantic salmon (*Salmo salar*) the VGLL3 locus (Barson et al., 2015), and in *D. melanogaster* the DDT-R locus (Rostant et al., 2015), the Ala-278-Thr polymorphism in the mitochondrial genome (Camus et al., 2015), and multiple candidate loci from a recent genome-wide association study (Ruzicka et al., 2019).

Although humans have also been shown to experience SA selection for some quantitative traits (Garver-Apgar et al., 2011; Stearns et al., 2012; Stulp et al., 2012), and there

are numerous reports of genetic loci with sex-specific effects (i.e. effects that differ in magnitude between the sexes) on multiple complex traits or disease phenotypes (Gilks et al., 2014; Winkler et al., 2015), there are apparently no clear examples of SA loci in humans, which is inconsistent with theoretical expectations (Connallon and Clark, 2014). One potential explanation is that sexual antagonism is a weak selective force in humans, with sexual dimorphism being rather limited (Dixson, 2009; Short, 1979). This may indicate that concordant selection pressures between the sexes have dominated our evolutionary history. But the recent quantitative genetic studies challenge this view (Stearns et al., 2012; Stulp et al., 2012), and indeed theory predicts there is an inevitability to SA loci occurring in organisms with separate sexes (Connallon and Clark, 2014; Parker et al., 1972). Thus, there is a clear disparity between, on the one hand, theoretical expectations and quantitative genetic evidence that SA selection does occur in humans, and on the other hand, a complete lack of specific examples of SA loci. All this in the context of decades of research into how individual genetic variants influence disease profiles.

An alternative explanation for the absence of documented SA loci in humans is that since biomedical science does not use the same terminology for SA effects that evolutionary biology uses, there may in fact be examples that have been misclassified and therefore remain hidden in the literature. For example, genetic loci with sex-specific effects are frequently referred to as *sex-dependent* effects in the biomedical literature. Other terms are also used for related concepts, including *sex-different* effects. Furthermore, *sex-opposite* effects (i.e. a positive effect in one sex and a negative in the other) could be SA effects if the phenotype in question is deleterious, as disease phenotypes often are, since they can influence an individual's survival or reproductive fitness. The biomedical literature also frequently uses the terms *sex* and *gender* interchangeably, with *males* and *females* more often than not referred to as *men and women*, or *boys and girls* (Khramtsova et al., 2019). There is therefore ample scope for discipline-specific terms to be used for describing the same concept, which may hinder communication between disciplines and obscure examples of SA loci. Moreover, although there has been some progress in introducing concepts from evolutionary biology into the field of medicine (Nesse and Williams, 1994), those inroads have been relatively recent. It is therefore likely that the concepts of (intralocus) sexual conflict and sexual antagonism are not generally well known within biomedical science, and variants that may have SA effects on disease risk may not be referred to as such. A secondary related hypothesis

is that SA effects, when discovered, are discounted as errors as they do not match
expectations that sex-differences in phenotype or genetic architecture are not important
(Clayton and Collins, 2014). As a result, that may also lead them to be misclassified or simply
go unreported, leading to a general publication bias.

For these reasons, we propose that a systematic review of the biomedical literature
to identify SA loci in humans requires a specific and targeted set of search terms that would
not normally be used within the field of evolutionary biology. We test this assumption by
dividing our systematic review into two stages. In the first stage, we search for papers
identifying specific genetic loci using terms directly relating to the concept of sexual
antagonism and others utilized in evolutionary biology. We supplement this search with a
second stage, where we develop a set of terms that we hypothesize may capture the same
concept as sexual antagonism within the biomedical literature, although not explicitly stated
as such, as well as additional terminology relating to possible alternatives for describing the
two sexes. From our searches we also extracted data on sex-specific effect sizes and allele
frequencies. We were then able to explore how effect sizes varied for the same alleles
between the sexes, and test the prediction that SA alleles experiencing net positive selection
between the sexes may achieve higher equilibrium frequencies than alleles that experience
more symmetric or net negative patterns of selection (Morrow and Connallon, 2013). Our
systematic review aims to advance our understanding of SA genes in humans, which show
considerable and largely unexplained diversity in sexual dimorphism for disease phenotypes
(Ober et al., 2006; Rigby and Kulathinal, 2015), but we also highlight how and why
interdisciplinary research (in this case between biology and medicine) can sometimes fail,
impeding scientific advances over significant periods of time.

Methods

For this systematic review we followed PRISMA guidance where possible (Moher et al., 2009).
PubMed (https://pubmed.ncbi.nlm.nih.gov/) was searched for articles on 2nd December 2020
with no time limit. The searches were carried out in two Stages, with the organism filter set
to human in both stages. In Stage 1, eligible studies were required to report specific genetic
variants or haplotypes that were referred to as sexually antagonistic or were an example of
intralocus sexual conflict. To achieve this, we conducted a Boolean search for articles that
used the terms "sexual antagonism" OR "sexually antagonistic" OR "intralocus sexual conflict"

AND "locus" OR "loci", "gene" OR "snp" OR "polymorphism" OR "variant" OR "allele" in their

abstract or title. The Stage 1 search returned 34 articles in total (full search term in the

supplementary   material;   search   output   is   accessible   at

https://pubmed.ncbi.nlm.nih.gov/collections/60255050/?sort=pubdate).

In Stage 2, studies were required to report specific genetic variants or haplotypes in

humans with opposite effects in the two sexes on either complex traits, the outcome of a

medical intervention, or disease risk/severity. We define complex traits as likely with a

polygenic genetic architecture but are not directly related to a disease phenotype. In this

second stage search terms were specifically designed to include papers from the biomedical

literature that may have been missed in the first stage because they do not report their

findings with terms normally found within the evolutionary biology literature. Again, we

conducted a Boolean search for articles that used terms in their title or abstract to describe

an opposite or different effect in the two sexes ("sex dependent", "sex different", "gender-

dependent", "sex AND opposite", or "gender AND opposite"), or that capture this concept

with alternative words for sex (("male AND female AND opposite" OR "men AND women AND

opposite" OR "boys AND girls AND opposite") AND ("locus" OR "loci" OR "gene" OR "snp" OR

"polymorphism" OR "variant" OR "allele")). Full details of the search terms used and the

numbers of articles returned are provided in the supplementary material. The Stage 2 search

returned 881 papers (Figure 1) (full search term in the supplementary material; search output

is accessible at https://pubmed.ncbi.nlm.nih.gov/collections/60254985/?sort=pubdate).

The abstracts of the papers from Stage 1 were then examined by EM and from Stage

2 by JH, and any papers that had the possibility of reporting an opposite effect of a specific

genetic locus on a complex trait, medical intervention or on disease risk/severity were

considered for further screening. From Stage 1, no articles passed the screening. From Stage

2,   this   screening   produced   a   shortlist   of   70   candidate   papers

(https://pubmed.ncbi.nlm.nih.gov/collections/57906298/?sort=pubdate) Both JH and EM

then reviewed all the full texts of these candidate papers in detail. Papers were included in

the final list if they described a sex-opposite or SA effect linked to a specific genetic locus or

loci, and reported the effect to be statistically significant (at a P-value cut-off of <0.05). One

additional paper was considered from an outside source.

We converted all reported sex-specific effects into a standard effect size (Cohen's d)

quantifying the magnitude of how a given variant affects the studied trait expressed in the

given sex. Specifically, Cohen's d was computed based on the reported descriptive statistics (N, mean, standard error) or by conversion from other effect sizes (Odds ratio) and test statistics (F-values, t-values) using formulas reported elsewhere (Borenstein, 2009; Gurevitch et al., 2013, pp. 195–206; Lajeunesse, 2013). We sought information directly from the authors where these metrics were not possible to extract from the papers themselves (12 authors contacted, 5 responded, 3 responded with data, all later excluded as they did not fulfil the criteria for eligibility). We also recorded the gene name, locus (with accession/rs number where possible), trait affected and the frequency of the focal allele having the effect, hereafter referred to as *effect allele frequency*. Identification for the variants were taken from the studies where possible, but others necessitated searching the National Centre for Biotechnology Information (NCBI) to find the Reference SNP cluster ID for the variants described. Where effect allele frequencies were not reported, we used genotype frequencies to calculate effect allele frequency. Not all studies reported allele or genotype frequencies and so we attempted to supplement these data with allele frequencies from the 1000 genomes database (Auton et al., 2015), since these show a strong correlation with effect allele frequencies reported in the studies reviewed (Pearson correlation: $N = 25$ $r = 0.94$, $P > 0.001$; Figure S1). However, using this approach we were only able to supplement our allele frequency data for one additional locus (rs7341475 in the *RELN* gene; Table 1).

     If b_f and b_m represent the estimated effect of the focal allele in females and males respectively, then positive values of b_f or b_m imply that the allele is positively associated with disease expression, whereas negative values imply that the allele is negatively associated with disease expression. SA alleles are therefore defined as those with opposite effects between the sexes (i.e. $b_f < 0 < b_m$ or $b_m < 0 < b_f$). An evolutionary model of evolution at SA loci predicts that as the magnitude of the positive effect that an SA allele has outweighs the negative effect, the higher the frequency that allele can achieve (Morrow and Connallon, 2013). We therefore expect a negative relationship between the ratio of positive to negative effect sizes (hereafter referred to as the *effect size ratio*) and observed allele frequency. To investigate this, we calculated the effect size ratio by dividing the positive standardized effect size by the negative standardized effect size; when $b_f < 0 < b_m$ effect size ratio was calculated as b_m / b_f, and when $b_m < 0 < b_f$ effect size ratio was calculated as b_f / b_m. Thus, SA alleles with a greater beneficial effect will have a smaller, more negative effect size ratio (<-1), while SA alleles that have a greater deleterious effect will have a larger effect size ratio (>-1), up to a maximum

value of 0 (positive values occur when the effect is positive or negative in both sexes, but the

allele would then no longer be defined as a SA allele).

We modelled how effect allele frequency changes with effect size ratio using a

generalized linear model (GLM), weighted by the inverse of the variance of the effect sizes

such that data points with smaller variance have a higher weight, since smaller sample sizes

were associated with larger and more variable effect sizes (Figure S2). We initially also

included trait class and its interaction with effect size ratio as a fixed factor with two levels

(complex trait and disease trait/severity), since alleles that influence complex traits in

opposite directions are not necessarily under SA selection and so may not behave in the way

predicted, whereas alleles influencing disease traits, unless very late acting, are more likely

to show a closer relationship with marginal effects on fitness. As a response variable, effect

allele frequency is limited between 0 and 1, so we looked at allele counts to allow allele

frequency to vary freely. GLMs with binomial error distribution showed substantial

overdispersion, so we used a quasibinomial function to address this issue (Payne et al., 2018).

We used a Chi-squared test to infer significance of the two predictor variables and their

interaction. The full model indicated that the interaction term and trait class have no

significant effect, so that we report the fit of the reduced model which only includes effect

size ratio. We also subsequently modelled the data for the two trait classes (complex traits

and disease risk/severity) separately, again using a quasibinomial distribution function, to see

if the result was replicated in these smaller subsets of the data. The raw data and R script is

available for replicating the analyses and figures we present (Supplementary Material).

Results

The Stage 1 search found no articles that described genetic loci in humans with effects that

were sexually antagonistic. In contrast, the Stage 2 search identified 32 articles that described

variants with statistically significant sex-opposite or SA effects (Figure 1;

https://pubmed.ncbi.nlm.nih.gov/collections/60278165/?sort=pubdate). From the studies

examined, 51 SA variants were identified (Table 1), affecting 21 different complex traits (30

loci) and 19 disease risk/severity traits (21 loci), with a large range of effect sizes. There were

no variants affecting medical intervention traits that passed the screening process. The

majority of alleles had effect sizes of similar absolute values in the two sexes, with larger

effects tending to show greater differences on absolute effect size (Figure S3).

We found 22 studies provided sufficient data to allow the relationship between effect

size ratio and effect allele frequency to be explored by statistical modelling. Trait class and its

interaction with effect size ratio were initially included in the model but neither were found

to have a significant effect (GLM: trait class dfs = 1,35, deviance = 15.5 × 10^3, P = 0.965;

interaction dfs = 1,34, deviance = 1476, P = 0.989) and were therefore excluded from the

subsequent model in which effect size ratio was found to relate negatively to effect allele

frequency, as predicted (GLM: estimate ± SE = -1.10 ± 0.50, dfs = 1,36, deviance = 38.7 × 10^6,

P = 0.024; Figure 2). For three loci there were data on effect sizes for more than one trait

(Table 1), which introduced a degree of non-independence between these values in the

dataset. However, resampling the data with effect sizes for only one trait for these three loci

in turn and modelling these datasets in the same way (18 individual GLMs, dfs = 1,31) did not

change the results qualitatively (minimum, maximum: model estimate = -1.0963, -1.0964;

deviance = 38.72 × 10^6, 38.73 × 10^6; P = 0.03618, 0.03620).

Although the interaction term between trait type and effect size ratio was not

significant we wanted to investigate whether the negative relationship between effect size

ratio and effect allele frequency across all loci was repeated when the data was divided

according to trait class, since while opposite effects on disease risk/severity may show a direct

relationship with SA fitness effects, complex traits may or may not be experiencing SA

selection, even if effects are in opposite directions in the two sexes. However, although the

trends were both again negative as predicted, the generalized linear models of these smaller

datasets were only marginally significant (complex traits model estimate ± SE = -1.10 ± 0.61,

dfs = 1,24, deviance = 38.7 × 106, P = 0.065; disease risk/severity model estimate ± SE = -1.34

± 0.77, dfs = 1,10, deviance = 5.14 × 104, P = 0.053; Figure S4). As for the full dataset, there

was some lack of independence for the complex traits subset which contained loci with effect

sizes for more than one trait. Again, resampling and modelling these datasets in the same

way (18 individual glms, dfs 1, 22) did not change the results quantitatively (minimum,

maximum: model estimate = -1.0960, -1.0961; deviance = 38.669 × 10^6, 38.675 × 10^6; P =

0.07774, 0.07776).

Discussion

280   The Stage 1 literature search of our systematic review focussed on finding reports of specific

281   genetic loci in humans with evidence for SA effects using terminology normally associated

with the evolutionary concepts of sexual conflict or sexual antagonism. The Stage 2 search sought equivalent evidence, but used search terms that we anticipated would be used by scientists outside the field of evolutionary biology, who may not use the same terminology. Although the Stage 1 search did not find any examples of SA loci occurring in humans, the Stage 2 search identified 51 genetic loci across 39 studies that had ostensibly SA effects, but were not described as such. Clearly then the Stage 1 review failed to identify multiple relevant reports going back some 20 years because the terminology used in those reports did not match the conceptual framework of the search. The Stage 2 search may also be a lower limit given searches of other larger databases may also harbour further examples (see below). Although not yet validated, these reports nonetheless represent a substantial body of evidence that humans, like other organisms with separate sexes (Bonduriansky and Chenoweth, 2009), inevitably experience SA selection for a wide range of complex traits (Connallon and Clark, 2014; Garver-Apgar et al., 2011; Stearns et al., 2012; Stulp et al., 2012), as well as for a range of diseases.

A key prediction from a population genetic model of SA genetic variation is that SA alleles are expected to achieve higher equilibrium frequencies, without necessarily going to fixation, as the relative magnitude of the positive effect in one sex outweighs the negative effect in the opposite sex i.e. an increasingly negative effect size ratio (Morrow and Connallon, 2013). We found clear support for this prediction, with a negative relationship between effect size ratio and allele frequency when examining all complex traits and diseases together (Fig 2), although this relationship was only marginally significant when complex or disease trait classes were modelled separately (Fig S3). These results lend support to the view that the loci identified are genuinely experiencing SA selection, and although it is difficult to discern how selection acts on trait size in males and females for complex traits we would not expect the frequencies of disease-causing alleles to be so high under a mutation-selection balance. We were not able to look at the evolutionary history or age of these alleles due to limited information (n = 8) (Albers and McVean, 2018), it would nonetheless be valuable to examine the population dynamics of these alleles over a broad timescale.

The loci showed a very broad range of effect sizes, from small to very large, with a generally symmetrical inverse relationship between effect size in one sex and the other (Figure S2), which is generally expected, since our screening process necessarily excluded studies reporting effects in the same direction across the two sexes. Nonetheless, it is striking

just how large some of the effects were, with very large negative effects in one sex
simultaneously occurring with similarly large positive effects. Larger and more variable effect
sizes are expected when sample sizes are small, a pattern we also found, which motivates
further investigation of the traits and loci included in this review to elucidate more accurate
estimates of real effect sizes.

The loci themselves fall within genes that influence a broad range of phenotypes,
including morphological, physiological and behavioural complex traits, and a similarly diverse
range of disease types, including various cancers, neurological disorders and immune system
processes (see Table 1). For the most part, the diseases appear to be early-acting rather than
late-onset, with the exception of perhaps bone mineral density. As such, it seems reasonable
that an allele that increases disease risk or severity in these cases will indeed have a
concomitant reduction in marginal fitness. Although a single large study identified several loci
related to BMI adjusted waist-to-hip ratio, there is generally no overall bias towards one
particular trait or disease class. The total number of loci is relatively small however, which
may limit power to identify such biases if they exist. We also did not find any examples of SA
loci on the X-chromosome, which may or may not be a hotspot of sexual antagonism (Fry,
2010; Rice, 1984; Ruzicka and Connallon, 2020). This was expected given that it is commonly
not included in genome-wide analyses (Wise et al., 2013), although most of the papers we
reviewed were candidate gene studies. Nonetheless, we anticipate further examples may be
identified should data from the X-chromosome be included systematically in association
studies.

We also found evidence that some traits are influenced by more than one genetic
variant in sex-opposite or SA ways (n=8), or that some specific variants have pleiotropic
effects on more than one trait or disease in a sex-opposite or SA way (n=6; see Table 1). This
complexity in the genetic architecture of traits or diseases may make conflict resolution
particularly difficult, as a change in the allele at a single genetic locus experiencing SA
selection may simultaneously influence multiple genes or phenotypes in both sexes in
divergent ways (Fitzpatrick, 2004; Pennell and Morrow, 2013). Consequently, this may explain
the persistence of intralocus sexual conflict at these loci. It may be that some of the remaining
loci also have as yet unidentified pleiotropic effects with other traits or there are as yet
unidentified SA loci influencing those same traits or diseases. Of the 51 variants identified, 4
were confirmed to be haplotypes. Such variants, consisting of two or more SNPs in linkage

disequilibrium having a joint effect, could also present a problem for conflict resolution, since

the ability of selection to act on any individual locus independent of the others in the linkage

block is reduced for linked loci. This issue may also extend to other single variants reported

here if they also occur in linkage blocks.

Although we have identified multiple genetic loci with either sex-opposite or SA

effects these are likely outnumbered by those with either sex-specific (same direction but

different magnitude) or sex-limited effects. For instance, Winkler et al. (2015) report 44 loci

with sex-specific or sex-limited effects but only 11 with sex-opposite effects. Several other

genome-wide association studies report sex-specific effects in several human diseases

(Khramtsova et al., 2019), and a recent (non-systematic) review identified 37 SNPs with sex-

dependent effects (Gilks et al., 2014). Detection of loci with SA versus sex-specific or sex-

limited effects may differ systematically since we expect SA loci are more likely to persist at

intermediate frequencies than loci with sex-specific effects (i.e. that differ between the sexes

in magnitude but not sign). A larger systematic review that targeted variants with sex-specific

or sex-dependent effects would therefore be a valuable contribution and enable us to more

clearly understand how important SA alleles are relative to the broader context of genes with

non-identical effects in the two sexes.

A key gap in our knowledge is whether the putative examples of sex-opposite or SA

alleles presented here can be validated using independently derived datasets in the same or

different sub-populations. For this, focussing on those variants associated with sex-

differential disease risk would naturally be the most fruitful for advancing our knowledge of

sexual antagonism in human disease. We therefore encourage specialists for the particular

disease groups presented in Table 1 to include in the future the potential for SA genetic

effects to occur when designing studies and analysing data. It is also important for all studies

to report results in sufficient detail so that effect sizes can be calculated (if not given) as well

as allele frequency data. Unfortunately, 17 studies failed to do this, so could not be included

in our review and subsequent test of the evolutionary model. More generally, our study

reinforces the view that sex is an important factor in shaping genetic associations with human

traits and disease, even in divergent and contradictory ways, and so should always be

considered when investigators examine genetic associations with phenotypes (Lee, 2018;

Ober et al., 2008).

There is a possibility that some bias exists in the PubMed database against papers
from the field of evolutionary biology, and if we had used another database then both stages
would have returned valid papers. We subsequently (14th Dec 2020) sought to investigate this
possibility with additional searches of Scopus and Web of Science. Repeating the Stage 1
search initially returned 85 papers in Web of Science (Basic search of Topic, refined by
"Human" within results) and 71 papers in Scopus (include keywords "Human", exclude "Non-
Human"), but again no papers made it through the screening process. Repeating the Stage 2
search returned a much larger number of studies in both Web of Science (4,573; Basic search
of Topic, refined by "Human" within results) and Scopus (6,616; Advanced search of Title-
Abstract, filter: include human and humans, exclude nonhumans, Articles only), with 8,604
unique records. This large number of records is too many to currently screen and may harbour
further examples of SA loci in humans, but it is noteworthy that the majority of the initial list
of studies found during Stage 2 using PubMed were also found in the other two databases
(712 out of 881) as well as 28 out of the screened list of 32 studies identified in the current
systematic review.

The disparity between the two stages of the review process provides a remarkable
example of how discipline-specific terminology and concepts can hinder scientific
communication between fields for substantial periods of time. For many of the studies
identified in this review, the focus of the research was not on quantifying or even identifying
sex-specific genetic effects in traits or disease, although they were tested for. The responses
to such findings varied. Some framed these results as a major finding, others merely made
note of them, with one suggesting that since the genetic associations were in opposite
directions in the two sexes then it should be regarded as a false positive (Wong et al., 2005).
It is possible, or even likely then that there exists a publication bias in the biomedical sciences
against studies with apparently incongruous sex-opposite or SA effects, perhaps in part
because the evolutionary framework that may provide a context for understanding those
results is not generally appreciated or known. Of course, for evolutionary biologists interested
in the role of SA selection on human traits and disease, this is clearly a problem. For many
years no specific examples of SA loci were known in humans, and yet the first putative
examples in fact appeared in the literature two decades ago. Our systematic review therefore
provides a concrete illustration of how advances to both fields have likely been hampered by

a limited understanding of the theoretical framework and terminology used by the
neighbouring field.

How can communication between the fields of medicine and evolutionary biology be
facilitated? We do not have a comprehensive answer but clearly training in, or an increased
awareness of, evolutionary medicine for scientists in both camps would probably be
beneficial. The topic of evolutionary biology now forms a key part of the education for some
biomedical scientists and clinicians. Similarly, evolutionary medicine is now frequently
included in fundamental textbooks on evolution, and is being taught to students of
evolutionary biology as part of their degree courses. These changes have been relatively
recent, are far from being universally employed, but they should be encouraged. Funders of
primary research are also now actively encouraging or obliging researchers to include sex as
a factor when designing their studies (Clayton and Collins, 2014), although this may simply
result in studies that aim to control for rather than investigate these effects explicitly.
Consequently even this remedy may still preclude a deeper understanding of the origins of
some human diseases.

Acknowledgements

425 We thank Jessica Abbott, Tim Connallon and Filip Ruzicka for comments on a previous version
426 of the MS. Funding was provided by the Swedish Research Council (Grant number: 2019-
427 03567) and by the Royal Society to EHM as a University Research Fellowship and
428 Enhancement Award. TJ was funded by the Centre national de la recherche scientifique
429 (CNRS) and the German Research Foundation (DFG grant number: JA 2653/2-1).

References

Albers, P.K., McVean, G., 2018. Dating genomic variants and shared ancestry in population-
scale sequencing data (preprint). Genomics. https://doi.org/10.1101/416610
Auton, A., Abecasis, G.R., Altshuler, D.M., Durbin, R.M., Abecasis, G.R., Bentley, D.R.,
Chakravarti, A., Clark, A.G., Donnelly, P., Eichler, E.E., Flicek, P., Gabriel, S.B., Gibbs,
R.A., Green, E.D., Hurles, M.E., Knoppers, B.M., Korbel, J.O., Lander, E.S., Lee, C.,
Lehrach, H., Mardis, E.R., Marth, G.T., McVean, G.A., Nickerson, D.A., Schmidt, J.P.,
Sherry, S.T., Wang, J., Wilson, R.K., Gibbs, R.A., Boerwinkle, E., Doddapaneni, H., Han,
Y., Korchina, V., Kovar, C., Lee, S., Muzny, D., Reid, J.G., Zhu, Y., Wang, J., Chang, Y.,
Feng, Q., Fang, X., Guo, X., Jian, M., Jiang, H., Jin, X., Lan, T., Li, G., Li, J., Li, Y., Liu, S.,
Liu, X., Lu, Y., Ma, X., Tang, M., Wang, B., Wang, G., Wu, H., Wu, R., Xu, X., Yin, Y.,

Zhang, D., Zhang, W., Zhao, J., Zhao, M., Zheng, X., Lander, E.S., Altshuler, D.M., Gabriel, S.B., Gupta, N., Gharani, N., Toji, L.H., Gerry, N.P., Resch, A.M., Flicek, P., Barker, J., Clarke, L., Gil, L., Hunt, S.E., Kelman, G., Kulesha, E., Leinonen, R., McLaren, W.M., Radhakrishnan, R., Roa, A., Smirnov, D., Smith, R.E., Streeter, I., Thormann, A., Toneva, I., Vaughan, B., Zheng-Bradley, X., Bentley, D.R., Grocock, R., Humphray, S., James, T., Kingsbury, Z., Lehrach, H., Sudbrak, R., Albrecht, M.W., Amstislavskiy, V.S., Borodina, T.A., Lienhard, M., Mertes, F., Sultan, M., Timmermann, B., Yaspo, M.-L., Mardis, E.R., Wilson, R.K., Fulton, L., Fulton, R., Sherry, S.T., Ananiev, V., Belaia, Z., Beloslyudtsev, D., Bouk, N., Chen, C., Church, D., Cohen, R., Cook, C., Garner, J., Hefferon, T., Kimelman, M., Liu, C., Lopez, J., Meric, P., O'Sullivan, C., Ostapchuk, Y., Phan, L., Ponomarov, S., Schneider, V., Shekhtman, E., Sirotkin, K., Slotta, D., Zhang, H., McVean, G.A., Durbin, R.M., Balasubramaniam, S., Burton, J., Danecek, P., Keane, T.M., Kolb-Kokocinski, A., McCarthy, S., Stalker, J., Quail, M., Schmidt, J.P., Davies, C.J., Gollub, J., Webster, T., Wong, B., Zhan, Y., Auton, A., Campbell, C.L., Kong, Y., Marcketta, A., Gibbs, R.A., Yu, F., Antunes, L., Bainbridge, M., Muzny, D., Sabo, A., Huang, Z., Wang, J., Coin, L.J.M., Fang, L., Guo, X., Jin, X., Li, G., Li, Q., Li, Y., Li, Z., Lin, H., Liu, B., Luo, R., Shao, H., Xie, Y., Ye, C., Yu, C., Zhang, F., Zheng, H., Zhu, H., Alkan, C., Dal, E., Kahveci, F., Marth, G.T., Garrison, E.P., Kural, D., Lee, W.-P., Fung Leong, W., Stromberg, M., Ward, A.N., Wu, J., Zhang, M., Daly, M.J., DePristo, M.A., Handsaker, R.E., Altshuler, D.M., Banks, E., Bhatia, G., del Angel, G., Gabriel, S.B., Genovese, G., Gupta, N., Li, H., Kashin, S., Lander, E.S., McCarroll, S.A., Nemesh, J.C., Poplin, R.E., Yoon, S.C., Lihm, J., Makarov, V., Clark, A.G., Gottipati, S., Keinan, A., Rodriguez-Flores, J.L., Korbel, J.O., Rausch, T., Fritz, M.H., Stütz, A.M., Flicek, P., Beal, K., Clarke, L., Datta, A., Herrero, J., McLaren, W.M., Ritchie, G.R.S., Smith, R.E., Zerbino, D., Zheng-Bradley, X., Sabeti, P.C., Shlyakhter, I., Schaffner, S.F., Vitti, J., Cooper, D.N., Ball, E.V., Stenson, P.D., Bentley, D.R., Barnes, B., Bauer, M., Keira Cheetham, R., Cox, A., Eberle, M., Humphray, S., Kahn, S., Murray, L., Peden, J., Shaw, R., Kenny, E.E., Batzer, M.A., Konkel, M.K., Walker, J.A., MacArthur, D.G., Lek, M., Sudbrak, R., Amstislavskiy, V.S., Herwig, R., Mardis, E.R., Ding, L., Koboldt, D.C., Larson, D., Ye, K., Gravel, S., The 1000 Genomes Project Consortium, Corresponding authors, Steering committee, Production group, Baylor College of Medicine, BGI-Shenzhen, Broad Institute of MIT and Harvard, Coriell Institute for Medical Research, European Molecular Biology Laboratory, E.B.I., Illumina, Max Planck Institute for Molecular Genetics, McDonnell Genome Institute at Washington University, US National Institutes of Health, University of Oxford, Wellcome Trust Sanger Institute, Analysis group, Affymetrix, Albert Einstein College of Medicine, Bilkent University, Boston College, Cold Spring Harbor Laboratory, Cornell University, European Molecular Biology Laboratory, Harvard University, Human Gene Mutation Database, Icahn School of Medicine at Mount Sinai, Louisiana State University, Massachusetts General Hospital, McGill University, National Eye Institute, N., 2015. A global reference for human genetic variation. Nature 526, 68–74. https://doi.org/10.1038/nature15393

Barson, N.J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G.H., Fiske, P., Jacq, C., Jensen, A.J., Johnston, S.E., Karlsson, S., Kent, M., Moen, T., Niemelä, E., Nome, T., Næsje, T.F., Orell, P., Romakkaniemi, A., Sægrov, H., Urdal, K., Erkinaro, J., Lien, S., Primmer, C.R., 2015. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. Nature 528, 405–408. https://doi.org/10.1038/nature16062

Bonduriansky, R., Chenoweth, S.F., 2009. Intralocus sexual conflict. Trends Ecol Evol 24,
280–288.
Borenstein, M., 2009. Converting Among Effect Sizes, in: Introduction to Meta-Analysis. John
Wiley & Sons, Ltd, pp. 45–49. https://doi.org/10.1002/9780470743386.ch7
Camus, M.F., Wolf, J.B.W., Morrow, E.H., Dowling, D.K., 2015. Single Nucleotides in the
mtDNA Sequence Modify Mitochondrial Molecular Function and Are Associated with
Sex-Specific Effects on Fertility and Aging. Curr. Biol. 25, 2717–2722.
https://doi.org/10.1016/j.cub.2015.09.012
Carlson, J.C., Nidey, N.L., Butali, A., Buxo, C.J., Christensen, K., Deleyiannis, F.W.-D., Hecht,
J.T., Field, L.L., Moreno-Uribe, L.M., Orioli, I.M., Poletta, F.A., Padilla, C., Vieira, A.R.,
Weinberg, S.M., Wehby, G.L., Feingold, E., Murray, J.C., Marazita, M.L., Leslie, E.J.,
2018. Genome-wide interaction studies identify sex-specific risk alleles for
nonsyndromic orofacial clefts. Genet. Epidemiol. 42, 664–672.
https://doi.org/10.1002/gepi.22158
Chippindale, A.K., Gibson, J.R., Rice, W.R., 2001. Negative genetic correlation for adult
fitness between sexes reveals ontogenetic conflict in Drosophila. Proc. Natl. Acad.
Sci. USA 98, 1671–1675.
Cicchetti, D., Rogosch, F.A., Hecht, K.F., Crick, N.R., Hetzel, S., 2014. Moderation of
maltreatment effects on childhood borderline personality symptoms by gender and
oxytocin receptor and FK506 binding protein 5 genes. Dev. Psychopathol. 26, 831–
849. https://doi.org/10.1017/S095457941400042X
Clayton, J.A., Collins, F.S., 2014. Policy: NIH to balance sex in cell and animal studies. Nature
509, 282–283. https://doi.org/10.1038/509282a
Connallon, T., Clark, A., G., 2014. Evolutionary inevitability of sexual antagonism. Proc. R.
Soc. B Biol. Sci. 281, 20132123. https://doi.org/10.1098/rspb.2013.2123
Connallon, T., Clark, A.G., 2013. Antagonistic Versus Nonantagonistic Models of Balancing
Selection: Characterizing the Relative Timescales and Hitchhiking Effects of Partial
Selective Sweeps. Evolution 67, 908–917. https://doi.org/10.1111/j.1558-
5646.2012.01800.x
Dauvilliers, Y., Neidhart, E., Lecendreux, M., Billiard, M., Tafti, M., 2001. MAO-A and COMT
polymorphisms and gene effects in narcolepsy. Mol. Psychiatry 6, 367–372.
https://doi.org/10.1038/sj.mp.4000911
Dixson, A., 2009. Dixson, A. F. (2009) Sexual selection and the Origins of Human Mating
Systems. Oxford University Press, UK.
Ellegren, H., Parsch, J., 2007. The evolution of sex-biased genes and sex-biased gene
expression. Nat Rev Genet 8, 689–698. https://doi.org/10.1038/nrg2167
Fillingim, R.B., Kaplan, L., Staud, R., Ness, T.J., Glover, T.L., Campbell, C.M., Mogil, J.S.,
Wallace, M.R., 2005. The A118G single nucleotide polymorphism of the mu-opioid
receptor gene (OPRM1) is associated with pressure pain sensitivity in humans. J. Pain
Off. J. Am. Pain Soc. 6, 159–167. https://doi.org/10.1016/j.jpain.2004.11.008
Fitzpatrick, M.J., 2004. Pleiotropy and the genomic location of sexually selected genes. Am.
Nat. 163, 800–808. https://doi.org/10.1086/386297
Foerster, K., Coulson, T., Sheldon, B.C., Pemberton, J.M., Clutton-Brock, T., Kruuk, L.E.B.,
2007. Sexually antagonistic genetic variation for fitness in red deer. Nature 447,
1107–1110.
Franck, N., Lanne, T., Astrand, O., Engvall, J., Lindstrom, T., Ostgren, C.J., Nystrom, F.H.,
2012. Cardiovascular risk factors related to the PPARgamma Pro12Ala polymorphism

in patients with type 2 diabetes are gender dependent. Blood Press. 21, 122–127.
https://doi.org/10.3109/08037051.2011.623349

Fry, J.D., 2010. The Genomic Location of Sexually Antagonistic Variation: Some Cautionary
Comments. Evolution 64, 1510–1516. https://doi.org/10.1111/j.1558-
5646.2009.00898.x

Garver-Apgar, C.E., Eaton, M.A., Tybur, J.M., Emery Thompson, M., 2011. Evidence of
intralocus sexual conflict: physically and hormonally masculine individuals have more
attractive brothers relative to sisters. Evol. Hum. Behav. 32, 423–432.
https://doi.org/10.1016/j.evolhumbehav.2011.03.005

Gastaldi, M., Diziere, S., Defoort, C., Portugal, H., Lairon, D., Darmon, M., Planells, R., 2007.
Sex-specific association of fatty acid binding protein 2 and microsomal triacylglycerol
transfer protein variants with response to dietary lipid changes in the 3-mo Medi-
RIVAGE primary intervention study. Am. J. Clin. Nutr. 86, 1633–1641.
https://doi.org/10.1093/ajcn/86.5.1633

Gilks, W.P., Abbott, J.K., Morrow, E.H., 2014. Sex differences in disease genetics: evidence,
evolution, and detection. Trends Genet. 30, 453–463.
https://doi.org/10.1016/j.tig.2014.08.006

Gloria-Bottini, F., Bottini, N., Renzetti, G., Bottini, E., 2007. ACP1 and Th class of
immunological disease: evidence of interaction with gender. Int. Arch. Allergy
Immunol. 143, 170–176. https://doi.org/10.1159/000099308

Grieshop, K., Arnqvist, G., 2018. Sex-specific dominance reversal of genetic variation for
fitness. PLOS Biol. 16, e2006810. https://doi.org/10.1371/journal.pbio.2006810

Gurevitch, J., Koricheva, J., Mengersen, K., 2013. Handbook of Meta-Analysis In Ecology and
Evolution, Handbook of Meta-analysis in Ecology and Evolution.
https://doi.org/10.1515/9781400846184

Herman, A.I., Conner, T.S., Anton, R.F., Gelernter, J., Kranzler, H.R., Covault, J., 2011.
Variation in the gene encoding the serotonin transporter is associated with a
measure of sociopathy in alcoholics. Addict. Biol. 16, 124–132.
https://doi.org/10.1111/j.1369-1600.2009.00197.x

Khramtsova, E.A., Davis, L.K., Stranger, B.E., 2019. The role of sex in the genomics of human
complex traits. Nat. Rev. Genet. 20, 173–190. https://doi.org/10.1038/s41576-018-
0083-1

Kim, B., Kim, H., Joo, Y.H., Lim, J., Kim, C.-Y., Song, K., 2010. Sex-different association of DAO
with schizophrenia in Koreans. Psychiatry Res. 179, 121–125.
https://doi.org/10.1016/j.psychres.2008.08.009

Kiy, A., Wilhelm, O., Hildebrandt, A., Reuter, M., Sommer, W., 2013. On the genetic basis of
face cognition and its relation to fluid cognitive abilities. Genes Brain Behav. 12, 438–
445. https://doi.org/10.1111/gbb.12034

Korner, A., Ma, L., Franks, P.W., Kiess, W., Baier, L.J., Stumvoll, M., Kovacs, P., 2007. Sex-
specific effect of the Val1483Ile polymorphism in the fatty acid synthase gene (FAS)
on body mass index and lipid profile in Caucasian children. Int. J. Obes. 2005 31,
353–358. https://doi.org/10.1038/sj.ijo.0803428

Lajeunesse, M., 2013. Recovering missing or partial data from studies: A survey of
conversions and imputations for meta-analysis. pp. 195–206.

Lande, R., 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic
characters. Evolution 34, 292–305.

Lang, U.E., Hellweg, R., Bajbouj, M., Gaus, V., Sander, T., Gallinat, J., 2008. Gender-
dependent association of a functional NGF polymorphism with anxiety-related
personality traits. Pharmacopsychiatry 41, 196–199. https://doi.org/10.1055/s-0028-
1082070

Lauritzen, L., Sorensen, L.B., Harslof, L.B., Ritz, C., Stark, K.D., Astrup, A., Dyssegaard, C.B.,
Egelund, N., Michaelsen, K.F., Damsgaard, C.T., 2017. Mendelian randomization
shows sex-specific associations between long-chain. Am. J. Clin. Nutr. 106, 88–95.
https://doi.org/10.3945/ajcn.117.152595

Lee, S.K., 2018. Sex as an important biological variable in biomedical research. BMB Rep. 51,
167–173. https://doi.org/10.5483/BMBRep.2018.51.4.034

Liu, Y.-H., Chen, Y.-J., Wu, H.-H., Wang, T.-Y., Tsai, F.-J., 2014. Single nucleotide
polymorphisms at the PRR3, ABCF1, and GNL1 genes in the HLA class I region are
associated with Graves' ophthalmopathy in a gender-dependent manner.
Ophthalmology 121, 2033–2039. https://doi.org/10.1016/j.ophtha.2014.04.027

Ljungberg, L.U., Ostgren, C.J., Nystrom, F.H., Lanne, T., 2014. Associations of genetic
polymorphisms in the renin-angiotensin system with central aortic and ambulatory
blood pressure in type 2 diabetic patients. J. Renin-Angiotensin-Aldosterone Syst.
JRAAS 15, 61–68. https://doi.org/10.1177/1470320312474052

McCarthy, J.J., Somji, A., Weiss, L.A., Steffy, B., Vega, R., Barrett-Connor, E., Talavera, G.,
Glynne, R., 2009. Polymorphisms of the scavenger receptor class B member 1 are
associated with insulin resistance with evidence of gene by sex interaction. J. Clin.
Endocrinol. Metab. 94, 1789–1796. https://doi.org/10.1210/jc.2008-2800

Mills, S.C., Koskela, E., Mappes, T., 2011. Intralocus sexual conflict for fitness: sexually
antagonistic alleles for testosterone. Proc. R. Soc. B Biol. Sci. rspb20112340.
https://doi.org/10.1098/rspb.2011.2340

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., Group, T.P., 2009. Preferred Reporting
Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLOS
Med. 6, e1000097. https://doi.org/10.1371/journal.pmed.1000097

Morea, A., Cameran, M., Rebuffi, A.G., Marzenta, D., Marangon, O., Picci, L., Zacchello, F.,
Scarpa, M., 2005. Gender-sensitive association of CFTR gene mutations and 5T allele
emerging from a large survey on infertility. Mol. Hum. Reprod. 11, 607–614.
https://doi.org/10.1093/molehr/gah214

Morrow, E.H., 2015. The evolution of sex differences in disease. Biol. Sex Differ. 6, 5.
https://doi.org/10.1186/s13293-015-0023-0

Morrow, E.H., Connallon, T., 2013. Implications of sex-specific selection for the genetic basis
of disease. Evol. Appl. 6, 1208–1217. https://doi.org/10.1111/eva.12097

Nesse, R.M., Williams, G.C., 1994. Why we get sick: The new science of Darwinian medicine.
Vintage, New York.

Ober, C., Loisel, D.A., Gilad, Y., 2008. Sex-specific genetic architecture of human disease. Nat
Rev Genet 9, 911–922. https://doi.org/10.1038/nrg2415

Ober, C., Pan, L., Phillips, N., Parry, R., Kurina, L.M., 2006. Sex-specific genetic architecture of
asthma-associated quantitative trait loci in a founder population. Curr. Allergy
Asthma Rep. 6, 241–246. https://doi.org/10.1007/s11882-006-0041-4

Ovadia, G., Shifman, S., 2011. The genetic variation of RELN expression in schizophrenia and
bipolar disorder. PloS One 6, e19955. https://doi.org/10.1371/journal.pone.0019955

Padmanabhan, S., Menni, C., Lee, W.K., Laing, S., Brambilla, P., Sega, R., Perego, R., Grassi,
G., Cesana, G., Delles, C., Mancia, G., Dominiczak, A.F., 2010. The effects of sex and

method of blood pressure measurement on genetic associations with blood pressure
in the PAMELA study. J. Hypertens. 28, 465–477.
https://doi.org/10.1097/HJH.0b013e32833594d7
Parker, G.A., 1979. SEXUAL SELECTION AND SEXUAL CONFLICT, in: Sexual Selection and
Reproductive Competition in Insects. Elsevier, pp. 123–166.
https://doi.org/10.1016/B978-0-12-108750-0.50010-0
Parker, G.A., Baker, R.R., Smith, V.G.F., 1972. The origin and evolution of gamete
dimorphism and the male-female phenomenon. J. Theor. Biol. 36, 529–553.
Payne, E.H., Gebregziabher, M., Hardin, J.W., Ramakrishnan, V., Egede, L.E., 2018. An
empirical approach to determine a threshold for assessing overdispersion in Poisson
and negative binomial models for count data. Commun. Stat. Simul. Comput. 47,
1722–1738. https://doi.org/10.1080/03610918.2017.1323223
Peng, Y.-M., Lei, S.-F., Guo, Y., Xiong, D.-H., Yan, H., Wang, L., Guo, Y.-F., Deng, H.-W., 2008.
Sex-specific association of the glucocorticoid receptor gene with extreme BMD. J.
Bone Miner. Res. Off. J. Am. Soc. Bone Miner. Res. 23, 247–252.
https://doi.org/10.1359/jbmr.071017
Pennell, T.M., Haas, F.J.H. de, Morrow, E.H., Doorn, G.S. van, 2016. Contrasting effects of
intralocus sexual conflict on sexually antagonistic coevolution. Proc. Natl. Acad. Sci.
113, E978–E986. https://doi.org/10.1073/pnas.1514328113
Pennell, T.M., Morrow, E.H., 2013. Two sexes, one genome: the evolutionary dynamics of
intralocus sexual conflict. Ecol. Evol. 3, 1819–1834.
https://doi.org/10.1002/ece3.540
Press, O.A., Zhang, W., Gordon, M.A., Yang, D., Haiman, C.A., Azuma, M., Iqbal, S., Lenz, H.-
J., 2011. Gender-related survival differences associated with polymorphic variants of
estrogen receptor-beta (ERbeta) in patients with metastatic colon cancer.
Pharmacogenomics J. 11, 375–382. https://doi.org/10.1038/tpj.2010.45
Price, J.S., Strong, J., Eliassen, J., McQueeny, T., Miller, M., Padula, C.B., Shear, P., Lisdahl, K.,
2013. Serotonin transporter gene moderates associations between mood, memory
and hippocampal volume. Behav. Brain Res. 242, 158–165.
https://doi.org/10.1016/j.bbr.2012.11.013
Rice, W.R., 1992. Sexually antagonistic genes: experimental evidence. Science 256, 1436–
1439. https://doi.org/10.1126/science.1604317
Rice, W.R., 1984. Sex chromosomes and the evolution of sexual dimorphism. Evolution 38,
735–742.
Rigby, N., Kulathinal, R.J., 2015. Genetic Architecture of Sexual Dimorphism in Humans. J.
Cell. Physiol. 230, 2304–2310. https://doi.org/10.1002/jcp.24979
Rostant, W.G., Kay, C., Wedell, N., Hosken, D.J., 2015. Sexual conflict maintains variation at
an insecticide resistance locus. BMC Biol. 13. https://doi.org/10.1186/s12915-015-
0143-3
Ruzicka, F., Connallon, T., 2020. Is the X chromosome a hot spot for sexually antagonistic
polymorphisms? Biases in current empirical tests of classical theory. Proc. R. Soc. B.
https://doi.org/10.1098/rspb.2020.1869
Ruzicka, F., Dutoit, L., Czuppon, P., Jordan, C.Y., Li, X.-Y., Olito, C., Runemark, A., Svensson,
E.I., Yazdi, H.P., Connallon, T., 2020. The search for sexually antagonistic genes:
Practical insights from studies of local adaptation and statistical genomics. Evol. Lett.
4–5, 398–415. https://doi.org/10.1002/evl3.192

Ruzicka, F., Hill, M.S., Pennell, T.M., Flis, I., Ingleby, F.C., Mott, R., Fowler, K., Morrow, E.H.,
Reuter, M., 2019. Genome-wide sexually antagonistic variants reveal long-standing
constraints on sexual dimorphism in fruit flies. PLoS Biol. 17, e3000244.
Sainz, J., Rudolph, A., Hoffmeister, M., Frank, B., Brenner, H., Chang-Claude, J., Hemminki,
K., Forsti, A., 2012. Effect of type 2 diabetes predisposing genetic variants on
colorectal cancer risk. J. Clin. Endocrinol. Metab. 97, E845-851.
https://doi.org/10.1210/jc.2011-2565
Sangalli, A., Orlandi, E., Poli, A., Maurichi, A., Santinami, M., Nicolis, M., Ferronato, S.,
Malerba, G., Rodolfo, M., Gomez Lira, M., 2017. Sex-specific effect of RNASEL
rs486907 and miR-146a rs2910164 polymorphisms' interaction as a susceptibility
factor for melanoma skin cancer. Melanoma Res. 27, 309–314.
https://doi.org/10.1097/CMR.0000000000000360
Schneider, K.K., Hule, L., Schote, A.B., Meyer, J., Frings, C., 2015. Sex matters! Interactions of
sex and polymorphisms of a cholinergic receptor gene (CHRNA5) modulate response
speed. Neuroreport 26, 186–191. https://doi.org/10.1097/WNR.0000000000000313
Schuurhof, A., Bont, L., Siezen, C.L.E., Hodemaekers, H., van Houwelingen, H.C., Kimman,
T.G., Hoebee, B., Kimpen, J.L.L., Janssen, R., 2010. Interleukin-9 polymorphism in
infants with respiratory syncytial virus infection: an opposite effect in boys and girls.
Pediatr. Pulmonol. 45, 608–613. https://doi.org/10.1002/ppul.21229
Shalev, I., Lerer, E., Israel, S., Uzefovsky, F., Gritsenko, I., Mankuta, D., Ebstein, R.P., Kaitz,
M., 2009. BDNF Val66Met polymorphism is associated with HPA axis reactivity to
psychological stress characterized by genotype and gender interactions.
Psychoneuroendocrinology 34, 382–388.
https://doi.org/10.1016/j.psyneuen.2008.09.017
Short, R.V., 1979. Sexual selection and its component parts, somatic and genital selection, as
illustrated by man and the great apes. Adv. Study Behav. 9, 131–158.
Sjoberg, R.L., Nilsson, K.W., Nordquist, N., Ohrvik, J., Leppert, J., Lindstrom, L., Oreland, L.,
2006. Development of depression: sex and the interaction between environment
and a promoter polymorphism of the serotonin transporter gene. Int. J.
Neuropsychopharmacol. 9, 443–449. https://doi.org/10.1017/S1461145705005936
Stearns, S.C., Govindaraju, D.R., Ewbank, D., Byars, S.G., 2012. Constraints on the
coevolution of contemporary human males and females. Proc. R. Soc. B Biol. Sci. 279,
4836–4844. https://doi.org/10.1098/rspb.2012.2024
Stulp, G., Kuijper, B., Buunk, A.P., Pollet, T.V., Verhulst, S., 2012. Intralocus sexual conflict
over human height. Biol. Lett. https://doi.org/10.1098/rsbl.2012.0590
Winkler, T.W., Justice, A.E., Graff, M., Barata, L., Feitosa, M.F., Chu, S., Czajkowski, J., Esko,
T., Fall, T., Kilpeläinen, T.O., Lu, Y., Mägi, R., Mihailov, E., Pers, T.H., Rüeger, S.,
Teumer, A., Ehret, G.B., Ferreira, T., Heard-Costa, N.L., Karjalainen, J., Lagou, V.,
Mahajan, A., Neinast, M.D., Prokopenko, I., Simino, J., Teslovich, T.M., Jansen, R.,
Westra, H.-J., White, C.C., Absher, D., Ahluwalia, T.S., Ahmad, S., Albrecht, E., Alves,
A.C., Bragg-Gresham, J.L., Craen, A.J.M. de, Bis, J.C., Bonnefond, A., Boucher, G.,
Cadby, G., Cheng, Y.-C., Chiang, C.W.K., Delgado, G., Demirkan, A., Dueker, N.,
Eklund, N., Eiriksdottir, G., Eriksson, J., Feenstra, B., Fischer, K., Frau, F., Galesloot,
T.E., Geller, F., Goel, A., Gorski, M., Grammer, T.B., Gustafsson, S., Haitjema, S.,
Hottenga, J.-J., Huffman, J.E., Jackson, A.U., Jacobs, K.B., Johansson, Å., Kaakinen, M.,
Kleber, M.E., Lahti, J., Leach, I.M., Lehne, B., Liu, Y., Lo, K.S., Lorentzon, M., Luan, J.,
Madden, P.A.F., Mangino, M., McKnight, B., Medina-Gomez, C., Monda, K.L.,

Montasser, M.E., Müller, G., Müller-Nurasyid, M., Nolte, I.M., Panoutsopoulou, K.,
Pascoe, L., Paternoster, L., Rayner, N.W., Renström, F., Rizzi, F., Rose, L.M., Ryan,
K.A., Salo, P., Sanna, S., Scharnagl, H., Shi, J., Smith, A.V., Southam, L., Stančáková, A.,
Steinthorsdottir, V., Strawbridge, R.J., Sung, Y.J., Tachmazidou, I., Tanaka, T.,
Thorleifsson, G., Trompet, S., Pervjakova, N., Tyrer, J.P., Vandenput, L., Laan, S.W.
van der, Velde, N. van der, Setten, J. van, Vliet-Ostaptchouk, J.V. van, Verweij, N.,
Vlachopoulou, E., Waite, L.L., Wang, S.R., Wang, Z., Wild, S.H., Willenborg, C., Wilson,
J.F., Wong, A., Yang, J., Yengo, L., Yerges-Armstrong, L.M., Yu, L., Zhang, W., Zhao,
J.H., Andersson, E.A., Bakker, S.J.L., Baldassarre, D., Banasik, K., Barcella, M.,
Barlassina, C., Bellis, C., Benaglio, P., Blangero, J., Blüher, M., Bonnet, F.,
Bonnycastle, L.L., Boyd, H.A., Bruinenberg, M., Buchman, A.S., Campbell, H., Chen,
Y.-D.I., Chines, P.S., Claudi-Boehm, S., Cole, J., Collins, F.S., Geus, E.J.C. de, Groot,
L.C.P.G.M. de, Dimitriou, M., Duan, J., Enroth, S., Eury, E., Farmaki, A.-E., Forouhi,
N.G., Friedrich, N., Gejman, P.V., Gigante, B., Glorioso, N., Go, A.S., Gottesman, O.,
Gräßler, J., Grallert, H., Grarup, N., Gu, Y.-M., Broer, L., Ham, A.C., Hansen, T., Harris,
T.B., Hartman, C.A., Hassinen, M., Hastie, N., Hattersley, A.T., Heath, A.C., Henders,
A.K., Hernandez, D., Hillege, H., Holmen, O., Hovingh, K.G., Hui, J., Husemoen, L.L.,
Hutri-Kähönen, N., Hysi, P.G., Illig, T., Jager, P.L.D., Jalilzadeh, S., Jørgensen, T.,
Jukema, J.W., Juonala, M., Kanoni, S., Karaleftheri, M., Khaw, K.T., Kinnunen, L.,
Kittner, S.J., Koenig, W., Kolcic, I., Kovacs, P., Krarup, N.T., Kratzer, W., Krüger, J.,
Kuh, D., Kumari, M., Kyriakou, T., Langenberg, C., Lannfelt, L., Lanzani, C., Lotay, V.,
Launer, L.J., Leander, K., Lindström, J., Linneberg, A., Liu, Y.-P., Lobbens, S., Luben, R.,
Lyssenko, V., Männistö, S., Magnusson, P.K., McArdle, W.L., Menni, C., Merger, S.,
Milani, L., Montgomery, G.W., Morris, A.P., Narisu, N., Nelis, M., Ong, K.K., Palotie,
A., Pérusse, L., Pichler, I., Pilia, M.G., Pouta, A., Rheinberger, M., Ribel-Madsen, R.,
Richards, M., Rice, K.M., Rice, T.K., Rivolta, C., Salomaa, V., Sanders, A.R., Sarzynski,
M.A., Scholtens, S., Scott, R.A., Scott, W.R., Sebert, S., Sengupta, S., Sennblad, B.,
Seufferlein, T., Silveira, A., Slagboom, P.E., Smit, J.H., Sparsø, T.H., Stirrups, K., Stolk,
R.P., Stringham, H.M., Swertz, M.A., Swift, A.J., Syvänen, A.-C., Tan, S.-T., Thorand,
B., Tönjes, A., Tremblay, A., Tsafantakis, E., Most, P.J. van der, Völker, U., Vohl, M.-C.,
Vonk, J.M., Waldenberger, M., Walker, R.W., Wennauer, R., Widén, E., Willemsen,
G., Wilsgaard, T., Wright, A.F., Zillikens, M.C., Dijk, S.C. van, Schoor, N.M. van,
Asselbergs, F.W., Bakker, P.I.W. de, Beckmann, J.S., Beilby, J., Bennett, D.A.,
Bergman, R.N., Bergmann, S., Böger, C.A., Boehm, B.O., Boerwinkle, E., Boomsma,
D.I., Bornstein, S.R., Bottinger, E.P., Bouchard, C., Chambers, J.C., Chanock, S.J.,
Chasman, D.I., Cucca, F., Cusi, D., Dedoussis, G., Erdmann, J., Eriksson, J.G., Evans,
D.A., Faire, U. de, Farrall, M., Ferrucci, L., Ford, I., Franke, L., Franks, P.W., Froguel, P.,
Gansevoort, R.T., Gieger, C., Grönberg, H., Gudnason, V., Gyllensten, U., Hall, P.,
Hamsten, A., Harst, P. van der, Hayward, C., Heliövaara, M., Hengstenberg, C., Hicks,
A.A., Hingorani, A., Hofman, A., Hu, F., Huikuri, H.V., Hveem, K., James, A.L., Jordan,
J.M., Jula, A., Kähönen, M., Kajantie, E., Kathiresan, S., Kiemeney, L.A.L.M., Kivimaki,
M., Knekt, P.B., Koistinen, H.A., Kooner, J.S., Koskinen, S., Kuusisto, J., Maerz, W.,
Martin, N.G., Laakso, M., Lakka, T.A., Lehtimäki, T., Lettre, G., Levinson, D.F., Lind, L.,
Lokki, M.-L., Mäntyselkä, P., Melbye, M., Metspalu, A., Mitchell, B.D., Moll, F.L.,
Murray, J.C., Musk, A.W., Nieminen, M.S., Njølstad, I., Ohlsson, C., Oldehinkel, A.J.,
Oostra, B.A., Palmer, L.J., Pankow, J.S., Pasterkamp, G., Pedersen, N.L., Pedersen, O.,
Penninx, B.W., Perola, M., Peters, A., Polašek, O., Pramstaller, P.P., Psaty, B.M., Qi,

L., Quertermous, T., Raitakari, O.T., Rankinen, T., Rauramaa, R., Ridker, P.M., Rioux, J.D., Rivadeneira, F., Rotter, J.I., Rudan, I., Ruijter, H.M. den, Saltevo, J., Sattar, N., Schunkert, H., Schwarz, P.E.H., Shuldiner, A.R., Sinisalo, J., Snieder, H., Sørensen, T.I.A., Spector, T.D., Staessen, J.A., Stefania, B., Thorsteinsdottir, U., Stumvoll, M., Tardif, J.-C., Tremoli, E., Tuomilehto, J., Uitterlinden, A.G., Uusitupa, M., Verbeek, A.L.M., Vermeulen, S.H., Viikari, J.S., Vitart, V., Völzke, H., Vollenweider, P., Waeber, G., Walker, M., Wallaschofski, H., Wareham, N.J., Watkins, H., Zeggini, E., Consortium, C., Consortium, D., Consortium, G., Consortium, G.-Bpg., Consortium, I., Consortium, M., Chakravarti, A., Clegg, D.J., Cupples, L.A., Gordon-Larsen, P., Jaquish, C.E., Rao, D.C., Abecasis, G.R., Assimes, T.L., Barroso, I., Berndt, S.I., Boehnke, M., Deloukas, P., Fox, C.S., Groop, L.C., Hunter, D.J., Ingelsson, E., Kaplan, R.C., McCarthy, M.I., Mohlke, K.L., O'Connell, J.R., Schlessinger, D., Strachan, D.P., Stefansson, K., Duijn, C.M. van, Hirschhorn, J.N., Lindgren, C.M., Heid, I.M., North, K.E., Borecki, I.B., Kutalik, Z., Loos, R.J.F., 2015. The Influence of Age and Sex on Genetic Associations with Adult Body Size and Shape: A Large-Scale Genome-Wide Interaction Study. PLOS Genet. 11, e1005378. https://doi.org/10.1371/journal.pgen.1005378

Wise, A.L., Gyi, L., Manolio, T.A., 2013. eXclusion: Toward Integrating the X Chromosome in Genome-wide Association Analyses. Am. J. Hum. Genet. 92, 643–647. https://doi.org/10.1016/j.ajhg.2013.03.017

Wong, S.Y.S., Lau, E.M.C., Li, M., Chung, T., Sham, A., Woo, J., 2005. The prevalence of Apo E4 genotype and its relationship to bone mineral density in Hong Kong Chinese. J. Bone Miner. Metab. 23, 261–265. https://doi.org/10.1007/s00774-004-0593-0

Wu, K., Chen, C., Moyzis, R.K., Greenberger, E., Yu, Z., 2016. Gender Interacts with Opioid Receptor Polymorphism A118G and Serotonin Receptor Polymorphism -1438 A/G on Speed-Dating Success. Hum. Nat. Hawthorne N 27, 244–260. https://doi.org/10.1007/s12110-016-9257-8

Youssef, S.S., Abd El Aal, A.M., Nasr, A.S., el Zanaty, T., Seif, S.M., 2013. Interleukin-12B gene polymorphism frequencies in Egyptians and sex-related susceptibility to hepatitis C infection. J. Interferon Cytokine Res. Off. J. Int. Soc. Interferon Cytokine Res. 33, 415–419. https://doi.org/10.1089/jir.2012.0161

Table 1. Genetic loci in humans showing sexually antagonistic effects on trait expression. EAF, effect allele frequency (* indicates allele frequency derived from 100 genomes database); M, male; F, female. Background colours denote instances where genes and/or variants appear more than once in the list.

Trait	Gene	Locus	EAF	M Effect	M Effect SE	F Effect	F Effect SE	Reference
Complex traits								
Concentration performance score	FADS1/2	rs1535	0.672	0.2275	0.1065	-0.2394	0.1116	(Lauritzen et al., 2017)
Reading speed, number of sentences	FADS1/2	rs174448	0.642	0.5511	0.1069	-0.6074	0.1129	(Lauritzen et al., 2017)
Reading speed, number of sentences	ELOVL5	rs2397142	0.695	-0.1963	0.1042	0.2364	0.1095	(Lauritzen et al., 2017)
Reading speed, correct sentences	FADS1/2	rs174448	0.642	0.6599	0.1077	-0.3451	0.1112	(Lauritzen et al., 2017)
Speed-dating success	OPRM1	rs1799971	0.922	0.4697	0.2275	-0.53	0.2856	(Wu et al., 2016)
Speed-dating success	HTR2A	rs6311	0.418	-0.4826	0.1982	0.4702	0.1964	(Wu et al., 2016)
Waist-hip-ratio (BMI adjusted)	SIM1	rs17185536	0.76	-0.0166	0.008	0.0264	0.0067	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	NMU	rs3805389	0.28	-0.0139	0.0067	0.0261	0.006	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	GNPNAT1	rs4898764	0.53	0.0208	0.0066	-0.0165	0.0059	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	SCGZ	rs17470444	0.71	0.0345	0.0081	-0.0149	0.0068	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	IQGAP2	rs2069664	0.53	0.0244	0.0078	-0.0183	0.0067	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	SLC2A3	rs741361	0.6	0.0272	0.0079	-0.0196	0.0067	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	IRS1	rs2673140	0.38	0.022	0.0066	-0.0209	0.0059	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	TTN	rs2042995	0.23	0.0198	0.0066	-0.022	0.0078	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	RXRA	rs10881574	0.07	0.0268	0.008	-0.0191	0.0067	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	PTPRD	rs7042428	0.98	0.0244	0.0071	-0.0166	0.0063	(Winkler et al., 2015)
Waist-hip-ratio (BMI adjusted)	CERC2	rs17809093	0.04	0.0294	0.0091	-0.024	0.0077	(Winkler et al., 2015)
Reaction time (RT) in the Stroop task, congruent condition	CHRNA5	rs3841324	0.406	-1.5421	0.3857	0.9945	0.2517	(Schneider et al., 2015)
Reaction time (RT) in the Stroop task, incongruent condition	CHRNA5	rs3841324	0.406	-1.6215	0.389	1.0215	0.2521	(Schneider et al., 2015)
Reaction time (RT) in the Negative priming task	CHRNA5	rs3841324	0.406	-1.675	0.3912	0.6299	0.247	(Schneider et al., 2015)

Trait	Gene	SNP	MAF	β1	SE1	β2	SE2	Reference
Borderline personality traits after maltreatment	OXTR	rs53576	0.738	-0.3429	0.0881	0.2066	0.0857	(Cicchetti et al., 2014)
Borderline personality traits after maltreatment	FKPB5	CATT haplotype	-	0.2041	0.0877	-0.195	0.0857	(Cicchetti et al., 2014)
fluid intelligence	C16orf62 (OXTR)	rs226849	0.46	-0.3949	0.1869	0.494	0.1814	(Kiy et al., 2013)
face perception	C16orf62 (OXTR)	rs226849	0.46	-0.5819	0.1909	0.3498	0.1788	(Kiy et al., 2013)
face memory	C16orf62 (OXTR)	rs226849	0.46	-0.3679	0.1864	0.3562	0.1789	(Kiy et al., 2013)
Hippocampal volume	SLC6A4	rs25531	0.39	-		-		(Price et al., 2013)
Sociopathy (in alcoholics)	SLC6A4	rs25531	0.51	-		-		(Herman et al., 2011)
systolic blood pressure S_NIGHT	PON1	rs3917542	-	-0.1047	0.0637	0.1129	0.0645	(Padmanabhan et al., 2010)
systolic blood pressure S_NIGHT	PON3	rs17884563	-	-0.1482	0.0637	0.1061	0.0645	(Padmanabhan et al., 2010)
systolic blood pressure S_CLIN	EDNRB	rs1335803	-	-0.1308	0.0637	0.1129	0.0645	(Padmanabhan et al., 2010)
diastolic blood pressure D_NIGHT	DRD1	rs265974	-	-0.1482	0.0637	0.1502	0.0646	(Padmanabhan et al., 2010)
diastolic blood pressure D_HOME	SLC4A2	rs2303933	-	-0.1482	0.0637	0.1325	0.0646	(Padmanabhan et al., 2010)
diastolic blood pressure D_CLIN	LPL	rs17410577	-	-0.1047	0.0637	0.1775	0.0647	(Padmanabhan et al., 2010)
Psychological Stress	BDNF	rs6265	0.149	-		-		(Shalev et al., 2009)
BMI standard deviation score (children)	FAS	rs2228305	0.0312	-0.552	0.2259	0.4427	0.2075	(Korner et al., 2007)
Pain sensitivity	OPRM1	rs1799971	0.8817	-		-		(Fillingim et al., 2005)
Central DBP	REN	rs6693954	0.22	-0.4444	0.2132	1.1052	0.3923	(Ljungberg et al., 2014)
Disease risk/severity								
Graves opthalmopathy in GD	PRR3	rs2074503	0.3586	0.4397	0.1682	-0.2185	0.0851	(Liu et al., 2014)
Colorectal cancer	LTA	rs1041981	0.32	-0.0722	0.0368	0.1027	0.0453	(Sainz et al., 2012)
Arterial blood pressure	PPAR	rs1801282	0.1213	-		-		(Franck et al., 2012)
Schizophrenia risk	RELN	rs7341475	0.851*	1.2396	0.2819	-0.762	0.3662	(Ovadia and Shifman, 2011)
Colon cancer survival	ESR2	(CA)n repeats	-	0.2272	0.0962	-0.2046	0.1205	(Press et al., 2011)
Schizophrenia risk	DAO	rs2070586	0.2729	0.1982	0.0888	-0.2537	0.0908	(Kim et al., 2010)

T2DM, fasting glucose	SCARB1	rs9919713	0.04	0.1865	0.0853	-0.2515	0.0697	(McCarthy et al., 2009)
Depression	SLC6A4	rs25531	0.4586	0.5404	0.2277	-0.3996	0.1859	(Sjoberg et al., 2006)
Infertility in cystic fibrosis	CFTR	rs213950	0.417931	-		-		(Morea et al., 2005)
Bone mineral density	APOE	rs429358	0.0798	-		-		(Wong et al., 2005)
Narcolepsy severity	COMT	Long allele	0.4588	1.3596	0.3878	-2.2152	0.4871	(Dauvilliers et al., 2001)
Nonsyndromic cleft lip	IMPK	rs11142081	0.863	1.2156	0.0937	-1.1161	0.0783	(Carlson et al., 2018)
Nonsyndromic cleft lip	IMPK	rs72804706	0.898	-0.816	0.0906	2.0665	0.0908	(Carlson et al., 2018)
Nonsyndromic cleft lip	IMPK	rs77590619	0.744	-0.8577	0.0904	1.2291	0.0804	(Carlson et al., 2018)
Allergy	ACP1	A* allele (CG haplotype)	-	-		-		(Gloria-Bottini et al., 2007)
Hepatitis C susceptability	IL-12B	rs3212227	0.3463	-		-		(Youssef et al., 2013)
RSV severity	IL-9	rs2069885	0.15	0.1888	0.0933	-0.2794	0.0957	(Schuurhof et al., 2010)
CV risk marker (plasma fatty acid composition)	FABP2	rs1799883	0.287	-		-		(Gastaldi et al., 2007)
Bone mineral density	NR3C1	GCAG haplotype	-	0.6854	0.3676	-0.9029	0.4311	(Peng et al., 2008)
Anxiety	NGF	rs6330	0.451	0.4562	0.18	-0.4674	0.1817	(Lang et al., 2008)
Melanoma risk	RNASEL, miR-146a	AG/AA and C haplotype	-	0.4525	0.1564	-0.3553	0.174	(Sangalli et al., 2017)

Figure legends

Figure 1. Figure 1. PRISMA flow diagram for systematic review of sexually antagonistic loci in humans.

Figure 2. The relationship between effect allele frequency and effect size ratio. Point size varies according to the variance of effect size ratio - larger points have smaller variance and therefore a larger weighting in the model. The vertical dotted line represents the switch point between a net-negative effect of a particular locus (effect size ratio >-1) and a net positive effect (effect size ratio <-1). The blue line represents the predicted values derived from the generalized linear model fitted to the data (see Results).