1 **Title** Research Letter: Reactive balance responses after mild traumatic brain injury
2 (mTBI): a scoping review
3

4 **Authors:** Amanda Morris[1*], Tallie Casucci[2], Mary M. McFarland[3], Benjamin Cassidy[1],
5 Ryan Pelo[4], Nicholas Kreter[1], Leland E. Dibble[4], Peter C. Fino[1]
6
[1]Department of Health and Kinesiology, University of Utah, 250 S 1850 E,Salt Lake City, UT,
USA
[2]J. Willard Marriot Library, University of Utah, 295 1500 E, Salt Lake City, UT, USA
[3]Eccles Health Sciences Library, 10 N 1900 E, Salt Lake City, UT, USA
[4]Department of Physical Therapy and Athletic Training, University of Utah, 520 S Wakara Way,
Salt Lake City, UT, USA
12
*Correspondence:
Amanda Morris
250 South 1850 East, HPER Room E 107 D
Salt Lake City, UT, 84112
amanda.morris@utah.edu
Author E-mails: TC: tallie.casucci@utah.edu; MMM: mary.mcfarland@utah.edu; BC:
cassidybp@mymail.vcu.edu; RP: ryan.pelo@hsc.utah.edu; NK: nick.kreter@utah.edu;
LD: lee.dibble@hsc.utah.edu; PF: peter.fino@utah.edu
21
ORCID: AM: 0000-0002-6297-5255; TC: 0000-0001-6311-9613; MMM: 0000-0002-
9062-2393; RP: 0000-0002-2109-366X; NK: 0000-0002-1567-6287, PF: 0000-0002-
8621-3706

Conflicts of Interest and Source of Funding: This project was supported in part by
the Pac-12 Conference's Student-Athlete Health and Well-Being Initiative (PI: Fino) and
by the Eunice Kennedy Shiver National Institute of Child Health & Human Development
of the National Institutes of Health under Award Number K12HD073945. The content of
this manuscript is solely the responsibility of the authors and does not necessarily
represent the official views of the Pac-12 Conference, or its members, or the National
Institute of Health. Sponsors had no role in the study design, in the collection, analysis
and interpretation of data; in the writing of the manuscript; or in the decision to submit
the manuscript for publication.

We affirm that we have no financial affiliation (including research funding) or
involvement with any commercial organization that has a direct financial interest in any
matter included in this manuscript, except as disclosed in an attachment and cited in the
manuscript. Any other conflict of interest (ie, personal associations or involvement as a
director, officer, or expert witness) is also disclosed in an attachment.

Publication history: This manuscript was previously posted to medRxiv, doi:
https://doi.org/10.1101/2020.12.17.20248433

Abstract

Objective: Balance testing after concussion or mild traumatic brain injury (mTBI) can be
useful in determining acute and chronic neuromuscular deficits that are unapparent from
symptom scores or cognitive testing alone. Current assessments of balance do not
comprehensively evaluate all three classes of balance: maintaining a posture, voluntary
movement, and reactive postural response. Despite the utility of reactive postural
responses in predicting fall risk in other balance impaired populations, the effect of
mTBI on reactive postural responses remains unclear. This review sought to (1)
examine the extent and range of available research on reactive postural responses in
people post-mTBI and (2) determine if reactive postural responses (balance recovery)
are affected by mTBI.

Design: PRISMA Scoping review.

Methods: Studies were identified using Medline, Embase, CINAHL, Cochrane Library,
Dissertations and Theses Global, PsycINFO, SportDiscus, and Web of Science.
Inclusion criteria were: injury classified as mTBI with no confounding central or
peripheral nervous system dysfunction beyond those stemming from the mTBI,
quantitative measure of reactive postural response, and a discrete, externally driven
perturbation was used to test reactive postural response.

Results: A total of 4,747 publications were identified and a total of three studies (5
publications) were included in the review.

Conclusion: The limited number of studies available on this topic highlight the lack of
knowledge on reactive postural responses after mTBI. This review provides a new
direction for balance assessments after mTBI and recommends incorporating all three
classes of postural control in future research.

Keywords: reactive balance, concussion, brain injury, return-to-play

1 Introduction

Control of balance, or postural control, can be categorized into three different

classes of activity: 1) maintaining posture/stability (e.g. standing), 2) voluntary

transitional movements (e.g. gait), and 3) reactive postural responses (RPR) to restore

stability after an unexpected disturbance [1]. The three classes of balance are related and

each involve complex interactions between motor and sensory processes [2, 3], but have

unique neuromechanical demands.

The classes of postural control exist on a continuum of feedback and feedforward

control [4-8] (Figure 1). Maintaining a posture requires controlling the center of mass

(COM) through active postural muscle activation and integration of sensory feedback

while maintaining a fixed base of support (BOS) [8]. Unlike static posture, transitional

movements involve voluntary movement of COM and changes in BOS through

incorporation of pre-planning and anticipatory control based on central set [8], and

sensory feedback for corrections [8, 9]. In contrast to static or transitional movements,

RPR involve involuntary, externally-driven motion of the COM or BOS that are corrected

using rapid, time-constrained, automatic postural responses [7, 8] to quickly stop the initial

falling motion by moving the COM or changing the BOS [10, 11] and stabilizing the body

through subsequent postural adjustments [8]. As RPR occur faster than voluntary

movement (70-100 ms versus 180-250 ms) [8, 12], there is little time for pre-planning or

feedback-driven control [13]. Because of differences in neuromechanical demands, each

class of postural control may yield unique, but complementary knowledge about the

neuromechanical and neurophysiological clinical information regarding individuals'

postural competence.

Multiple reviews have concluded that concussion, hereafter mild traumatic brain

injury (mTBI), impairs balance [14-19]. The vast majority of studies on postural control after

mTBI examine static posture [20-23] and a smaller, but significant, number of studies

investigate voluntary transitional movements such as gait and turning [24-29]. Yet, a

comprehensive review of the literature surrounding RPR is lacking and the effect of

mTBI on RPR remains unclear. Therefore, the purpose of this scoping review was to

examine the extent and range of available research on RPR post-mTBI. We aimed to

determine if RPR are affected by mTBI. Objectives, inclusion criteria, and methods for

this review were specified in advance and documented in a protocol (Supplemental

Digital Content 1) [30].

102   *Insert Figure 1 here*

2   Methods

A scoping review was conducted using Arksey's five stages [31] (detailed methods

in Supplemental Digital Content 1). Covidence (Veritas Health Innovation, Melbourne,

AU) was used to screen and select studies.

2.1 Inclusion Criteria

Articles were included if: 1) the injury was classified as mTBI (loss of

consciousness <30 minutes, Glasgow Coma Scale >13, posttraumatic amnesia <24

hours), 2) the outcome was a measure of RPR (Supplemental Digital Content 1), 3) the

perturbation was an externally driven change in support structure, visual surround, or

vestibular information (sway referenced postural control tests such as the Sensory

Organization Test (SOT), were not considered RPR paradigms as these perturbations

were not externally driven), and 4) the perturbation was discrete.

Articles were excluded if there were confounding population factors such as:

central or peripheral nervous system dysfunction beyond those stemming from the

mTBI, pregnancy, or orthopedic injuries that affect postural control and gait.

2.2 Literature searching

Literature searching was conducted by a librarian (TC) and peer reviewed by an

information specialist (MMM) with Peer Review of Electronic Search Strategies

(PRESS) guidelines [32]. The following databases were used: Medline, Embase, CINAHL,

Cochrane Library, Dissertations and Theses Global, PsycINFO, SportDiscus, and Web

of Science (Supplemental Digital Content 1). No date limits, nor search filters, were

applied. The search was conducted January 31st - February 4, 2020, and updated on

March 9th , 2021. Search strategies were composed using database subject terms and

multiple keywords for the main concepts of (brain injuries OR concussions) AND

(balance) AND (posture OR postural OR perturbations) (Supplemental Digital Content 2

contains detailed search strategies). References of included studies were checked for

relevancy.

Titles and abstracts were reviewed independently by four reviewers (AM, BC,

RP, NK). Publications that did not meet the inclusion criteria were excluded from the

review. Full-text screening and extraction of results was completed independently by

three reviewers (AM, BC, RP). Approval by at least two reviewers was required for

inclusion; if reviewers disagreed, all reviewers came to a consensus based on inclusion

and exclusion criteria.

3   Results

3.1 Extent of Research Available

After duplicates were removed, 4,747 publications were identified (Figure 2).

After title and abstract screening, 4,710 studies were excluded due to incorrect study

type, population, or relevance to the aim of this review. Thirty-five full-text articles were

assessed for eligibility. Of 35 articles, five articles were extracted, three of the articles [33-

35] were from the same study (M.Walker, Personal Communication, June 5, 2020). The

three selected studies and their characteristics are summarized in Table 1. Excluded

studies are summarized in Supplemental Digital Content 3.

146   *Insert Figure 2 here*
147   *Insert Table 1 here*

3.2 Perturbation Method and Postural Response Measures

Only three studies met our outlined criteria examining RPR to discrete

perturbations after mTBI. Study one delivered expected and unexpected upper-

extremity perturbations during a standing reaching task using the KINARM robot (BKIN

Technologies Ltd., Kingston, ON)[36]. Perturbations were delivered after the reach started

and consisted of a 12 N square pulse with a 10 second rise time applied to the left or

right side of the participant (C.Mang, personal communication, June 9, 2020).

Anticipatory and reactive postural adjustments were measured. RPR were quantified

using the maximum absolute COP velocity from 100ms after reach onset to end of the

reach.

Study two perturbed static stance by delivering unexpected forces (10% of

participant's body weight) to a waist-belt via a computer controlled motor. Perturbations

were delivered for 0.5 seconds in two directions, forward and right [35]. Motion capture

was used to quantify postural sway length, displacement, and postural oscillations

(frequency of marker acceleration) as RPR outcomes.

Study three perturbed static balance and walking using eight conditions on a

Computer-Assisted Rehabilitation Environment (CAREN) system: 1) single, rapid (<1s)

forward surface translations 2) slow rotational perturbations (platform rotated downward

over 5s), 3) visual scene tilt, 4) visual scene translation, 5) visual scene congruent with

tilt, 6) visual scene incongruent with tilt, 7) visual scene congruent with translation, and

8) visual scene incongruent with translation [37]. Discrete perturbations were delivered

with an interstimulus interval of 8-12 seconds. RPR outcomes were recorded using

motion capture to quantify joint angles and spatiotemporal features (i.e. stride length),

an accelerometer to quantify acceleration, and electromyography to record muscle

activity.

3.3 Effect of mTBI on Reactive Postural Responses

Study 1 tested RPR in athletes at baseline and within ten days of mTBI [36]. Before

mTBI, anticipatory postural adjustments were seen only with an expected perturbation.

After mTBI, anticipatory postural adjustments were seen in both expected and

unexpected perturbations. However, there was no effect of mTBI on RPR.

Study 2 tested ten non-veterans without mTBI history and veterans with a history

of mTBI (7 months – 7 years post mTBI)[35]. Veterans were divided into two groups,

chronic complaints of imbalance (N=8) and no balance complaints (N=6). Those with

chronic complaints of imbalance had significantly more upper trunk sway in response to

perturbation than non-veterans and veterans without balance complaints. Additionally,

those with complaints of imbalance had high-frequency postural oscillations (frequency

of marker acceleration) after perturbations.

Study three tested ten healthy controls and eleven participants with mTBI who

reported lingering balance deficits. Incongruent visual scene with translation or rotation,

platform movement only, and visual only perturbations eliciting RPR were more

sensitive than standard clinical tests (e.g.Berg Balance Scale, Dynamic Gait Index, and

High Level Mobility Assessment tool, Balance Evaluations Systems Test) to detecting

whether an individual reported lingering balance deficits post mTBI. Walking

perturbations provided greater ability to discriminate than standing perturbations.

4 Conclusion

This scoping review presents the available research on RPR after mTBI. The

small number of available studies prohibits a definitive conclusion. Instead, our results

highlight the lack of knowledge on this topic, and demonstrate methodological variability

between existing studies examining reactive postural control post–mTBI.

Two of the three studies included in this review presented results indicating that

objective measures of RPR can detect lingering balance deficits after mTBI, while one

study reported no change in RPR after mTBI. There was no effect of mTBI on RPR in

collegiate athletes in response to upper body perturbations [36], nor were there reactive

postural deficits in veterans who did not report disequilibrium after perturbation at the

waist [35]. However, both veterans who reported chronic imbalance after mTBI[35] and

adults with previous mTBI who reported lingering balance deficits had impaired RPR

when perturbed. Therefore, it is possible that abnormal RPR are evident in individuals

with chronic, persisting symptoms from mTBI, but absent in asymptomatic individuals.

Impaired RPR suggest increased fall and injury risk, however, the available literature

examining reactive balance deficits and their clinical consequences are severely limited

and more research is needed.

The studies presented here used different methodology and populations, limiting

the ability to compare results. The perturbations were different in magnitude, direction,

and method; one study used right and left upper body perturbations through a hand-

held apparatus, another study used a forward and rightward perturbation delivered to a

waist-belt, while the third study used surface and visual perturbations. Task differences

in type and perturbation profile (force and acceleration) likely resulted in different

strengths of perturbation, necessitating different magnitudes of postural response [38, 39],

making it difficult to compare the magnitude of balance deficit after mTBI between the

studies. Different muscle activation patterns, joint torques, and sensory feedback may

have been used depending on where the perturbation was applied or the magnitude of

the perturbation. The populations also varied in age, symptom severity, and time since

injury making comparisons between the three studies difficult (Table 1). It is likely that

different age groups have inherent differences in RPR; there are significant differences

in RPR between healthy young and old adults [40] and between children of different motor

developmental levels [41]. Finally, all studies tested outside of what is considered the

acute period after mTBI (<48 hours) and did not perform any prospective testing, limiting

the ability to determine the immediate and longitudinal effects of mTBI on RPR. Due to

conflicting results, inconsistent methods, and extremely limited number of studies, the

effects of mTBI on RPR to physical perturbations remains unclear.

4.1 Clinical Implications

Balance testing can be useful in determining acute and chronic neuromuscular

impairments after mTBI that may be unapparent from symptom scores or cognitive

testing alone. Studies of maintenance of posture [21-23, 42] and voluntary transitional

movement [17, 19, 24-26, 43, 44] after mTBI suggest these domains of balance can be impaired

both acutely and chronically after injury. Consequently, the quantification of static

balance using the Balance Error Scoring System (BESS) [45], SOT [46], or dynamic

balance using the tandem gait test [47] have become valuable clinical tools. However,

these tests do not examine RPR. Given the critical importance of RPR in the prevention

of falls and injury, their inclusion in the comprehensive examination of postural control

after mTBI is warranted.

RPR have been used to predict fall risk in balance impaired populations such as:

older adults [48], stroke [49] and Parkinson's Disease [50-52]. These studies, using consistent

location of perturbation and systematic manipulation of perturbation magnitude, have

allowed the characterization of rapid postural response to unpredictable body

movement [53]. These responses involve early automatic postural responses [12] and

coordination of multiple body segments via neural pathways encompassing spinal,

brainstem, and cortical regions [7]. While traditionally considered a cortical injury, deficits

in RPR may indicate mTBI-related dysfunction in subcortical regions and throughout the

central nervous system. Increasingly, subtype-based models of care for patients with

mTBI [54, 55] acknowledge the heterogeneous consequences of mTBI and the need for

targeted, specialized treatment strategies. Future research examining RPR as well as

the other classes of balance may provide more specificity than single-classed

assessments of balance about the neurophysiological processes underlying motor

deficits, and appropriate targets for rehabilitative care, in individuals post-mTBI.

5 References

1.   Pollock AS, Durward BR, Rowe PJ, Paul JP. What is balance? *Clinical*

Rehabilitation. 2000;14(4):402-406. doi:10.1191/0269215500cr342oa

2.   Woollacott MH, Shumway-Cook A. *Motor Control: Theory and Practical*

Applications. Second ed. Lippincott Williams and Wilkins; 2001:164-165.

3.   Horak FB. Postural orientation and equilibrium: what do we know about neural

control of balance to prevent falls? *Age and Ageing*. 2006;35(S2):ii7-ii11.

4.   Nashner LM, Shupert CL, Horak FB, Black FO. Organization of posture controls:

An analysis of sensory and mechanical constraints. *Progress in Brain Research*.

1989;80(C):411-418. doi:10.1016/S0079-6123(08)62237-2

5.   Dietz V, Trippel M, Ibrahim IK, Berger W. Human stance on a sinusoidally

translating platform: balance control by feedforward and feedback mechanisms.

Experimental Brain Research. 1993;93(2):352-362. doi:10.1007/BF00228405

6.   Pavol MJ, Pai Y-C. Feedforward adaptations are used to compensate for a

potential loss of balance. *Exp Brain Res*. 2002;145:528-538. doi:10.1007/s00221-002-

1143-4

7.   Jacobs JV, Horak FB. Cortical control of postural responses. *Journal of Neural*

Transmission. 2007/10// 2007;114:1339-1348. doi:10.1007/s00702-007-0657-0

8.   Horak FB. Postural Control. Springer Berlin Heidelberg; 2008:3212-3219.

9.   Reimann H, Fettrow T, Thompson ED, Jeka JJ. Neural Control of Balance During

Walking. *Frontiers in Physiology*. 2018;9(SEP):1271-1271.

doi:10.3389/fphys.2018.01271

10.   Maki BE, Edmondstone MA, McIlroy WE. Age-related differences in laterally

directed compensatory stepping behavior. *Journals of Gerontology - Series A Biological*

Sciences and Medical Sciences. 2000;55(5):270-277. doi:10.1093/gerona/55.5.M270

11.   Maki BE, McIlroy WE. Cognitive demands and cortical control of human balance-

recovery reactions. *Journal of Neural Transmission*. 2007/10// 2007;114:1279-1296.

doi:10.1007/s00702-007-0764-y

12.   Horak FB, Henry SM, Shumway-Cook A. Postural Perturbations: New Insights for

Treatment of Balance Disorders. *Physical Therapy*. 1997;77(5)

13.   Rasman BG, Forbes PA, Tisserand R, Blouin JS. Sensorimotor manipulations of

the balance control loop-beyond imposed external perturbations. *Frontiers in Neurology*.

2018;9(OCT)doi:10.3389/fneur.2018.00899

14.   Guskiewicz KM. Balance Assessment in the Management of Sport-Related

Concussion. *Clinics in Sports Medicine*. 2011;30(1):89-102.

doi:10.1016/j.csm.2010.09.004

15.   Valovich McLeod TC, Hale TD. Vestibular and balance issues following sport-

related concussion. *Brain Injury*. 2015;29(2):175-184.

doi:10.3109/02699052.2014.965206

16.   Buckley TA, Oldham JR, Caccese JB. Postural control deficits identify lingering

post-concussion neurological deficits. *Journal of sport and health science*. 2016;5(1):61-

69. doi:10.1016/j.jshs.2016.01.007

17.   Fino PC, Parrington L, Pitt W, et al. Detecting gait abnormalities after concussion

or mild traumatic brain injury: A systematic review of single-task, dual-task, and complex

gait. *Gait and Posture*. 2018;62:157-166. doi:10.1016/j.gaitpost.2018.03.021

18.   Martini DN, Broglio SP. Long-term effects of sport concussion on cognitive and

motor performance: A review. *International Journal of Psychophysiology*. 2018;132:25-

30. doi:10.1016/j.ijpsycho.2017.09.019

19.   Wood TA, Hsieh KL, An R, Ballard RA, Sosnoff JJ. Balance and Gait Alterations

Observed More Than 2 Weeks After Concussion. *American Journal of Physical*

Medicine & Rehabilitation. 2019;98(7):566-576. doi:10.1097/PHM.0000000000001152

20.   Guskiewicz KM, Ross SE, Marshall SW. Postural Stability and

Neuropsychological Deficits after Concussion in Collegiate Athletes. *Journal of Athletic*

Training. 2001;36(3):263-273.

21.   DeBeaumont L, Mongeon D, Tremblay S, et al. Persistent motor system

abnormalities in formerly concussed athletes. *Journal of Athletic Training*.

2011;46(3):234-240. doi:10.4085/1062-6050-46.3.234

22.   Quatman-Yates CC, Bonnette S, Hugentobler JA, et al. Postconcussion Postural

Sway Variability Changes in Youth: The Benefit of Structural Variability Analyses.

Pediatric Physical Therapy. 2015;27(4):316-327. doi:10.1097/PEP.0000000000000193

23.   Fino PC, Nussbaum MA, Brolinson PG. Decreased high-frequency center-of-

pressure complexity in recently concussed asymptomatic athletes. *Gait and Posture*.

2016;50:69-74. doi:10.1016/j.gaitpost.2016.08.026

24.   Buckley TA, Munkasy BA, Tapia-Lovler TG, Wikstrom EA. Altered gait

termination strategies following a concussion. *Gait and Posture*. 2013;38(3):549-551.

doi:10.1016/j.gaitpost.2013.02.008

25.   Howell DR, Osternig LR, Chou LS. Dual-task effect on gait balance control in

adolescents with concussion. *Archives of Physical Medicine and Rehabilitation*.

2013;94(8):1513-1520. doi:10.1016/j.apmr.2013.04.015

26.   Oldham JR, Munkasy BA, Evans KM, Wikstrom EA, Buckley TA. Altered dynamic

postural control during gait termination following concussion. *Gait and Posture*.

2016;49:437-442. doi:10.1016/j.gaitpost.2016.07.327

27.   Buckley T, Murray NG, Munkasy BA, Oldham JR, Evans KM, Clouse B.

Impairments in Dynamic Postural Control Across Concussion Clinical Milestones.

Journal of Neurotrauma. 2020:neu.2019.6910-neu.2019.6910.

doi:10.1089/neu.2019.6910

28.   Buckley TA, Munkasy BA, Krazeise DA, Oldham JR, Evans KM, Clouse B.

Differential Effects of Acute and Multiple Concussions on Gait Initiation Performance.

Archives of Physical Medicine and Rehabilitation. 2020;101(8):1347-1354.

doi:10.1016/j.apmr.2020.03.018

29. Lynall RC, Campbell KR, Mauntel TC, Blackburn JT, Mihalik JP. Single-Legged

Hop and Single-Legged Squat Balance Performance in Recreational Athletes With a

History of Concussion. *Journal of athletic training*. 2020;55(5):488-493.

doi:10.4085/1062-6050-185-19

30. The Joanna Briggs I. Methodology for JBI Scoping Reviews. The Joanna Briggs

Institute; 2015.

31. Arksey H, O'Malley L. Scoping studies: Towards a methodological framework.

International Journal of Social Research Methodology: Theory and Practice.

2005;8(1):19-32. doi:10.1080/1364557032000119616

32. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C.

PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement.

Journal of Clinical Epidemiology. 2016;75:40-46. doi:10.1016/j.jclinepi.2016.01.021

33. Walker M, Pan T, Liao K. Dynamic postural instability in OEF/OIF veterans with

mild TBI and disequilibrium. *Journal of Head Trauma Rehabilitation*. 2012;27(5):E34-35.

34. Walker M, Pan T, Liao K, Roenigk K, Daly J. Postural Stability in Veterans with

Mild Traumatic Brain Injury and Disequilibrium. *Neurology*. 2015;84(14):7.161-7.161.

35.   Pan T, Liao K, Roenigk K, Daly JJ, Walker MF. Static and dynamic postural

stability in veterans with combat-related mild traumatic brain injury. *Gait and Posture*.

2015;42(4):550-557. doi:10.1016/j.gaitpost.2015.08.012

36.   Mang CS, Whitten TA, Dukelow SP, Benson BW. Upper-extremity and postural

responses to perturbed reaching movements for assessment of sport-related

concussion. *Journal of Neurotrauma*. 2018;35(16):A245-A245.

37.   Rao HM, Talkar T, Ciccarelli G, et al. Sensorimotor conflict tests in an immersive

virtual environment reveal subclinical impairments in mild traumatic brain injury.

Scientific Reports. 2020/09/08 2020;10(1):14773. doi:10.1038/s41598-020-71611-9

38.   Brown LA, Jensen JL, Korff T, Woollacott MH. The translating platform paradigm:

perturbation displacement waveform alters the postural response. *Gait and Posture*.

2001;14(3):256-263. doi:10.1016/S0966-6362(01)00131-X

39.   Freyler K, Gollhofer A, Colin R, Brüderlin U, Ritzmann R. Reactive balance

control in response to perturbation in unilateral stance: Interaction effects of direction,

displacement and velocity on compensatory neuromuscular and kinematic responses.

PLoS ONE. 2015;10(12)doi:10.1371/journal.pone.0144529

40.   Thelen DG, Wojcik LA, Schultz AB, Ashton-Miller JA, Alexander NB. Age

Differences in Using a Rapid Step To Regain Balance During a Forward Fall | The

Journals of Gerontology: Series A | Oxford Academic. *The Journals of Gerontology:

Series A*. 1997;52A(1):M8-M13.

41.   Sundermier L, Woollacott M, Roncesvalles N, Jensen J. The development of

balance control in children: Comparisons of EMG and kinetic variables and

chronological and developmental groupings. *Experimental Brain Research*.

2001;136(3):340-350. doi:10.1007/s002210000579

42.   King LA, Horak FB, Mancini M, et al. Instrumenting the balance error scoring

system for use with patients reporting persistent balance problems after mild traumatic

brain injury. *Archives of Physical Medicine and Rehabilitation*. 2014;95(2):353-359.

doi:10.1016/j.apmr.2013.10.015

43.   Register-Mihalik JK, Littleton AC, Guskiewicz KM. Are Divided Attention Tasks

Useful in the Assessment and Management of Sport-Related Concussion? :

Neuropsychol Rev; 2013. p. 300-313.

44.   Kleiner M, Wong L, Dubé A, Wnuk K, Hunter SW, Graham LJ. Dual-Task

Assessment Protocols in Concussion Assessment: A Systematic Literature Review.

Journal of Orthopaedic and Sports Physical Therapy. 2018;48(2):87-103.

doi:10.2519/jospt.2018.7432

45.   Bell DR, Guskiewicz KM, Clark MA, Padua DA. Systematic review of the balance

error scoring system. *Sports Health*. 2011;3(3):287-295.

doi:10.1177/1941738111403122

46.   Broglio SP, Ferrara MS, Sopiarz K, Kelly MS. Reliable change of the sensory

organization test. *Clinical Journal of Sport Medicine*. 2008;18(2):148-154.

doi:10.1097/JSM.0b013e318164f42a

47.   Oldham JR, Difabio MS, Kaminski TW, Dewolf RM, Howell DR, Buckley TA.

Efficacy of Tandem Gait to Identify Impaired Postural Control after Concussion.

Medicine and Science in Sports and Exercise. 2018;50(6):1162-1168.

doi:10.1249/MSS.0000000000001540

48.   Carty C, Cronin N, Nicholson D, et al. Reactive stepping behaviour in response

to forward loss of balance predicts future falls in community-dwelling older adults | Age

and Ageing | Oxford Academic. *Age and Ageing.* 2015;44(1):109-115.

49.   Mansfield A, Wong JS, McIlroy WE, et al. Do measures of reactive balance

control predict falls in people with stroke returning to the community? *Physiotherapy*

(United Kingdom). 2015;101(4):373-380. doi:10.1016/j.physio.2015.01.009

50.   Valkovic P, Broz ová H, Bötzel K, Ru z ic ka Ee, Benetin J. Push and

release test predicts better Parkinson fallers and nonfallers than the pull test:

Comparison in OFF and ON medication states. *Movement Disorders.*

2008;23(10):1453-1457. doi:10.1002/mds.22131

51.   Mak MKY, Auyeung MM. The mini-bestest can predict parkinsonian recurrent

fallers: A 6-month prospective study. *Journal of Rehabilitation Medicine.*

2013;45(6):565-571. doi:10.2340/16501977-1144

52.   Schlenstedt C, Brombacher S, Hartwigsen G, Weisser B, Möller B, Deuschl G.

Comparison of the Fullerton Advanced Balance Scale, Mini-BESTest, and Berg Balance

Scale to Predict Falls in Parkinson Disease. *Physical Therapy.* 2016;96(4):494-501.

doi:10.2522/ptj.20150249

53.   Maki BE, McIlroy WE. Control of rapid limb movements for balance recovery:

Age-related changes and implications for fall prevention. *Age and Ageing*. 2006/09//

2006;35:12-18. doi:10.1093/ageing/afl078

54.   Kontos AP, Collins MW, Holland CL, et al. Preliminary Evidence for Improvement

in Symptoms, Cognitive, Vestibular, and Oculomotor Outcomes Following Targeted

Intervention with Chronic mTBI Patients. *Mil Med*. Mar 1 2018;183(suppl_1):333-338.

doi:10.1093/milmed/usx172

55.   Möller MC, Lexell J, Wilbe Ramsay K. Effectiveness of specialized rehabilitation

after mild traumatic brain injury: A systematic review and meta-analysis. *J Rehabil Med*.

Feb 5 2021;53(2):jrm00149. doi:10.2340/16501977-2791

56.   Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred Reporting Items for

Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Medicine*.

2009;6(7):e1000097-e1000097. doi:10.1371/journal.pmed.1000097

6 Tables

Table 1. Characteristics of selected studies using discrete perturbations to examine

reactive postural response after mTBI.

Study	Population	N , groups	Testing Timeline	Perturbation Used	Outcome measures	Interpretation	Type
Mang et al. [a 36]	Athletes 17 ± 4 years	**430** BL: 430 Post-mTBI: 19	BL and ≤10 days post injury	KINARM robot, Unanticipated and anticipated upper-extremity perturbations during a reaching task	Response time, hand path length, anticipatory (mean COP velocity from 50 ms prior to reach onset to 100 ms after reach onset) and reactive postural adjustments (maximum absolute COP velocity from 100 ms after reach onset to end of reach)	Anticipatory postural adjustments, but not reactive postural adjustments, are impaired in some athletes ≤10 days post mTBI.	Abstract
Pan et al. [b 35]	Veterans 22-40 years	**24** Control: 10 No Disequilibrium: 6 Chronic Disequilibrium: 8	7 months to 7 years post injury	Motor controlled rope attached to belt at waist, unanticipated forward and rightward perturbations	Sway length and displacement for 2.5 seconds after onset of perturbation	Veterans with chronic disequilibrium after mTBI have measurable deficits in reactive balance control months to years after injury.	Article
Rao et al.[37]	mTBI: 39.1 ± 17.18 years Control: 32.4 ± 14 years	**21** mTBI: 11 Control: 10		Single, translational treadmill perturbations while standing, and while walking	Spatiotemporal (e.g. stride length), kinematic (e.g. joint angles), accelerations, EMG features (e.g.	Compared to standard clinical testing, features from sensorimotor perturbation can detect	Article

	amplitude)	lingering balance impairments after mTBI

Abbreviations: BL, testing at baseline

[a]Methods information provided via personal communication with author.

[b]Two other publications were confirmed to be part of the same study via personal communication with the author.

432 **7 Figure List**

434 **Figure 1.** Similarities and differences between the three classes of postural control.

435 BOS = Base of support; COM = Center of mass

Figure 2. PRISMA Flow Diagram [56]

439 **8   List of Supplemental Digital Content**

440 **Supplemental Digital Content 1.** Scoping review protocol.

442 **Supplemental Digital Content 2.** Detailed search strategies. Microsoft Word Doc

444 **Supplemental Digital Content 3.** Table of excluded studies with study characteristics.
445 Microsoft Word Doc