

Title	Transforming Obesity Prevention for CHILDren (TOPCHILD) Collaboration: protocol for a systematic review with individual participant data meta-analysis of behavioural interventions for the prevention of early childhood obesity
Author names	Kylie E Hunter ¹ , Brittany J Johnson ² , Lisa Askie ¹ , Rebecca K Golley ² , Louise A Baur ³ , Ian C Marschner ¹ , Rachael W Taylor ⁴ , Luke Wolfenden ⁵ , Charles T Wood ⁶ , Seema Miharshahi ⁷ , Alison J Hayes ⁸ , Chris Rissel ⁸ , Kristy P Robledo ¹ , Denise A O'Connor ^{9,10} , David Espinoza ¹ , Lukas P Staub ¹ , Paul Chadwick ¹¹ , Sarah Taki ^{8,12} , Angie Barba ¹ , Sol Libesman ¹ , Mason Aberoumand ¹ , Wendy A Smith ^{13,14} , Michelle Sue-See ¹⁴ , Kylie D Hesketh ¹⁵ , Jessica L Thomson ¹⁶ , Maria Bryant ¹⁷ , Ian M Paul ¹⁸ , Vera Verbestel ¹⁹ , Cathleen Odar Stough ²⁰ , Li Ming Wen ^{8,12} , Junilla K Larsen ²¹ , Sharleen L O'Reilly ²² , Heather M Wasser ²³ , Jennifer S Savage ²⁴ , Ken K Ong ²⁵ , Sarah-Jeanne Salvy ²⁶ , Mary Jo Messito ²⁷ , Rachel S Gross ²⁷ , Levie T Karssen ²¹ , Finn E Rasmussen ²⁸ , Karen Campbell ¹⁵ , Ana Maria Linares ²⁹ , Nina Cecilie Øverby ³⁰ , Cristina Palacios ³¹ , Kaumudi J Joshipura ^{32,33} , Carolina González Acero ³⁴ , Rajalakshmi Lakshman ²⁵ , Amanda L Thompson ^{23,35} , Claudio Maffei ³⁶ , Emily Oken ³⁷ , Ata Ghaderi ³⁸ , Maribel Campos Rivera ³⁹ , Ana B Perez-Exposito ⁴⁰ , Jinan C Banna ⁴¹ , Kayla de la Haye ⁴² , Michael Goran ⁴² , Margrethe Røed ³⁰ , Stephanie Anzman-Frasca ⁴³ , Anna Lene Seidler ¹ , on behalf of the Transforming Obesity Prevention for CHILDren (TOPCHILD) Collaboration
Author affiliations	¹ National Health and Medical Research Council Clinical Trials Centre, University of Sydney, Camperdown, NSW, Australia ² Caring Futures Institute, College of Nursing and Health Sciences, Flinders University, Adelaide, SA, Australia ³ Children's Hospital Westmead Clinical School, The University of Sydney, Westmead, NSW, Australia ⁴ Department of Medicine, University of Otago, Dunedin, New Zealand ⁵ School of Medicine and Public Health, The University of Newcastle, Newcastle, NSW, Australia ⁶ Duke University School of Medicine, Durham, NC, USA ⁷ Department of Health Systems and Populations, Faculty of Medicine, Health and Human Sciences, Macquarie University, NSW, Australia ⁸ School of Public Health, Faculty of Medicine and Health, University of Sydney, Camperdown, NSW, Australia ⁹ Monash Department of Clinical Epidemiology, Cabrini Institute, Malvern, VIC, Australia ¹⁰ Department of Epidemiology and Preventive Medicine, School of Public Health and Preventive Medicine, Monash University, Melbourne, VIC, Australia ¹¹ Centre For Behaviour Change, UCL, London, UK ¹² Population Health Research and Evaluation Hub, Sydney Local Health District, Camperdown, NSW, Australia ¹³ Canterbury Community Health Centre, Campsie, NSW Australia ¹⁴ Consumer Representative, Sydney, NSW Australia ¹⁵ Institute for Physical Activity and Nutrition, Deakin University, Geelong, VIC, Australia ¹⁶ USDA Agricultural Research Service, Stoneville, MS, USA ¹⁷ Department of Health Sciences and the Hull York Medical School, University of York, York, UK ¹⁸ Penn State College of Medicine, Hershey, PA, USA ¹⁹ Department of Rehabilitation Sciences, Ghent University, Ghent, Belgium ²⁰ Department of Psychology, University of Cincinnati, Cincinnati, OH, USA ²¹ Behavioural Science Institute, Radboud University, Nijmegen, the Netherlands ²² School of Agriculture and Food Science, University College Dublin, Dublin, Ireland ²³ Department of Nutrition, University of North Carolina at Chapel Hill, NC, USA ²⁴ Department of Nutritional Sciences & Center for Childhood Obesity Research,

	<p>Pennsylvania State University, State College, PA, USA</p> <p>²⁵ MRC Epidemiology Unit, University of Cambridge, Cambridge, UK</p> <p>²⁶ Research Center for Health Equity, Cedars-Sinai Medical Center, West Hollywood, CA, USA</p> <p>²⁷ New York University Grossman School of Medicine, New York, NY, USA</p> <p>²⁸ Department of Global Public Health, Karolinska Institute, Stockholm, Sweden</p> <p>²⁹ College of Nursing, University of Kentucky, Lexington, KY, USA</p> <p>³⁰ Faculty of Health and Sport Sciences, Department of Nutrition and Public Health University of Agder, Kristiansand, Norway</p> <p>³¹ Department of Dietetics and Nutrition, Robert Stempel College of Public Health & Social Work, Florida International University, Miami, FL, USA</p> <p>³² Center for Clinical Research and Health Promotion, University of Puerto Rico Medical Sciences Campus, San Juan, Puerto Rico, USA</p> <p>³³ Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA, USA</p> <p>³⁴ Inter-American Development Bank, Santo Domingo, Distrito Nacional, Dominica Republic</p> <p>³⁵ Department of Anthropology, University of North Carolina at Chapel Hill, NC, USA</p> <p>³⁶ Pediatric Diabetes and Metabolic Disorders Unit, University of Verona, Verona, Italy</p> <p>³⁷ Division of Chronic Disease Research Across the Lifecourse, Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA, USA</p> <p>³⁸ Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden</p> <p>³⁹ Dental and Craniofacial Genomics Core, Endocrinology Section School of Medicine, University of Puerto Rico, San Juan, Puerto Rico, USA</p> <p>⁴⁰ Inter-American Development Bank, Washington, DC, USA</p> <p>⁴¹ Department of Human Nutrition, Food and Animal Sciences, University of Hawaii, Honolulu, HI, USA</p> <p>⁴² Department of Preventive Medicine, University of Southern California, Los Angeles, CA, USA</p> <p>⁴³ Department of Pediatrics, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY, USA</p>
Corresponding author	<p>Kylie E Hunter NHMRC Clinical Trials Centre, University of Sydney Locked Bag 77, Camperdown NSW 1450, Australia kylie.hunter@ctc.usyd.edu.au</p>

TITLE

Transforming Obesity Prevention for CHILDren (TOPCHILD) Collaboration: protocol for a systematic review with individual participant data meta-analysis of behavioural interventions for the prevention of early childhood obesity

ABSTRACT

Introduction: Behavioural interventions in early life appear to show some effect in reducing childhood overweight and obesity. However, uncertainty remains regarding their overall effectiveness, and whether effectiveness differs among key subgroups. These evidence gaps have prompted an increase in very early childhood obesity prevention trials worldwide. Combining the individual participant data (IPD) from these trials will enhance statistical power to determine overall effectiveness and enable examination of intervention-covariate interactions. We present a protocol for a systematic review with IPD meta-analysis to evaluate the effectiveness of obesity prevention interventions commencing antenatally or in the first year after birth, and to explore whether there are differential effects among key subgroups.

Methods and analysis: Systematic searches of Medline, Embase, CENTRAL, CINAHL, PsycInfo, and trial registries for all ongoing and completed randomised controlled trials evaluating behavioural interventions for the prevention of early childhood obesity have been completed up to March 2020 and will be updated annually to include additional trials. Eligible trialists will be asked to share their IPD; if unavailable, aggregate data will be used where possible. An IPD meta-analysis and a nested prospective meta-analysis (PMA) will be performed using methodologies recommended by the Cochrane Collaboration. The primary outcome will be body mass index (BMI) z-score at age 24 +/- 6 months using World Health Organisation Growth Standards, and effect differences will be explored among pre-specified individual and trial-level subgroups. Secondary outcomes include other child weight-related measures, infant feeding, dietary intake, physical activity, sedentary behaviours, sleep, parenting measures and adverse events.

Ethics and dissemination: Approved by The University of Sydney Human Research Ethics Committee (2020/273) and Flinders University Social and Behavioural Research Ethics Committee (project no. HREC CIA2133-1). Results will be relevant to clinicians, child health services, researchers, policy-makers and families, and will be disseminated via publications, presentations, and media releases.

Registration: Prospectively registered on PROSPERO: CRD42020177408

KEYWORDS

Early childhood overweight/obesity, prevention, individual participant data, prospective meta-analysis, behavioural interventions, diet, physical activity, sleep, sedentary behaviours, infancy

STRENGTHS AND LIMITATIONS OF THIS STUDY

- This will be the largest individual participant data (IPD) meta-analysis evaluating behavioural interventions for the prevention of early childhood obesity to date, and will provide the most reliable and precise estimates of early intervention effects to inform future decision-making.
- IPD meta-analysis methodology will enable unprecedented exploration of important individual and trial-level characteristics that may be associated with childhood obesity or that may be effect modifiers.
- The proposed innovative methodologies are feasible and have been successfully piloted by members of our group.
- It may not be possible to obtain IPD from all eligible trials; in this instance, aggregate data will be used where available, and sensitivity analyses will be conducted to assess inclusion bias.
- Outcome measures may be collected and reported differently across included trials, potentially increasing imprecision; however, we will harmonise available data where possible, and encourage those planning or conducting ongoing trials to collect common core outcomes following prospective meta-analysis methodology.

INTRODUCTION

Childhood obesity is one of the most serious public health issues of the 21st century, and requires urgent action.^{1, 2} Globally, an estimated 38 million (6%) children aged under five years were living with overweight or obesity in 2019,³ and prevalence is increasing across every continent as environments become more obesity conducive.^{4, 5} While childhood obesity affects all sections of society, it disproportionately affects racial and ethnic minority groups^{6, 7} and populations with a lower socioeconomic position (SEP), and thus is also a major health equity issue.⁴ Children with obesity are much more likely to have obesity across the lifecourse,^{8, 9} and are at increased risk of short and long term negative health sequelae, such as poor mental and musculoskeletal health, type 2 diabetes, asthma, and cardiovascular disease.^{10, 11} This places a large burden on healthcare systems,¹² and has significant economic consequences arising from increased disability and decreased productivity and life expectancy.¹³ Thus, identifying modifiable behaviours for the early prevention of childhood obesity is critical to inform the development of early intervention strategies.

There are a variety of modifiable behaviours that may influence energy balance and therefore may be implicated in childhood obesity prevention, namely, feeding practices, dietary intake, physical activity, sedentary behaviours and sleep. For instance, appropriate responsive feeding has been identified as promising for obesity prevention,¹⁴⁻¹⁶ while consumption of sugar sweetened beverages is associated with severe obesity in children aged less than 5 years.¹⁷ Data are mixed on the protective benefits of breastfeeding for the prevention of obesity, though some studies suggest that longer duration of exclusive breastfeeding may provide modest protection.¹⁸⁻²³ Similarly, there may be an association between age at introduction of solids and growth,²⁴ with mixed results surrounding the direction of this association and the underlying causal mechanisms. Previous systematic reviews have reported significant inverse associations between physical activity and measures of adiposity in children.²⁵⁻²⁷ Conversely, sedentary behaviours such as television viewing or screen time are associated with higher body mass index (BMI) levels^{28, 29} and greater adiposity³⁰ in young children. There is now also a large body of observational evidence supporting the relationship between short sleep duration and an increased risk of obesity across all age groups, including infants and young children,³¹⁻³⁵ though a recent systematic review found inconsistent evidence of an association between longer infant sleep duration and healthier body composition up to age 24 months.³⁶

In addition to these behaviours, individual-level covariates known or hypothesised to be predictive for childhood obesity include pre-pregnancy maternal and paternal BMI, age, race, ethnicity, socioeconomic position, excess gestational weight gain, parity, smoking during pregnancy, gestational

diabetes, birth mode of delivery (caesarean, vaginal), birthweight, gestational age at birth, baby's sex, and childcare attendance.^{6, 7, 22, 37} Some of these covariates may also be individual-level effect modifiers, predicting how effective an intervention is likely to be, e.g. socioeconomic position and race/ethnicity. Trial-level characteristics such as timing of intervention onset, setting and the level of well-child health care available in the community may also modify intervention effectiveness.³⁸

Limitations and evidence gaps identified in previous reviews

In the past five years there have been numerous reviews of childhood obesity prevention trials encompassing a variety of intervention types, settings and age groups.^{14, 39-45} Few of these focused solely on infancy, and many spanned multiple life stages from the prenatal period to 18 years of age. One review found that family-based childhood obesity prevention interventions most frequently targeted children 2-10 years of age (78%), with fewer targeting infants aged 0-1 year (24%) or the prenatal period (8%).³⁹ Most reviews highlighted the urgent need for further rigorous evidence to inform obesity prevention interventions in the very early childhood years.^{14, 39-43, 45} Given the consequences of rapid early life weight gain, associated epigenetic changes and early onset of obesity in many children,^{3, 46, 47} there is strong rationale to start preventive interventions early when biology is most amenable to change, and before negative obesity-conducive behavioural patterns are established.²

Most of the childhood obesity prevention reviews to date have used qualitative methodology such as narrative reviews, content analysis, and systematic reviews without meta-analysis to describe variations in study design, setting, population, interventions, and outcomes, and to hypothesise that certain individual and trial-level characteristics may enhance effectiveness via proposed conceptual frameworks and intervention models.^{14, 39, 41-45} Yet, quantitative evaluation is required to formally test these hypotheses. Recently, Brown et al⁴⁰ updated a Cochrane systematic review and aggregate data meta-analysis on obesity prevention in children aged 0-18 years, and found that interventions focusing on diet and physical activity combined can lead to a small reduction in BMI z-score in children aged 0-5 years of age (mean difference -0.07, 95% CI -0.14 to -0.01). However, a huge variety in intervention approaches limited their ability to conduct meaningful comparisons, and many multicomponent interventions were originally reported as a whole package, precluding evaluation of discrete intervention characteristics. Moreover, the aggregated data were insufficient to derive conclusions on effect differences by individual-level characteristics such as ethnicity and socioeconomic position.

The Early Prevention of Obesity in Children (EPOCH) Collaboration conducted a world-first IPD prospective meta-analysis (PMA) of four randomised controlled trials (RCTs) of behavioural interventions for the prevention of early childhood obesity.³⁸ They found that, compared with usual

care, early childhood interventions were modestly effective in reducing BMI z-score 18-24 months after birth by 0.12 standard deviations (which translates to a 2% decrease in obesity prevalence). However, when accounting for missing data this difference was no longer significant. There was some heterogeneity across trials, and interventions appeared to be more effective in populations with limited publicly funded existing health care programs, in this instance defined as a maximum of one postnatal home visit.³⁸ However, this finding needs to be confirmed in analyses including more than four studies. EPOCH's predictive analyses of individual and trial-level factors did not have sufficient power to detect reliable differences in BMI z-score. Thus, the overall effectiveness of early obesity prevention interventions remains uncertain, as does whether there may be differential effects amongst subgroups.

Need for individual participant data meta-analysis

The limitations and evidence gaps described above highlight the need for more powerful and in-depth analyses focusing on preventive interventions in very early childhood. Since the EPOCH PMA,³⁸ we have identified more than 60 additional ongoing or completed very early obesity prevention trials worldwide with a combined sample size of more than 50,000 participants. While most trials are powered to detect some important differences in key outcomes, individually they have limited power to detect a difference in our primary outcome, BMI z-score at 24 +/- 6 months of age. In order to reliably detect a reduction in BMI z-score similar to that seen in EPOCH (-0.12),³⁸ 2920 participants are required (90% power, two-sided 5% level of significance). Moreover, usually about four times that sample size (n~12,000) is required to detect differences in subgroups.⁴⁸ The expected total sample size for TOPCHILD will exceed these estimates (as by December 2020, 37 trials including 34,370 participants have already agreed to share their IPD).

Conducting a trial of this size would be time and resource intensive. A more efficient method is to combine IPD from trials in a pooled analysis to increase the sample size and therefore statistical power. This strengthens the chance of detecting intervention effect differences, and enables us to determine the size of such effects with greater certainty,⁴⁹ while also allowing variation in study designs and population which heightens generalisability and allows a greater diversity to study effect modification for different subgroups of individuals or trial characteristics.⁵⁰ Moreover, this collaborative approach maximises the use of existing data, thereby reducing research waste.

Thus, we will conduct an IPD meta-analysis with detailed subgroup analyses of all available trials to confirm whether early obesity prevention interventions commencing antenatally or in the first year after birth are effective, and whether effectiveness varies across subgroups defined by individual- or trial-level characteristics. The knowledge generated from this study can be used to inform decision-

making around the design and implementation of more effective, efficient, equitable and targeted interventions for the prevention of childhood obesity and its sequelae.

Objectives

This IPD meta-analysis will address the following research questions:

- 1) Compared with usual care/attentional control, what are the effects of parent/caregiver-focused behavioural obesity prevention interventions commencing during pregnancy or infancy on:
 - a. child BMI z-score at age 24 months (+/- 6 months)? (primary outcome)
 - b. child BMI z-score at alternative timepoints, other child weight-related measures, infant feeding, dietary intake, physical activity, sedentary behaviours, sleep, parenting measures and adverse events? (secondary outcomes)
- 2) Do intervention effects vary across individual-level characteristics (e.g. parental BMI, parity, socioeconomic position, birthweight)?
- 3) Do intervention effects vary across trial-level characteristics (e.g. access to existing well-child health care programs, intervention mode of delivery, timing of intervention onset)?

METHODS AND ANALYSIS

We will conduct a systematic review with IPD meta-analysis and a nested PMA according to the methods recommended by the Cochrane Collaboration.^{51, 52} A nested PMA enables integration of prospective evidence into a retrospective meta-analysis, and harmonisation among planned/ongoing studies.⁵² Lead investigators of eligible trials will be invited to share their IPD and join the Transforming Obesity Prevention for CHILDren (TOPCHILD) Collaboration (www.topchildcollaboration.org). This protocol adheres to the Preferred Reporting Items for Systematic Reviews and Meta-Analysis extension for protocols (PRISMA-P)⁵³ (Supplementary Appendix 1).

Eligibility criteria

Types of studies

This systematic review will include randomised controlled trials only, including feasibility studies, pilot trials and definitive trials. Randomisation may occur at the individual level or by cluster (e.g. child care, community), including stepped-wedge designs. Quasi-randomised trials are excluded as they may introduce bias. There are no language or date restrictions.

Trial participants

Participants will be parents/caregivers (including pregnant women) and their infant(s) aged 0-12 months (at baseline). Caregiver is defined as the person with primary responsibility for care of the child, and excludes secondary sources of support, such as child care providers and early childhood teachers. Women may be primipara or multipara, and both singletons and multiples are eligible.

Types of interventions

Interventions must be behavioural interventions targeting parents/caregivers, and include at least one component related to modifiable child behaviours that may influence overweight/obesity risk (e.g. infant feeding, dietary intake, physical activity, sedentary behaviours, sleep). They may commence in the pre-conception or antenatal phase but must include intervention exposure targeting the birth to 12 months infancy stage, as pregnancy-only interventions are considered distinct and are currently being examined by Dodd et al in a separate IPD meta-analysis.⁵⁴ Only childhood obesity prevention-focused trials will be included; these are defined as trials that clearly state childhood obesity prevention as a key aim/objective. Interventions focused only on improving an obesity related behaviour (e.g. sleep, delayed introduction of solid foods), as well as those focused on treatment of obesity, stunting or underweight will be excluded. Trials with a dual focus to prevent obesity and undernutrition are eligible, though we will carefully consider and pre-specify how their data will be incorporated in the statistical analysis plan. Interventions focused solely on nutritional supplements will be excluded, as they are not considered to be behavioural interventions.

Types of comparator/control

Eligible trials must have either 1) a usual care control arm, defined as existing local child health care, or 2) no intervention (including waitlist control) or 3) attention control (e.g. child safety education).

Types of outcome measures

To be included, trials must collect at least one of the child weight-related outcomes listed in Table 1 post intervention (at any age), i.e. BMI/BMI z-score, prevalence of overweight/obesity, percent fat content/adiposity, skinfold thickness, abdominal circumference, waist-to-height ratio. This is considered a legitimate and pragmatic approach given our review is of multi-component public health interventions focusing on obesity.⁵⁵

Eligibility for nested prospective meta-analysis

In accordance with PMA methodology,⁵² only planned/ongoing trials will be eligible for the nested PMA if trial results were not yet known to the investigator/s at the time the main components of the TOPCHILD protocol (i.e. aims and objectives, hypotheses, eligibility criteria, main outcomes, subgroup

and sensitivity analyses) were initially agreed in December 2020. We encourage investigators of planned/ongoing studies to collect the outcomes and covariates listed in Table 1 where possible, to facilitate data harmonisation and synthesis.

Information sources and search strategy

In March 2020 we undertook an initial systematic search for eligible trials using the following databases from their inception: Medline (Ovid), Embase (Ovid), Cochrane Central Register of Controlled Trials (CENTRAL), CINAHL (EBSCO), PsycInfo, ClinicalTrials.gov and the World Health Organization's International Clinical Trials Registry Platform's (WHO ICTRP) Search Portal. The full search strategy is available in Supplementary Appendix 2. This search will be updated annually for the duration of the TOPCHILD Collaboration (currently funded until end 2023). Collaborators and contacts will also be asked to notify us of any planned, ongoing or completed trials of which they are aware that may meet the eligibility criteria.

Selection of studies for inclusion in the review

Two members of the TOPCHILD Steering Group will independently screen all retrieved records against eligibility criteria. Any discrepancies will be resolved by discussion or, if required, adjudication by a third reviewer from the Steering Group. The Principal Investigator and/or corresponding author of eligible trials will be invited by email to join the TOPCHILD Collaboration. If there is no response to initial emails and reminders, we will contact co-authors and/or other contacts listed in registration records and consult our existing networks to see if they can reach out to those they may know. If IPD cannot be obtained for an eligible trial, we will use aggregate data sourced from publications where available.

Supplementary Appendix 3 lists eligible trials identified up to March 2020.

Data collection, management, and confidentiality

Data receipt / extraction

Trialists of all eligible studies will be invited to share de-identified individual participant data via secure data transfer platforms or via an institutional-secure email using password-protected zip files. Data will be provided according to a pre-specified coding template where possible. Otherwise, data will be accepted in any format and recoded as necessary. The data management team (within the TOPCHILD Steering Group) will receive and store the data in perpetuity in a secure, customised database at the NHMRC Clinical Trials Centre, University of Sydney, and data management will follow the *University of Sydney Data Management Policy 2014*. Each trial will also be asked to provide metadata (i.e. data that

provides information about their trial dataset), such as questionnaires, data collection forms, and data dictionaries to aid understanding of the dataset. Trial-level data, such as setting, intervention timing, mode of delivery, comparator/control details, method of sequence generation, allocation concealment, geographical location, sample size, outcome measures and definitions will be extracted into a database and cross-checked against any published reports, trial protocols, registration records and data collection sheets.

Data processing

Data from each trial will be checked with respect to range, internal consistency, consistency with published reports and missing items. Integrity of the randomisation process will be examined by reviewing the chronological randomisation sequence and pattern of assignment, as well as the balance of participant characteristics across intervention and control groups. Any inconsistencies or missing data will be discussed with trialists and/or data managers and resolved by consensus. Once finalised, data from each of the trials will be combined into a single TOPCHILD Collaboration database.

Risk of bias assessment and certainty of evidence appraisal

Included studies will be assessed for risk of bias by two independent reviewers from the TOPCHILD Steering Group using Version 2 of the Cochrane risk-of-bias tool for randomised trials (RoB 2).⁵⁶ This tool includes five domains encompassing bias arising from: the randomisation process, deviations from intended interventions, missing outcome data, measurement of the outcome, and selection of the reported result. For cluster-randomised trials, bias arising from identification or recruitment of individual participants within clusters will also be assessed.⁴⁵ The certainty of evidence will be assessed according to Cochrane procedures⁵⁷ using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.⁵⁸ Any differences will be resolved by consensus or with a third reviewer from the TOPCHILD Steering Group.

Primary outcome

The primary outcome will be BMI z-score at age 24 months (+/- 6 months). BMI z-score will be determined in accordance with World Health Organization (WHO) growth standards.⁴⁷

Secondary outcomes

All outcomes are detailed in Table 1. Where possible, definitions will be standardised, otherwise outcomes will be used as defined within each trial. Secondary outcomes include BMI z-score at other

timepoints, other measures of child weight, infant feeding (including breastfeeding and introduction of solid foods), dietary intake, sedentary behaviours, physical activity, and sleep, as well as parent/caregiver-related measures. We will also assess any adverse events, such as underweight or poor weight gain.

Covariates/subgroups

All included covariates are listed in Table 1. Individual-level and trial-level subgroup analyses will be conducted for the primary outcome of BMI z-score at age 24 months (+/- 6 months). Those of primary interest at the individual level include socioeconomic position, race/ethnicity, pre-pregnancy maternal/paternal BMI, maternal age, gestational weight gain and parity, and at the trial level include timing of intervention onset, current level of background care in the community, recruitment country and mode of delivery.

Where possible, outcomes and covariates will be collected as continuous variables to maximise power to detect intervention effects and interactions, and enable exploration of any non-linear relationships.⁵⁹ Dichotomous and categorical variables will also be collected to aid interpretation, and if data are insufficient for the pre-specified subgroup analyses, categories will be collapsed prior to any analyses being conducted.

Table 1. Outcomes and covariates/subgroups

<i>Variable</i>	<i>Definition/explanatory text/examples</i>
Primary outcome	
BMI z-score at age 24 months (+/- 6 months)	determined in accordance with WHO growth standards ⁶⁰
Secondary outcomes	
BMI z-score at 12 months (+/- 3 months)	determined in accordance with WHO growth standards ⁶⁰
BMI z-score at 48 months (+/- 12 months)	determined in accordance with WHO growth standards ⁶⁰
Other weight-related measures	e.g. prevalence of overweight/obesity (defined as BMI z-score of at least 2 standard deviations above the WHO reference), percent fat content/adiposity, skinfold thickness, abdominal circumference, waist-to-height ratio, velocity of weight gain
Infant feeding	e.g. breastfeeding initiation and duration, exclusivity of breastfeeding, age at introduction of solid foods (complementary feeding)
Dietary intake	e.g. energy intake, intake of fruit, vegetables, energy dense nutrient poor (EDNP) foods, and sugar-sweetened beverages (SSBs)
Sedentary behaviours	e.g. screen time, restrained time while awake (in prams/strollers, high-chairs, strapped on a caregiver's back or chest)
Physical activity	e.g. active play duration, prone play ('tummy time'), device assessed physical activity time
Sleep	e.g. sleep duration, measures of sleep quality such as frequency and duration of

It is made available under a [CC-BY-NC 4.0 International license](https://creativecommons.org/licenses/by-nc/4.0/) .

	waking at night
Parental/caregiver measures	general and domain-specific parenting styles and practices, e.g. parenting self-efficacy, parenting styles, parent feeding practices, parent physical activity practices, parent sleep practices, stress
Adverse events	e.g. underweight, injuries, infection
Individual-level covariates /subgroups	
Socioeconomic position	e.g. household income/country median household income, parent/caregiver highest education level, employment status
Parental weight status	e.g. maternal pre-pregnancy BMI, paternal BMI
Race/ethnicity	trialist defined
Maternal age	at recruitment
Maternal gestational weight gain	in kilograms
Parity	primipara, multiparous
Mode of delivery at birth	caesarean, vaginal
Birthweight	in grams
Weight for gestational age	small-for-gestational-age (SGA), appropriate-for-gestational-age (AGA), large-for-gestational-age (LGA)
Sex	female, male, uncertain/other
Gestational age at birth	preterm, term
Household composition	e.g. 2 vs 1 adult household, siblings, marital status
Type of pregnancy	singleton, multiple
Maternal diabetes	gestational, type 1, type 2
Smoking during pregnancy	yes/no
Infant's age at enrolment	in months
Child's age at final assessment	in months
Child care attendance	yes/no
Trial-level covariates/subgroups	
Delivery mode (intervention)	e.g. face-to-face, letter, mobile digital device, individual vs group
Intervention setting	e.g. household residence, community healthcare facility
Intervention dose/intensity	e.g. total number of contacts, frequency of contact, duration of contact
Fidelity	planned, actual
Timing of intervention onset	pre-conception, antenatal, postnatal
Timing of intervention completion	child age in months
Current level of background care in the community	descriptive, categorisation to be determined, e.g. expected number of health contacts between birth and 1 year, expectation of attending prenatal programs (yes/no), etc.
Country	low, middle, high income
Behavioural +/- other intervention type	behavioural intervention(s) alone vs behavioural + other intervention type (e.g. supplement)

Data analysis

A detailed statistical analysis plan will be prepared and agreed upon by the TOPCHILD Collaboration members prior to any analyses being undertaken. Analyses will follow the intention-to-treat principle and include all randomised infant-parent/caregiver dyads for which data are available (including any that were excluded from the original study analysis). For cluster RCTs, correlated data will be taken into account by fitting the models using generalised estimating equations to derive appropriate standard errors. Correlations between multiples also will be accounted for in the analyses.

The primary analysis for all outcomes will be conducted using a one-stage approach combining all available IPD and aggregate data (where IPD are unavailable) to reduce the risk of availability bias.^{61, 62} The combined dataset will be analysed including trial as a random effect. Models will be chosen appropriate to the outcome type. Generalised linear models with appropriate distributions and link functions will be used for continuous and binary outcomes, while Cox proportional hazards regression will be used to analyse time-to-event outcomes subject to censoring. For example, linear models will be used for the primary outcome while relative risk binomial regression with log link function will be used for prevalence of overweight/obesity, and Cox models will be used for breastfeeding duration. Where possible, continuous covariates and outcomes will be analysed on their continuous scale to maximise utility of available data.⁵⁹

Heterogeneity of intervention effects across trials will be investigated using quantitative measures (I^2) supplemented by graphical presentations as recommended in the Cochrane Handbook.⁶³ Any notable heterogeneity identified will be explored further to ascertain if the combination of trials is appropriate.

Results will be reported using appropriate estimates of intervention effect (relative risks, mean differences or hazard ratios) with 95% confidence intervals and associated two-sided p values. For trials with multiple intervention arms, we will present the data for each intervention arm compared to the control arm, with the number of participants in the control arm adjusted to ensure no double counting.⁴⁰ Missing data will be explored in sensitivity analyses using appropriate methods. All analyses will be performed using the open-source software R.⁶⁴

Differences in intervention effect between the pre-specified subgroups will be examined by testing a treatment by subgroup interaction term within the 1-stage-model. Findings of subgroup analyses will be reported as exploratory,⁶⁵ and summarised using a 1-stage-approach supplemented by graphical presentation in a forest plot using a 2-stage-approach. Non-linear relationships will be explored for continuous covariates using a multivariate meta-analysis of the trend.⁵⁹

Other exploratory analyses for the primary outcome will include graphical presentation of BMI z-score distributions to investigate any differences beyond mean differences and examine any non-linear relationships. The potential for mediation and moderator analyses using parent/caregiver measures will be explored and detailed in a statistical analysis plan after we have extracted information about relevant variables collected by included trials.

Assessment of selection or publication bias

Potential selection bias and publication bias will be investigated by conducting a nested prospective meta-analysis (PMA) and comparing prospectively versus retrospectively included trials in a sensitivity analysis.⁵² We will also seek to include any unreported outcomes sourced from each trial's IPD, which may alleviate selective outcome reporting bias.⁵¹ Lastly, contour-enhanced funnel plots will be used to examine whether there are differences in results between more and less precise studies.

Adjustments for multiple testing

Only one primary outcome was selected for this study (Table 1). For secondary outcomes and subgroup analyses, no formal adjustments will be made for the potential inflation of type 1 error rates due to multiple testing. Instead, we will follow Schulz and Grimes' approach⁵⁴ and recommendations of the Cochrane Collaboration.⁶³ This involves cautious interpretation of the magnitudes of effect, patterns and consistency of results across related outcomes and clinical/biological plausibility rather than focusing on any single statistically significant result in isolation which can be extremely misleading.^{63, 65}

Planned sensitivity analyses

Where possible, the following sensitivity analyses will be conducted for the primary outcome:

- Two stage approach
- Including IPD only, i.e. excluding trials without IPD available⁵⁵

- Including prospectively included trials only (nested PMA), i.e. planned/ongoing trials for which results were not yet known to investigator/s at the time the main components of the TOPCHILD protocol were agreed⁵²
- Adjusting for birthweight as a covariate
- Excluding trials with a high risk of bias for sequence generation and/or allocation concealment and/or loss to follow-up
- Excluding trials with a significant conflict of interest (e.g. funded by industry)
- The impact of missing data on conclusions about the intervention effect (if appropriate)

Project management

Membership of the TOPCHILD Collaboration includes trial representatives from each of the trials contributing IPD to the project, a Steering Group and an Advisory Group. Trial representatives have the opportunity to contribute their expert knowledge to the TOPCHILD Collaboration and provide input into the protocols, statistical analysis plan, and final results manuscript. The Steering Group will be responsible for data collection, management and analysis, as well as communication within the Collaboration, including newsletter updates, maintenance of the TOPCHILD website and organisation of virtual or face-to-face collaborator meetings. The Advisory Group will comprise invited experts in childhood obesity prevention, IPD meta-analysis, statistics, behaviour change theory/methods and policy implementation.

Patient and public involvement

The TOPCHILD Collaboration involves a broad range of stakeholders including health professionals, policy-makers and trialists. In addition, the Advisory Group includes a parent representative and intervention facilitator/nurse. They have given input into and feedback on this protocol and will be involved in discussion and interpretation of results.

ETHICS AND DISSEMINATION

Ethical considerations

IPD will be provided by each included trial on the stipulation that ethical approval has been provided by their respective Human Research Ethics Committees (or equivalent), and participants gave informed consent before enrolment to participate in the initial individual trials. Trialists remain the custodians of their own data, which will be de-identified before being shared with the TOPCHILD Collaboration. Ethical approval for this project has been granted by The University of Sydney Human

Research Ethics Committee (2020/273) and Flinders University Social and Behavioural Research Ethics Committee (project no. HREC CIA2133-1).

Publication policy

TOPCHILD manuscripts will be prepared by the Steering Group in consultation with the Advisory Group, and circulated to the full Collaboration for comment, revision and approval prior to submission for publication. Any reports of the results of this study will be published either in the name of the collaborative group, or by representatives of the collaborative group on behalf of the TOPCHILD Collaboration, as agreed by members of the collaborative group.

DISCUSSION

This will be the largest IPD meta-analysis to date of trials evaluating behavioural obesity prevention interventions commencing in very early childhood. The findings will inform next generation obesity prevention initiatives that are effective, efficient, and equitable. Such interventions could set children on a better health trajectory early on and reduce the potentially life-long burden of disease associated with obesity.

The main strengths of this study arise from use of IPD meta-analysis methodology, which is considered the 'gold standard'.⁶⁶ It involves collecting the raw line-by-line data for each participant in each study from the original trialists. This can improve the quality of data, and enables more in-depth and precise analyses than would be possible using only published aggregate data.⁵¹ In particular, IPD meta-analysis will enable thorough exploration of individual-level and trial-level subgroups, so that we may quantify any differential effects and uncover the key determinants of successful outcomes. This addresses the limitations identified in previous reviews of childhood obesity prevention,^{38, 40, 45} where such detailed and sufficiently powered analyses were simply not possible.

A potential limitation of this study is the risk of not obtaining IPD from all eligible studies, resulting in inclusion bias. Where available, we will include aggregate data from these studies, and conduct sensitivity analyses with inclusion of IPD only to explore potential bias.⁶⁷ Further, there may be variations across studies in how measures are collected and reported, which may lead to some imprecision and difficulties pooling the data. We will seek to address this using nested prospective meta-analysis methodology, whereby researchers of planned or ongoing trials are encouraged to harmonise their trial design and collect core outcome measures to facilitate meta-analysis and

interpretation.⁵² For completed studies, we will derive common outcome variables by cleaning, re-coding, and converting existing measures where possible.

We plan to complete the first round of study identification and IPD collection by early 2021, then conduct the primary analyses and disseminate the results by the end of 2022. Trials that are not completed in time to provide data for this cycle will remain a part of the TOPCHILD Collaboration, and their data will be included in future updates of TOPCHILD.

This IPD meta-analysis will be conducted in parallel with a complementary TOPCHILD project (Johnson et al, unpublished), which aims to deconstruct childhood obesity interventions into their components (i.e. delivery features, target behaviours and behaviour change techniques) using systematic, internationally recognised frameworks and both published and unpublished trial materials. In future, the resulting dataset curated from these two projects will be used for predictive modelling of intervention component effectiveness at an individual participant level, facilitating a personalised or precision medicine approach to public health prevention.

The TOPCHILD Collaboration will maximise use of existing trial data that will enable us to understand and utilise the most effective intervention components for specific population groups and contexts. It will provide urgently needed evidence to inform development and implementation of effective, efficient and equitable interventions for the prevention of early childhood obesity. The results will be of prime importance for guideline developers, policymakers, consumers and the research community. Further information and updates on the TOPCHILD Collaboration can be found at www.topchildcollaboration.org.

Acknowledgements

The authors thank Slavica Berber for guidance and input into the database search strategy. We would also like to acknowledge the NHMRC Centre of Research Excellence in Early Prevention of Obesity in Childhood (EPOCH CRE), who supported the pilot and foundational work for this project. Lastly, we thank Maria Luisa Garmendia and Camila Corvalan for their feedback on an earlier version of this protocol.

TOPCHILD Collaboration members

Steering group: Anna Lene Seidler, Kylie Hunter, Brittany Johnson, Rebecca Golley, Angie Barba, Lisa Askie, Mason Aberoumand, Sol Libesman

Advisory group: Louise Baur, Paul Chadwick, David Espinoza, Alison Hayes, Ian Marschner, Seema Mhrshahi, Denise O'Connor, Chris Rissel, Kristy Robledo, Lee Sanders, Wendy Smith, Lukas Staub, Michelle Sue-See, Sarah Taki, Rachael Taylor, Luke Wolfenden, Charles Wood, Shonna Yin

Trial representatives (to date): Stephanie Anzman-Frasca, Jinan Banna, Maria Bryant, Karen Campbell, Maribel Campos Rivera, Eva Corpeleijn, Lynne Daniels, Kayla de la Haye, Ata Ghaderi, Carolina González Acero, Michael Goran, Rachel Gross, Christine Helle, Kylie Hesketh, Kaumudi Joshipura, Alison Karasz, Levie Karssen, Rajalakshmi Lakshman, Junilla Larsen, Ana Maria Linares, Claudio Maffei, Mary Jo Messito, Emily Oken, Ken Ong, Sharleen O'Reilly, Nina Øverby, Cristina Palacios, Ian Paul, Ana Perez-Exposito, Eliana Perrin, Hein Raat, Finn Rasmussen, Margrethe Røed, Russell Rothman, Sarah-Jeanne Salvy, Jennifer Savage, Cathleen Odar Stough, Barry Taylor, Rachael Taylor, Amanda Thompson, Jessica Thomson, Vera Verbestel, Heather Wasser, Li Ming Wen

Author contributions

ALS together with KEH, BJJ, LA, RKG conceived the idea for the study.

KEH, BJJ, ALS, LA, RKG developed the research question and protocol registration.

KEH wrote the first draft of the manuscript.

KEH, BJJ, ALS, RKG, LA developed the eligibility criteria and search strategy.

KEH, MA, AB, BJJ, SL, ALS performed the search and screening.

AJH, CR, CTW, DE, DAO'C, ICM, KPR, LAB, LPS, LW, MS-S, PC, RWT, SM, ST, WS provided critical review and feedback at each stage of the process.

All authors critically revised the manuscript for intellectual content, and agreed and approved the final manuscript. KEH is the guarantor of the review.

Funding statement

This work was supported by the Australian National Health and Medical Research Council (NHMRC) Ideas Grant TOPCHILD (Transforming Obesity Prevention for CHILDren): Looking into the black box of interventions (GNT1186363). Funders had no role in developing this protocol. Individual authors declare the following funding: ALT reports funding from NIH R01HD073237; AML reports funding from NIH National Center for Advancing Translational Science through grant # UL1TR000117 and UL1TR001998; BJT reports funding from Health Research Council of New Zealand; CGA reports funding from The PepsiCo Foundation; CP reports funding from National Institutes of Health, Robert Wood Johnson Foundation, World Health Organization, US Department of Agriculture; COS reports funding from University of Cincinnati University Research Council; DAOC is supported by an Australian National

Health and Medical Research Council (NHMRC) Translating Research into Practice Fellowship (APP1168749); EO reports the PROBIT study was supported by grant MOP-53155 from the Canadian Institutes of Health Research and grant R01 HD050758 from the US National Institutes of Health; IMP reports funding from grant R01DK088244 from the United States National Institute of Diabetes and Digestive and Kidney Diseases; JCB reports funding from US Department of Agriculture; JKL and LTK reports funding from Fonds NutsOhra awarded (100.939); JSS reports funding from NIH NIDDK, NIH NHLBI, PCORI; JLT is an employee of USDA ARS and the Agency did fund the Delta Healthy Sprouts Trial (Project 6401-5300-003-00D); KdlH reports funding from 1R01HD092483-01 (MPI: de la Haye, Salvy), NIH/NICHD; KDH is supported by an Australian Research Council Future Fellowship (FT130100637); LMW reports funding from NHMRC (#393112; #1003780; #1169823); LW is supported by a NHMRC Career Development and NHF Future Leader Fellowship; MB reports HAPPY was funded by a UK NIHR Programme Grant for Applied Research (project number RP-PG-0407-10044); MCR reports The Baby Act Trial was sponsored by the Center for Collaborative Research in Health Disparities under grant U54 MD007600 of the National Institute on Minority Health and Health Disparities from the National Institutes of Health; MJM reports funding from USDA AFRI 2011-68001-30207; NCØ reports their original study was partly funded by the Norwegian Women's Public Health Association, who had no influence on any part of the study design, implementation and evaluation; RKG is a Chief Investigator on the Early Prevention of Obesity in Childhood, NHMRC Centre for Research Excellence (1101675); RSG is supported by the National Institute of Food and Agriculture/US Department of Agriculture, award number 2011-68001-30207, and the National Institutes of Health/National Institute of Child Health and Human Development through a K23 Mentored Patient-Oriented Research Career Development Award (K23HD081077; principal investigator: Rachel S. Gross); RL reports funding from UK NPRI (National Prevention Research Initiative), MRC PHIND (Public Health Intervention Development programme); SLOR reports funding from European Unions Horizon 2020 Research and Innovation Programme (Grant Agreement no. 847984) and Australia National Health and Medical Research Council (NHMRC) (Grant application no. APP1194234); S-JS reports funding from NIMHD U54MD000502 (MPI: Salvy & Dutton Project #2), NICHD R01HD092483 (MPI: Salvy & de la Haye); SA-F is a co-investigator on the current INSIGHT grant which follows participants to ages 6 and 9: NIH 2R01DK088244; ABP-E reports the SPOON program in Guatemala is funded by the Inter-American Development Bank with donations of The Government of Japan and The PepsiCo Foundation.

Competing interests statement

Authors listed as Trial Representatives are investigators of eligible trials. All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: no support from

any organisation for the submitted work; AB, ALS, BJJ, KEH, MA, RKG, SL and LPS reports grants from NHMRC Ideas Grant TOPCHILD (Transforming Obesity Prevention for CHILDren) (GNT1186363); APE and CGA reports grants administered by the Inter-American Development Bank from The Government of Japan and The PepsiCo Foundation; ALT reports grants from National Institute of Health; BJT reports grants from NZ Health Research Council; EO reports grants from the US National Institutes of Health, and the Canadian Institutes for Health Research; IMP reports grants from NIH/NIDDK; JSS reports grants from PCORI, NIH NIDDK and NHLBI, and personal fees from Danone Organic, American Academy of Pediatrics, and Lets Move Maine; LTK and JKL reports grants from Fonds NutsOhra; MCR reports grants from National Institute on Minority Health and Health Disparities-National Institutes of Health/Center for Collaborative Research in Health Disparities, and personal fees from Rhythm Pharmaceuticals; RSG reports grants from US Department of Agriculture and NIH/NICHD; no other relationships or activities that could appear to have influenced the submitted work.

References

1. World Health Organisation. *Taking action on childhood obesity*. WHO: Geneva, Switzerland 2018.
2. World Health Organisation. *Report of the Commission on Ending Childhood Obesity*. WHO: Geneva, Switzerland 2016.
3. United Nations Children's Fund (UNICEF), World Health Organization, International Bank for Reconstruction and Development/The World Bank. *Levels and trends in child malnutrition: Key Findings of the 2020 Edition of the Joint Child Malnutrition Estimates*. Geneva: World Health Organization. 2020.
4. UNICEF. *The State of the World's Children 2019. Children, Food and Nutrition: Growing well in a changing world*. UNICEF, New York 2019.
5. Di Cesare M, Soric M, Bovet P, et al. The epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. *BMC Medicine*. 2019;17(1):212.
6. Guerrero AD, Mao C, Fuller B, Bridges M, Franke T, Kuo AA. Racial and Ethnic Disparities in Early Childhood Obesity: Growth Trajectories in Body Mass Index. *Journal of Racial and Ethnic Health Disparities*. 2016;3(1):129-137.
7. Kumanyika SK. Unraveling common threads in obesity risk among racial/ethnic minority and migrant populations. *Public Health*. Jul 2019;172:125-134.
8. Simmonds M, Llewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obesity Reviews*. 2016;17(2):95-107.
9. Geserick M, Vogel M, Gausche R, et al. Acceleration of BMI in Early Childhood and Risk of Sustained Obesity. *New England Journal of Medicine*. 2018;379(14):1303-1312.
10. Halfon N, Larson K, Slusser W. Associations Between Obesity and Comorbid Mental Health, Developmental, and Physical Health Conditions in a Nationally Representative Sample of US Children Aged 10 to 17. *Academic Pediatrics*. 2013;13(1):6-13.
11. Mihrshahi S, Gow ML, Baur LA. Contemporary approaches to the prevention and management of paediatric obesity: an Australian focus. *Med J Aust*. 2018;209(6).
12. Wolfenden L, Ezzati M, Larijani B, Dietz W. The challenge for global health systems in preventing and managing obesity. *Obesity Reviews*. 2019;20(S2):185-193.
13. Finkelstein EA, Graham WCK, Malhotra R. Lifetime Direct Medical Costs of Childhood Obesity. *Pediatrics*. 2014;133(5):854-862.
14. Redsell SA, Edmonds B, Swift JA, et al. Systematic review of randomised controlled trials of interventions that aim to reduce the risk, either directly or indirectly, of overweight and obesity in infancy and early childhood. *Matern Child Nutr*. 2016;12(1):24-38.
15. Hurley KM, Cross MB, Hughes SO. A systematic review of responsive feeding and child obesity in high-income countries. *The Journal of nutrition*. 2011;141(3):495-501.
16. Spill MK, Callahan EH, Shapiro MJ, et al. Caregiver feeding practices and child weight outcomes: a systematic review. *The American Journal of Clinical Nutrition*. 2019;109(Supplement_1):990S-1002S.
17. Porter R, Tindall A, Gaffka B, et al. A Review of Modifiable Risk Factors for Severe Obesity in Children Ages 5 and Under. *Childhood Obesity*. 2018;14(7):468-476.
18. Arenz S, Ruckerl R, Koletzko B, von Kries R. Breast-feeding and childhood obesity - a systematic review. *Int J Obes Relat Metab Disord*. 2004;28(10):1247-1256.
19. Lefebvre CM, John RM. The effect of breastfeeding on childhood overweight and obesity: A systematic review of the literature. *Journal of the American Association of Nurse Practitioners*. 2014;26(7):386-401.
20. Qiao J, Dai L-J, Zhang Q, Ouyang Y-Q. A Meta-Analysis of the Association Between Breastfeeding and Early Childhood Obesity. *Journal of Pediatric Nursing*. 2020;53:57-66.

21. Weng SF, Redsell SA, Swift JA, Yang M, Glazebrook CP. Systematic review and meta-analyses of risk factors for childhood overweight identifiable during infancy. *Archives of Disease in Childhood*. 2012;97(12):1019-1026.
22. Woo Baidal JA, Locks LM, Cheng ER, Blake-Lamb TL, Perkins ME, Taveras EM. Risk Factors for Childhood Obesity in the First 1,000 Days: A Systematic Review. *American journal of preventive medicine*. 2016/06/01/ 2016;50(6):761-779.
23. Yan J, Liu L, Zhu Y, Huang G, Wang PP. The association between breastfeeding and childhood obesity: a meta-analysis. *BMC Public Health*. 2014/12/13 2014;14(1):1267.
24. Vail B, Prentice P, Dunger DB, Hughes IA, Acerini CL, Ong KK. Age at Weaning and Infant Growth: Primary Analysis and Systematic Review. *The Journal of pediatrics*. 2015;167(2):317-324.e311.
25. Jiménez-Pavón D, Kelly J, Reilly JJ. Associations between objectively measured habitual physical activity and adiposity in children and adolescents: Systematic review. *International Journal of Pediatric Obesity*. 2010;5(1):3-18.
26. Pate RR, Hillman CH, Janz KF, et al. Physical Activity and Health in Children Younger than 6 Years: A Systematic Review. *Medicine and science in sports and exercise*. Jun 2019;51(6):1282-1291.
27. Timmons BW, Leblanc AG, Carson V, et al. Systematic review of physical activity and health in the early years (aged 0-4 years). *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2012;37(4):773-792.
28. Wen LM, Baur LA, Rissel C, Xu H, Simpson JM. Correlates of body mass index and overweight and obesity of children aged 2 years: Findings from the healthy beginnings trial. *Obesity*. 2014;22(7):1723-1730.
29. Cox R, Skouteris H, Rutherford L, Fuller-Tyszkiewicz M, Dell'Aquila D, Hardy LL. Television viewing, television content, food intake, physical activity and body mass index: a cross-sectional study of preschool children aged 2–6 years. *Health Promotion Journal of Australia*. 2012;23(1):58-62.
30. Jochem C, Schmid D, Leitzmann MF. Sedentary Behaviour and Adiposity. In: Leitzmann MF, Jochem C, Schmid D, eds. *Sedentary Behaviour Epidemiology*. Cham: Springer International Publishing; 2018:155-178.
31. Carter PJ, Taylor BJ, Williams SM, Taylor RW. Longitudinal analysis of sleep in relation to BMI and body fat in children: the FLAME study. *BMJ*. 2011;342:d2712.
32. Felső R, Lohner S, Hollódy K, Erhardt É, Molnár D. Relationship between sleep duration and childhood obesity: Systematic review including the potential underlying mechanisms. *Nutrition, Metabolism and Cardiovascular Diseases*. 2017;27(9):751-761.
33. Miller MA, Krusbrink M, Wallace J, Ji C, Cappuccio FP. Sleep duration and incidence of obesity in infants, children, and adolescents: a systematic review and meta-analysis of prospective studies. *Sleep*. 2018;41(4).
34. Morrissey B, Taveras E, Allender S, Strugnell C. Sleep and obesity among children: A systematic review of multiple sleep dimensions. *Pediatric Obesity*. 2020;15(4):e12619.
35. Sluggett L, Wagner SL, Harris RL. Sleep Duration and Obesity in Children and Adolescents. *Canadian Journal of Diabetes*. 2019;43(2):146-152.
36. Harskamp-van Ginkel MW, Chinapaw MJM, Harmsen IA, Anujoo KO, Daams JG, Vrijkotte TGM. Sleep during Infancy and Associations with Childhood Body Composition: A Systematic Review and Narrative Synthesis. *Childhood Obesity*. 2020/03/01 2019;16(2):94-116.
37. Ziauddeen N, Wilding S, Roderick PJ, et al. Predicting the risk of childhood overweight and obesity at 4–5 years using population-level pregnancy and early-life healthcare data. *BMC Medicine*. 2020/05/11 2020;18(1):105.
38. Askie L, Espinoza D, Martin A, et al. Interventions commenced by early infancy to prevent childhood obesity - the EPOCH Collaboration: an individual participant data prospective meta-

- analysis of four randomised controlled trials. *Pediatric Obesity*. 2020:e12618.
<https://doi.org/10.1111/ijpo.12618>.
39. Ash T, Agaronov A, Young TL, Aftosmes-Tobio A, Davison KK. Family-based childhood obesity prevention interventions: a systematic review and quantitative content analysis. *International Journal of Behavioral Nutrition and Physical Activity*. 2017;14(1):113.
 40. Brown T, Moore T, Hooper L, et al. Interventions for preventing obesity in children. *Cochrane Database of Systematic Reviews*. 2019(7).
<https://doi.org/10.1002/14651858.CD001871.pub4>.
 41. Narzisi K, Simons J. Interventions that prevent or reduce obesity in children from birth to five years of age: A systematic review. *Journal of Child Health Care*. 2020:1367493520917863.
 42. Blake-Lamb TL, Locks LM, Perkins ME, Baidal JAW, Cheng ER, Taveras EM. Interventions for childhood obesity in the first 1,000 days: a systematic review. *American journal of preventive medicine*. 2016;50(6):780-789.
 43. Bleich SN, Vercammen KA, Zatz LY, Frelier JM, Ebbeling CB, Peeters A. Interventions to prevent global childhood overweight and obesity: a systematic review. *Lancet Diabetes & Endocrinology*. 2018;6(4):332-346.
 44. Grobler L, Visser M, Siegfried N. Healthy Life Trajectories Initiative: Summary of the evidence base for pregnancy-related interventions to prevent overweight and obesity in children. *Obesity Reviews*. 2019;20(S1):18-30.
 45. Hennessy M, Heary C, Laws R, et al. The effectiveness of health professional-delivered interventions during the first 1000 days to prevent overweight/obesity in children: A systematic review. *Obesity Reviews*. 2019;20(12):1691-1707.
 46. Haire-Joshu D, Tabak R. Preventing Obesity Across Generations: Evidence for Early Life Intervention. *Annu Rev Public Health*. 2016;37:253-271.
 47. Lillycrop KA, Burdge GC. Epigenetic changes in early life and future risk of obesity. *International Journal of Obesity*. 2011;35(1):72-83.
 48. Brookes ST, Whitely E, Egger M, Smith GD, Mulheran PA, Peters TJ. Subgroup analyses in randomized trials: risks of subgroup-specific analyses; power and sample size for the interaction test. *J Clin Epidemiol*. 2004;57(3):229-236.
 49. Egger M, Smith GD. Meta-Analysis. Potentials and promise. *BMJ* 1997;315(7119):1371.
 50. Anna Lene Seidler, Kylie E Hunter, David Espinoza, Seema Miharshahi, Lisa Askie. Quantifying the advantages of conducting a prospective meta-analysis (PMA): a case study of early childhood obesity prevention. *PREPRINT (Version 1) 2020*:available at Research Square [<https://doi.org/10.21203/rs.21203.rs-36734/v21201+>].
 51. Tierney J, Stewart L, Clarke M. Chapter 26: Individual participant data. In: Higgins J, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.0*. Cochrane, 2019.
 52. Seidler A, Hunter K, Cheyne S, Ghersi D, Berlin J, Askie L. A guide to prospective meta-analysis. *BMJ*. 2019(367):l5342.
 53. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews*. 2015;4(1):1.
 54. Dodd JM, Grivell RM, Louise J, et al. The effects of dietary and lifestyle interventions among pregnant women who are overweight or obese on longer-term maternal and early childhood outcomes: protocol for an individual participant data (IPD) meta-analysis. *Systematic Reviews*;6(1):51.
 55. McKenzie JE, Brennan SE, Ryan RE, Thomson HJ, Johnston RV, Thomas J. Chapter 3: Defining the criteria for including studies and how they will be grouped for the synthesis. In: Higgins JPT, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020)*. Cochrane, Available from www.training.cochrane.org/handbook2020.

56. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:l4898.
57. Schünemann HJ, Higgins JPT, Vist GE, et al. Chapter 14: Completing ‘Summary of findings’ tables and grading the certainty of the evidence. In: Higgins JPT, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020)*. Cochrane, 2020. Available from www.training.cochrane.org/handbook2020.
58. Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology*. 2011/04/01/2011;64(4):383-394.
59. Riley RD, Debray TPA, Fisher D, et al. Individual participant data meta-analysis to examine interactions between treatment effect and participant-level covariates: Statistical recommendations for conduct and planning. *Statistics in Medicine*. 2020;39(15):2115-2137.
60. World Health Organization. The WHO Child Growth Standards. 2010. <http://www.who.int/childgrowth/en/>.
61. Burke DL, Ensor J, Riley RD. Meta-analysis using individual participant data: one-stage and two-stage approaches, and why they may differ. *Statistics in Medicine*. 2017;36(5):855-875.
62. Riley RD, Steyerberg EW. Meta-analysis of a binary outcome using individual participant data and aggregate data. *Res Synth Methods*. 2010;1(1):2-19.
63. Deeks JJ, Higgins JPT, Altman DG. Chapter 10: Analysing data and undertaking meta-analyses. In: Higgins JPT, Thomas J, Chandler J, et al., eds. *Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020)*2020:Available from www.training.cochrane.org/handbook.
64. R Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical Computing*. 2020: Vienna, Austria. URL <https://www.R-project.org/>.
65. Schulz KF, Grimes DA. Multiplicity in randomised trials II: subgroup and interim analyses. *Lancet*. May 7-13 2005;365(9471):1657-1661.
66. Stewart LA, Parmar MK. Meta-analysis of the literature or of individual patient data: is there a difference? *Lancet* 1993;341:418-422.
67. Tierney JF, Fisher DJ, Burdett S, Stewart LA, Parmar MKB. Comparison of aggregate and individual participant data approaches to meta-analysis of randomised trials: An observational study. *PLOS Medicine*. 2020;17(1):e1003019.