Identifying Risk Groups for SARS-CoV-2 Vaccination "By the Numbers"

Clement J. McDonald, MD[1], Seo H. Baik, PhD[1], Zhaonian Zheng, MS[1], Liz Amos, MLIS[1]

Author Affiliations: [1]Lister Hill National Center for Biomedical Communications, National Library of Medicine, U.S. National Institutes of Health, Bethesda, Maryland, USA

Address reprint request to Dr. Clement J. McDonald (corresponding author) at National Library of Medicine, U.S. National Institutes of Health, 8600 Rockville Pike, Building 38A, Room 07N710, Bethesda, Maryland 20894, USA

Abstract (WC 249)

Background

Given the limited supply of two COVID-19 vaccines, it will be important to choose which risk groups to prioritize for vaccination in order to get the most health benefits from that supply.

Method

In order to help decide how to get the maximum health yield from this limited supply, we implemented a logistic regression model to predict COVID-19 death risk by age, race and sex and did the same to predict COVID-19 case risk.

Results

Our predictive model ranked all demographic groups by COVID-19 death risk. It was highly concentrated by demographic group, e.g. 85+ year old Black, Non-Hispanic patients suffered 2,160 deaths per 100,000. The comparable risks did not reach the level of our 43 highest risk demographic groups (92% of all COVID 19 deaths), the risks of nursing home resident exceeded them at 5,200 deaths/100,000.

We saw minimal concentration of COVID-19 case risk by demographic groups compared to the high concentration of COVID-19 death risk we saw among demographic groups.

Conclusion

SARS-CoV-2 vaccines protect against severe COVID infection and thus against COVID-19 death per vaccine studies. Allocating at least some of the early vaccine supplies to high risk demographic groups could maximize lives saved. Our model predicts that allocating only 12 million vaccine courses to the 17 demographic groups at highest COVID-19 death risk (table 3 row 1-17 in our report) could eliminate 47% of COVID deaths with by investing only 3.7% the vaccine courses needed to protect all US residents -- a highly leveraged payoff.

Introduction

As of December 14, SARS-CoV-2 had infected more than 16 million persons and caused more than 298,000 deaths[1] in the U.S. It has infected and killed ten times the number in the 2018 flu epidemic (35.5 million cases and 34,200 deaths[2]) and nearly 100 times the number in the 1952 polio epidemic (57,879 cases and 3,145 deaths[3]).

Two COVID-19 vaccines with 95% efficacy and no serious adverse events[4] have Food and Drug Administrative (FDA) Emergency Use Authorization. However, vaccine supply is limited and may remain so for many months. Given the increasing number of cases and death, methods for prioritizing the risk groups who will receive the most benefit from the available supply are important. In this report, we propose a method to rank the vaccination benefit per risk group numerically.

Methods

Two data sets for defining COVID-19 risk based on demographic categories

The first is National Center for Health Statistics (NCHS)[5] which includes vetted data from death certificates about a selected set of deaths including COVID-19 deaths, coded to ICD-10 code U07.1 This information is reported by age range, race-ethnicity and sex. It includes no individual level data. We considered only COVID-19 deaths from this database, dated November 25, 2020. We only had COVID-19 data from April 1 (when the COVID-19 ICD code was created) through October 30, 2020, the last available data. Age ranges begin at 0-4 years and increase in 10-year increments until age 84, beyond which all cases are lumped in one, 85+ group. Cells containing fewer than 10 individuals are flagged as

such and delivered with no content. This data came grouped by age/race/sex. We used this data set for our analyses about COVID-19 death.

Our second data set comes from the Centers for Disease Control and Prevention (CDC)'s COVID-19 Surveillance Public Use Data[6], updated December 4, 2020 with data up to November 19, 2020. It carries patient level data including age group, race, ethnicity, sex, hospitalization death status (and other variables). Ages are binned differently from the NCHS data. Age ranges start at 0-9 and step in 10-year intervals until 70-79; above which all records are classed as 80+ years. We grouped the individual records from this data set into combinations of age range, sex and combined race-ethnic categories. We used this data for our case rate analyses.

In both data sets, race-ethnicity is combined into five terms using categories defined by the 2019 NCHS Bridged-Race Population Estimates[7]: American Indian/Alaska Native, Non-Hispanic; Asian, Non-Hispanic; Black, Non-Hispanic; Native Hawaiian/Other Pacific Island, Non-Hispanic, Multiple/Other, Non-Hispanic; and Hispanic/Latino. To simplify, the tables and discussion we use "Black" to mean Black, Non-Hispanic", Hispanic to mean Hispanic/Latino; "American Indian" to mean American Indian/Alaska Native, Non-Hispanic, "Asian" to mean: Asian, Non-Hispanic, and "White" to mean White, Non-Hispanic. We will use "race" to mean the combined race-ethnicity terms.

We obtained estimated population counts from the 2019 NCHS Bridged-Race Population Estimates[7] for each demographic category in each of the data sets, as approximates for 2020 counts, and used them to compute death and case rates per 100,000 U.S. inhabitants for each demographic category. We only used de-identified and publicly available data in this study.

Data about special risk groups

We obtained information about nursing home cases and deaths from the Centers for Medicare & Medicaid Service (CMS).[8] It reported COVID-19 cases and deaths per nursing home, without patient level data or demographics breakdown. The total number of nursing home residents (1.2 million) came from Kaiser Family Foundation's analysis.[9]

We obtained information about health care workers from a number of incomplete sources.[10,11] For example, one source with information on 11.9 million COVID-19 cases lacked health care worker status for 80%.[11]

Statistical methods

We generated cross-tabulations[12] by race, age, and sex for both data sets, the first with death as the outcome, the second with the occurrence of a COVID-19 case as the outcome and reported both. We also ran a logistic regression analysis[13,14] on the first data set with death as the outcome, and the second with the occurrence of COVID-19 cases as outcome. We ran our case analysis on the CDC data records with complete age/sex/race data and a secondary analysis on a larger set with complete sex/age data. Logistic regression produces an equation based on characteristics (variables) in a set of cases with known outcomes. The equation will then calculate the probability of that outcome when fed the same variables from a new case. Logistic regression it has the advantage that it can be used on grouped data or on individual level data, can accommodate continuous as well as categorical variables, and provides confidence limits on its estimates. It was necessary to include all pair wise interactions between age, sex and race to obtain stable statistic in these analyses.

The input to for these analyses were tables with a row for each demographic category e.g. age >80, sex = male, race = Black. Each row also carries a count of the number of individuals within category who suffered the outcome (a COVID-19 death or infection) and a count of the number of U.S. inhabitants within that category from NCHS Bridged-Race Population.[7]

In their phase II trials, SARS-CoV-2 vaccine prevented severe cases and thus death. We assume vaccination of a given risk group will prevent 100% of the COVID-19 death predicted for that group. For simplicity sake, we also assume 100% of effectiveness against cases, a slight exaggeration over their reported 95% effectiveness.

Because our goal is to develop a prioritization algorithm to save the maximum number of lives, we counted all deaths equally. Other weightings, such as life years lost are possible, but the WHO SAGE report[15] argues for use of a simple unweighted count, as we did. [Statistical analyses performed by SH and ZZ].

RESULTS

Break down of death rates

The NCHS data set dated November 25, 2020[5] carried results for 220,582 COVID-19 deaths occurring between April 1 and October 30. No age or race data was missing but 2,234 (1.01%) of the groups specified "other" as the race. We excluded these records leaving 218,348 for our analysis shown in Table

1. Note the empty cells in the age range up to range 24-35 had values from 0-9 but they were hidden in this public data set. We show these as "<10" in our tables.

Ignoring age and sex, overall death rates were highest at 99 deaths/100,000 for Black, and they decrease in order to 82 for American Indian, 75, for Hispanic, 59 for White and 41 for Asian per 100,000. Overall, deaths rated climbed almost geometrically with age (see Table 1), so much so, that seventy percent of all COVID-19 deaths occurred in individual over age 70. The highest COVID-19 death rates occurred in males over age 85. For Black, it was 1,953, Hispanic 1,886, American Indian 1,120, White 1,029 and Asians 925 deaths per 100,000. In every age/race category, the male COVID-19 death rate exceeded that of the females.

We used a more recent set of death records from weeks 44 through 47 (10/21/2020 to11/21/2020) which did not yet have race information, to assess the death rate during the current surge in COVID-19 cases.[16] (Supplement Table 1) During four weeks of that surge, COVID-19 killed 22,000 elderly older than 74, highlighting need to urgency to save lives in this age group.

Breakdown of case rates

The CDC case surveillance data set[6] contained 8,242,333 COVID-19 cases from April 1 when ICD code for COVID-19 became available, until October 30, the date of the most recent data. Age and sex data were missing in trivial amounts (0.14%) and (1.17%). However, definitive race status was absent from 43.6% of these. We used the 4,646,942 records with complete sex age and race data to report case rates, broken down by age sex and race (Table 2).

Ignoring age and sex, American Indian had the highest case rates at 2,133 per 100,000. The rate for

Hispanic, Black, White and Asian followed in decreasing order of 1935, 1655, 1267 and 744, per 100,000,

respectively. The Asian case rate was one third rate of the highest case rate.

Among Blacks, the case rates peak in the 80+ age range, among Hispanics in the 40-49 age range, and

among Whites and Asians, in the 20-29-year age group. American Indian case rates peaked in the 30-39

age range to the highest level of any age/race category.

In Blacks, the female case rates exceeded the male rate until age sixty after which the relationship

reverses. Among Hispanics and Whites, the transition from more females to more males occurs at age

50, among American Indians at age 80. Ignoring race, the highest case prevalence of any age/sex

category (2,167/100, 000) was in 20-29-year-old women, a rate almost 20% greater than the next

closest age/sex category.

The difference in the COVID-19 case prevalence in all 10 pairwise comparisons between racial categories

was significant at $p < 0.001$ after Bonferroni's correction for multiple comparisons.

Logistic regression analysis

Table 3 presents the logistic regression estimate for likelihood of a COVID-19 death for each

demographic group sorted by risk, with 85+ year old black men at the highest risk (see row 1). The first

five columns show the NCHS data regarding each risk category. Columns A-C carry the values of race,

age and sex that define the category. Column D carries the total number of people in each category, and

column E, the number of COVID-19 deaths in that category. The next three columns display the logistic

regression's estimated risk of death (Column E) and the upper and lower confidence intervals (Columns F and G) for that estimate. We expressed these risk in terms of number per 100,000, rather than pure probabilities, to fit public health reporting conventions. In a given row, Columns I and J are cumulative sums of all of the deaths and all of the people in the rows at or above that row. One can think of the cumulative sum of people (Column J) to be the number of people to vaccinate to prevent the number of deaths in Column I.

For a given vaccine supply one would read down to the first row Table 3 to find the first row that had account in Column J that that equaled or exceeded the vaccine supply, and vaccinate all risk groups at or above that row and possibly some people in the next category. For example, if one had a supply for 12 million people one could vaccinate all risk groups in from row 1 to 17 of table 3, and few people in row 18 and thereby cover 3.7% of the U.S. population to prevent 47% if all COVID deaths -- a highly leveraged use of the supply.

Table 4 shows the results of an analogous logistic analysis of cases (infections) as the outcome. The table shows demographic categories ranked from highest to lowest case risk. Our primary analysis was on 4,646,942 records with complete age/sex/race data.

The highest risk occurs among 30-39-year old female American Indians. Most of the top risk categories were American Indian or Hispanic females of different age ranges. If we dedicated the a vaccine supply of 14 million to individual with the highest infection risk rather than the highest death risk we would vaccinate all of the demographic categories in Table 4 from rows 1 to row 7, which would include 5.7% of the total population but would only prevent 11% of the COVID-19 cases. The results for our analysis of the 8,120,644, with complete age and sex data were similar. Vaccinating 6.7% of the individuals with

highest case risk would prevent only 10.4% of case. In both cases the payoff for vaccinating based on case risk was a fifth of the payoff for vaccinating based on death rates.

Cases and deaths among special populations

Any allocation rule for SARS-CoV-2 vaccine will want to consider the infection and death rates in some special population for prioritization. Among nursing home residents, the cumulative number of deaths per 100,000 is staggering. (See row 1, Table 5).

Data collected from the Department of Corrections websites are principally represented by prisons, with exception of a few states that include jails.[17] For prisoners, the death rate was modest at the average of 55-65-year-old men and would not reach a priority until vaccine supplies were enough to cover 78 million people (see open arrow at row 45, Table 3).

Data about COVID-19 deaths cases is limited. However, we only have data for total healthcare workers (HCWs) (14 to 20 million). Based on the data about HCW deaths, their risk would be 10/100,000 deaths (table 5) and would have a have priority would be less than about 60% of our demographic groups ranked by death risk (Table 3, Row 59). However, other factors apply to vaccination decisions about HCWs.

DISCUSSION

We created two simple statistical models using logistic regression, one for predicting COVID-19 deaths and another for predicting COVID-19 cases. Babus, Das, and Lee[18] report a more complicated model but only considered infections (cases) as outcome.

We used NCHS's vetted COVID death records using only age ranges, race and sex as predictors of COVID-19 deaths. Our model distinguished death risk by demographic categories quite over death risk across a two orders of magnitude range. It would probably produce more accurate predictions and more granular categories if more detailed data were available. For example, integer ages, (which are available to some agencies), rather than 10-year ranges, would produce more refined predictions. Including other variable such as geographic location case rates would also likely improve predictions. New York State plans to weighing location risk by simply prioritizing high risk locations over low ones.[19] Location risk could be included formally in a logistic model to improve its accuracy, if demographic breakdowns and coved death case rates were known for those locations.

Demographic breakdowns were not available for special risk groups, so we obtained simple overall risk figures (Table 5). The risk of COVID-19 deaths among 1.3 million nursing home residents[20] is extreme at 5,200 per 100,000;[8] this is more than twice of risk of even 85+ year old black males and puts them in front of the line in most vaccination plans. The death risks for prison residents and HCWs are below that of the 78 million people in the 41 highest risk demographics.

Front line HCWs are widely recognized to deserve special priority because they bravely enter the COVID-19 hot zone every day. However, many of the nominal 14-20 million HCWs, are not front line. The risk of administrators and pharmacist preparing IV's in the basement is exponentially less than a nurse anesthetist or emergency room physician intubating COVID-19 patients. It will be important to

distinguish carefully between HCW with, and those without, direct COVID-19 patient contact to preserve some early vaccine supply for the high-risk elderly.

Some argue for allocating vaccine supply by occupation class e.g. essential workers, or teachers with the goal of protecting education, food supplies, etc. and reducing the case rate.[19] New York State's vaccination program places their senior citizens (age 65 and above) in phase 3 *behind* teachers, grocery store, pharmacists, transit workers, and others who will be vaccinated in in phase 2.[21]

Our data indicated that small early supplies of vaccines will not be enough to vaccinate large occupational groups, or to have an important effect on case rates. On the other hand, small early supplies could have huge effect on COVID-19 death rates among high risk elderly because COVID-19 deaths are so concentrated in these relatively small demographic categories. For example, our model would allocate 12 million vaccine regimens (2 per person) allocated by risk numbers would cover everybody over age 85, all Blacks and Hispanics over 75 and all Black males over 65, and could prevent 47% of COVID deaths. Allocation by risk number also delivers vaccines equitably by race. Sixty-five percent of 12 million regimens assumed by demographic risk, would go to racial minorities. Finally, COVID-19 is presently is killing 75+ year old seniors at rate of 20,000 lives per month. Our model predicts opportunities to save many of these lives through COVID vaccination, if delivered by the risk numbers and in a timely fashion. After analyzing their numbers, Great Britain has committed their first tranches of vaccines of its vaccine supply to their 80+ year old elders.[22]

Acknowledgements: This research was supported by the Intramural Research Program of the National Library of Medicine, National Institutes of Health.

REFERENCES

1.   CDC COVID Data Tracker. 2020. (Accessed December 14, 2020, at https://covid.cdc.gov/covid-data-tracker/#cases_casesper100klast7days.)

2.   Estimated Influenza Illness, Medical visits, Hospitalizations, and Deaths in the United States - 2018-2019 influenza season. 2020. (Accessed December 10, 2020, at https://www.cdc.gov/flu/about/burden/2018-2019.html#:~:text=CDC%20estimates%20that%20influenza%20was,the%202018%E2%80%932019%20influenza%20season.)

3.   Polio. 2017. (Accessed December 10, 2020, at https://ourworldindata.org/polio.)

4.   Jackson LA, Anderson EJ, Rouphael NG, et al. An mRNA Vaccine against SARS-CoV-2 — Preliminary Report. New England Journal of Medicine 2020;383:1920-31.

5.   Monthly provisional counts of death by age group, sex, and race/ethnicity for select cause of death. 2020. (Accessed December 10, 2020, at https://data.cdc.gov/NCHS/Monthly-provisional-counts-of-deaths-by-age-group-/65mz-jvh5.)

6.   COVID-19 Case Surveillance Public Use Data. 2020. (Accessed December 10, 2020, at https://data.cdc.gov/Case-Surveillance/COVID-19-Case-Surveillance-Public-Use-Data/vbim-akqf.)

7.   Vintage 2019 postcensal estimates of the resident population of the United States (April 1, 2010, July 1, 2010-July 1, 2019), by year, county, single-year of age (0, 1, 2, .., 85 years and over), bridged race, Hispanic origin, and sex. Prepared under a collaborative arrangement with the U.S. Census Bureau. . 2020. (Accessed December 10, 2020, at https://www.cdc.gov/nchs/nvss/bridged_race.htm.)

8.   Nursing Home COVID-19 Public File. 2020. (Accessed December 10, 2020, at https://data.cms.gov/stories/s/COVID-19-Nursing-Home-Data/bkwz-xpvg.)

9.   Total Number of Residents in Certified Nursing Facilites. 2019. (Accessed December 10, 2020, at

https://www.kff.org/other/state-indicator/number-of-nursing-facility-

residents/?currentTimeframe=0&selectedRows=%7B%22wrapups%22:%7B%22united-

states%22:%7B%7D%7D%7D&sortModel=%7B%22colId%22:%22Number%20of%20Nursing%20Facility%

20Residents%22,%22sort%22:%22asc%22%7D.)

10.   National Nurses United. Sins of Omission: How Government Failtures to Track COVID-19 Data

Have Led to More Than 1,700 Health Care Worker Deaths and Jeopardize Public Health2020.

11.   Cases & Deaths Among Healthcare Personnel. 2020. (Accessed December 14, 2020, at

https://covid.cdc.gov/covid-data-tracker/#health-care-personnel.)

12.   SAS Version 9.4 PROC FREQ. 2020.

13.   SAS Version 9.4 PROC LOGISTIC. 2020.

14.   Agresti A, Kateri M. Categorical Data Analysis. International Encyclopedia of Statistical Science

Berlin, Heidelberg: Springer Berlin Heidelberg; 2011:206-8.

15.   World Health Organization [WHO]. WHO SAGE values framework for the allocation and

prioritization of COVID-19 vaccination2020 September 14, 2020.

16.   Provisional COVID-19 Death Counts by Sex, Age, and Week. 2020. (Accessed December 14, 2020,

at https://data.cdc.gov/NCHS/Provisional-COVID-19-Death-Counts-by-Sex-Age-and-W/vsak-wrfu.)

17.   UCLA Law COVID-19 Behind Bars Data Project,. 2020. (Accessed December 10, 2020, at

https://law.ucla.edu/centers/criminal-justice/criminal-justice-program/related-programs/covid-19-

behind-bars-data-project/.)

18.   Babus A, Das S, Lee S. The Optimal Allocation of Covid-19 Vaccines. medRxiv

2020:2020.07.22.20160143.

19.   New York State Department of Health [NYSDOH]. New York State's COVID-19 Vaccination

Program2020 October 2020.

20. National Center for Health Statistics [NCHS]. Long-term Care Providers and Services Users in the United States 2015-20162019.

21. Goodnough A, Hoffman J. The Elderly vs. Essential Workers: Who Should Get the Coronavirus Vaccine First? The New York Times 2020 December 5, 2020.

22. Department of Health & Social Care. Priority groups for coronavirus (COVID-19) vaccination: advice from the JCVI, 2 December 20202020 December 3, 2020.

23. Mass Incarceration: The Whole Pie 2020. Prison Policy Initiative, 2020. (Accessed December 10, 2020, at

https://www.prisonpolicy.org/reports/pie2020.html#:~:text=The%20American%20criminal%20justice%20system,civil%20commitment%20centers%2C%20state%20psychiatric-accessed.)

24. Total Health Care Employment. 2018. (Accessed December 10, 2020, at https://www.kff.org/other/state-indicator/total-health-care-employment/?currentTimeframe=0&sortModel=%7B%22colId%22:%22Location%22,%22sort%22:%22asc%22%7D.)

25. Health Care Still Largest U.S. Employer. Census Bureau,, 2020. (Accessed December 10, 2020, at https://www.census.gov/library/stories/2020/10/health-care-still-largest-united-states-employer.html.)

Table 1: COVID-19 death rates/100,000 from the NCHS COVID-19 data set broken down by age, sex and race

Age	Sex	Black # Deaths /100,000	Black # in category	Hispanic # Deaths /100,000	Hispanic # in category	American Indian # Deaths /100,000	American Indian # in category	Asian # Deaths /100,000	Asian # in category	White # Deaths /100,000	White # in category	All # Deaths /100,000	All # in category
	All	<10	2.99 M	<10	5.09 M	<10	0.19 M	<10	1.13 M	<10	10.16 M	<10	19.58 M
0-4	Female	<10	1.47 M	<10	2.50 M	<10	0.10 M	<10	0.55 M	<10	4.95 M	<10	9.57 M
	Male	<10	1.52 M	<10	2.60 M	<10	0.10 M	<10	0.58 M	<10	5.21 M	<10	10.01 M
	All	<10	6.23 M	<10	10.57 M	<10	0.41 M	<10	2.42 M	<10	21.36 M	<10	40.99 M
5-14	Female	<10	3.07 M	<10	5.18 M	<10	0.20 M	<10	1.19 M	<10	10.42 M	<10	20.05 M
	Male	<10	3.16 M	<10	5.39 M	<10	0.21 M	<10	1.24 M	<10	10.95 M	<10	20.94 M
	All	1.2	6.42 M	1.3	9.88 M	<10	0.41 M	<10	2.67 M	<10	23.30 M	0.6	42.69 M
15-24	Female	1.1	3.16 M	0.7	4.83 M	<10	0.20 M	<10	1.33 M	<10	11.35 M	<10	20.88 M
	Male	1.2	3.25 M	2.0	5.05 M	<10	0.21 M	<10	1.34 M	<10	11.95 M	0.8	21.81 M
	All	6.6	6.81 M	7.5	9.53 M	8.7	0.43 M	<10	3.54 M	1.1	25.65 M	3.3	45.94 M
25-34	Female	5.3	3.42 M	4.7	4.55 M	5.7	0.21 M	<10	1.80 M	0.8	12.60 M	2.3	22.58 M
	Male	7.9	3.38 M	10.1	4.98 M	11.6	0.21 M	0.8	1.74 M	1.5	13.04 M	4.3	23.36 M
	All	21.0	5.48 M	24.6	8.66 M	29.3	0.34 M	3.2	3.29 M	2.8	23.90 M	10.0	41.66 M
35-44	Female	17.8	2.88 M	12.9	4.20 M	20.4	0.17 M	0.7	1.74 M	2.0	11.88 M	6.4	20.87 M
	Male	24.5	2.60 M	35.6	4.46 M	38.5	0.17 M	6.0	1.55 M	3.6	12.01 M	13.5	20.79 M
45-54	All	57.0	5.21 M	70.0	7.13 M	86.7	0.32 M	15.0	2.80 M	10.1	25.41 M	27.5	40.87 M

	Female	42.3	2.77 M	36.1	3.54 M	55.7	0.17 M	7.6	1.49 M	7.4	12.73 M	17.4	20.70 M
	Male	73.7	2.44 M	103.4	3.59 M	119.7	0.15 M	23.5	1.31 M	12.8	12.68 M	37.8	20.17 M
	All	149.0	5.04 M	177.3	5.08 M	148.0	0.33 M	50.2	2.28 M	31.1	29.73 M	64.5	42.45 M
55-64	Female	115.4	2.73 M	103.7	2.60 M	119.0	0.18 M	25.1	1.24 M	22.4	15.21 M	44.5	21.95 M
	Male	188.8	2.31 M	254.4	2.48 M	180.8	0.16 M	80.0	1.04 M	40.2	14.52 M	85.9	20.50 M
	All	367.3	3.15 M	395.3	2.82 M	283.5	0.21 M	126.0	1.58 M	91.9	23.72 M	149.6	31.48 M
65-74	Female	269.1	1.80 M	258.9	1.53 M	240.1	0.11 M	78.0	0.88 M	69.0	12.46 M	109.4	16.78 M
	Male	498.1	1.35 M	556.7	1.29 M	334.2	0.10 M	186.9	0.70 M	117.1	11.26 M	195.6	14.70 M
	All	784.8	1.39 M	756.0	1.31 M	540.0	0.09 M	292.6	0.75 M	281.5	12.44 M	366.1	15.97 M
75-84	Female	619.6	0.85 M	557.4	0.76 M	469.2	0.05 M	203.1	0.42 M	227.9	6.89 M	293.2	8.97 M
	Male	1,046.0	0.54 M	1,033.1	0.55 M	630.2	0.04 M	407.4	0.33 M	348.0	5.55 M	459.5	7.00 M
	All	1,609.2	0.52 M	1,495.0	0.51 M	927.0	0.03 M	841.8	0.30 M	943.4	5.24 M	1,033.5	6.60 M
85+	Female	1,458.1	0.36 M	1,271.3	0.32 M	813.7	0.02 M	792.0	0.19 M	894.3	3.33 M	966.3	4.23 M
	Male	1,953.3	0.16 M	1,885.9	0.19 M	1,120.4	0.01 M	924.8	0.11 M	1,029.0	1.91 M	1,152.9	2.38 M
All		99.4	43.24 M	75.3	60.57 M	82.2	2.77 M	40.6	20.76 M	59.3	200.91 M	66.5	328.24 M

Table 2 COVID-19 Case rates/100,000 from CDC COVID-19 public surveillance data broken down by age, sex and race

Age	Sex	Black # Cases /100,000	Black # in category	Hispanic # Cases /100,000	Hispanic # in category	American Indian # Cases /100,000	American Indian # in category	Asian # Cases /100,000	Asian # in category	White # Cases /100,000	White # in category	All # Cases /100,000	All # in category
	All	458	6.10 M	618	10.00 M	898	0.40 M	273	2.30 M	341	20.60 M	432	39.80 M
0 - 9	Female	457	3.00 M	614	5.10 M	934	0.20 M	259	1.10 M	338	10.10 M	429	19.40 M
	Male	459	3.10 M	621	5.30 M	864	0.20 M	286	1.20 M	343	10.60 M	435	20.30 M
	All	938	6.30 M	1,285	10.40 M	1,738	0.40 M	524	2.50 M	1,161	22.30 M	1,126	41.90 M
10 - 19	Female	983	3.10 M	1,352	5.10 M	1,841	0.20 M	526	1.20 M	1,269	10.90 M	1,207	20.50 M
	Male	893	3.20 M	1,220	5.30 M	1,637	0.20 M	522	1.30 M	1,059	11.40 M	1,048	21.40 M
	All	1,871	6.90 M	2,474	9.80 M	2,505	0.40 M	1,006	3.10 M	1,890	24.80 M	1,958	45.10 M
20 - 29	Female	2,177	3.40 M	2,684	4.70 M	2,814	0.20 M	1,028	1.60 M	2,099	12.10 M	2,167	22.10 M
	Male	1,572	3.50 M	2,277	5.10 M	2,203	0.20 M	984	1.60 M	1,692	12.70 M	1,758	23.10 M
	All	2,097	6.00 M	2,504	9.10 M	2,895	0.40 M	843	3.50 M	1,371	25.10 M	1,674	44.20 M
30 - 39	Female	2,334	3.10 M	2,647	4.30 M	3,141	0.20 M	834	1.80 M	1,477	12.50 M	1,790	21.90 M
	Male	1,844	2.90 M	2,373	4.80 M	2,647	0.20 M	852	1.70 M	1,268	12.70 M	1,559	22.20 M
	All	2,098	5.20 M	2,667	8.00 M	2,789	0.30 M	799	3.10 M	1,399	23.70 M	1,706	40.30 M
40 - 49	Female	2,261	2.80 M	2,750	3.90 M	2,912	0.20 M	802	1.60 M	1,495	11.80 M	1,798	20.30 M
	Male	1,914	2.50 M	2,586	4.00 M	2,660	0.20 M	795	1.40 M	1,304	11.90 M	1,612	20.00 M
50 - 59	All	2,087	5.20 M	2,517	6.10 M	2,386	0.30 M	870	2.50 M	1,345	28.10 M	1,588	42.40 M

Age	Sex												
	Female	2,160	2.80 M	2,521	3.10 M	2,471	0.20 M	850	1.30 M	1,372	14.20 M	1,615	21.60 M
	Male	2,004	2.40 M	2,514	3.10 M	2,292	0.20 M	892	1.20 M	1,317	13.90 M	1,559	20.70 M
	All	1,948	4.20 M	2,083	3.90 M	1,978	0.28 M	758	2.00 M	1,109	27.70 M	1,289	38.00 M
60 - 69	Female	1,881	2.30 M	1,959	2.00 M	2,011	0.15 M	706	1.10 M	1,070	14.30 M	1,243	20.00 M
	Male	2,032	1.90 M	2,222	1.80 M	1,939	0.13 M	823	0.90 M	1,151	13.30 M	1,341	18.10 M
	All	2,014	2.10 M	1,819	2.00 M	1,842	0.15 M	636	1.10 M	1,116	18.20 M	1,237	23.70 M
70 - 79	Female	1,838	2.30 M	1,633	1.10 M	1,916	0.08 M	573	0.60 M	1,057	9.80 M	1,165	12.90 M
	Male	2,264	0.90 M	2,056	0.90 M	1,753	0.07 M	715	0.50 M	1,184	8.50 M	1,324	10.80 M
	All	2,622	1.10 M	2,053	1.00 M	2,115	0.06 M	900	0.60 M	1,782	10.10 M	1,834	12.90 M
80+	Female	2,495	0.70 M	1,953	0.60 M	2,179	0.04 M	881	0.40 M	1,831	6.10 M	1,858	7.90 M
	Male	2,869	0.40 M	2,214	0.40 M	2,017	0.03 M	928	0.20 M	1,708	4.00 M	1,795	5.10 M
All		1,655	43.30 M	1,935	60.60 M	2,133	2.80 M	744	20.80 M	1,267	200.90 M	1,416	328.20 M

Table 3: Logistic regression on NCHS COVID death data showing the risk of COVID-19 death for each age/race/sex risk group sorted in order by death risk/100,000 US inhabitants

	A	B	C	D	E	F	G	H	I	J
						Death risk	Risk CI		Cum. # of	Cum. # in
	Race	Age	Sex	# Death	# in Population	/100,000	Lower	Risk CI Upper	deaths	population
1	Black	85+	Male	3,096	0.16 M	2,160.7	2,114.3	2,208.2	3,096	0.16 M
2	Hispanic	85+	Male	3,495	0.19 M	1,951.6	1,907.9	1,996.2	6,591	0.34 M
3	Black	85+	Female	5,261	0.36 M	1,367.0	1,337.8	1,396.8	11,852	0.70 M
4	Hispanic	85+	Female	4,116	0.32 M	1,233.7	1,206.0	1,262.0	15,968	1.03 M
5	White	85+	Male	19,625	1.91 M	1,233.3	1,221.3	1,245.4	35,593	2.94 M
6	American Indian	85+	Male	125	0.01 M	1,208.7	1,075.9	1,357.8	35,718	2.95 M
7	Asian	85+	Male	1,058	0.11 M	1,094.8	1,053.3	1,138.0	36,776	3.06 M
8	Black	75-84	Male	5,624	0.54 M	1,014.9	995.6	1,034.5	42,400	3.60 M
9	Hispanic	75-84	Male	5,641	0.55 M	963.9	944.8	983.3	48,041	4.14 M
10	White	85+	Female	29,820	3.33 M	777.5	770.1	785.1	77,861	7.48 M
11	American Indian	85+	Female	155	0.02 M	762.0	677.9	856.4	78,016	7.50 M
12	Asian	85+	Female	1,509	0.19 M	689.9	663.6	717.2	79,525	7.69 M
13	American Indian	75-84	Male	250	0.04 M	681.6	623.8	744.7	79,775	7.73 M
14	Black	75-84	Female	5,267	0.85 M	639.3	627.1	651.7	85,042	8.58 M
15	Hispanic	75-84	Female	4,246	0.76 M	607.0	594.9	619.4	89,288	9.34 M
16	Hispanic	65-74	Male	7,195	1.29 M	494.9	485.7	504.3	96,483	10.63 M
17	Black	65-74	Male	6,728	1.35 M	466.2	457.6	474.9	103,211	11.98 M
18	American Indian	75-84	Female	237	0.05 M	428.8	392.4	468.7	103,448	12.03 M

19	Asian	75-84	Male	1,334	0.33 M	369.8	354.6	385.7	104,782	12.36 M
20	American Indian	65-74	Male	327	0.10 M	354.5	327.3	383.9	105,109	12.46 M
21	White	75-84	Male	19,305	5.55 M	354.5	350.5	358.4	124,414	18.01 M
22	Hispanic	65-74	Female	3,961	1.53 M	311.2	305.2	317.2	128,375	19.54 M
23	Black	65-74	Female	4,841	1.80 M	293.1	287.6	298.6	133,216	21.34 M
24	Asian	75-84	Female	853	0.42 M	232.4	222.8	242.4	134,069	21.76 M
25	American Indian	65-74	Female	275	0.11 M	222.8	205.6	241.3	134,344	21.87 M
26	White	75-84	Female	15,703	6.89 M	222.7	220.2	225.3	150,047	28.76 M
27	Hispanic	55-64	Male	6,305	2.48 M	219.0	214.4	223.6	156,352	31.24 M
28	Black	55-64	Male	4,353	2.31 M	186.7	182.5	191.0	160,705	33.54 M
29	American Indian	55-64	Male	281	0.16 M	184.4	168.8	201.5	160,986	33.70 M
30	Asian	65-74	Male	1,305	0.70 M	159.1	152.3	166.3	162,291	34.40 M
31	Hispanic	55-64	Female	2,695	2.60 M	137.5	134.6	140.5	164,986	36.99 M
32	Black	55-64	Female	3,152	2.73 M	117.2	114.6	120.0	168,138	39.73 M
33	American Indian	55-64	Female	209	0.18 M	115.8	106.0	126.5	168,347	39.90 M
34	White	65-74	Male	13,191	11.26 M	114.2	112.6	115.8	181,538	51.16 M
35	American Indian	45-54	Male	185	0.15 M	107.3	95.4	120.7	181,723	51.32 M
36	Asian	65-74	Female	690	0.88 M	99.9	95.6	104.4	182,413	52.20 M
37	Hispanic	45-54	Male	3,714	3.59 M	85.9	83.5	88.3	186,127	55.79 M
38	White	65-74	Female	8,596	12.46 M	71.7	70.7	72.7	194,723	68.25 M
39	Black	45-54	Male	1,796	2.44 M	71.0	68.5	73.6	196,519	70.69 M
40	American Indian	45-54	Female	92	0.17 M	67.4	59.9	75.8	196,611	70.85 M
41	Asian	55-64	Male	831	1.04 M	62.9	59.4	66.7	197,442	71.89 M

42	Hispanic	45-54	Female	1,278	3.54 M	53.9	52.4	55.5	198,720	75.43 M
43	Black	45-54	Female	1,173	2.77 M	44.6	43.0	46.2	199,893	78.20 M
44	Asian	55-64	Female	311	1.24 M	39.5	37.3	41.9	200,204	79.44 M
45	White	55-64	Male	5,837	14.52 M	38.4	37.6	39.2	206,041	93.96 M
46	American Indian	35-44	Male	64	0.17 M	36.1	29.7	44.0	206,105	94.13 M
47	Hispanic	35-44	Male	1,587	4.46 M	30.0	28.8	31.3	207,692	98.59 M
48	Black	35-44	Male	638	2.60 M	26.1	24.6	27.7	208,330	101.19 M
49	White	55-64	Female	3,402	15.21 M	24.1	23.6	24.6	211,732	116.40 M
50	American Indian	35-44	Female	35	0.17 M	22.7	18.6	27.6	211,767	116.57 M
51	Hispanic	35-44	Female	542	4.20 M	18.8	18.0	19.7	212,309	120.76 M
52	Asian	45-54	Male	308	1.31 M	18.7	17.0	20.6	212,617	122.07 M
53	Black	35-44	Female	513	2.88 M	16.4	15.5	17.4	213,130	124.95 M
54	White	45-54	Male	1,629	12.68 M	12.5	12.0	13.0	214,759	137.63 M
55	Asian	45-54	Female	113	1.49 M	11.8	10.7	12.9	214,872	139.12 M
56	American Indian	25-34	Male	25	0.21 M	10.7	7.7	14.7	214,897	139.34 M
57	Hispanic	25-34	Male	501	4.98 M	9.1	8.5	9.8	215,398	144.32 M
58	Black	25-34	Male	269	3.38 M	8.2	7.5	9.0	215,667	147.70 M
59	White	45-54	Female	947	12.73 M	7.8	7.5	8.1	216,614	160.43 M
60	American Indian	25-34	Female	12	0.21 M	6.7	4.8	9.2	216,626	160.65 M
61	Hispanic	25-34	Female	214	4.55 M	5.7	5.3	6.2	216,840	165.19 M
62	Black	25-34	Female	183	3.42 M	5.1	4.7	5.6	217,023	168.61 M
63	Asian	35-44	Male	93	1.55 M	4.0	3.3	4.8	217,116	170.16 M
64	White	35-44	Male	433	12.01 M	3.5	3.2	3.7	217,549	182.18 M

65	Asian	35-44	Female	12	1.74 M	2.5	2.1	3.0	217,561	183.92 M
66	White	35-44	Female	242	11.88 M	2.2	2.0	2.3	217,803	195.80 M
67	Hispanic	15-24	Male	99	5.05 M	1.6	1.4	1.9	217,902	200.85 M
68	Black	15-24	Male	40	3.25 M	1.4	1.1	1.8	217,942	204.10 M
69	White	25-34	Male	190	13.04 M	1.4	1.2	1.6	218,132	217.15 M
70	Hispanic	15-24	Female	32	4.83 M	1.0	0.9	1.2	218,164	221.98 M
71	Black	15-24	Female	35	3.16 M	0.9	0.7	1.1	218,199	225.14 M
72	White	25-34	Female	103	12.60 M	0.9	0.8	1.0	218,302	237.75 M
73	Asian	25-34	Male	14	1.74 M	0.5	0.3	0.8	218,316	239.48 M
74	Asian	25-34	Female	*	1.80 M	0.3	0.2	0.5	218,316	241.28 M
75	White	15-24	Male	32	11.95 M	0.2	0.1	0.2	218,348	253.23 M
76	White	15-24	Female	*	11.35 M	0.1	0.1	0.1	218,348	264.58 M
77	Black	0-4	Male	*	1.52 M	†	†	†	†	266.11 M
78	American Indian	0-4	Male	*	0.10 M	†	†	†	†	266.20 M
79	Hispanic	0-4	Male	*	2.60 M	†	†	†	†	268.80 M
80	White	0-4	Male	*	5.21 M	†	†	†	†	274.01 M
81	Asian	0-4	Male	*	0.58 M	†	†	†	†	274.59 M
82	White	5-14	Male	*	10.95 M	†	†	†	†	285.54 M
83	Black	5-14	Male	*	3.16 M	†	†	†	†	288.70 M
84	American Indian	15-24	Male	*	0.21 M	†	†	†	†	288.91 M
85	American Indian	5-14	Male	*	0.21 M	†	†	†	†	289.12 M
86	Hispanic	5-14	Male	*	5.39 M	†	†	†	†	294.51 M
87	Asian	15-24	Male	*	1.34 M	†	†	†	†	295.85 M

88	Asian	5-14	Male	*	1.24 M	†	†	†	†	297.09 M
89	Black	0-4	Female	*	1.47 M	†	†	†	†	298.56 M
90	American Indian	0-4	Female	*	0.10 M	†	†	†	†	298.66 M
91	Hispanic	0-4	Female	*	2.50 M	†	†	†	†	301.15 M
92	White	0-4	Female	*	4.95 M	†	†	†	†	306.11 M
93	Asian	0-4	Female	*	0.55 M	†	†	†	†	306.66 M
94	White	5-14	Female	*	10.42 M	†	†	†	†	317.07 M
95	Black	5-14	Female	*	3.07 M	†	†	†	†	320.14 M
96	American Indian	15-24	Female	*	0.20 M	†	†	†	†	320.34 M
97	American Indian	5-14	Female	*	0.20 M	†	†	†	†	320.55 M
98	Hispanic	5-14	Female	*	5.18 M	†	†	†	†	325.73 M
99	Asian	15-24	Female	*	1.33 M	†	†	†	†	327.05 M
100	Asian	5-14	Female	*	1.19 M	†	†	†	†	328.24 M

* Source data was suppressed for >10 deaths
† Could not estimate because of suppressed data

Table 4: Logistic regression on CDC COVID case data showing the risk of COVID-19 case for each age/race/sex risk group sorted in order by case risk/100,000 US inhabitants

	A	B	C	D	E	F	G	H	I	J
							Risk CI	Risk CI	Cum. # of	Cum. # in
	Race	Age	Sex	# of Cases	# in Population	Case risk/100,000	Lower	Upper	cases	population
1	American Indian	30 - 39	Female	5,933	0.19 M	3,163.4	3,101.0	3,226.9	5,933	0.19 M
2	American Indian	40 - 49	Female	4,735	0.16 M	2,997.2	2,932.5	3,063.2	10,668	0.35 M
3	American Indian	20 - 29	Female	6,085	0.22 M	2,821.4	2,766.0	2,877.8	16,753	0.57 M
4	Hispanic	40 - 49	Female	107,877	3.92 M	2,764.2	2,750.5	2,778.0	124,630	4.49 M
5	Hispanic	20 - 29	Female	127,038	4.73 M	2,689.4	2,677.0	2,701.9	251,668	9.22 M
6	Black	80+	Female	17,529	0.70 M	2,667.5	2,635.8	2,699.7	269,197	9.93 M
7	Hispanic	30 - 39	Female	114,899	4.34 M	2,642.4	2,629.7	2,655.3	384,096	14.27 M
8	American Indian	30 - 39	Male	4,940	0.19 M	2,624.0	2,570.4	2,678.8	389,036	14.45 M
9	Hispanic	40 - 49	Male	104,477	4.04 M	2,572.6	2,559.6	2,585.6	493,513	18.49 M
10	American Indian	40 - 49	Male	4,126	0.16 M	2,571.0	2,513.8	2,629.4	497,639	18.65 M
11	Black	80+	Male	10,322	0.36 M	2,531.6	2,498.5	2,565.2	507,961	19.01 M
12	Hispanic	50 - 59	Male	77,349	3.08 M	2,530.9	2,516.6	2,545.3	585,310	22.09 M
13	Hispanic	50 - 59	Female	77,743	3.08 M	2,503.7	2,489.5	2,518.0	663,053	25.17 M
14	American Indian	50 - 59	Female	4,326	0.18 M	2,466.5	2,410.6	2,523.8	667,379	25.34 M
15	Hispanic	30 - 39	Male	112,841	4.76 M	2,377.1	2,365.5	2,388.7	780,220	30.10 M
16	American Indian	50 - 59	Male	3,671	0.16 M	2,297.2	2,243.9	2,351.7	783,891	30.26 M
17	Black	30 - 39	Female	72,451	3.10 M	2,286.5	2,272.6	2,300.4	856,342	33.36 M
18	Hispanic	20 - 29	Male	114,993	5.05 M	2,271.8	2,261.0	2,282.7	971,335	38.41 M

19	Black	40 - 49	Female	62,765	2.78 M	2,248.7	2,234.3	2,263.3	1,034,100	41.19 M
20	Hispanic	60 - 69	Male	40,695	1.83 M	2,229.6	2,212.8	2,246.4	1,074,795	43.02 M
21	American Indian	20 - 29	Male	4,894	0.22 M	2,196.0	2,151.1	2,241.8	1,079,689	43.24 M
22	American Indian	80+	Female	841	0.04 M	2,159.3	2,047.6	2,277.0	1,080,530	43.28 M
23	Black	50 - 59	Female	60,604	2.81 M	2,155.8	2,141.8	2,169.8	1,141,134	46.09 M
24	Black	70 - 79	Male	20,068	0.89 M	2,148.3	2,126.0	2,170.8	1,161,202	46.98 M
25	White	20 - 29	Female	254,471	12.12 M	2,112.7	2,105.4	2,120.1	1,415,673	59.10 M
26	Black	20 - 29	Female	74,736	3.43 M	2,108.4	2,095.8	2,121.1	1,490,409	62.53 M
27	Hispanic	80+	Male	8,852	0.40 M	2,090.2	2,060.3	2,120.5	1,499,261	62.93 M
28	American Indian	80+	Male	514	0.03 M	2,047.5	1,940.0	2,160.7	1,499,775	62.96 M
29	American Indian	60 - 69	Male	2,503	0.13 M	2,031.8	1,976.6	2,088.6	1,502,278	63.09 M
30	Hispanic	80+	Female	12,433	0.64 M	2,030.4	2,002.7	2,058.6	1,514,711	63.72 M
31	Hispanic	70 - 79	Male	17,741	0.86 M	2,019.1	1,996.9	2,041.5	1,532,452	64.59 M
32	Black	50 - 59	Male	48,846	2.44 M	2,008.5	1,994.9	2,022.3	1,581,298	67.02 M
33	Black	60 - 69	Male	38,149	1.88 M	2,003.9	1,988.4	2,019.4	1,619,447	68.90 M
34	American Indian	70 - 79	Male	1,171	0.07 M	1,954.3	1,880.5	2,030.9	1,620,618	68.97 M
35	Hispanic	60 - 69	Female	39,904	2.04 M	1,951.9	1,937.1	1,966.8	1,660,522	71.00 M
36	American Indian	60 - 69	Female	3,021	0.15 M	1,931.1	1,879.0	1,984.6	1,663,543	71.16 M
37	Black	40 - 49	Male	47,022	2.46 M	1,928.1	1,914.8	1,941.4	1,710,565	73.61 M
38	Black	70 - 79	Female	23,241	1.26 M	1,919.2	1,900.0	1,938.5	1,733,806	74.88 M
39	Black	60 - 69	Female	44,100	2.34 M	1,903.3	1,889.1	1,917.6	1,777,906	77.22 M
40	American Indian	10 - 19	Female	3,735	0.20 M	1,897.2	1,851.8	1,943.8	1,781,641	77.42 M
41	Black	30 - 39	Male	53,717	2.91 M	1,894.9	1,882.7	1,907.2	1,835,358	80.34 M

42	White	80+	Female	112,207	6.13 M	1,804.4	1,794.4	1,814.5	1,947,565	86.47 M
43	White	80+	Male	68,808	4.03 M	1,748.6	1,736.7	1,760.5	2,016,373	90.49 M
44	American Indian	70 - 79	Female	1,520	0.08 M	1,746.6	1,680.7	1,815.0	2,017,893	90.57 M
45	White	20 - 29	Male	215,143	12.72 M	1,678.2	1,671.9	1,684.5	2,233,036	103.29 M
46	Hispanic	70 - 79	Female	17,947	1.10 M	1,661.7	1,643.5	1,680.1	2,250,983	104.39 M
47	Black	20 - 29	Male	55,129	3.51 M	1,639.4	1,629.1	1,649.9	2,306,112	107.90 M
48	American Indian	10 - 19	Male	3,406	0.21 M	1,582.0	1,543.3	1,621.7	2,309,518	108.11 M
49	White	40 - 49	Female	176,915	11.83 M	1,493.0	1,486.9	1,499.2	2,486,433	119.94 M
50	White	30 - 39	Female	183,972	12.46 M	1,487.3	1,481.3	1,493.3	2,670,405	132.39 M
51	White	50 - 59	Female	195,134	14.22 M	1,377.9	1,372.4	1,383.4	2,865,539	146.62 M
52	Hispanic	10 - 19	Female	68,694	5.08 M	1,350.0	1,341.7	1,358.4	2,934,233	151.70 M
53	White	50 - 59	Male	183,066	13.90 M	1,310.9	1,305.5	1,316.3	3,117,299	165.60 M
54	White	40 - 49	Male	155,308	11.91 M	1,306.4	1,300.7	1,312.0	3,272,607	177.50 M
55	White	30 - 39	Male	160,821	12.69 M	1,257.5	1,252.2	1,262.9	3,433,428	190.19 M
56	White	10 - 19	Female	137,951	10.87 M	1,257.1	1,251.1	1,263.0	3,571,379	201.06 M
57	Hispanic	10 - 19	Male	64,451	5.28 M	1,221.9	1,214.3	1,229.6	3,635,830	206.35 M
58	White	70 - 79	Male	100,689	8.50 M	1,197.2	1,190.5	1,204.0	3,736,519	214.85 M
59	White	60 - 69	Male	153,468	13.34 M	1,151.1	1,145.9	1,156.3	3,889,987	228.18 M
60	White	10 - 19	Male	120,966	11.42 M	1,070.4	1,065.1	1,075.7	4,010,953	239.61 M
61	White	60 - 69	Female	153,478	14.34 M	1,069.7	1,064.8	1,074.5	4,164,431	253.95 M
62	Asian	20 - 29	Female	16,083	1.57 M	1,051.9	1,039.2	1,064.7	4,180,514	255.52 M
63	White	70 - 79	Female	103,470	9.79 M	1,045.8	1,039.9	1,051.7	4,283,984	265.30 M
64	Black	10 - 19	Female	30,480	3.10 M	1,024.2	1,015.3	1,033.1	4,314,464	268.41 M

65	Asian	20 - 29	Male	15,467	1.57 M	959.8	948.1	971.7	4,329,931	269.98 M
66	Asian	80+	Male	2,256	0.24 M	959.4	933.4	986.2	4,332,187	270.22 M
67	American Indian	0 - 9	Female	1,827	0.20 M	913.1	882.6	944.6	4,334,014	270.42 M
68	Asian	50 - 59	Male	10,315	1.16 M	912.1	899.0	925.3	4,344,329	271.57 M
69	American Indian	0 - 9	Male	1,750	0.20 M	884.3	854.8	914.8	4,346,079	271.77 M
70	Asian	80+	Female	3,170	0.36 M	860.2	837.3	883.8	4,349,249	272.13 M
71	Asian	30 - 39	Female	15,356	1.84 M	853.9	843.4	864.5	4,364,605	273.97 M
72	Black	10 - 19	Male	28,505	3.19 M	853.3	845.7	861.0	4,393,110	277.17 M
73	Asian	60 - 69	Male	7,267	0.88 M	849.1	834.9	863.6	4,400,377	278.05 M
74	Asian	50 - 59	Female	11,349	1.34 M	832.9	821.1	844.9	4,411,726	279.38 M
75	Asian	30 - 39	Male	14,458	1.70 M	830.5	820.1	841.1	4,426,184	281.08 M
76	Asian	40 - 49	Male	11,394	1.43 M	801.4	790.5	812.5	4,437,578	282.52 M
77	Asian	40 - 49	Female	13,098	1.63 M	796.1	785.5	806.9	4,450,676	284.15 M
78	Asian	70 - 79	Male	3,572	0.50 M	735.1	717.8	752.8	4,454,248	284.65 M
79	Asian	60 - 69	Female	7,713	1.09 M	685.1	673.6	696.8	4,461,961	285.74 M
80	Hispanic	0 - 9	Male	32,740	5.27 M	633.1	627.4	638.9	4,494,701	291.01 M
81	Hispanic	0 - 9	Female	31,107	5.07 M	601.4	595.9	607.1	4,525,808	296.08 M
82	Asian	70 - 79	Female	3,619	0.63 M	557.1	544.0	570.6	4,529,427	296.71 M
83	Asian	10 - 19	Female	6,504	1.24 M	529.3	519.8	538.9	4,535,931	297.95 M
84	Asian	10 - 19	Male	6,559	1.26 M	518.6	509.3	528.1	4,542,490	299.20 M
85	Black	0 - 9	Female	13,654	2.99 M	465.2	459.3	471.2	4,556,144	302.19 M
86	Black	0 - 9	Male	14,169	3.09 M	450.8	445.0	456.6	4,570,313	305.28 M
87	White	0 - 9	Female	33,967	10.06 M	342.4	339.4	345.4	4,604,280	315.34 M

88	White	0 - 9	Male	36,325	10.58 M	339.0	336.0	342.0	4,640,605	325.91 M
89	Asian	0 - 9	Male	3,410	1.19 M	290.0	282.7	297.4	4,644,015	327.11 M
90	Asian	0 - 9	Female	2,927	1.13 M	254.2	247.7	260.8	4,646,942	328.24 M

Table 5: Deaths and population sized of COVID special risk categories

Special risk category	Number of COVID-19 deaths	Number in the category	# Deaths/100,000
Nursing home (long term care)	72,642[8]	1,400,000[9,20]	5,188
Prison	1,494[17]	2,300,000[23]	64.9
Health Care Workers	1,700[10] 847[11]	16,866,020[24] 20,000,000[25]	10