SARS-CoV-2 neutralizing antibodies; longevity, breadth, and evasion by

emerging viral variants

Fiona Tea[1*], Alberto Ospina Stella[2*], Anupriya Aggarwal[2*], David Ross Darley[3,4*], Deepti

Pilli[1], Daniele Vitale[5], Vera Merheb[1], Fiona X. Z. Lee[1], Philip Cunningham[6], Gregory J.

Walker[7], David A. Brown[5,7], William D. Rawlinson[7,8], Sonia R. Isaacs[7], Vennila

Mathivanan[2], Markus Hoffman[9,10], Stefan Pöhlmann[9,10], Dominic E. Dwyer[7,11,12], Rebeca

Rockett[11,12], Vitali Sintchenko[7,11,12,13], Veronica C. Hoad[14], David O. Irving[14,15], Gregory J.

Dore[2,3], Iain B. Gosbell[14,16], Anthony D. Kelleher[2**], Gail V. Matthews[2,3**], Fabienne

Brilot[1,12,13,17,18**], Stuart G Turville[2**]

*First author: contributed equally

**Senior author: contributed equally

Affiliations

[1]Brain Autoimmunity Group, Kids Neuroscience Centre, Kids Research at the Children's

Hospital at Westmead, Sydney, New South Wales, Australia

[2]The Kirby Institute, The University of New South Wales, Sydney, New South Wales,

Australia

[3]St Vincent's Hospital, Sydney, New South Wales, Australia

[4]School of Medicine, St Vincent's Clinical School, The University of New South Wales,

Sydney, New South Wales, Australia

[5]Westmead Institute for Medical Research, Sydney, New South Wales, Australia

[6]St Vincent's Applied Medical Research, Sydney, New South Wales, Australia

[7]New South Wales Health Pathology, Sydney, Australia

[8]School of Medical Sciences, Biotechnology and Biomolecular Sciences, and School of

Women's and Children's Health, The University of New South Wales Sydney, New South

Wales, Australia

[9]German Primate Center Infection Biology Unit, Georg-August-University, Göttingen,

Germany

[10]Faculty of Biology and Psychology, Georg-August-University, Göttingen, Göttingen,

Germany

[11]Centre for Infectious Diseases & Microbiology - Public Health, New South Wales Health

Pathology - Institute of Clinical Pathology & Medical Research (ICPMR), Westmead,

Sydney, New South Wales, Australia

[12]Marie Bashir Institute for Biosecurity, Faculty of Medicine and Health, The University of

Sydney, Sydney, New South Wales, Australia

[13]Faculty of Medicine and Health, The University of Sydney, Sydney, New South Wales,

Australia

[14]Australian Red Cross Lifeblood, Melbourne, Victoria, Australia

[15]Faculty of Health, University of Technology, Sydney, New South Wales, Australia

[16]School of Medicine, Western Sydney University, Sydney, New South Wales, Australia

[17]School of Medical Sciences, Discipline of Applied Medical Science, Faculty of Medicine

and Health, The University of Sydney, Sydney, New South Wales, Australia

[18]Brain and Mind Centre, The University of Sydney, Sydney, New South Wales, Australia

Correspondence:

Fabienne Brilot

Brain Autoimmunity Group, Kids Neuroscience Centre, Kids Research at the Children's

Hospital at Westmead

School of Medical Sciences, Discipline of Applied Medical Science, Faculty of Medicine and

Health, The University of Sydney, Sydney, New South Wales, Australia,

Fabienne.brilot@sydney.edu.au

Number of Figures: 5

Number of tables: 2

Number of Supplementary Figures: 5

Word count: 8020

One Sentence Summary:

Neutralizing antibody responses to SARS-CoV-2 are sustained, associated with COVID19

severity, and evaded by emerging viral variants

Short title: Neutralizing SARS-CoV2 antibody response

Abstract

The SARS-CoV-2 antibody neutralization response and its evasion by emerging viral variants are unknown. Antibody immunoreactivity against SARS-CoV-2 antigens and Spike variants, inhibition of Spike-driven virus-cell fusion, and infectious SARS-CoV-2 neutralization were characterized in 807 serial samples from 233 RT-PCR-confirmed COVID-19 individuals with detailed demographics and followed up to seven months. A broad and sustained polyantigenic immunoreactivity against SARS-CoV-2 Spike, Membrane, and Nucleocapsid proteins, along with high viral neutralization were associated with COVID-19 severity. A subgroup of 'high responders' maintained high neutralizing responses over time, representing ideal convalescent plasma therapy donors. Antibodies generated against SARS-CoV-2 during the first COVID-19 wave had reduced immunoreactivity and neutralization potency to emerging Spike variants. Accurate monitoring of SARS-CoV-2 antibody responses would be essential for selection of optimal plasma donors and vaccine monitoring and design.

Introduction

Control of the SARS-CoV-2 pandemic relies on population resistance to infection due to a post-infection and vaccination-induced immunity. Current questions relate to the level, breadth, and longevity of generated immunity, and whether mutation of the virus will compromise immunity. Previous studies reported varying results in longitudinal changes of the virus-specific antibody response. Some detected stable antibody titers 4-6 months after diagnosis (*1, 2*), while others reported waning of the antibody response 2-3 months after infection (*3, 4*). Differences in assay sensitivity and antigen targets may account for these

discrepancies, with Spike and nucleocapsid being the main antigens investigated.

Immunoreactivity to other abundant antigens, such as Membrane or Envelope, are unknown.

Neutralization of SARS-CoV-2 has been reported for antibodies that bind to Spike, a large

homo-trimeric glycoprotein studded across the viral surface (*5, 6*), whereas Membrane and

Envelope proteins, although exposed on the viral surface, remain to be identified as

neutralizing antibody targets. Rapid development of neutralizing antibody response to Spike

correlates with viral immunity, and individuals who seroconvert may develop a lasting

neutralization response (*7*).

The SARS-CoV-2 virus has accumulated many polymorphisms across its genome, especially

within the Spike gene (*8*). Shortly after the introduction of SARS-CoV-2 into the human

population, many early and dominant amino acid polymorphisms were associated with viral

entry fitness, such as D614G (*9, 10*). However, the pressure of the neutralising antibody

response might select for escape mutations in Spike that limit post-infectious immunity or

vaccine protection (*11*). One example is the S477N/D614G Spike variant which appeared in

Australia during July and August 2020 and was traced to a single event from Australian hotel

quarantine (*12*). The S477N/D614G Spike variant currently represents greater than 5% of

Spike variants worldwide, 15% in Europe, and 58% in Oceania (*13*).

Using the lessons learned from research of other viral pathogens and neuroimmunological

autoantibodies (*14, 15*), we have developed a suite of novel high-content assays that

sensitively assess antibody responses against the native oligomeric structure of Spike and its

emerging variants (*16*). To measure the neutralizing capacity, we have also developed a

Biosafety Level 2 surrogate Spike-driven virus-cell fusion assay that has been cross-validated

with a novel high content, machine-scored, Biosafety Level 3 authentic SARS-CoV-2

neutralization assay.

Herein we characterize the longevity, polyantigenic breadth, and neutralization capacity of

the SARS-CoV-2 antibody response in individuals and their responses to globally emerging

SARS-CoV-2 variants. Using two longitudinal SARS-CoV-2 community- and hospital-based

Australian cohorts representative of the broad spectrum of disease severity at acute infection,

we showed that the polyantigenic and neutralizing responses to SARS-CoV-2 are sustained,

associated with COVID19 severity, and are evaded by emerging viral variants.

This work provides a community snapshot of humoral immunity in those recovering from

infection and sheds light on important considerations for vaccine design and selection of

donors for convalescent plasma therapy. Additionally, the modular assays used herein can be

adapted for novel viral pathogens to respond rapidly to emerging pathogens.

122   Results

123   SARS-CoV-2 antibody responses are sustained for up to seven months
124   post-infection and are focused on Spike

SARS-CoV-2 antibodies were assessed in RT-PCR-confirmed COVID-19 convalescent

adults in two Australian cohorts; ADAPT, a hospital-based cohort of patients recruited during

the first and second wave of infection in Australia (n=83 and n=17), and LIFE, a cohort of

plasma donors (n=159) (Table 1, Fig. 1A). Antibody immunoreactivity to SARS-CoV-2

antigens, inhibition of virus-cell fusion, live SARS-CoV-2 neutralization, and

immunoreactivity to Spike emerging variants were assessed and antibody features were

compared with demographic data (Fig. 1A). At first date of collection post-infection, 96%

(81/83 ADAPT, median 71 days, mean 74 days, post first PCR positivity) and 98% (152/159

LIFE, median 59 days, mean 61 days) of infected patients were Spike IgG-positive, and 81%

(66/83 ADAPT) and 91% (139/152 LIFE) were Spike IgM-positive (Table 2). A broad range

of Spike IgG levels was observed. No differences in Spike IgG and IgM levels were observed

between females and males, but higher IgG and IgM levels were associated with older age

(P<0.0001) (Fig. S1). Detection of convalescent positive serostatus was more sensitive when

Spike IgG were detected by live cell flow cytometry compared to Nucleocapsid IgG or Spike

IgG using commercial assays (Table 2).

The longevity of antibody responses was assessed in 807 Spike IgG-positive serial samples

from 233 individuals (n=162 ADAPT, n=645 LIFE), spanning up to 205 days post-PCR

positivity (Fig. 1A and Table 1). There was a range of Spike IgG titers at first collection date,

and among all Spike IgG-positive individuals, no individual seroreverted, even up to 205

days post-PCR positivity. The majority of ADAPT patients had stable IgG responses (85%),

whereas most LIFE donors exhibited decreased IgG over time (59%), where a decrease was

defined as >30% change from first collected sample (Fig. 1B) (*14*). A two-phase decay in

those with decreasing responses characterized by an initial high rate of decay followed by

stabilization, and the breakpoint between the two phases was estimated at 85 days post-PCR

positivity (Fig. 1B). The level at which Spike IgG stabilized was dependent on intial antibody

response. High Spike IgG levels decayed to mid-level reactivity and mid-low level reactivity

to low level (Fig. 1B). In Spike IgM-positive patients, the majority had decreased IgM levels

over time (68% ADAPT; 84% LIFE), in which levels initially decreased and then stabilised

at lower levels, but did not serorevert up to 205 days (Fig. 1C). Only five ADAPT (6%) and

fourteen LIFE (9%) individuals sero-reverted for Spike IgM at median 146 days post-PCR

positivity (Fig. 1C). The breakpoint between the two phases of IgM decay was at 93 days

post-PCR positivity.

The polyantigenic breadth of Spike IgG-positive individuals against the virus was examined

by detecting IgG targeting the SARS-CoV-2 Membrane, Envelope, and Nucleocapsid

proteins (Table 2). 54% (45/83 ADAPT) and 57% (87/152 LIFE) individuals harboured IgG

targeting the SARS-CoV-2 Membrane protein, whereas 78% had antibody targeting the

Nucleocapsid protein (65/83 ADAPT, 118/152 LIFE) (Fig. 1D and 1E). Antibody titers

toward the Membrane protein remained stable over the period of observation in most

individuals (91%, 41/45 ADAPT; 95%, 83/87, LIFE), whereas responses toward the

Nucleocapsid protein differed between ADAPT and LIFE, and were reminiscent of the Spike

IgG response, i.e. mostly stable in ADAPT, and mostly decreased in LIFE over time (Fig. 1D

and 1E). Across both cohorts, reactivity to the Envelope protein was very limited with only

two ADAPT patients (2%) positive for Envelope IgG (Table 2). Antibody responses to

SARS-CoV-2 were highly focused on Spike, followed by the Nucleocapsid and Membrane

proteins. Individuals with higher Spike IgG had also high levels of Nucleocapsid and

Membrane IgG (Fig. S1).

The overall decay of SARS-CoV-2 antibodies between both cohorts behaved similarly for

Spike IgM, but not for Spike IgG, Membrane IgG, and Nucleocapsid IgG, with LIFE donors

exhibiting a higher proportion of decreased profiles (Fig. 1). The first collected sample in

ADAPT started later post-PCR positivity, and the time duration between paired samples was

shorter than for LIFE samples, therefore some ADAPT patients may have been captured

during the 2^{nd}, more stable, phase (Fig. 1 and Table 1). Furthermore, few ADAPT patients

underwent plasmapheresis, whereas all LIFE donors underwent plasmapheresis as part of

convalescent plasma donations (median 6 donations, IQR 3-9, max 14). However, donors

with more than 10 donations (n=30) had decay profiles similar to the whole cohort, in which

donors stabilized at mid-low level, and none of these highly recurrent donors became

seronegative (Fig. S2).

Neutralization of SARS-CoV-2 is correlated with Spike antibody levels and is maintained over time

The neutralization capacity of these individual responses was assessed on a Spike-driven virus-cell fusion assay and a whole-virus neutralization assay (Fig. 1A). Most sera were capable of inhibiting virus-cell fusion (82%, 68/83 ADAPT; 68%, 104/152 LIFE) and mediating viral neutralization (88%, 73/83 ADAPT; 94%, 143/152 LIFE) (Fig. 2A, Table 2). In both cohorts, the virus-cell fusion assay was more stringent than the SARS-CoV-2 neutralization assay as a proportion of individual sera with lower titers in the SARS-CoV-2 neutralization assay were negative in the virus-cell fusion assay (7%, 6/83 ADAPT; 27%, 41/152 LIFE), and most individuals had higher titers in the neutralization assay (Fig. 2A). To understand the discrepancy between both viral assays, live SARS-CoV-2 viral particles were enumerated and directly compared to Spike-pseudotyped lentiviral particles. Cell permeable RNA-specific staining of live virions detected viral particles that were Nucleocapsid-positive (Fig. 2B). The particle to transduction ratios from the fusion assay were 1.03×10^5, consistent with the low specific infectivity of lentiviruses such as HIV-1[17]. In contrast, the SARS-CoV-2 particle to infectivity ranged from 58 (HekAT14) to 578 (VeroE6), consistent with the ratio reported for influenza virus [18]. However, the absolute viral particle number was 74-fold higher in Spike-pseudotyped particle preparation (1.64×10^8 particles per ml) compared to authentic SARS-CoV-2 (2.22×10^6 particles per ml). Thus, the specific infectivity of SARS-CoV-2 was higher than that of Spike-expressing lentiviral particles, which may account for the higher sensitivity of the SARS-CoV-2-based neutralization assay.

Most ADAPT patients had stable virus-cell fusion inhibition (99%) and neutralization (89%) titers over time (Fig. 2C and 2D). Most of LIFE donors had decreased virus-cell fusion inhibition (82%) and neutralization (56%) capacity over time, and the majority exhibited a

single-phase decay in both assays (Fig. 2C and 2D). The greater number of samples per LIFE

donor enabled finer characterization of the decay profile in 34 donors in the virus-cell fusion

and 44 donors in the neutralization assay). Most donors had a single-phase decay, whilst a

two-phase decay was observed in those with >1:320 titers at first collection. These rapidly

dropped and then stabilized over time at 1:80 to 1:160 (28% and 25% of LIFE donors in the

virus-cell fusion and neutralization assays, respectively). Individuals with two-phase decay

had much higher starting titers than individuals with a single-phase decay (Fig. 2D). In the

neutralization assay, LIFE donors with decreased profile had a similar median follow up as

the stable profile (~63 days and 56 days, respectively). In both cohorts, the neutralization and

fusion profiles were similar to the Spike IgG profiles, in which ADAPT had more stable

responses than LIFE. Indeed, Spike IgG and IgM titers were strongly correlated with virus-

cell fusion inhibition and SARS-CoV-2 neutralization (Fig. 2E).

A broad antigenic repertoire and high neutralization capacity against

SARS-CoV-2 is associated with COVID-19 severity

Approximately half of individuals (55% ADAPT and 49% LIFE) had broad polyantigenic

immunoreactivity as defined by IgG responses against each of SARS-CoV-2 Spike,

Membrane, and Nucleocapsid proteins (Fig. 3A). Around a third of individuals exhibited

antibodies against only two proteins (27% and 30%, Nucleocapsid and Spike; 2% and 9%

toward Membrane and Spike in ADAPT and LIFE respectively), and a smaller proportion

had responses against Spike alone (12 and 17%) (Fig. 3A). Polyantigenic immunoreactivity

did not change overtime in most individuals (82%, 15/81 ADAPT, 83%, 41/152 LIFE, data

not shown). No individual developed IgG to new antigens at any point of follow up, but

instead, lost immunoreactivity to one antigen, either Nucleocapsid or Membrane.

Patients had broader responses across the spectrum of severity in ADAPT (Fig. 3B). ADAPT

and LIFE hospitalized patients with more severe symptoms were more likely to exhibit a

broader antibody response to SARS-CoV-2, i.e. polyreactive toward the three antigens (Fig.

3B). Interestingly, two of seven hospitalized LIFE patients who had a short 24 hour

hospitalization harboured non-broad responses, and Spike-only responses were exclusively

observed in non-hospitalized, mild, and moderate individuals (Fig. 3B). Higher IgG titers

against Membrane and Nucleocapsid proteins were also associated with disease severity in

both cohorts (Fig. 3C). Patients with broader SARS-CoV-2 responses and higher disease

severity had greater viral neutralization and virus-cell fusion inhibition (Fig. 3D, E). This

polyreactive, high severity subgroup was populated almost exclusively by older males (Fig.

3F). Similarly, higher neutralization and virus-cell fusion inhibition titers were more enriched

in older males with moderate disease and who were hospitalized (Fig. 3G and H).

High responders with strong and broad SARS-CoV-2 antibody responses

are ideal plasma donors

A small subgroup of individuals were "high responders" characterized by high Spike IgG,

Spike IgM-positive, broad polyantigenic immunoreactivity (binding to Nucleocapsid, Spike,

and Membrane), virus-cell fusion inhibition (>1:160), and neutralization (>1:320). They

maintained this high response over time (n=14, 17% ADAPT, n=19, 12% LIFE). High

responders were more likely to be male, hospitalized, and were of older age (Fig. 4A).

Further characterization was performed on a series of increasingly permissive cell lines,

VeroE6, HekAT14, HekAT10, and HekAT24 (Fig. 4B and Fig. S3). Low, i.e. non-high

responders, and high responders sera neutralized live SARS-CoV-2 in VeroE6, HekAT14,

and HekAT10 cell lines, whereas limited neutralization was observed in the hyper-permissive

HekAT24 cell line (Fig. 4B). Using the HekAT24 cell line, two elite responders were

identified in LIFE (Fig. 4C), with high Spike IgG and IgM levels, and neutralization titers 30-

to 4-fold greater than other individuals (Fig. 4D). Interestingly, elite responders had the

highest detectable IgM levels, and early IgM decay coincided with a decrease in

neutralization titers, whereas Spike IgG remained stable overtime (Fig. 4E). This association

between decreased IgM and neutralization titers was observed in ~10% of individuals in both

cohorts (data not shown).

Spike IgG antibody binding and neutralizing capacity are dependent on

Spike mutations in emerging new variants

Numerous Spike polymorphisms have evolved over the course of the pandemic (*11*) with the

most attention given to the transmission fitness gain variants, such as D614G (*10, 11*). To test

the breadth of the antibody response, Spike IgG immunoreactivity to several Spike variants

implicated in the Receptor Binding Domain (RBD) and S1 was assessed (Fig. 1A).

Expression of all Spike variants was similar across each transfected cell line used in the flow

cytometry antibody assays (Fig. S4A). Compared to the Wuhan-1 D614 variant, most patients

had similar binding and were able to recognise the Spike RBD variants G476F, V483A, and

V367S (Fig. 5A). However, across both cohorts, there was an overall reduced binding to

D614G, a prominent non-RBD S1 variant present during the Australian first wave (Fig. 5A).

65% of ADAPT and 91% of LIFE individuals, infected from the first world-wide wave,

generated antibodies that bound broadly to G476F, V483A, V367S, and D614G Spike,

whereas 35% of ADAPT and 9% of LIFE had more restricted Spike recognition; i.e. they

recognized G476F, V483A, V367S, but had a decreased binding to D641G (Fig. 5A).

Immunoreactivity toward all Spike variants was stable overtime in most patients (data not

shown).

Importantly, sera with reduced D614G IgG binding also had lower neutralization and virus-

cell fusion inhibition compared to those who recognised D614G Spike (Fig. 5B), suggesting

implications for blocking infection in patients who cannot induce robust Spike antibody

recognition. Furthermore, patients who bound D614G Spike had broad SARS-CoV-2

polyantigenic immunoreactivity, whereas patients who displayed reduced binding to D614G

had more limited antigenic recognition, with 36% recognising Spike only (Fig. 5C). In a

D614G virus-cell fusion assay, patients who maintained binding to D614G showed enhanced

virus-cell fusion inhibition, compared to when parental Wuhan-1 D614 Spike was used (Fig.

5D). Individuals with lower IgG binding to D614G, i.e. restricted variant recognition had

limited D614G Spike virus-cell fusion inhibition, and most (8/11) were unable to prevent

Spike fusion (data not shown), emphasizing the need to maintain robust binding to Spike

variants for efficient viral neutralization. Patients with restricted Spike variant recognition

were not distinguished by age and severity, but were more likely to be female (Fig. S5).

Although D614G Spike remains a predominant variant globally (Fig. 5E), in the second wave

of Australian infection between July to September, an isolate with additional polymorphisms,

primarily S477N, and in some cases an additional V1068F, was identified. These variants

were not detected during the first Australian wave (Fig. 5F), which included the original

Wuhan-1 D614 or the D614G variant equally. To assess the antibody binding capacity

between original and emerging variants, patients infected by two Spike variants,

S477N/D614G and S477N/D614G/V1068F, were recruited during the second wave in

Australia (n=17, from the ADAPT cohort, Table 1). All ADAPT patients from the first and

second wave had detectable IgG against all Spike variants (Fig. S4B). Compared to the

D614G variant, a strong decrease in immunoreactivity to S477N/D614G and

S477N/D614G/V1068F was observed in all ADAPT patients from the second wave, whereas

the third mutation within the Spike S2 domain V1068F did not have an additive effect (Fig.

5G). This decrease was also observed irrespective of the virus variant that had infected the

ADAPT patients (Fig. 5G). Importantly ADAPT patients from the first wave, who had not

encountered the new variants, had reduced binding to S477N/D614G and

S477N/D614G/V1068F, suggesting a global decrease of immunoreactivity toward both new

variants (Fig. 5G).

Discussion

The current study characterizes the breadth, longevity, and neutralizing capacity of SARS-

CoV-2 antibody response in two Australian cohorts, encompassing a wide range of

demographics and disease states, up to seven months after COVID-19 diagnosis. We show

the development of broad and sustained immunoreactivity against SARS-CoV-2 antigens,

and found high titers of Spike-binding and virus-neutralizing antibodies were associated with

COVID-19 severity. A group of high responders were identified with high, broad, and

sustained neutralizing responses, who may represent ideal donors for convalescent plasma

donations. Most importantly, although most patients seroconverted, antibodies generated

after early infection displayed a significantly reduced antibody binding and neutralization

potency to emerging evasive variants. Our data has important implications on hyperimmune

therapy, monoclonal antibody treatments, and vaccine development strategies against

emerging viral variants.

The longevity of the immune response against SARS-CoV-2 is a fundamental yet currently

unresolved question. Like others, we observed a strong correlation between Spike IgG levels

and neutralization capacity (*19, 20*). Although reports on neutralization prevalence and

average titers vary widely depending on sampling and detection assay strategies (*21, 22*), our

results expand on previous findings by comparing neutralization levels with antigen-specific

response over a longer follow-up period with more timepoints than most previous studies.

The decline in IgG titers and neutralization often stabilized at different levels later into

convalescence, addressing whether decreasing IgG levels eventually plateau. Especially in

LIFE whose samples were collected later post-infection and with a longer follow-up period

than ADAPT. Spike IgM levels decreased more rapidly than IgG, but were still detectable up

to 205 days after diagnosis, much later than previously reported (*1, 19*), and consistent with

mathematical modelling of decline of IgM titers in a smaller convalescent cohort (*23*). While

our results reveal widely different magnitudes of initial responses and a decrease in

neutralizing antibodies titres, most patients have detectable Spike IgG and neutralizing

responses more than 5 months after diagnosis, suggesting extended humoral protection, even

in those with mild manifestations of the disease.

IgG and IgM against conformational Spike antibody assays have been seldom used, and

Spike IgM detection has been challenging. Although many serological assays have reported

100% sensitivity at ~15 days post-infection (*24*), prevalence studies, vaccine efficacy, and

assessment for convalescent COVID-19 plasma donors may not recruit so early post-

infection or -vaccination. In this context, and future seroprevalence studies, more sensitive

antibody assays will be essential. Flow cytometry assays are used in clinical diagnostics,

mainly in the sensitive and specific detection of neuroimmunological autoantibodies in which

antigen conformation and discrimination of seropositive patients from healthy controls are

critical (*14, 15*). Within the follow up time, the detection of Nucleocapsid and Spike IgG by

high capacity commercial assays was significantly less sensitive compared to the flow

cytometry assay. Integration of the flow cytometry assay to detect Spike IgG would be

valuable to include in the diagnostic pipeline in addition to resource-intensive whole virus

neutralization. Given the sensitivity of the flow cytometry assay, this methodology would be

ideally suited towards seroprevalence in populations to reveal the true rates of community

infection.

The majority of individuals in both cohorts were treated in the community. COVID-19

severity, from mild to hospitalization, was associated with an antibody immune response

against SARS-CoV-2 that was reactive toward an increasing number of SARS-CoV-2

antigens, as recently reported (25). As our cohorts included only convalescent individuals, the

role of broad polyantigenic immunoreactivity in the acute response of hospitalized patients

remains unknown. Indeed, reports of patients with absent humoral immune responses have

hinted at the role of T cells and innate immune response during the acute disease.

Nonetheless, the presence of a broad polyantigenic viral immunoreactivity can be useful to

monitor the quality of the antibody response after vaccination.

Whilst the correlation of Spike IgG levels with viral neutralization was strong, high Spike

IgM levels were also associated with high viral neutralization in some, especially during the

early convalescent days. A lack of somatic mutations was observed in hundreds of cloned

neutralizing human antibodies from convalescent patients (26). In addition, many antibody

precursor sequences were observed in naïve B cells from pre-pandemic patient samples,

highlighting the importance of pre-existing germline antibody sequences in the neutralization

response. The lack of somatic mutations observed in IgG may be consistent with IgM being

potent in a neutralization response as both isotypes could have similar affinity binding sites

for Spike, but with multiple binding sites per molecule on IgM, the avidity for Spike would

be higher.

Full virus neutralization and prevention of virus-cell fusion were associated. Whilst many

assays aim to assess neutralization surrogates outside of level 3 biosafety laboratories, key

differences were observed between Spike-driven virus-cell fusion and the authentic SARS-

CoV-2 assay. In our study, the particle to transduction ratio in the virus-cell fusion assay was

much higher than the SARS-CoV-2 neutralization assay. This is consistent with the

respective infectivity of HIV-1 compared to respiratory viruses such as SARS, non-SARS

coronaviruses, and influenza (*17, 18*). The virus-cell fusion assay involves a single round of

infection, whereas the full virus in the neutralization assay is replication-competent and

undergoes multiple rounds of replication over a three day culture. Therefore, the spread of the

virus must be considered alongside the capacity of antibodies to inhibit the initial single

particle entry and blocking of the virus spread between cells. Although the pseudotyping

fusion assay had lower sensitivity, most individuals across both cohorts had titers in this

assay with potency ranking similar to full virus neutralization.

Transfusion of convalescent COVID-19 plasma has been proposed as a therapy, with >70

ongoing randomized controlled trials. The few clinical trials to date have supported an

acceptable safety profile, but evidence regarding efficacy is mixed (*27*). Most unsuccessful

trials included donors with unknown neutralizing status or FDA-classified low titers, whereas

successful trials either used high-titer or convalescent COVID-19 plasma delivered within

three days of hospitalization (*28*). Indeed, utility of convalescent plasma could be improved

by donations during early disease stages and by selecting donors with high neutralization

antibody titers (*29*). Our findings that the immunological response to SARS-CoV-2 is widely

heterogeneous, with large variations in SARS-CoV-2 antibodies and neutralization,

polyantigenic immunoreactivity, and longitudinal responses complement these assertions. To

take into account the first phase of decay observed during early convalescence, we propose

an optimal window for plasmapheresis, up to 100 days post-diagnosis. Furthermore, the

occurrence of a small group of individuals, termed "high and elite responders" with highly

neutralising, broad, and sustained SARS-CoV-2 antibody responses over time, may be due to

the rapid and lasting generation of memory B cells (*30*). These patients were likely to be

hospitalized older males. Alongside appropriate serology screening programs, the targeted

recruitment for plasma donations could help to identify optimal convalescent resources

available within affected communities.

A clear advantage of the methodologies used in this study is the capacity of both level 2

biosafety pseudotyped fusion and flow cytometry assays to monitor the effects of viral

polymorphisms in real time. Indeed with acceleration of global viral spread, we are now

observing evolution of viral fitness and/or immune escape across millions of infected people.

The SARS-CoV-2 fitness gain of D614G (*9*) appeared very early in the pandemic and still

represents the majority of viral infections globally (>80%)(*10*). Zoonosis of a virus is often

followed by finer tuning of replication, as observed in the 2014 Ebola outbreak, in which the

variant A82V enabled more efficient receptor NPC1 usage (*31*). Although D614G is a

polymorphism outside of the RBD, it significantly impacts the RBD positioning and Spike

quaternary structure. The release of hydrogen bonds leading to structural changes are

proposed to expose Spike to increase ACE2-dependent fusion (*32*). RBD exposure in the

D614G variant may explain the association with great inhibition of virus-cell fusion in

patients who recognized the D614G Spike variant. These results are consistent with recent

studies in hamsters (*9*) and preliminary data on protection from the ongoing vaccine human

trials in areas where the D614G Spike variant remains prevalent. However, our data also

highlighted that a subgroup of patients who displayed limited antibody binding to D614G

Spike also had reduced virus neutralization irrespective of the viral variant that had infected

them. This could be a major concern for vaccine candidate design, especially given the

emergence of the S477N/D614G polymorphism in the majority of patients that were infected

in the Australian second wave and in Europe (*12, 13*). Seroconversion was observed in all

patients from the first and second wave, and good antibody binding to Wuhan-1 D614 and

D614G, but there also was a significant decrease in binding to S477N/D614G Spike

independent of the variant that had infected individuals. Therefore the emergence of the

additional polymorphism S477N/D614G could represent an immune evasive variant leading

to less antibody binding and a resistance to virus neutralization, which could imply a need for

periodic variation in vaccine design, as for the influenza vaccine (*33*). Indeed, the mapping of

Spike monoclonal antibody escape *in vitro* has recently shown that S477N/D614G is broadly

resistant to many neutralizing antibody clones (*34*). Whilst the mechanism behind these

observations is unknown, the appearance of a N-glycosylation site within Spike RBD could

lead to glycan shielding, as in HIV (*35*), and our evidence that S477N/D614G-infected

patients have a similar binding to this variant, albeit reduced, compared to first wave patients,

may suggest changing the Wuhan-1 D614 Spike to the S477N/D614G variant in vaccine

generation may not overcome the resistance of this variant to the neutralising antibody

response. As antibodies against Spike harness the majority of neutralizing activity, selecting

the optimal Spike variants in monovalent or multivalent vaccine strategies may be critical.

Our study has important translatable implications to understand the natural history of

COVID-19, and post-infection and vaccination-induced immunity. We have highlighted that

molecular epidemiology and sero-surveillance will both be required to detect emerging

polymorphisms. Furthermore, sensitive monitoring of antibody binding and neutralization

capacity will be paramount in vaccine design strategy and convalescent plasma therapy, and

in seroprevalence studies, and this would require involvement of more rapidly adaptive

methodologies to characterize the magnitude of the neutralization antibody responses against

emerging variants.

Materials and Methods

Subjects

This study investigated two cohorts of RT-PCR-confirmed convalescent individuals recruited from February to October 2020 in Australia (Table 1 and Fig. 1A). The Adapting to Pandemic Threats (ADAPT) cohort included 83 patients diagnosed at a community-based fever clinic whose sera was collected at two time points post PCR-positivity during the first wave (March-August, n=166 samples). The second wave included sera from 17 patients recruited between July to October. The Australian Red Cross Lifeblood (Lifeblood) cohort (LIFE) included 645 sera samples from 159 donors collected at multiple timepoints post-PCR-positivity (at least 28 days post-recovery) from volunteers presenting to Lifeblood for whole blood or plasma donation. The disease severity of ADAPT patients ranged from mildly symptomatic (mild), community-managed (moderate) to critically unwell and hospitalized (hosp), whereas the self-reported disease severity of LIFE donors included community-managed (non-hosp) and hospitalized (hosp) (Table 1). A healthy adult non-infected pre-pandemic cohort was collected in Australia and consisted of healthy and non-inflammatory neurological disorder donors (n=24). No re-exposure to SARS-CoV-2 and no re-infection was reported. Ethics approval for this study was granted by St Vincent's Hospital (2020/ETH00964) and Lifeblood (30042020) Research Ethics Committees. Written consent was obtained from all ADAPT patients. In LIFE, the donor consent form included a statement that blood donation may be used in research.

Flow cytometry cell-based assay for detection of SARS-CoV-2 antibodies

A flow cytometry cell-based assay detected patient serum antibodies against SARS-CoV-2

antigens as for neuroimmunological autoantibodies (*14, 15*). SARS-CoV-2 full-length Spike

(Wuhan-1 D614, V367F, G476S, V483A, D614G, S477N/D614G, and

S477N/D614G/V1068F) (*10, 11*), Membrane, and Envelope proteins were expressed on

transfected HEK293 cells. Serum (1:80) was added to live Spike-expressing cells, and

Membrane-, and Envelope-expressing cells were treated with 4% paraformaldehyde and

0.2% saponin, followed by AlexaFluor 647-conjugated anti-human IgG (H+L)

(ThermoFisher Scientific) or anti-human IgM (A21249, ThermoFisher Scientific). Cells were

acquired on the LSRII flow cytometer (BD Biosciences). Patients were SARS-CoV-2

antibody-positive if their delta median fluorescence intensity (ΔMFI $=$ MFI transfected

cells – MFI untransfected cells) was above the positive threshold (mean ΔMFI+4SD of 24

pre-pandemic controls) in at least two of three quality-controlled experiments (*14*). Binding

to Spike variants was expressed as a percentage of reduced binding compared to Spike. Data

was analysed using FlowJo 10.4.1 (TreeStar, USA), Excel (Microsoft, USA), and GraphPad

Prism (GraphPad Software, USA).

Commercial SARS-CoV-2 ELISA

Nucleocapsid IgG assay on the ARCHITECT-I (Abbott Diagnostics, USA), quantitative

Spike-1/Spike-2 (S1/S2) IgG on LIASON-155 XL (DiaSorin S.p.A, Italy), and Spike (S1)

IgG immunoassay (EUROIMMUN, Germany) were performed. Samples were reported

positive if the signal was greater than the published cut-off value (>1.4). Signal to cut-off

ratios were used.

SARS-CoV-2 viral-cell fusion assay

The hACE2 ORF (Addgene# 1786) was cloned into a 3rd generation lentiviral expression

vector and clonal stable ACE2-expressing Hek293T cells were generated by lentiviral

transductions (*36*). Lentiviral particles pseudotyped with SARS-CoV-2 Spike envelope were

produced by co-transfecting Hek293T cells with a GFP encoding lentiviral plasmid HRSIN-

CSGW (*37*), psPAX2, and plasmid expressing C-terminal truncated Spike (pCG1-SARS-2-S

Delta18) (*38*) including D614 or D614G (*36*). Neutralization activity of sera was measured

using a single round infection of ACE2-HEK293T with Spike-pseudotyped lentiviral

particles. Virus particles were incubated with serially diluted donor sera for 1 hour at 37°C.

Virus-serum mix was then added onto ACE2-HEK293T cells (2.5×10^3/well) in a 384-well

plate. Following spinoculation at 1200g for 1 hour at 18°C, the cells were moved to 37°C for

72 hours. Entry of Spike particles was imaged by GFP-positive cells (InCell Analyzer)

followed by enumeration with InCarta software (Cytiva, USA). Neutralization was measured

by reduction in GFP expression relative to control group infected with the virus particles

without any serum treatment.

The virus entry pathway in VeroE6, used in live virus neutralization assays, is primarily

endosomal (*39*). In contrast, cells derived from nasopharyngeal tissues, express ACE2 in

addition to the surface serine protease TMPRSS2 which drives virus-cell membrane fusion

and can signficantly enhance viral entry (*38*). To address viral neutralization in the presence

of ACE2 and TMPRSS2, a portfolio of Hek293T expressing clonal cell lines with ACE2 and

TMPRSS2 (HekAT) was generated. The coexpression of ACE2 and TMPRSS2 led to a series

of increasingly permissive cell lines that were readily susceptible to SARS-CoV-2 cytopathic

effects, VeroE6, HekAT14, HekAT10, HekAT24 (Fig. S3).

High content fluorescdent live SARS-CoV-2 neutralization assay

Sera were serially diluted and mixed in duplicate with an equal volume of virus solution at

$1.5x10^3$ TCID50/mL. After 1 hour of virus-serum coincubation at 37°C, 40μL were added to

equal volume of freshly-trypsinized VeroE6 cells, and three clonal HekAT cells in 384-well

plates ($5x10^3$/well) selected on SARS-CoV-2 permissiveness. After 72h, cells were stained

with NucBlue (Invitrogen, USA) and the entire well was imaged with InCell Analyzer.

Nuclei counts, proxy for resulting cytopathic effect, were compared between convalescent

sera, mock controls (defined as 100% neutralisation), and infected controls (defined as 0%

neutralization) using the formula; % viral neutralization = (D-(1-Q))x100/D, where Q =

nuclei count normalized to mock controls, and D = 1-Q for average of infection controls

(InCarta software).

Enumeration of SARS-CoV-2 particles

Live SARS-CoV-2 and lentiviral particles were stained using SYTO™ RNASelect™ Green

Fluorescent cell Stain (Invitrogen, USA) at a final concentration of 10μM for 30 minutes at

37°C in freshly thawed unpurified viral particles. Particles were then diluted 1/10 and 1/100

in sterile PBS and then adhered to Poly-L-Lysine coated glass bottom 96-well Greiner

Sensoplates (Sigma Aldrich, USA) through spinoculation at 1200g for 1 hour at 18°C.

Particles were either imaged live or immune-fluorescently counter-stained using a rabbit

polyclonal SARS-CoV-2 Nucleocapsid antibody, followed by Alexa647-conjugated goat

anti-rabbit IgG (Novus Biologicals, USA). Viral particles were then imaged and quantified as

previously described (*40*). Particle to infectivity ratios were determined by dividing the total

particle count per ml with the calculated TCID50/ml. Particle to GFP transduction ratios were

used for lentiviruses.

SARS-CoV-2 Spike sequencing and analysis

Clinical respiratory samples were sequenced using an existing amplicon-based Illumina sequencing approach. The raw sequence data were subjected to an in-house quality control procedure before further analysis as reported in (*41*). Non-synonymous SARS-CoV-2 Spike mutations (read frequency >0.8, minimum coverage 10x) were inferred from variant calling files during bioinformatic analysis using phylogenetic assignment of named global outbreak lineages (PANGOLIN)(*11*). All consensus SARS-CoV-2 genomes identified have been uploaded to GISAID (www.gisaid.org).

Statistics

Statistical analyses were performed in R v4.0.3. Loess curves were generated using ggplot2 v3.3.2. For categorical variables, a log-linear model was fitted and Pearson residuals plotted in a mosaic plot (MASS v7.3-51.6). Shapiro-Wilk test was used to test for normality in continuous variables and a Dwass-Steel-Critchlow-Fligner test was used to test for significance between continuous and categorical variables. Correlations were measured using the Spearman method (psych v2.0.8). Virus-cell fusion and neutralization data were fitted using an exponential decay curve (Origin Lab). Patient curves unable to be fitted, <3 collection dates or low viral fusion and neutralization, were undetermined. Statistical significance was determined as $p<0.05$.

References

1. D. F. Gudbjartsson *et al.*, Humoral Immune Response to SARS-CoV-2 in Iceland. *N Engl J Med* **383**, 1724-1734 (2020).
2. B. Isho *et al.*, Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci Immunol* **5**, (2020).

3.   F. J. Ibarrondo *et al.*, Rapid Decay of Anti–SARS-CoV-2 Antibodies in Persons with
Mild Covid-19. *New England Journal of Medicine* **383**, 1085-1087 (2020).

4.   M. Pollán *et al.*, Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide,
population-based seroepidemiological study. *Lancet* **396**, 535-544 (2020).

5.   T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and
protection from disease in a small animal model. *Science* **369**, 956-963 (2020).

6.   M. Hoffmann *et al.*, Camostat mesylate inhibits SARS-CoV-2 activation by
TMPRSS2-related proteases and its metabolite GBPA exerts antiviral activity.
bioRxiv, (2020).

7.   A. Wajnberg *et al.*, Robust neutralizing antibodies to SARS-CoV-2 infection persist
for months. *Science* **370**, 1227-1230 (2020).

8.   Q. Li *et al.*, The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and
Antigenicity. *Cell* **182**, 1284-1294 e1289 (2020).

9.   Y. J. Hou *et al.*, SARS-CoV-2 D614G variant exhibits efficient replication ex vivo
and transmission in vivo. *Science*, (2020).

10.   B. Korber *et al.*, Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G
Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819 (2020).

11.   Z. Liu *et al.*, Landscape analysis of escape variants identifies SARS-CoV-2 spike
mutations that attenuate monoclonal and serum antibody neutralization. *bioRxiv*,
2020.2011.2006.372037 (2020).

12.   A. T. Chen, K. Altschuler, S. H. Zhan, Y. A. Chan, B. E. Deverman, COVID-19 CG:
Tracking SARS-CoV-2 mutations by locations and dates of interest. *bioRxiv*, (2020).

13.   E. B. Hodcroft *et al.*, Emergence and spread of a SARS-CoV-2 variant through
Europe in the summer of 2020. *medRxiv*, 2020.2010.2025.20219063 (2020).

14.   F. Tea *et al.*, Characterization of the human myelin oligodendrocyte glycoprotein
antibody response in demyelination. *Acta Neuropathol Commun* **7**, 145 (2019).

15.   F. Graus *et al.*, A clinical approach to diagnosis of autoimmune encephalitis. *Lancet*
Neurol **15**, 391-404 (2016).

16.   D. R. Burton, L. Hangartner, Broadly Neutralizing Antibodies to HIV and Their Role
in Vaccine Design. *Annu Rev Immunol* **34**, 635-659 (2016).

17.   P. J. Klasse, Molecular determinants of the ratio of inert to infectious virus particles.
Prog Mol Biol Transl Sci **129**, 285-326 (2015).

18.   K. Martin, A. Helenius, Transport of incoming influenza virus nucleocapsids into the
nucleus. *J Virol* **65**, 232-244 (1991).

19.   A. S. Iyer *et al.*, Persistence and decay of human antibody responses to the receptor
binding domain of SARS-CoV-2 spike protein in COVID-19 patients. *Sci Immunol* **5**,
(2020).

20.   S. L. Klein *et al.*, Sex, age, and hospitalization drive antibody responses in a COVID-
19 convalescent plasma donor population. *J Clin Invest* **130**, 6141-6150 (2020).

21.   D. F. Robbiani *et al.*, Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442 (2020).

22.   X. Wang *et al.*, Neutralizing Antibodies Responses to SARS-CoV-2 in COVID-19
Inpatients and Convalescent Patients. *Clin Infect Dis*, (2020).

23.   A. K. Wheatley *et al.*, Evolution of immunity to SARS-CoV-2. *medRxiv*,
2020.2009.2009.20191205 (2020).

24.   A. Bryan *et al.*, Performance Characteristics of the Abbott Architect SARS-CoV-2
IgG Assay and Seroprevalence in Boise, Idaho. *J Clin Microbiol* **58**, (2020).

25.   E. Shrock *et al.*, Viral epitope profiling of COVID-19 patients reveals cross-reactivity
and correlates of severity. *Science* **370**, eabd4250 (2020).

26.   C. Kreer *et al.*, Longitudinal Isolation of Potent Near-Germline SARS-CoV-2-
Neutralizing Antibodies from COVID-19 Patients. *Cell* **182**, 843-854 e812 (2020).
27.   K. L. Chai *et al.*, Convalescent plasma or hyperimmune immunoglobulin for people
with COVID-19: a living systematic review. *Cochrane Database Syst Rev* **10**,
Cd013600 (2020).
28.   H. Abolghasemi *et al.*, Clinical efficacy of convalescent plasma for treatment of
COVID-19 infections: Results of a multicenter clinical study. *Transfus Apher Sci* **59**,
102875 (2020).
29.   E. Salazar *et al.*, Significantly Decreased Mortality in a Large Cohort of Coronavirus
Disease 2019 (COVID-19) Patients Transfused Early with Convalescent Plasma
Containing High-Titer Anti-Severe Acute Respiratory Syndrome Coronavirus 2
(SARS-CoV-2) Spike Protein IgG. *Am J Pathol*, (2020).
30.   A. Arunasingam *et al.*, Long-Term Persistence of Neutralizing Memory B Cells in
Sars-Cov-2. *SSRN*, (2020).
31.   W. E. Diehl *et al.*, Ebola Virus Glycoprotein with Increased Infectivity Dominated the
2013-2016 Epidemic. *Cell* **167**, 1088-1098 e1086 (2016).
32.   L. Yurkovetskiy *et al.*, Structural and Functional Analysis of the D614G SARS-CoV-
2 Spike Protein Variant. *Cell* **183**, 739-751 e738 (2020).
33.   K. Houser, K. Subbarao, Influenza vaccines: challenges and solutions. *Cell Host*
Microbe **17**, 295-300 (2015).
34.   L. Liu *et al.*, Potent Neutralizing Antibodies Directed to Multiple Epitopes on SARS-
CoV-2 Spike. *bioRxiv*, 2020.2006.2017.153486 (2020).
35.   T. Zhou *et al.*, Quantification of the Impact of the HIV-1-Glycan Shield on Antibody
Elicitation. *Cell Rep* **19**, 719-732 (2017).
36.   A. Aggarwal *et al.*, Mobilization of HIV spread by diaphanous 2 dependent filopodia
in infected dendritic cells. *PLoS Pathog* **8**, e1002762 (2012).
37.   M. G. Toscano *et al.*, Efficient lentiviral transduction of Herpesvirus saimiri
immortalized T cells as a model for gene therapy in primary immunodeficiencies.
Gene Ther **11**, 956-961 (2004).
38.   M. Hoffmann *et al.*, SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and
Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280 e278 (2020).
39.   J. Wei *et al.*, Genome-wide CRISPR Screens Reveal Host Factors Critical for SARS-
CoV-2 Infection. *Cell*, (2020).
40.   S. G. Turville, M. Aravantinou, H. Stossel, N. Romani, M. Robbiani, Resolution of de
novo HIV production and trafficking in immature dendritic cells. *Nat Methods* **5**, 75-
85 (2008).
41.   R. J. Rockett *et al.*, Revealing COVID-19 transmission in Australia by SARS-CoV-2
genome sequencing and agent-based modeling. *Nat Med* **26**, 1398-1404 (2020).

Acknowledgements: We thank all the patients and donors who participated in this study. We

thank Drs. Suat Dervish, Edwin Lau, and Maggie Wang for providing advices at the Flow

Cytometry Core Facility of the Westmead Research Hub. We thank Ms Rebecca Rielly and

the Operation Team at Kids Research for providing acces to the PC2 facility; **Funding:** This

work was supported by Snow Medical (Australia), The University of New South Wales

Rapid Response grant (Australia), the University of Sydney Research Excellence Initiative

grant (Australia), and the MRRF NHRMC COVID-19 grant. The Australian Governments

fund Australian Red Cross Lifeblood for the provision of blood, blood products, and services

for the Australian community; **Author contributions:** FB, SGT, AK, and GM designed the

study. FT, DP, DV, AOS, AA, VM, FXL, GW, RR, VS, WR, PC conducted and analysed

experiments. GD, GM, DRD, IG, VH, DI enrolled and managed patients. FT, AOS, AA, and

DRD wrote the manuscript first draft and FT prepared Figs and tables. FB, SGT, AK, and

GM designed and coordinated research and verified results. All authors reviewed the draft

before submission; **Competing Interests:** FB has received honoraria from Biogen Idec and

Merck Serono as invited speaker. All other authors declare no competing interests; **Data and**

materials availability: Correspondence and requests for data should be addressed to FB and

may be limited due to ethical considerations. Plasmids transfer should be obtained through a

MTA.

Figure Legends

Fig 1. SARS-CoV-2 antibody responses are sustained and are predominantly focused on Spike. (**A**) The first wave of Australian infections were from D614 and D614G Spike, and the S477N/D614G Spike variant emerged during the second wave. Convalescent patient sera from ADAPT (first and second waves) and LIFE (first wave) were examined for SARS-CoV-2 antibodies. Mean time and range of PCR-positivity (red), and dates of first and last sample collection (blue) are shown. Seropositive patients with at least three weeks between first and last samples were examined over time. (**B**) 96-98% (grey) of patients were Spike IgG+. Most ADAPT patients had stable levels overtime, whereas most of LIFE Spike IgG levels

decreased. No patients seroreverted. **(C)** 81-91% (grey) were Spike IgM+, most had

decreasing levels over time and Spike IgM+ individuals started with and maintained low IgM

levels. **(D)** 54-57% (grey) of sera were Membrane IgG+, and most ADAPT had stable levels,

whereas a larger proportion of LIFE had decreasing levels. **(E)** 78% of sera were

Nucleocapsid IgG+, most were stable in ADAPT, whereas most decreased in LIFE. Loess

curves with 95% confidence intervals are shown.

Fig 2. **Viral neutralization and inhibition of viral-cell fusion are strongly correlated to**

Spike antibody titers, and sustained overtime. **(A)** 68-82% of convalescent sera inhibited

virus-cell fusion, whereas 88-94% sera neutralized live authentic SARS-CoV-2. **(B)** ~75% of

virus particles were SARS-CoV-2 Nucleocapsid- and RNA-positive (overlay, yellow) **(C)** All

but one ADAPT patient had stable responses over time, whereas most LIFE donors (82%)

had a decreased virus-cell fusion over time, with the majority (71%) exhibiting a single-phase

decay. **(D)** In sera capable of viral neutralization, most ADAPT sera were stable (89%),

whereas most LIFE sera (56%) had a decreased score over time, with the majority (75%)

exhibiting a single-phase decay. Serum curves unable to be fitted were classified as

undetermined. **(E)** Spike IgG and IgM levels were correlated to inhibition of virus-cell fusion

and neutralization scores. R^2 values are shown and * indicates significance.

Fig 3. The antibody responses of patients with more severe COVID-19 disease have

broader SARS-CoV-2 polyantigenicity. **(A)** ~ half of patients (49-55%) had broad SARS-

CoV-2 antibodies (blue). Some had responses to two antigens (light and dark green), and a

few reacted to Spike only (red). **(B)** Hospitalized patients were more likely to have broad

SARS-CoV-2 polyantigenic immunoreactivity, whereas patients with only Spike reactivity

exhibited mild-moderate symptoms. **(C)** Hospitalized patients exhibited higher Spike IgG,

IgM, Membrane IgG, and Nucleocapsid IgG levels. High virus-cell fusion inhibition and

neutralization titers were observed in patients with broad polyantigenic immunoreactivity **(D)**

and in hospitalized patients **(E).** Older males were more likely to present with broader

polyantigenic immunoreactivity **(F)**, higher virus-cell fusion inhibition **(G),** and

neutralization scores **(H)**. Younger females were more enriched in mild to moderate disease

severity, with narrow antigenicity **(F)**, and lower virus-cell fusion inhibition **(G)** and

neutralization scores **(H)**.

Fig 4. High and elite responders are discriminated with SARS-CoV-2-permissive cells.

(A) Patients with high and robust SARS-CoV-2 responses were more likely male,

hospitalised (left), and of older age (right). **(B)** Low and high responders to SARS-CoV-2

showed limited neutralization in hyperpermissive HekAT24 clonal cells. Permissiveness is

indicated by +. **(C)** Only Elite responders showed neutralization in HekAT24 cells. **(D)**

Serum titration curves from an Elite responder (blue) showed IgG and IgM levels greater

than low (red) and high (green) responders, and incredibly high neutralization titers

(≥10,000) that decreased and stabilized at high levels (≥1280). **(E)** The elite donor

demonstrated stable high Spike IgG, but the early decrease in viral neutralization was parallel

to IgM decline before stabilization (at high titer).

Fig 5. SARS-CoV-2 antibody responses show evasion to emerging Spike variants. (A)

Most patients had broad recognition of Spike variants (blue), whereas a smaller group had

restricted Spike variant recognition and did not have a strong immunoreactivity to D614G

Spike (red). Patients with reduced binding to D614G Spike had lower virus-cell fusion and

neutralization scores **(B)**, and presented with less broad polyantigenic SARS-CoV-2

recognition **(C)**. **(D)** D614G Spike-binding sera had greater inhibition of D614G Spike-

pseudotyped virus-cell fusion. **(E, F)** In Australia, D614G Spike was the predominant variant

during the first wave and acquired additional mutations during the second wave (S477N,

V1068F). **(G)** All patients had decreased immunoreactivity to S477N/D614G and

S477N/D614G/V1068F Spike, while V1068F did not have an additive effect. The level of

decrease binding was irrespective of the virus that infected patients during the second wave.

Tables

Table 1. Demographics of the convalescent SARS-CoV-2 ADAPT and LIFE cohorts

	ADAPT		LIFE
Timeline*	First wave	Second wave	First wave
RT-PCR confirmed patients *n*	83	17**	159
Serial samples *n*	166	-	645
Gender *Male:female (ratio)*	35:48 (0.7)	6:11 (0.5)	72:79 (0.9)
Age at RT-PCR positivity *Median years (IQR, min, max)*	48 (35-59, 20, 79)	44 (34-64)	51 (30-63, 19, 78)
Days after RT-PCR positivity at first sample collection *Median days (IQR, min, max)* *Mean days (SD)*	71 (64-86, 36, 122) 74 (16)	31 (26-39, 21, 47) 33 (8)	59 (52-67, 33, 100) 61 (12)
Days after RT-PCR positivity at latest sample collection *Median days (IQR, min, max)* *Mean days (SD)*	118 (115-132, 114, 139) 123 (12)	n/a	95 (77-126, 55, 205) 127 (39)
Disease severity at acute infection***			
Non-hospitalized *n (% total)*	73 (88)	17 (100)	145 (95)#
Mild *n (%)*	31(42)	8 (47)	n/a
Moderate *n (%)*	42 (58)	9 (53)	n/a
Hospitalized *n (%)*	10 (12)	0 (0)	7 (5) [‡]
Admitted to ICU *n (%)*	3 (30)	0 (0)	-

*Timeline according to Fig. 1A

**n=8 infected and PCR-positive for S477N/D614G, n=9 infected with and PCR-positive for

S477N/D614G/V1068F

***Non-hospitalized Mild: community-managed with minor, mostly upper respiratory tract

viral symptoms including sore throat, rhinorrhoea, headache, and anosmia/ageusia. Non-

hospitalized Moderate: community-managed with fever/chills and one, or ≥ 2 of the

following organ-localizing symptoms; cough, haemoptysis, shortness of breath, chest pain,

nausea/vomiting, diarrhoea, or altered consciousness/confusion. Hospitalized: inpatient ward

care. Hospitalized and admitted to intensive care unit (ICU): care in the ICU for acute

respiratory distress syndrome.

no data on disease severity or symptoms were collected in LIFE.

‡ information on hospitalization was self-reported in LIFE.

Table 2. Comparison of the sensitivity of SARS-CoV-2 antibody detection assays.

	ADAPT n=166 samples		LIFE			
			All samples n=645		First and last samples n=302*	
	Positive samples, n (%)	Sensitivity % (95% CI)	Positive samples, n (%)	Sensitivity % (95% CI)	Positive samples, n (%)	Sensitivity % (95% CI)
Spike IgG						
Flow cytometry assay	162 (98)	98 (94-99)	645	n/a	302	n/a
Euroimmun	121 (73)	73 (65-80)	n/a	n/a	n/a	n/a
Spike IgM						
Flow cytometry assay	127 (77)	76** (70-83)	608 (94)	94[^§] (92-96)	276 (91)	91[^§] (87-94)
S1/S2 Spike IgG						
DiaSorin Liason SARS-CoV-2 S1/S2 IgG assay	134 (81)	81 (74-86)	n/a	n/a	n/a	n/a
Nucleocapsid IgG						
Abbott Architect SARS-CoV-2 assay	116 (70)	70 (62-77)	472 (73)	73 (70-77)	222 (74)	74 (68-78)
Euroimmun	121 (73)	73 (65-80)	n/a	n/a	n/a	n/a
Nucleocapsid/Spike IgG*	n/a	n/a	577 (89)	89 (87-92)	271 (90)	90 (86-93)
Membrane IgG						
Flow cytometry assay	87 (52)	52 (45-60)	n/a	n/a	173 (57)	57 (51-63)
Envelope IgG	4 (2)	2 (0.8-6)	0	0	0	0

754   *Only first and last samples of LIFE cohort were tested for Membrane IgG

755   **Sensitivity is influenced by IgM sero-reversion in 5 ADAPT and 14 LIFE donors

756   ***Positivity determined using a two-step clinical diagnostic testing with the Abbott

757   Architect SARS-CoV-2 Nucleocapsid IgG assay, followed by the Euroimmun Spike IgG

758   assay.

759
760

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5