

1 **Estimating the risk of incident SARS-CoV-2 infection among healthcare**
2 **workers in quarantine hospitals: the Egyptian example**

3
4 *Sofía Jijón*^{1,2†} (PhD), *Ahmad Al Shafie*^{3†} (MSc), *Essam Hassan*⁴ (MD), *EMAE-MESuRS*
5 *working group on nosocomial SARS-CoV-2 modeling*, *Laura Temime*^{1,2} (PhD), *Kévin*
6 *Jean*^{1,2,5*} (PhD), *Mohamed El-Kassas*^{3*} (MD)

7
8 ¹ Laboratoire Modélisation, épidémiologie et surveillance des risques sanitaires (MESuRS),
9 Conservatoire national des Arts et Métiers, Paris, France

10 ² Unité PACRI, Institut Pasteur, Conservatoire national des Arts et Métiers, Paris, France

11 ³ Endemic Medicine Department, Faculty of Medicine, Helwan University, Cairo, Egypt

12 ⁴ Tropical Medicine Department, Faculty of Medicine, Fayoum University, Fayoum, Egypt

13 ⁵ MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease
14 Epidemiology, Imperial College London, United Kingdom

15
16 [†] *Contributed equally to the work*

17 ^{*} *Contributed equally to the work*

18
19 **Corresponding author:**

20 Sofía Jijón, sofia.jijon_alban@sorbonne-universite.fr

21 Current address: Sorbonne Université, Campus Pierre et Marie Curie, 4 Place Jussieu, 75005
22 Paris, France.

23 **Running title:** SARS-CoV-2 infection risk among HCW in quarantine hospitals

24 **Word count.** Abstract: 297 words; Main text: 3240 words (3964 including captions and tables)

25 **ABSTRACT**

26 In response to the COVID-19 epidemic, Egypt established a unique care model based on
27 quarantine hospitals where only externally-referred confirmed COVID-19 patients were
28 admitted, and healthcare workers resided continuously over 1- to 2-week working shifts.
29 While the COVID-19 risk for HCWs has been widely reported in standard healthcare settings,
30 it has not been evaluated yet in quarantine hospitals.

31 Here, we relied on longitudinal data, including results of routine RT-PCR tests, collected
32 within three quarantine hospitals located in Cairo and Fayoum, Egypt. Using a model-based
33 approach that accounts for the time-since-exposure variation in false-negative rates of RT-
34 PCR tests, we computed the incidence of SARS-CoV-2 infection among HCWs. Over a total
35 follow-up of 6,064 person-days (PD), we estimated an incidence rate (per 100 PD) of 1.05
36 (95% CrI: 0.58–1.65) at Hospital 1, 1.92 (95% CrI: 0.93–3.28) at Hospital 2 and 7.62 (95%
37 CrI: 3.47–13.70) at Hospital 3. The probability for an HCW to be infected at the end of a
38 shift was 13.7% (95% CrI: 7.8%–20.8%) and 23.8% (95% CrI: 12.2%–37.3%) for a 2-week
39 shift at Hospital 1 and Hospital 2, respectively, which lies within the range of risk levels
40 previously documented in standard healthcare settings, whereas it was >3-fold higher for a 7-
41 day shift at Hospital 2 (42.6%, 95%CrI: 21.9%–64.4%). Our model-based estimates unveil a
42 proportion of undiagnosed infections among HCWs of 46.4% (95% CrI: 18.8%–66.7%),
43 45.0% (95% CrI: 5.6%–70.8%) and 59.2% (95% CrI: 34.8%–78.8%), for Hospitals 1 to 3,
44 respectively.

45 The large variation in SARS-CoV-2 incidence we document here suggests that HCWs from
46 quarantine hospitals may face a high occupational risk of infection, but that, with sufficient
47 anticipation and infection control measures, this risk can be brought down to levels similar to
48 those observed in standard healthcare settings.

49
50 **Keywords.** COVID-19; SARS-CoV-2; Nosocomial infection; Healthcare workers;
51 Quarantine hospitals.

52 **WHAT THIS PAPER ADDS**

53 **What is already known on this topic**

54 Previous studies conducted in standard care settings have documented that frontline
55 healthcare workers (HCWs) face high risk of COVID-19. Whether risk levels differ in
56 alternative care models, such as COVID-19 quarantine hospitals in Egypt where HCWs
57 resided in the hospital days and nights for various durations, is unknown.

58 **What this study adds**

59 COVID-19 risk for HCWs in quarantine hospitals varies substantially between facilities, from
60 risk levels that are in the range of those documented in standard healthcare settings to levels
61 that were approximatively 3 times higher.

62 **How this study might affect research, practice or policy**

63 With sufficient anticipation and infection control measures, occupational COVID-19 risk for
64 HCWs working in quarantine hospitals can be brought down to levels similar to those
65 observed in standard healthcare settings.

66 INTRODUCTION

67 Healthcare settings have faced multiple challenges related to the currently ongoing severe
68 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic. They have notably had
69 to deal with influxes of infected patients across successive epidemic waves while controlling
70 the risk of nosocomial spread to other patients and staff. As a result, healthcare workers
71 (HCWs) have been a population of interest in terms of risk assessment and control measure
72 implementation [1].

73 At early stages of the pandemic, Egypt was identified as one of the African countries most
74 vulnerable to SARS-CoV-2 importation [2]. On February 14, 2020, Egypt reported the first
75 confirmed case of coronavirus disease 2019 (COVID-19) in Africa and remained among the
76 five African countries most assigned by the COVID-19 epidemic up to the end of 2020 [3].
77 From February 14 to August 31, 2020 (the period to which we refer to as the first wave), the
78 COVID-19 epidemic in Egypt resulted in about 9,700 confirmed infections and 5,500 deaths
79 reported nationally [4]. These numbers most certainly reflect underreporting of the real
80 number of infections, due to the high proportion of asymptomatic SARS-CoV-2 infection
81 [5,6] and limited testing.

82 To mitigate the high risk of SARS-CoV-2 spreading, Egypt established a unique care model
83 under supervision from the World Health Organization, whereby specific hospitals were
84 assigned as quarantine hospitals for patients with COVID-19, and where dedicated medical
85 teams stayed in the hospital days and nights during working shifts of various durations [7].

86 While the quarantine-hospital strategy has the potential to be highly efficient in terms of
87 patient care, as well as in limiting the potential spread of the virus from hospitals into the
88 community, its impact on infection risk for HCWs remains understudied. One theoretical
89 modelling study assessing healthcare working force organization found that alternating HCW
90 teams by 1-week periods may reduce the overall number of infected HCWs [8]. On the other
91 hand, to the best of our knowledge, previous epidemiological studies focusing on the infection
92 risk faced by Egyptian HCWs have been conducted in non-quarantine settings exclusively [9–
93 11]. Yet, the quarantine strategy was adopted again to face the second wave of the COVID-19
94 epidemic at the national level, and could be adopted for future epidemic waves in Egypt
95 and/or in other countries [12].

96 Here, we used mathematical modelling to estimate the risk of SARS-CoV-2 infection among
97 HCWs participating in quarantine-hospital interventions, relying on detailed longitudinal data
98 collected in three Egyptian healthcare facilities during the first wave of the COVID-19
99 epidemic.

100 **METHODS**

101 *Study settings*

102 Data was collected within three Egyptian hospitals (hereafter denoted by Hosp1, Hosp2 and
103 Hosp3) located in Cairo (Hosp1 and Hosp3), and Fayoum (middle Egypt, Hosp2), that were
104 temporarily transformed into quarantine hospitals during the first COVID-19 wave. During
105 the quarantine-organization period (Hosp1: March 14th to August 1st, 2020; Hosp2: April 1st
106 to July 31th; Hosp3: June 6th to July 11th, 2020), only externally-referred COVID-19
107 confirmed patients were admitted to these hospitals for medical care. Multidisciplinary
108 medical teams fully dedicated to COVID-19 patient management worked in total isolation,
109 organized by shifts (Hosp1: 2-week shifts; Hosp2: 1 to 2-week shifts; Hosp3: 1-week shifts).
110 During their shift, HCWs were assigned either to intensive care units (ICU) or non-ICU.
111 HCWs were screened for SARS-CoV-2 infection before starting a working shift using rapid
112 serological IgM/IgG antibody tests (Artron laboratories Burnaby, Canada; sensitivity: 83.3%,
113 specificity: 100% [13]). Only HCWs with no SARS-CoV-2 antibodies were allowed to start
114 working in the hospital; except for the last shift in Hosp3, where staff recruitment relied
115 exclusively on HCWs who had been previously infected (positive serological tests), and were
116 additionally tested using reverse transcription-polymerase chain reaction (RT-PCR) on
117 nasopharyngeal swabs, obtaining negative results. During the working shifts, HCWs were
118 tested for SARS-CoV-2 infection using RT-PCR tests: i) routinely at the end of the shift, ii)
119 upon symptoms, and iii) in case of outbreak suspicion (>2 positive tests among HCWs).
120 HCWs testing negative at the end of the shift were then released for self-isolation at home for
121 two weeks. HCWs testing positive before or during their shift self-isolated at home in the case
122 of presenting no or mild symptoms, or were admitted to the same quarantine hospital for
123 medical care in the case of presenting moderate to severe symptoms.

124 *Observed risk of SARS-CoV-2 infection*

125 Crude incidence rates were obtained from the number of incident SARS-CoV-2 infections
126 observed (ie, diagnosed during or at the end of a shift) among HCWs screened as seronegative
127 for SARS-CoV-2 before their working shift. Incidence rate ratios were obtained through a
128 Poisson regression adjusted by hospital and by type of care unit (ICU/ non-ICU). In addition,
129 we computed the attack rates over each working shift, based on the observed infections
130 among HCWs having a negative screening test at the beginning of the working shift.

131 Observed per-shift attack rates of were considered to be healthcare-associated outbreaks
132 among HCWs.

133 *Model-based estimates of the incidence rate of SARS-CoV-2 infection*

134 An important feature arising from COVID-19 surveillance within quarantine hospitals is
135 potential right-truncation of data: HCWs infected a short time before the end of a shift are
136 likely to remain undiagnosed despite the systematic testing at the end of shifts, especially if
137 the test is performed early in the incubation period. We thus developed the following
138 mathematical model to estimate the incidence rate of both diagnosed and undiagnosed SARS-
139 CoV-2 infections.

140 We simulated the daily number of incident infections among HCWs, $X_{k,i}^A$ based on an
141 unobserved binomial process:

$$X_{k,i}^A \sim \text{Bin}(n = S_{k,i}^A, p = \lambda)$$

142 where k denotes the working shift, i denotes the day within the shift, A is the hospital unit of
143 HCWs' assignments ($A \in \{\text{ICU}, \text{Non-ICU}\}$), $S_{k,i}^A$ denotes the number of susceptible HCWs
144 assigned to A , at the beginning of the i -th day of shift k and λ denotes the constant daily
145 probability of SARS-CoV-2 infection faced by HCWs. The number of susceptible HCWs was
146 updated on a daily basis:

$$S_{k,i+1}^A = S_{k,i}^A - X_{k,i}^A$$

147 To model right-truncation in the surveillance data, we simulated an observation process
148 representing the systematic RT-PCR testing of HCWs at the end of each shift, accounting for
149 documented variation in test sensitivity as a function of time since infection [14]. The period
150 of time since infection was computed as $j_k - i$, where j_k denotes the day at the end of shift k
151 (i.e., the day when testing was performed) and i denotes the day of infection. Then, the daily
152 number of incident infections eventually diagnosed at the end of the -th shift was given by

$$I_{k,i}^A = X_{k,i}^A \theta_{j_k - i}$$

153 where $\theta_{j_k - i}$ denotes the probability of testing positive to RT-PCR, $j_k - i$ days after
 154 infection.

155 We used a Bayesian Markov Chain Monte-Carlo (MCMC) approach to estimate the
 156 parameter λ that best fitted the observed number of infections. We assumed that λ was
 157 constant over the study period. We considered a non-informative uniform prior distribution
 158 $\sim U(0,1)$ for λ and the following binomial likelihood function:

$$L(\lambda | \text{data}) = \prod_A \left(\frac{\sum_{k,i} N_{k,i}^{S,A}}{\sum_{k,i} N_{k,i}^{I,A}} \right) \left(\frac{\sum_{k,i} I_{k,i}^A}{\sum_{k,i} S_{k,i}^A} \right)^{\sum_{k,i} N_{k,i}^{I,A}} \left(1 - \frac{\sum_{k,i} I_{k,i}^A}{\sum_{k,i} S_{k,i}^A} \right)^{\sum_{k,i} N_{k,i}^{S,A} - \sum_{k,i} N_{k,i}^{I,A}},$$

159 where $\sum_{k,i} N_{k,i}^{S,A}$ and $\sum_{k,i} N_{k,i}^{I,A}$ denote the total number of susceptible HCWs (denoted by the
 160 superscript S) and the total number of HCW infected and diagnosed with COVID-19
 161 (denoted by the superscript I), observed in hospital unit A , during the study period,
 162 respectively. The model was thus fitted to the total number of observed infections in each
 163 hospital unit.

164 Medians and 95% credibility intervals (CrI) for λ were computed from posterior samples
 165 obtained after 10,000 model runs. The chains were visually inspected for convergence. For
 166 each hospital, we further estimated the probability for a susceptible HCW to be infected at the
 167 end of a shift as

$$168 \quad P(\text{Infected at shift end}) = 1 - (1 - \lambda)^d,$$

169 where d is the length of the working shift, in days.

170 Finally, to assess the proportion of undetected infections among HCWs, we ran our model on
 171 each hospital, using the posterior distributions for λ , to obtain an estimate for the real number
 172 of SARS-CoV-2 infections in each hospital (both diagnosed and undiagnosed), over the study
 173 period.

174 The numeric implementation of the model was coded in R 4.0.3 [15], using the FME package
 175 [16] for parameter estimation.

176

177 *Ethical approval*

178 This study was approved by the Research Ethics Committee (REC) of the Central Directorate
179 of Research and Health Development and Reviews at the Egyptian Ministry of Health and
180 Population (Serial: 25-2020/16), and by REC for human subject research at the Faculty of
181 Medicine, Helwan University (Serial: 50-2020).

182

183 **RESULTS**

184 The study period covered ten 2-week shifts in Hosp1, nine ~2-week shifts in Hosp2 and five
185 1-week shifts in Hosp3. The mean (min–max) number of HCWs per shift was 46 (34–63) in
186 Hosp1, 15 (5–26) in Hosp2 and 19 (16–20) in Hosp3 (Table 1). Over a total follow-up of
187 8,733 person-days (PD), 54 SARS-CoV-2 infections were observed (ie., diagnosed by RT-
188 PCR) among 722 HCWs showing no evidence of SARS-CoV-2 antibodies at the beginning of
189 their shifts, across the three hospitals. This represented an overall incidence rate of 0.62 (95%
190 CI: 0.45–0.78) diagnosed SARS-Cov-2 infections per 100 PD (Table 2). A significantly
191 higher incidence rate was observed for HCWs working at Hosp3 as compared to the other two
192 hospitals; and HCWs working in non-ICU units tended to be more at risk than HCWs
193 working in ICU, though this difference was not statistically significant (Table 2).

194

195 **Table 1. Hospital characteristics.**

196 Quarantine organization period, location and mean number of HCWs and patients by hospital (Hosp1, Hosp2
197 and Hosp3). Abbreviations: HCWs = Healthcare workers, PD = person-days.

	Hosp1	Hosp2	Hosp3
Quarantine organization period	March 14 – August 1	April 1 – July 31	June 6 – July 11
Location	Cairo	Fayoum	Cairo
Mean daily number of patients (min–max)	62 (0–108)	37 (0–103)	8 (0–20)
Mean number of HCWs per shift (min–max)	46 (34–63)	15 (5–26)	19 (16–20)
Shift duration	14 days	7–14 days	7 days

198

199

200

201 **Table 2. Observed risk of SARS-CoV-2 infection among HCWs, by hospital and by hospital unit.**
 202 A total of 54 infections were observed (ie., diagnosed before or at the end of a shift) over 8,733 person-days, in
 203 the three hospitals (Hosp1, Hosp1 and Hosp3). Crude rates of SARS-CoV-2 infection among HCWs by hospital
 204 and by hospital unit: intensive and non-intensive care units and 95% confidence intervals. A Poisson regression
 205 adjusted by hospital and by hospital unit was performed on the observed infections to obtain the incidence rate
 206 ratios and 95% confidence intervals. Abbreviations: CI = confidence interval, ICU = intensive care unit, IRR =
 207 incidence rate ratio, PD = person-days.

		Observations		Crude rates		Adjusted Poisson Regression		
		Events	PD	Rate per 100 PD	95% CI	IRR	95% CI	<i>p</i>
Hospital	Hosp1	28	6258	0.45	0.28–0.61	1	ref	–
	Hosp2	11	1808	0.61	0.25–0.97	1.49	0.74–2.87	0.28
	Hosp3	15	667	2.25	1.1–3.39	5.59	3.13–10.01	<0.001
Hospital Unit	ICU	15	2628	0.57	0.28–0.86	1	ref	–
	Non-ICU	39	6105	0.64	0.44–0.84	1.49	0.85–2.59	0.16

208

209 Healthcare-associated outbreaks among HCWs (observed attack rates of) represented ~70%
 210 (38/54) of all SARS-CoV-2 infections observed over the study period (Figure 1). Two of
 211 these outbreaks occurred in Hosp2 with 30% (3/10) and 24% (5/21) of HCWs being infected
 212 over two different working shifts. An outbreak where 36% of HCWs were infected (16/44)
 213 occurred in Hosp1 around the same time period. Two outbreaks took place later on in Hosp3,
 214 leading to, respectively, 30% (6/20) and 40% (8/20) of susceptible HCWs being infected over
 215 two different shifts. Of note, each outbreak resulted in infections in both ICU and non-ICU.
 216 For the last shift in Hosp3, staff recruitment relied exclusively on HCWs who had been
 217 infected during the first or the second shifts. This likely prevented HCWs in Hosp3 from
 218 becoming infected during the last shift (cf. Figure 1).




Figure 1. Per-week number of infections by hospital unit.

For a given working shift, healthcare workers were assigned to either the intensive care unit (ICU, yellow) or a non-ICU (blue). Dates are given in weeks, denoted by the prefix W followed by the week number of year 2020. Hosp1 was established as a quarantine hospital from March 14th to August 1st (W11 to W31), Hosp2 from April 1st to July 31st (W14 to W31) and Hosp3 from June 6th to July 11th (W23 to W28). In Hosp1 and Hosp2, most infections occurred in non-ICU units. Per-shift attack rates are shown above the bars. Of note, only previously infected HCWs worked during the last two shifts (W27 and W28) in Hosp3.

219

220

221

222

223

224

225

226

To estimate the mean incidence rate of both observed and unobserved infections in each hospital, we ran our model considering a constant risk, λ , over the whole study period, in each hospital unit. The chains for λ , for the three hospitals, are depicted in the Supplementary Figure 1. We estimated significantly different risk levels between the three hospitals (Kolmogorov-Smirnov test; p-value < 0.001), with an incidence rate of 1.05 (95% CrI: 0.58–1.65) per 100 PD in Hosp1, 1.92 (95% CrI: 0.93–3.28) per 100 PD in Hosp2 and 7.62 (95% CrI: 3.47–13.70) per 100 PD in Hosp3.

227

228

229

230

231

232

233

234

The probability for a HCW to be infected at the end of a working shift was estimated at 13.7% (95% CrI: 7.8%–20.8%) and 23.8% (95% CrI: 12.2%–37.3%) for 2-week shifts at Hosp1 and Hosp2, respectively, whereas a much higher probability of 42.6% (95% CrI: 21.9%–64.4%) was found for a 7-day shift at Hosp3 (see Table 3). These probabilities become 7.1% (95% CrI: 4.0%–11.0%) and 12.7% (95% CrI: 6.3%–20.8%), for Hosp1 and Hosp2, respectively, if 7-day shifts are considered as well, for easier comparison (Table 3). Given the notable differences between hospitals (Figure 1), we did not assess infection risk by care unit across hospitals, but within each hospital.

235

236

237

238

239

The posterior distributions for the incidence rate λ yield the following model-based median numbers of total SARS-CoV-2 infections: 59 (95% CrI: 45–74) at Hosp1, 31 (95% CrI: 14–52) at Hosp2 and 41 (95% CrI: 19–63) at Hosp3 (Table 3), which unveils a proportion of undiagnosed infections among HCWs of 46.4% (95% CrI: 18.8%–66.7%), 45.0% (95% CrI: 5.6%–70.8%) and 59.2% (95% CrI: 34.8%–78.8%), respectively.

240 We further ran our model on each hospital unit, on each hospital, which confirmed a notable
 241 difference between Hosp1–2 and Hosp3: our results suggest that HCWs working in Hosp1
 242 and Hosp2 face a higher risk when assigned to non-ICU, as compared to those assigned to the
 243 ICU; whereas HCWs in the ICU of Hosp3 face higher risk (see Fig. 1 and Table 3).

244 **Table 3. Model-based estimates of all SARS-CoV-2 infections**

245 The model-based estimates of the incidence rate, the probability of infection for a 7-day working shift and the
 246 estimated number of SARS-CoV-2 infections in each hospital, over the study period. Running our model for
 247 each type of care unit (ICU and non-ICU) in each hospital unveils the difference in the risk faced by HCWs
 248 between the three hospitals of our study. Indeed, HCWs working in Hosp1 and Hosp2 face a higher risk when
 249 assigned to non-ICU, as compared with those working in the ICU; whereas HCWs in Hosp3 face a slightly
 250 higher risk when assigned to the ICU. All estimates concern both diagnosed and undiagnosed infections (all),
 251 unless otherwise stated (diagnosed). Abbreviations: CrI = credibility interval, ICU = intensive care unit, PD =
 252 person-days.

	Median estimated incidence rate per 100 PD (95% CrI)			Probability of infection for a 7- day shift (95% CrI)	Median number of SARS-CoV-2 infections (95% CrI)		
	ICU	Non-ICU	Overall		All	Diagnosed	Proportion of undiagnosed infections
Hosp1	0.59 (0.11–1.61)	1.27 (0.68–2.01)	1.05 (0.58–1.65)	7.1% (4.0%–11.0%)	59 (45–74)	31 (15-51)	46.4% (18.8%–66.7%)
Hosp2	1.33 (0.33–3.33)	2.75 (1.12–5.02)	1.92 (0.93–3.28)	12.7% (6.3%–20.8%)	31 (14–52)	17 (7-31)	45.0% (5.6%–70.8%)
Hosp3	9.37 (3.48–19.66)	7.12 (2.06–18.71)	7.62 (3.47–13.70)	42.6% (21.9%–64.4%)	41 (19–63)	16 (6-30)	59.2% (34.8%–78.8%)

253

254

255 **DISCUSSION**

256 Here, we studied the risk of incident SARS-CoV-2 infection among HCWs residing days and
257 nights in quarantine hospitals, relying on detailed longitudinal data collected during the first
258 wave of COVID-19 in Egypt. We found that most diagnosed infections (70%) occurred
259 during what we defined as healthcare-associated outbreaks (as compared to isolated
260 infections). We observed high variability in nosocomial incidence, ranging from 0.45 to 2.25
261 infections per 100 PD across the three hospitals. Using a model-based approach, we
262 estimated the risk of both diagnosed and undiagnosed SARS-CoV-2 infections among HCWs,
263 by hospital and by hospital unit (ICU and non-ICU). We further estimated that a substantial
264 proportion of infections may have remained undetected, ranging from 45.0% to 59.2% across
265 the three hospitals.

266 Our study design has several limitations. First, screening HCWs using serological tests before
267 starting their working shifts may not detect active infections at the time of testing. Indeed,
268 antibodies may be detected by serological tests in less than 40% of infected individuals within
269 7 days since symptoms onset, reaching higher detection levels at day 16 after symptom onset
270 [17]. Hence, serological screening may allow recently infected HCWs –who are probably
271 infectious– to start a working shift. However, the Egyptian public health authorities' choice of
272 a protocol relying on serological tests before working shifts was mainly driven by material
273 constraints. Indeed, the use of rapid serological tests as a diagnostic tool was and remains of
274 particular interest in limited-resource contexts due to their lower cost, the minimal equipment
275 required, and faster results, as compared with RT-PCR tests [18]. Second, the assumption of a
276 constant overall risk of infection among HCWs disregards dynamic fluctuations in risk driven
277 by variation in the number of HCWs and/or patients in the hospitals, as well as the frequency
278 and nature of HCWs' contacts during their working shifts. This simplification was made in
279 order to estimate average transmission rate first; fine reproduction of transmission dynamics
280 were left for future work. Third, despite the quarantine strategy being adopted nationally, the
281 differences between the organizations of each of the three hospitals (working shifts lengths,
282 quarantine-organization period) resulted in some difficulties in comparing the risk estimations
283 across the three hospitals. Data from other hospitals that have adopted a quarantine strategy
284 will be beneficial to better understand occupational risks and eventually study the
285 performance of such a strategy in comparison to standard care settings.

286 We found a ~5-fold higher observed risk for Hosp3 as compared with the risk found in
287 Hosp1; whereas the risk of infection at Hosp2 was just slightly higher. The higher risk of
288 SARS-CoV-2 infection we report for Hosp3 and the higher risk faced by HCWs assigned to
289 the ICU of Hosp3 may be partly explained by the short period over which this hospital
290 adopted the quarantine organization, which coincided with the country's highest epidemic
291 activity of the first wave [4]. This may have led to a high proportion of severe and thus highly
292 contagious COVID-19 patients referred to quarantine hospitals together with a higher
293 workload for HCWs. Contrarily, Hosp1 adopted the quarantine organization in mid-March
294 2020, and thus experienced several weeks of lower epidemic intensity that may have
295 improved preparedness for intense COVID-19 activity and implementation of infection
296 control measures for invasive procedures (e.g., intubation in the ICU).

297 Our model-based approach, accounting for false-negativity rates of testing, as well as the
298 right-truncation of our data, allowed us to estimate the risk of both diagnosed and
299 undiagnosed SARS-CoV-2 infection among HCWs, by hospital and by hospital care unit. We
300 estimated that a rigorous quarantine organization with systematic testing of HCWs at the end
301 of their working shifts captures only about half of all HCW infections (depending on the
302 length of the working shift). Of note, the higher proportion of infections that remained
303 undetected in Hosp3 (59.2%, versus 46.4% and 45.0% in Hosp1 and Hosp2, respectively)
304 may be explained by the shorter duration of the working shifts (7 days), which may result in
305 false-negative tests for HCWs infected a short time before the end of their shift. Moreover,
306 these results suggest that, in the absence of the quarantine-hospital organization and, more
307 specifically, in the absence of systematic testing at the end of quarantine working shifts, an
308 even larger proportion of infections among HCWs may remain undetected, thus putting
309 HCWs' close contacts at risk of infection [19] and, consequently, putting themselves at risk
310 for adverse psychological symptoms [20].

311 The model-based estimates we found for the SARS-CoV-2 infection risk in Hosp1 and Hosp2
312 are consistent with infection point-prevalence reported in earlier studies performed in non-
313 quarantine Egyptian hospitals (varying from 4.2% to 14.3%) and with incidence estimates
314 reported among front-line HCWs in the UK (13% infection rate after one month) [10,11,21].
315 For comparison, a summary of previous results obtained early in the Covid-19 pandemic
316 and/or specifically in the Egyptian context is presented in Table 4. Moreover, our results
317 suggest that HCWs assigned to non-ICU face a higher risk than those assigned to the ICU, in

318 Hosp1 and Hosp2, which is in line with what was observed in previous studies addressing the
319 occupational risk of SARS-CoV-2 infection for HCWs in non-quarantine settings [22].
320 Overall, our findings on Hosp1 and Hosp2 suggest that, providing sufficient preparedness,
321 HCWs working in quarantine hospitals may not face a higher infection risk, and thus
322 highlight the benefits of implementing a quarantine-hospital strategy.

323

324 **Table 4. Our results in a context.**

325 Our estimation of the per-shift probability of infection in Hosp1–3, along with estimates of the point-prevalence
326 in other Egyptian studies and the observed cumulative incidence rate among HCWs in an English study.

Indicator	Estimate (%)	Context	Study
Per-shift probability of infection (95% CrI)	12.8 (7.6–19.5)	Hosp1 (14-days shift)	Our results
	17.3 (7.5–30.7)	Hosp2 (14-days shift)	
	48.2 (23.8–74.5)	Hosp3 (7-days shift)	
Point-prevalence	4.2	Symptomatic and asymptomatic HCWs in 12 healthcare facilities (N=4040).	[23]
Point-prevalence	14.3	Asymptomatic HCWs in emergency departments of tertiary care facility (N=203).	[10]
Incidence rate	13.0	Front-line HCWs testing negative at enrollment.	[21]

327

328 A notable strength of our study lies in the specific nature of the quarantine hospital set-up:
329 because HCWs resided continuously in the hospitals over their entire working shifts, we were
330 able to exclude risk of infection in the community and specifically quantify the nosocomial
331 risk for HCWs. Conversely, results reported from non-quarantine hospitals worldwide are
332 generally unable to distinguish between the nosocomial vs. community risk of infection.
333 However, as most studies conducted in healthcare settings, we were unable to distinguish
334 between patient-to-HCW and HCW-to-HCW routes of transmission. A previous study on
335 several nosocomial Covid-19 healthcare-associated outbreaks in Germany reported that
336 HCW-to-HCW transmission could represent an outsized risk as compared to the one due to
337 infected patients [24]. Still, assessing the relative contribution between patient-to-HCW
338 versus HCW-to-HCW SARS-Cov-2 transmission was left for future work. However, it is
339 worth noting that during the outbreaks observed in our study, infections occurred in different
340 care units, which may prime HCW-to-HCW transmission, rather than simultaneous
341 independent events of patient-to-HCW transmission. In early stages of the pandemic, personal
342 protective equipment (PPE) placed a focus on the risk induced by patients, but could have

343 contributes to underestimate the risk of infection from –infected and undiagnosed–
344 colleagues. This may be especially true for quarantine hospitals in which HCWs share resting
345 and conviviality rooms for longer times than in standard care settings. Adapting the use of
346 PPE and social distancing between colleagues while ensuring sufficient social interaction and
347 support to maintain HCWs’ mental health thus constitutes a specific challenge for quarantine
348 hospitals [20].

349 Preventing nosocomial SARS-CoV-2 infection remains an urgent need, especially in settings
350 where preparedness does not match the risk faced by HCWs. The large variation in infection
351 risk we found between hospitals suggests that HCWs from quarantine hospitals may face a
352 high risk of infection, but that, with sufficient anticipation and infection control measures, this
353 risk can be brought down to levels similar to those observed in standard COVID-19 care
354 settings. A comprehensive assessment of quarantine hospital care models should also include
355 their impact on HCWs’ mental health as well as the potential benefits of earlier infection
356 diagnosis, which is likely to reduce further hospital, household and community transmission
357 [20].

358

359 **EMAE-MESuRS working group on nosocomial SARS-CoV-2 modeling**

360 Audrey Duval^{6,7,8}, Kenza Hamzi¹, Niels Hendrickx⁶, Kévin Jean^{1,2,5}, Sofía Jijón^{1,2}, Ajmal

361 Oodally^{6,7,1}, Lulla Opatowski^{6,7}, George Shirreff^{6,7,8}, David RM Smith^{6,7,1}, Cynthia

362 Tamandjou⁷ and Laura Temime^{1,2}

363 ⁶ Institut Pasteur, Epidemiology and Modelling of Antibiotic Evasion (EMAE), Paris, France

364 ⁷ Université Paris-Saclay, UVSQ, Inserm, CESP, Anti-infective evasion and

365 pharmacoepidemiology team, Montigny-Le-Bretonneux, France

366 ⁸ IAME, UMR 1137, Université Paris 13, Sorbonne Paris Cité, France

367

368 **Declaration of interest**

369 The authors declare no conflicts of interest.

370

371 **Funding**

372 This study was supported by INSERM-ANRS (France Recherche Nord and Sud Sida-HIV

373 Hépatites, grant number ANRS-COV-19).

374

375 **Acknowledgments**

376 The authors thank all the dedicated HCWs who contributed to this study.

377 **REFERENCES**

- 378 1. Lai X, Wang M, Qin C, et al. Coronavirus Disease 2019 (COVID-2019) Infection
379 Among Health Care Workers and Implications for Prevention Measures in a Tertiary
380 Hospital in Wuhan, China. *JAMA Netw Open* **2020**; 3:e209666. Available at:
381 <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2766227>.
- 382 2. Gilbert M, Pullano G, Pinotti F, et al. Preparedness and vulnerability of African
383 countries against importations of COVID-19: a modelling study. *Lancet* **2020**;
384 395:871–877.
- 385 3. Salyer SJ, Maeda J, Sembuche S, et al. The first and second waves of the COVID-19
386 pandemic in Africa: a cross-sectional study. *Lancet* **2021**; 397:1265–1275. Available
387 at: [http://dx.doi.org/10.1016/S0140-6736\(21\)00632-2](http://dx.doi.org/10.1016/S0140-6736(21)00632-2).
- 388 4. WHO. Coronavirus Disease (COVID-19) Dashboard. Egypt. 2020; Available at:
389 <https://covid19.who.int/region/emro/country/eg>.
- 390 5. Buitrago-Garcia D, Egli-Gany D, Counotte MJ, et al. Occurrence and transmission
391 potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living
392 systematic review and meta-analysis. *PLOS Med* **2020**; 17:e1003346. Available at:
393 <https://dx.plos.org/10.1371/journal.pmed.1003346>.
- 394 6. Yanes-Lane M, Winters N, Fregonese F, et al. Proportion of asymptomatic infection
395 among COVID-19 positive persons and their transmission potential: A systematic
396 review and meta-analysis. *PLoS One* **2020**; 15:e0241536. Available at:
397 <https://dx.plos.org/10.1371/journal.pone.0241536>.
- 398 7. Medhat MA, El Kassas M. COVID-19 in Egypt: Uncovered figures or a different
399 situation? *J Glob Health* **2020**; 10. Available at:
400 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7303805>.
- 401 8. Sánchez-Taltavull D, Castelo-Szekely V, Candinas D, Roldán E, Beldi G. Modelling
402 strategies to organize healthcare workforce during pandemics: Application to COVID-
403 19. *J Theor Biol* **2021**; 523.
- 404 9. Meguid A, Talaat H, Shawky S, et al. SARS-CoV-2 infection among healthcare
405 workers of a gastroenterological service in a tertiary care facility. *Arab J Gastroenterol*
406 **2020**; 2020. Available at: <https://doi.org/10.1016/j.ajg.2020.07.005>.
- 407 10. Abdelmoniem R, Fouad R, Shawky S, et al. SARS-CoV-2 infection among
408 asymptomatic healthcare workers of the emergency department in a tertiary care
409 facility. *J Clin Virol* **2021**; 134:104710. Available at:
410 <https://doi.org/10.1016/j.jcv.2020.104710>.
- 411 11. Mostafa A, Kandil S, El-Sayed MH, et al. Universal COVID-19 screening of 4040
412 health care workers in a resource-limited setting: an Egyptian pilot model in a
413 university with 12 public hospitals and medical centers. *Int J Epidemiol* **2020**; :1–12.
414 Available at: [https://academic.oup.com/ije/advance-](https://academic.oup.com/ije/advance-article/doi/10.1093/ije/dyaa173/5936053)
415 [article/doi/10.1093/ije/dyaa173/5936053](https://academic.oup.com/ije/advance-article/doi/10.1093/ije/dyaa173/5936053).
- 416 12. Egypt reopens 21 coronavirus quarantine hospitals. 2020. Available at:

- 417 [https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-](https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-hospitals/)
418 [hospitals/](https://www.egyptindependent.com/egypt-reopens-21-coronavirus-quarantine-hospitals/). Accessed 21 November 2021.
- 419 13. Lassaunière R, Frische A, Harboe ZB, et al. Evaluation of nine commercial SARS-
420 CoV-2 immunoassays. *medRxiv* **2020**; :2020.04.09.20056325. Available at:
421 <http://medrxiv.org/content/early/2020/04/10/2020.04.09.20056325.abstract>.
- 422 14. Kucirka LM, Lauer SA, Laeyendecker O, Boon D, Lessler J. Variation in False-
423 Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-
424 CoV-2 Tests by Time Since Exposure. *Ann Intern Med* **2020**; 173:262–267.
- 425 15. R Core Team. R: A language and environment for statistical computing. 2020;
426 Available at: <https://www.r-project.org/>.
- 427 16. Soetaert K, Petzoldt T. Inverse modelling, sensitivity and monte carlo analysis in R
428 using package FME. *J Stat Softw* **2010**; 33:1–28.
- 429 17. Zhao J, Yuan Q, Wang H, et al. Antibody Responses to SARS-CoV-2 in Patients With
430 Novel Coronavirus Disease 2019. *Clin Infect Dis* **2020**; 71:2027–2034. Available at:
431 <https://academic.oup.com/cid/article/71/16/2027/5812996>.
- 432 18. Jiang J-C, Zhang Y. Serological antibody testing in the COVID-19 pandemic: their
433 molecular basis and applications. *Biochem Soc Trans* **2020**; 48:2851–2863. Available
434 at: [https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-](https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-antibody-testing-in-the-COVID-19)
435 [antibody-testing-in-the-COVID-19](https://portlandpress.com/biochemsoctrans/article/48/6/2851/226924/Serological-antibody-testing-in-the-COVID-19).
- 436 19. Krastinova E, Garrait V, Lecam M-T, et al. Household transmission and incidence of
437 positive SARS-CoV-2 RT-PCR in symptomatic healthcare workers, clinical course and
438 outcome: a French hospital experience. *Occup Environ Med* **2020**; :oemed-2020-
439 106866. Available at: <https://oem.bmj.com/lookup/doi/10.1136/oemed-2020-106866>.
- 440 20. Youssef N, Mostafa A, Ezzat R, Yosef M, El Kassas M. Mental health status of health-
441 care professionals working in quarantine and non-quarantine Egyptian hospitals during
442 the COVID-19 pandemic. *East Mediterr Heal J* **2020**; 26:1155–1164. Available at:
443 [https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-](https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-eng.pdf)
444 [eng.pdf](https://applications.emro.who.int/emhj/v26/10/1020-3397-2020-2610-1155-1164-eng.pdf).
- 445 21. Houlihan CF, Vora N, Byrne T, et al. Pandemic peak SARS-CoV-2 infection and
446 seroconversion rates in London frontline health-care workers. *Lancet* **2020**; 0.
447 Available at: [https://www.thelancet.com/journals/lancet/article/PIIS0140-](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31484-7/abstract)
448 [6736\(20\)31484-7/abstract](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(20)31484-7/abstract).
- 449 22. Eyre DW, Lumley SF, O'Donnell D, et al. Differential occupational risks to healthcare
450 workers from SARS-CoV-2 observed during a prospective observational study. *Elife*
451 **2020**; 9. Available at: <https://elifesciences.org/articles/60675>.
- 452 23. Mostafa A, Kandil S, El-Sayed MH, et al. SARS-CoV-2 seroconversion among 4040
453 Egyptian healthcare workers in 12 resource-limited healthcare facilities: A prospective
454 cohort study. *Int J Infect Dis* **2021**; Available at:
455 <https://doi.org/10.1016/j.ijid.2021.01.037>.

- 456 24. Schneider S, Piening B, Nouri-Pasovsky PA, Krüger AC, Gastmeier P, Aghdassi SJS.
457 SARS-Coronavirus-2 cases in healthcare workers may not regularly originate from
458 patient care: lessons from a university hospital on the underestimated risk of healthcare
459 worker to healthcare worker transmission. *Antimicrob Resist Infect Control* **2020**;
460 9:192. Available at: [https://aricjournal.biomedcentral.com/articles/10.1186/s13756-](https://aricjournal.biomedcentral.com/articles/10.1186/s13756-020-00848-w)
461 020-00848-w.