

1 **TITLE:** Home confinement during COVID-19 pandemic reduced physical activity but not health-
2 related quality of life in previously active older women

3

4 **RUNNING TITLE:** Physical activity, quality of life and COVID-19 home confinement

5

6 **AUTHORS:** Vanessa Teixeira do Amaral, Isabela Roque Marçal, Thiago da Cruz Silva, Fernanda
7 Bianchi Souza, Yacco Volpato Munhoz, Pedro Henrique Camprigher Witzler, Matheus Monge
8 Soares Correa, Bianca Fernandes, Emmanuel Gomes Ciolac

9

10 **AFFILIATIONS:** São Paulo State University (UNESP), School of Sciences, Department of Physical
11 Education, Exercise and Chronic Disease Research Laboratory (ECCR), Bauru, Brazil.

12

13 **CORRESPONDING AUTHOR:** Emmanuel Gomes Ciolac – Universidade Estadual Paulista “Júlio
14 de Mesquita Filho” (UNESP), Faculdade de Ciências, Departamento de Educação Física.

15 Av. Engenheiro Luiz Edmundo Carrijo Coube 14-06, Bauru – SP, Brazil, 17033-360.

16 E-mail: emmanuel.ciolac@unesp.br

17

18 **WORD COUNT:** 2.178

19 **ABSTRACT WORD COUNT:** 223

20 **NUMBER OF TABLES/FIGURES:** 1/3

21 **NUMBER OF REFERENCES:** 19

22 **ABSTRACT**

23

24 **Background:** To investigate the effect of COVID-19 home confinement on levels of physical
25 activity, sedentary behavior and health-related quality of life (HRQL) in older women previously
26 participating in exercise and educational programs.

27 **Methods:** 64 older women (age = 72 ± 5 ys) who participated in a physical exercise/educational
28 program and adhered to government home confinement recommendations have their levels of
29 physical activity, sedentary behavior and HRQL assessed before and during (11 to 13 weeks after
30 introduction of government recommendations to reduce virus transmission) COVID-19 pandemic.

31 **Results:** There were significant reductions in total physical activity (-259 METs/week, $P = 0.02$), as
32 a result of a ~17.0 % reduction in walking (-30.8 min/week, $P = 0.004$) and ~41.8 % reduction in
33 vigorous-intensity activity (-29.6 min/week, $P < 0.001$). Sedentary behavior also increased (2.24
34 h/week, $P < 0.001$; 1.07 h/week days, $P < 0.001$; and 1.54 h/weekend days, $P < 0.001$). However, no
35 significant change occurred in moderate-intensity physical activity, and HRQL domains and facets,
36 except for an improvement in environment domain.

37 **Conclusion:** Home confinement due to COVID-19 pandemic decreased levels of physical activity
38 and increased levels of sedentary behavior in previously active older women who participated in an
39 educational program. However, there were no significant changes in HRQL. These results suggest
40 that educational programs promoting healthy behaviors may attenuate the impact of home
41 confinement in older women.

42

43 **Key Words:** aging; educational program; exercise; quarantine; social isolation.

44

45 **1. Introduction**

46 The outbreak of coronavirus disease 2019 (COVID-19) is a global public health emergency,
47 responsible for 75,479,471 confirmed cases and 1,686,267 deaths worldwide (data as of December
48 21, 2020)¹. Older individuals and those with diabetics, hypertension, and cardiovascular and
49 respiratory diseases are at higher risk of COVID-19 complications and mortality ².

50 Although the measures to contain the spread of the virus (i.e.: social distancing, home
51 confinement, quarantine and/or lockdown) are effective for reducing incidence and mortality of
52 COVID-19³, they may result in negative behaviors that have important repercussions for both
53 individual health and global burden of noncommunicable chronic disease (NCDs). For example, the
54 adoption of these measures has been associated with reduced levels of physical activity ^{4,5}, increased
55 levels of sedentary behavior ^{4,5}, and a more unhealthy food consumption⁴. Few weeks of these
56 negative behaviors may result in several deleterious cardiometabolic and functional consequences
57 that impact management of NCDs ⁶, mainly in older and high-risk populations ⁶⁻⁸.

58 However, little is known about the effect of measures to contain COVID-19 spread on physical
59 activity levels and health-related quality life (HRQL) of older individuals that are physically active,
60 particularly those who had their group exercise routine interrupted, which may also result in social
61 relation impairments. Thus, our aim was to investigate the effect of COVID-19 home confinement
62 on levels of physical activity, sedentary behavior and HRQL in older women that were previously
63 participating in community-based exercise programs. We hypothesized that home confinement would
64 result in impairments on physical activity, sedentary behavior and HRQL.


65 **2. Methods**

66 *2.1 Population and Study Design*

67 The present study is a longitudinal, mixed-methods, observational study, assessing older
68 women (age > 65 years) that were previously participating in the *Ativa Melhor Idade* project. The
69 *Ativa Melhor Idade* is a university extension program (São Paulo State University [UNESP] at Bauru)
70 that promotes twice-weekly community-based exercise programs and a health-related educational
71 program (i.e.: importance and how to meet the recommended levels of physical activity, to avoid
72 sedentary behavior, and to have good nutritional habits). Participants have their levels of physical
73 activity, sedentary behavior and HRQL assessed before (between January and February 2020) and 11
74 to 13 weeks after introduction of societal shutdown, self-isolation and social-distancing measures to
75 reduce virus transmission (June 2020) by State of Sao Paulo (Brazil) government. Inclusion criteria
76 included (1) age > 65 ys, (2) regular participation (at least 60% of compliance) in the *Ativa Melhor*
77 *Idade* project for a minimum of six months prior inclusion in the study, (3) to have responded the
78 physical activity/sedentary behavior and HRQL questionnaires during the project annual assessment
79

80 before beginning of COVID-19 pandemic, and (4) to adhere the government home confinement
81 recommendations to reduce virus transmission. Participants with cognitive impairment who were
82 unable to answer the questionnaires accurately were not included. Participants who did not answer
83 any questionnaire in the follow-up assessment were excluded. A total of 130 older women
84 volunteered to participate in the study, ninety women met the inclusion criteria and were included in
85 the study. However, only 64 women (age = 72 ± 5 ys; body mass index = 29.0 ± 5.2 kg/m²) responded
86 the questionnaires during follow-up assessment and were included in final analysis (Figure 1).
87

88 **Figure 1.** Study design.



89

90 Baseline assessment were performed in the Exercise and Chronic Disease Research
91 Laboratory at UNESP (Bauru–SP, Brazil), during the annual assessment of *Ativa Melhor Idade*
92 project (between January and February 2020), and included anamnesis (health status and medications
93 used), and measures of physical activity/sedentary behavior (questionnaire), HRQL (questionnaires),
94 anthropometrics (weight, height and body mass index [BMI]), hemodynamics (blood pressure, heart
95 rate and arterial stiffness), functional levels (flexibility, balance, muscle strength and endurance
96 performance). Follow-up assessment was performed by a telephone call interview (average duration
97 of 50 min), and included anamnesis (health and behavioral status, medications used, compliance to
98 social isolation/home confinement measures, and participants' perception about the effects of
99 pandemic) and measures of physical activity/sedentary behavior and HRQL. All procedures were

100 approved by the Research Ethics Committee of School of Sciences of UNESP (CAEE
101 21220919.0.0000.5398) and all participants provided written informed consent.

102

103 *2.2 Measurements*

104 Baseline and follow-up assessments were performed by trained evaluators, during the
105 afternoon. Anamnesis and questionnaires were applied in the following sequence: (1) anamnesis and
106 physical evaluations (only at baseline) to characterize the sample profile; (2) levels of physical
107 activity and sedentary behavior (International Physical Activity Questionnaire [IPAQ] short version)
108 ⁹; (3) a questionnaire assessing compliance to home confinement and to other preventive measures of
109 virus spread (only during follow-up); (4) a questionnaire assessing participants' perception about the
110 effects of pandemic on daily living activities, health and anxiety, sleep quality, pain, medication use,
111 eating habits, weight, and alcohol and cigarettes consumption (only during follow-up); and (5) HRQL
112 questionnaires (WHOQOL-BREF and WHOQOL-OLD)^{10,11}. Weekly time and metabolic equivalents
113 (METs) ¹² levels of physical activity, as well as the total, domains and facets scores of HRQL were
114 compared between baseline and follow-up.

115

116 *2.3. Statistical Analysis*

117 The Shapiro-Wilk and Levene's tests were applied to test normality and homoscedasticity of
118 the data assessed before and during COVID-19 pandemic (IPAQ and WHOQOL). Paired samples t-
119 test and Wilcoxon signed-rank test were used to analyze the differences between pre and during
120 COVID-19 variables in parametric and non- parametric data, respectively. Data are presented in mean
121 \pm standard deviation (SD), median (interquartile range) or median (minimum–maximum).
122 Participants' perception about the effects of COVID-19 pandemic are presented descriptively.
123 Statistical software SPSS 17.0™ (SPSS Inc., Chicago, IL, USA) was used to perform the statistical
124 analyses. The statistically significant level was considered when $P < 0.05$.

125

126 **3. Results**


127 All participants had at least one comorbidity (hypertension: $N = 44$; musculoskeletal disorders:
128 $N = 28$; type 2 diabetes Mellitus: $N = 23$; dyslipidemia: $N = 13$; thyroid diseases: $N = 6$; and heart
129 diseases: $N = 6$) and were ~81 days in social isolation/home confinement. During the pandemic, all
130 participants reported to leave their homes once a week or once every two weeks to buy food and/or
131 medications. Participants reported several behavioral alterations as a consequence of the social

132 isolation/home confinement, including sleeping quality (24 participants reported that their sleep
133 worsened, and 28 participants reported to wake up later than normal), and eating habits (22
134 participants reported worsening of eating habits, 31 participants reported weight gain and 11
135 participants reported weight loss). Among the participants who consumed alcoholic beverages (15
136 participants), three participants reported decreasing weekly alcohol consumption, and 12 participants
137 reported no change. Among smokers (10 participants), two participants reported an increase in
138 cigarette consumption, and eight participants reported no change. Sixty participants reported to have
139 some kind of pain in their body, where 15 participants reported an increase and 11 participants
140 reported a decrease during COVID-19 pandemic. Fifty-eight participants reported to maintain routine
141 medication consumption, and six participants reported an increase ($N = 3$) or decrease ($N = 3$) in
142 routine medication consumption.

143 Participants also reported that COVID-19 pandemic affected very much or completely their
144 physical activity routine ($N = 44$). Fifty-one participants reported that were exercising regularly
145 during this period, where 44 participants were exercising at home and seven participants were
146 exercising outdoor (public spaces). Among those who were exercising, 29 participants reported to
147 receive professional guidance remotely. The pre vs. during COVID-19 pandemic assessment showed
148 significant reductions in total physical activity (-259 METs/week, $Z = -2.32$, $P = 0.02$), as a result of
149 a ~17.0 % reduction in walking (-30.8 min/week; $Z = -2.90$, $P = 0.004$) and ~41.8 % reduction in
150 vigorous-intensity activity (-29.6 min/week, $Z = -3.93$, $P < 0.001$); however, no significant change
151 occurred in moderate-intensity activity (Figure 2). Sedentary behavior (total sitting time) also
152 increased (2.24 h/week, $Z = -3.70$, $P < 0.001$) during COVID-19 pandemic in both week days (1.07
153 h/week, $Z = -3.64$, $P < 0.001$) and weekend days (1.54 h/week, $Z = -3.48$, $P < 0.001$) (Figure 2).

154

155 **Figure 2.** Levels of physical activity and sedentary behavior before and during home confinement.



*: significant difference between pre and post ($P < 0.05$).

156

157

158 However, HRQL domains and facets did not change during COVID-19 pandemic, except for
159 an improvement in environment domain (Table 1).

160

161

Table 1. Quality of life before and during home confinement/quarantine.

WHOQOL-BREF			
Domains	PRE	POST	<i>P</i>
Physical Health	14.3 ± 2.5	14.4 ± 2.8	0.61
Psychological	15.4 ± 2.2	15.6 ± 2.5	0.39
Social Relationship	15.7 ± 2.7	15.3 ± 2.7	0.25
Environment	14.0 ± 2.2	14.6 ± 2.2	0.01 *
Global QOL	15.2 ± 2.9	15.5 ± 2.9	0.38

WHOQOL-OLD			
Facets	PRE	POST	<i>P</i>
Sensory Abilities	14.8 ± 3.2	15.5 ± 3.2	0.09
Autonomy	14.1 ± 2.9	14.6 ± 3.1	0.13
Past, Present and Future Activities	14.5 ± 2.7	14.9 ± 2.1	0.34
Social Participation	15.4 ± 2.8	14.7 ± 2.7	0.09
Death and Dying	14.3 ± 4.1	14.1 ± 4.4	0.67
Intimacy	14.7 ± 3.5	15.6 ± 3.6	0.08

162 *: Significant difference between pre and post.

163

164 **4. Discussion**

165 The main findings of present study were that older women who were physically active prior
166 to COVID-19 pandemic showed substantial reduction in physical activity and increase in sedentary
167 behavior during the first months of home confinement. However, there were no significant changes
168 in HRQL, except for a small improvement in environment domain. According to our knowledge, this
169 is the first longitudinal study that investigated the effect of COVID-19 home confinement on physical
170 activity, sedentary behavior and HRQL in older women that were physically active prior to pandemic.

171 Cross-sectional studies has shown that social isolation and home confinement imposed during
172 COVID-19 pandemic reduced the vigorous activities and walking in reduced levels of physical
173 activity and increased levels of sedentary behavior in individuals aged < 65 ys ^{4,13} In addition, a small
174 qualitative study also suggested that home confinement affected the number of older people who
175 participated in group physical activity programs¹⁴. The reduced levels of physical activity and
176 increased levels of sedentary behavior in the present study, as well as the high prevalence (69 %) of
177 participants reporting that COVID-19 pandemic affected very much or completely their physical
178 activity routine study, is in accordance with previous cross-sectional ^{4,13} and qualitative ¹⁴ studies.
179 However, it is important to note that the reduction in walking found in the present study (17.1 %) was
180 lower than the reductions in the previous studies with participants aged < 65 ys (34 % to 58.2 %) ^{4,13}.
181 In addition, the previous cross-sectional studies showed reductions in substantial reductions in
182 moderate-intensity physical activity ^{4,13}, which did not change in the present study. Although
183 differences in age of the population and in study design may not be discharged as reasons for the
184 lower physical activity reduction in the present than previous studies ^{4,13}, the fact that participants of
185 present study were physically active and participated in an educational program to promote healthy
186 behaviors prior to pandemic should be considered. Educational programs promoting healthy
187 behaviors has shown to be effective in improving physical activity ^{15,16}. Thus, it is possible that the
188 participation in the educational program prior to COVID-19 pandemic may be attenuated the walking
189 and moderate-intensity activity reduction in the present study. Future studies are welcome to confirm
190 this hypothesis.

191 The positive effects of physical activity on HRQL in older individuals are well known ^{17,18} .
192 In this context, we hypothesized that home confinement due to COVID-19 pandemic would result in
193 reduction of physical activity and, consequently, HRQL in participants of present. However, despite
194 reduction in physical activity, there were no significant changes in 10 out of 11 domains/facets of
195 HRQL. The small reduction in physical activity and the prior participation in the educational program
196 may be associated with the no change in HRQL in the present study. It is important to note that, in
197 contrast to our hypothesis, there was a significant increase in environment domain of HRQL during

198 follow-up. This improvement may have occurred because participants felt more secure and protected
199 at home during a pandemic. In addition, as participants of present study were retired women who
200 have no change in retirement salary, it is possible that home confinement resulted in greater savings
201 in financial resources, affecting positively HRQL environment domain during follow-up.

202 The lack of a comparison group with older individuals who were not participating in physical
203 activity programs and/or an educational program prior to COVID-19 pandemic is a limitation of
204 present study, because it does not allow us to affirm that the present small reduction in physical
205 activity and the no change in HRQL was due to prior physical exercise and/or educational program
206 participation. The measure of physical activity levels by questionnaire, instead of more accurate tools
207 such as accelerometers is also a limitation that should be acknowledged.

208 The present study has also important implications. Low levels of physical activity is associated
209 with innumerable age-related cardiometabolic and functional impairments ^{6,7,19}. For example, few
210 weeks of reduction in levels of physical activity may result in deleterious effects on glycemic control,
211 body composition, inflammatory cytokines, blood pressure, vascular function,
212 cardiorespiratory/muscle fitness, balance and agility ^{6,7,19}. In addition, reductions in levels of physical
213 activity, as the observed in the present study, may result in increased incidence of NCDs and mortality
214 if persisting for long periods ⁶. Finally, it is also important to note that we found substantial increases
215 in the levels of sedentary behavior, which may potentiate the above-mentioned deleterious effects of
216 reduced levels of physical activity ⁶. In this context, strategies for promoting adequate levels of
217 physical activity and reducing sedentary behavior during COVID-19 home confinement should be
218 urgently implemented.

219

220 **5. Conclusion**

221 In summary, home confinement due to COVID-19 pandemic resulted in decreased levels of
222 physical activity and increased levels of sedentary behavior in previously active older women who
223 participate in an educational program promoting healthy behaviors. However, there were no
224 significant changes in HRQL. As the reduction in physical activity was lower in the present study
225 than in previous cross-sectional studies ^{4,13}, it can be suggested that educational programs promoting
226 healthy behaviors may attenuate the impact of home confinement in physical activity. Future studies
227 addressing this hypothesis are welcome.

228

229 **Acknowledgements**

230 VTA and BF were supported by *Coordenação de Aperfeiçoamento de Pessoal de Nível*
231 *Superior (CAPES)*. IRM and EGC were supported by *Fundação de Amparo à Pesquisa do Estado de*
232 *São Paulo (FAPESP # 2018/09695-5)* and *Conselho Nacional de Desenvolvimento Científico e*
233 *Tecnológico (CNPq #303399/2018-0)*, respectively. The results of the present study are presented
234 clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation.

235

236 **Author's Contributions**

237 VTA participated in study design, literature review, data collection and analysis, and
238 manuscript writing. TCS, FBS, YVM, PHCW and MMSC participated in data collection and
239 manuscript writing. IRM and BF participated in data analysis and manuscript writing. EGC prepared
240 the study, supervised the division of tasks, and participate in data analysis and manuscript writing.
241 All authors approved the final version of the manuscript.

242

243 **Competing interests**

244 The authors declare that they have no competing interests.

245

246

247 **References**

248

- 249 1. World Health Organization. Coronavirus disease (COVID-19). Accessed November 16,
250 2020. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
- 251 2. Jiménez-Pavón, D, Carbonell-Baeza, A, Lavie, C J. Physical exercise as therapy to fight
252 against the mental and physical consequences of COVID-19 quarantine: Special focus in
253 older people. *Prog Cardiovasc Dis*. Published online 2020:386-388.
254 doi:<https://doi.org/10.1016/j.pcad.2020.03.009>
- 255 3. Nussbaumer-Streit B, Mayr V, Dobrescu AI, et al. Quarantine alone or in combination with
256 other public health measures to control COVID-19: a rapid review. *Cochrane Database Syst*
257 *Rev*. 2020;2020(9). doi:10.1002/14651858.CD013574.pub2
- 258 4. Ammar A, Brach M, Trabelsi K, et al. Effects of COVID-19 Home Confinement on Eating
259 Behaviour and Physical Activity: Results of the ECLB-COVID19 International Online
260 Survey. *Nutrients*. 2020;12(6):1583. doi:10.3390/nu12061583
- 261 5. Visser M, Schaap LA, Wijnhoven HAH. Self-Reported Impact of the COVID-19 Pandemic
262 on Nutrition and Physical Activity Behaviour in Dutch Older Adults Living Independently.
263 *Nutrients*. 2020;12. doi:10.3390/nu12123708
- 264 6. Marçal IR, Fernandes B, Viana AA, Ciolac EG. The Urgent Need for Recommending
265 Physical Activity for the Management of Diabetes During and Beyond COVID-19 Outbreak.
266 *Front Endocrinol (Lausanne)*. 2020;11. doi:10.3389/fendo.2020.584642
- 267 7. Ciolac EG, Rodrigues da Silva JM, Vieira RP. Physical Exercise as an Immunomodulator of
268 Chronic Diseases in Aging. *J Phys Act Heal*. 2020;17(6):662-672. doi:10.1123/jpah.2019-

- 269 0237
- 270 8. McGlory C, von Allmen MT, Stokes T, et al. Failed Recovery of Glycemic Control and
271 Myofibrillar Protein Synthesis With 2 wk of Physical Inactivity in Overweight, Prediabetic
272 Older Adults. *Journals Gerontol Ser A*. 2018;73(8):1070-1077. doi:10.1093/gerona/glx203
- 273 9. Questionnaire G for data processing and analysis of the IPA, Forms (IPAQ)—Short and
274 Long. Guidelines for data processing and analysis of the International Physical Activity
275 Questionnaire (IPAQ)-short and long forms. IPAQ Web site. Published 2005. Accessed
276 December, 21, 2020. <https://sites.google.com/site/theipaq/scoring-protocol>
- 277 10. The Whoqol Group. Development of the World Health Organization WHOQOL-BREF
278 Quality of Life Assessment. *Psychol Med*. 1998;28(3):551-558.
279 doi:10.1017/S0033291798006667
- 280 11. Power M, Quinn K, Schmidt S. Development of the WHOQOL-Old module. *Springer*.
281 2005;14(10):2197-2214. doi:10.1007/s11136-005-7380-9
- 282 12. Pardini R, Matsudo S, Araújo T, et al. Validation of the international questionnaire of
283 physical activity level (IPAQ—version 6): pilot study in young Brazilian adults. *Rev Bras*
284 *Ciê n e Mov Brasília*. 2001;9:45-51.
- 285 13. Castañeda-Babarro A, Arbillaga-Etxarri A, Gutiérrez-Santamaria B, Coca A. Impact of
286 COVID-19 confinement on the time and intensity of physical activity in the Spanish
287 population. *Research Square*. Published online 2020. doi:10.21203/rs.3.rs-26074/v1
- 288 14. Goethals L, Barth N, Guyot J, Hupin D, Celarier T, Bongue B. Impact of Home Quarantine
289 on Physical Activity Among Older Adults Living at Home During the COVID-19 Pandemic:
290 Qualitative Interview Study. *JMIR Aging*. 2020;3(1):e19007. doi:10.2196/19007
- 291 15. Da Silva JMR, De Rezende MU, Spada TC, et al. Educational program promoting regular
292 physical exercise improves functional capacity and daily living physical activity in subjects
293 with knee osteoarthritis. *BMC Musculoskelet Disord*. 2017;18(1). doi:10.1186/s12891-017-
294 1912-7
- 295 16. Da Silva JMR, De Rezende MU, Spada TC, et al. Muscular and functional capacity in
296 subjects under treatment for knee osteoarthritis: Role of physical activity status. *J Phys Act*
297 *Heal*. 2019;16(5):362-367. doi:10.1123/jpah.2018-0318
- 298 17. Marquez DX, Aguiñaga S, Vásquez PM, et al. A systematic review of physical activity and
299 quality of life and well-being. *Transl Behav Med*. 2020;10(5):1098-1109.
300 doi:10.1093/tbm/ibz198
- 301 18. Scarabottolo CC, Cyrino ES, Nakamura PM, et al. Relationship of different domains of
302 physical activity practice with health-related quality of life among community-dwelling older
303 people: A cross-sectional study. *BMJ Open*. 2019;9(6):1-10. doi:10.1136/bmjopen-2018-
304 027751
- 305 19. Ciolac EG. Exercise training as a preventive tool for age-related disorders: A brief review.
306 *Clinics*. 2013;68(5):710-717. doi:10.6061/clinics/2013(05)20
- 307