The EEG multiverse of schizophrenia

Authors: Janir Ramos da Cruz[1,2,a], Dario Gordillo[1,a]*, Eka Chkonia[3,4], Wei-Hsiang Lin[1], Ophélie Favrod[1], Andreas Brand[1], Patrícia Figueiredo[2], Maya Roinishvili[4,5], and Michael H. Herzog[1]

Affiliations:

[1] Laboratory of Psychophysics, Brain Mind Institute, École Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland

[2] Institute for Systems and Robotics – Lisboa (LARSyS), Department of Bioengineering, Instituto Superior Técnico, Universidade de Lisboa, Lisbon, Portugal

[3] Department of Psychiatry, Tbilisi State Medical University, Tbilisi, Georgia

[4] Institute of Cognitive Neurosciences, Free University of Tbilisi, Tbilisi, Georgia

[5] Laboratory of Vision Physiology, Beritashvili Centre of Experimental Biomedicine, Tbilisi, Georgia

[a] These authors contributed equally

*Corresponding author:

Dario Gordillo, Laboratory of Psychophysics, Brain Mind Institute, School of Life Sciences, École Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

Phone number: +41 76 633 40 47

Email: dario.gordillolopez@epfl.ch

Keywords: Electroencephalography, Schizophrenia, Psychosis, Neuroimaging, Resting-State, Psychiatry

***ABSTRACT:** 190 WORDS*

***MAIN TEXT:** 3981 WORDS*

Abstract

Research on schizophrenia typically focuses on one paradigm, for which clear-cut differences between patients and controls are established. Great care is taken to understand the underlying genetical, neurophysiological, and cognitive mechanisms, which eventually may explain the clinical outcome. One tacit assumption of these *deep rooting* approaches is that paradigms tap into common and representative aspects of the disease. Here, we analyzed the resting-state electroencephalogram (EEG) of 121 schizophrenia patients and 75 controls. Using multiple signal processing methods, we extracted 194 EEG features. Sixty-nine out of the 194 EEG features showed a significant difference between patients and controls indicating that these features capture an important aspect of the disease. Surprisingly, the correlations between these features were very low, potentially because each feature picks up a different aspect of the disease. Our results are an invitation to rethink the current approach in schizophrenia research, which is centered on drawing general conclusions of schizophrenia based on single features extracted from one paradigm. We propose that complementing *deep* with *shallow* rooting approaches, where many roughly independent features are extracted from one paradigm (or several paradigms), will strongly improve diagnosis and potential treatment of schizophrenia.

Introduction

Schizophrenia patients show strong abnormalities in many domains including personality, cognition, perception, and even immunology. In many experimental paradigms, the differences between patients and controls have large effect sizes, indicating that important aspects of the disease are detected. This provokes two questions: what do these abnormalities have in common and how representative are they for the disease? For example, patients exhibit strong deficits in cognition, such as in working memory tasks [1], which are attributed to abnormalities of cortico-cerebellar-thalamic-cortical circuits [2]. Patients show

also diminished skin flushing with the niacin skin test [3], which is attributed to dysfunctional phospholipase

A2 arachidonic acid signaling [4]. How do the working memory deficits correspond to deficits in skin

functioning? Very few studies have correlated deficits with each other [5–10]. The Consortium on the

Genetics of Schizophrenia studied neurocognitive and neurophysiological abnormalities in schizophrenia

patients with a battery of 15 paradigms [9]. They found that neurocognitive measures shared a significant

amount of variance while neurophysiological measures were almost entirely independent. Price and

colleagues [8] studied four candidate electrophysiological endophenotypes of schizophrenia (mismatch

negativity, P50, P300, and antisaccades). Even though patients *and* their family members showed deficits

in each of these endophenotypes, the features were largely uncorrelated.

Here, we took another road. Instead of comparing different paradigms, we analyzed the very same data

of the very same patients with different electroencephalogram (EEG) analysis methods. Schizophrenia

patients show atypical patterns in multiple resting-state EEG features [11–14,15,16–18]. Resting-state EEG

features revealing significant differences between schizophrenia patients and controls are thought to tap

into brain functions linked to important aspects of the disease. For example, schizophrenia patients

exhibit reduced long-range temporal correlations (LRTC) in the alpha and beta frequency bands [16]. These

responses were proposed to reflect excessive switching of neuronal states in patients. Patients also have

shown atypical patterns in the dynamics of EEG microstates classes C and D [13,19]. These responses were

proposed to correspond to imbalances in attentional and information processing in schizophrenia.

Schizophrenia patients have shown increased power in the delta, theta, and beta frequency bands [20].

Increased beta power was suggested to reflect cortical hyperexcitability, increased power in the delta and

theta bands were proposed to relate to atypical dopaminergic function, to name a few examples. All these

results, individually, suggest that each EEG feature captures important aspects of schizophrenia. But how

representative are these abnormalities of the disorder? Does a patient showing abnormal microstate

dynamics also show deficits in LRTC, or in other EEG features? Even though all these atypical patterns in

patients are obtained from the same EEG data, no study has evaluated how these EEG features relate to

each other. This is not only the case for resting-state EEG studies, but constitutes the conventional

approach in schizophrenia research. This approach is centered on drawing general conclusions about

schizophrenia based on one paradigm, assumed to unveil common and representative aspects of the

disease.

Aiming to shed light on this EEG *multiverse* of schizophrenia, in this work, we analyzed the resting-state

EEG data of 121 schizophrenia patients and 75 healthy controls with multiple methods. This allowed us to

extract 194 EEG features, such as time-domain features, frequency-domain and connectivity features both

in electrode and source space, and nonlinear dynamical features, including many that have shown strong

atypical patterns in patients [11–14,15,16–18]. Then, we correlated the features revealing group differences to

evaluate how these abnormalities/deficits relate to each other. We also examined whether EEG features

show adequate predictive power to clinical scales measuring key symptoms of schizophrenia. Finally, we

propose that future studies in schizophrenia research should consider multiple features extracted from

the same and/or different paradigms in order to improve diagnosis and potential treatment.

Methods

Participants

Two groups of participants joined the experiment: schizophrenia patients (n = 121) and healthy controls

(n = 75). All participants took part in a battery of tests comprising perceptual and cognitive tasks as well

as EEG recordings. Patients were recruited from the Tbilisi Mental Health Hospital or the psycho-social

rehabilitation center. Patients were invited to participate in the study when they had recovered

sufficiently from an acute psychotic episode. Thirty-five were inpatients and 86 were outpatients. Patients

were diagnosed using the Diagnostic and Statistical Manual of Mental Disorders Fourth Edition (DSM-IV)

by means of an interview based on the Structured Clinical Interview for DSM-IV, Clinical Version,

information from staff, and study of patients' records. Psychopathology of patients was assessed by an

experienced psychiatrist using the Scale for the Assessment of Negative Symptoms (SANS) and the Scale

for the Assessment of Positive Symptoms (SAPS). Out of the 121 patients, 106 were receiving neuroleptic

medication. Chlorpromazine (CPZ) equivalents are indicated in **Table 1**. Controls were recruited from the

general population in Tbilisi, aiming to match patients' demographics as closely as possible. All controls

were free from psychiatric axis I disorders and had no family history of psychosis. General exclusion

criteria were alcohol or drug abuse, severe neurological incidents or diagnoses, developmental disorders

(autism spectrum disorder or intellectual disability), or other somatic mind-altering illnesses, assessed

through interview by certified psychiatrists. All participants were no older than 55 years. Group

characteristics are presented in **Table 1**. All participants signed informed consent and were informed that

they could quit the experiment at any time. All procedures complied with the Declaration of Helsinki

(except for pre-registration) and were approved by the Ethical Committee of the Institute of Postgraduate

Medical Education and Continuous Professional Development (Georgia). Protocol number: 09/07. Title:

"Genetic polymorphisms and early information processing in schizophrenia".

Table 1 - Group average statistics (±SD)

	Patients	Controls	Statistics
Gender (F/M)	22/99	39/36	$\chi^2(1) = 24.702$, $p = 6.690e\text{-}7$[b]
Age (years)	35.8 ± 9.2	35.1 ± 7.7	$t(194) = 0.519$, $p = 0.604$[c]
Education (years)	13.3 ± 2.6	15.1 ± 2.9	$t(194) = -4.418$, $p = 1.657e\text{-}5$[c]
Handedness (L/R)	6/115	4/71	$\chi^2(1) = 0.013$, $p = 0.908$[b]
Illness duration (years)	10.8 ± 8.7		
SANS	10.1 ± 5.2		
SAPS	8.6 ± 3.2		
CPZ equivalent[a]	561.1 ± 389.4		
SANS - Scale for the Assessment of Negative Symptoms, SAPS - Scale for Assessment of Positive, CPZ - chlorpromazine			
[a]Average CPZ equivalents calculated over the 106 Patients receiving neuroleptic medication			

EEG recording and data processing

Participants were sitting in a dim lit room. They were instructed to keep their eyes closed and to relax for

5 minutes. Resting-state EEG was recorded using a BioSemi Active Two Mk2 system (Biosemi B.V., The

Netherlands) with 64 Ag-AgCl sintered active electrodes, referenced to the common mode sense

electrode. The recording sampling rate was 2048 Hz. Offline data were downsampled to 256 Hz and

preprocessed using an automatic pipeline [21]. Preprocessed EEG data were analyzed using multiple signal

processing methods in the electrode and source space. In total, 194 EEG features were extracted (See

Supplementary Table 1). Out of the 194 EEG features, 50 were obtained in the source space and 144 in

the electrode space. For source space analysis, we defined 80 brain regions (40 per hemisphere) according

to the AAL atlas (See Supplementary Table 4). See Supplementary Methods for details.

Group comparisons

We compared patients' and controls' scores for each of the 194 EEG features. For each of the *J* variables

(i.e., 64 electrodes, 80 brain regions, or 12 microstate parameters, depending on the number of variables

of each EEG feature) of a given feature, we performed a two-way ANCOVA, with Group (patients and

controls) and Gender (male and female) as factors and Education as a covariate. *P*-values for the effect of

Group were corrected for *J* comparisons using False Discovery Rate (FDR; with an error rate of 5%). Group

effects' η^2 were converted to Cohen's *d*.

Pearson and partial least squares correlations

First, for each EEG feature that contained at least one variable showing a significant difference between patients and controls (after correcting for multiple comparisons), we selected the variable (i.e., electrode, brain region, or microstate parameter) with the biggest effect size to be the representative variable for that feature. Then, for patients and controls separately, we computed pairwise Pearson correlations between the representative variables of each significant EEG feature. Second, to quantify the overall relationship, i.e., the amount of *shared information*, between pairs of multivariate EEG features, we used Partial Least Squares Correlation (PLSC). PLSC generalizes correlations between two variables to two matrices [22,23]. The *shared information* can be quantified as the *inertia* common to the two features [24]. The statistical significance of the inertia was assessed using a permutation test [25,26]. The inertia values were normalized. Hence, the normalized inertias ($\Im_{relative}$) ranged from 0 (the two EEG features are completely unrelated) to 1 (the two EEG features contain the same information). PLSC analysis was done for patients and controls separately. See Supplementary Methods for details.

Regression analysis

To evaluate whether EEG features predict psychopathology scores (SAPS and SANS) adequately, we used elastic net regression models [27]. Elastic nets can handle regression problems where the number of predictors is relatively large compared to the number of samples as well as multicollinearity (i.e., the predictors are not linearly independent) by combining the L1 and L2 penalties to achieve regularization. For each of the 194 EEG features (with all its variables), we built two regression models, one to predict SAPS scores and one to predict SANS scores. We performed 20 repetitions of a 3-fold nested cross-validation procedure. First, one third of the data (1 fold) was left out for validation (test set), while the remaining data (2 folds; train set) were used to find the optimal parameters, namely the amount of penalization and the compromise between L1 and L2 penalties, using 3-fold cross-validation. The model

with the parameters leading to best performance in the train set was tested on the left-out data (test set).

The entire procedure was repeated 20 times, with different allocations of the patients in the train and

test sets. Using the same cross-validation procedure, i.e., 20 repetitions of a 3-fold cross-validation, we

also evaluated predictive performance using a nonlinear random forest regression model, setting the

maximum depth of the tree to 10 and the number of trees to 100. Random forests are meta estimators

that average several decision trees trained on subsets of the dataset to improve accuracy and avoid

overfitting. Prediction performance was calculated using the coefficient of determination (R^2) and the root

mean squared error (*RMSE*). The distribution of the prediction performance values was obtained from the

60 aggregated RMSE and R^2, across repetitions of the procedure.

155 Results

156 *Multiple EEG features reveal significant group effects*

For 121 patients (22 females, 35.8 ± 9.2 years old, 13.3 ± 2.6 years of education) and 75 age-matched

healthy controls (39 females, 35.1 ± 7.7 years old, 15.1 ± 2.9 years of education; Table 1), we extracted in

total 194 features from the resting-state EEG recordings, including time-domain, frequency-domain,

connectivity, and nonlinear dynamical features (Supplementary Table 1). Among the 194 EEG features, 69

(35.57%) showed significant differences between patients and controls with medium to large effect sizes

(Cohen's *d* varied from 0.463 to 1.037, **Figure 1**). Patients showed significantly reduced values in 24 out

of the 69 EEG features revealing significant group differences (illustrated as negative effect size in **Figure**

1). Patients exhibited significantly higher values than controls in 45 EEG features.

Figure 1. Effect size (Cohen's *d*) of the group differences between patients and controls

for each of the 194 EEG features. We took the values of the electrode, brain region, or

microstate parameter with the largest effect size according to Cohen's *d* (η^2 values were

converted to Cohen's *d*) to be the representative variable for each feature. Significant

group differences, after correction for multiple comparisons (using FDR), are depicted in

red, with dotted red vertical lines serving as a guide to their labels. > and < were added

to the feature labels to indicate if patients had significantly higher or lower values than

controls, respectively. The non-significant effects are shown in blue. Error bars represent

95% confidence intervals (C.I.). A list with the abbreviations and the corresponding name

of each feature is presented in Supplementary Table 1.

Correlations between EEG features

To evaluate to what extent features that showed significant group differences are sensitive to the same

aspects of the disorder we computed Pearson's correlations between pairs of features (**Figure 2**). As the

representative variable for each feature, we took the values of the electrode, brain region, or microstate

parameter that showed the largest group difference according to Cohen's *d* (**Figure 1**). Surprisingly, we

found that in the patients group only 36.49% of the pairwise correlations were significant at a level of 0.05

(without correcting for multiple comparisons). For the control group, only 26.73% of the correlations were

significant. Since significance depends on the sample size, here, we focus on the magnitude of the

correlation coefficients (r-values). In general, the magnitudes of the r-values were very low in both

patients (0.055, 0.122, 0.251, for the 25th, 50th, and 75th percentiles, respectively) and controls (0.059,

0.129, 0.242, for the 25th, 50th, and 75th percentiles, respectively; **Figure 2**). Strong correlations were found

mainly for pairs of very closely related features, such as between waiting-time statistics of gamma bursts

(*waiting time gamma*) and life-time statistics of gamma bursts (*life time gamma*; r=0.836 and r=0.926, in

patients and controls, respectively).

Figure 2. Pairwise correlations between the 69 EEG features that showed significant group

differences between patients and controls. Patients' r-values are presented in the upper

triangle and controls' r-values are shown in the lower triangle. Strong negative and

positive r-values are depicted in red and blue, respectively, and r-values around 0 in

yellow. For each feature, we used the values of the electrode, brain region, or microstate

parameter that showed the largest effect size as the representative variable for the

correlations. A list with the abbreviations and corresponding name of each feature is

shown in Supplementary Table 1.

To quantify the overall *shared information* between pairs of EEG features, which showed significant group

differences, by taking not only variables with the largest effect size into account but all variables of the

features we used partial least squares correlation (PLSC). For the patients, 55.92% of the pairwise inertias

were significant (without correcting for multiple comparisons) and for controls, 40.28%. In general,

relative inertias were not very high in both patients (0.254, 0.329, 0.409, for the 25th, 50th, and 75th

percentiles, respectively) and controls (0.305, 0.387, 0.472, for the 25th, 50th, and 75th percentiles,

respectively; **Figure 3**). As in the Pearson's correlation results, features that showed strong associations

were mainly similar features, such as the same network statistics for different connectivity measures in

the theta band, for example, at the electrode level: clustering coefficient connectivity estimated with

phase locking value (*clust coeff e-plv theta*) and with imaginary part of coherence (*clust coeff e-icoh theta*;

$\mathfrak{I}_{relative}$=0.804 and $\mathfrak{I}_{relative}$=0.826, in patients and controls, respectively).

Figure 3. Shared information between the 69 EEG features that showed significant group

differences, as measured by the relative inertia ($\mathfrak{I}_{relative}$) computed with partial least

squares correlations (PLSC). The relative inertia ranges from 0 (two features are

completely unrelated) to 1 (the two features' values move together by the exact same

percentage). Patients' relative inertias are presented in the upper triangle and controls'

relatives are shown in the lower triangle. A list with the abbreviations and corresponding

name of each feature is shown in Supplementary Table 1.

Prediction of psychopathology scores

We evaluated whether EEG features were adequate predictors of psychopathology scores determined by

the Scale for the Assessment of Positive Symptoms (SAPS) and the Scale for the Assessment of Negative

Symptoms (SANS), which target positive (hallucinations, delusions, bizarre behavior, and positive formal

thought disorder) and negative (affective flattening, alogia, apathy, anhedonia, and attention) symptoms,

respectively. All 194 EEG features exhibited very weak out-of-sample predictive ability to both the SANS

and SAPS scores. Results were very similar for both the linear (i.e., elastic net) and nonlinear (i.e., random

forest) models. See Supplementary Table 2 and Supplementary Table 3 for details.

Discussion

Traditionally, schizophrenia research focuses on a single experimental paradigm and analysis method

showing significant differences between patients and controls, and then tries to derive the underlying

genetic or neurophysiological causes of the disorder. This approach has been quite successful in the

formulation of hypotheses, such as the dopamine hypothesis [28], the social brain hypothesis [29], the

glutamate hypothesis [30], or the dysconnection hypothesis [31], just to name a few. Here, we examined to

what extent abnormalities quantified by different EEG features correlate with each other. Many of the

investigated features were previously linked to different abnormalities of brain processes in

schizophrenia, and, here, we reproduced many of these results, such as imbalance in microstates

dynamics [13,19], decreased long-range temporal correlations in the alpha and beta bands [16], decreased life-

and waiting-times in the beta band [17], increased spectral amplitude in the theta band [12], increased

connectivity in the theta band at the source level [11,14], decreased Lyapunov exponent [15], among others.

With our systematic analysis, we also found abnormalities in EEG features, which, to the best of our

knowledge, have not been reported yet, namely, delta-phase gamma-amplitude coupling, range EEG

coefficient of variation and asymmetry in the theta and alpha bands, etc. In some way, deeper analysis of

each feature may have warranted an in-depth study and a potential publication. However, we did not

want to elaborate on these methods individually because we wanted to understand how all EEG features

relate to each other in their entirety. The surprising insight from our analysis is that, even though we are

probing the same signals from the same participants, we found only weak correlations between the 69

significant features. The only strong correlations were between features that are similar from the outset,

thereby resembling test-retests. This suggests that, even though EEG features reveal clear-cut and

reproducible atypical patterns in patients, none of the features is truly representative for the disease, but

rather that all these features pick up more or less independent aspects of schizophrenia. Hence, the

traditional approach of focusing on a single experimental paradigm and analysis method has its

limitations. These results remind us that schizophrenia is indeed a very heterogeneous disease, a well-

known fact, which is however not always taken seriously enough because, as mentioned above, most

research tries to find the one or a few causes of schizophrenia within one well described paradigm by

digging as deep as possible into the underlying neurophysiological and genetic mechanisms. In analogy to

botany, one may call these approaches "deep rooting" approaches.

We propose that it may be useful to complement these deep rooting approaches with "shallow rooting"

approaches, representing schizophrenia within a high-dimensional space, where many tests and analysis

outcomes are the basis variables. The outcomes should ideally have large effect sizes, low mutual

correlations, and a "flat" factor structure. Whether this is possible is an open question and depends very

much on the underlying causes of schizophrenia. On the lowest complexity level, there may be only a few

independent causes (or even only one), which were not found yet. Given the heterogeneity of the disease,

including abnormalities in the cognitive [2] but also the skin functioning domain [4], the causes need to be on

a rather general level, likely subcellular, present in all human functioning. Alternatively, schizophrenia

may be an approximatively "additive" disease, where many small abnormalities add up to severe

symptoms. In an even more complex scenario, only certain combinations of redundant functions, each

coming with at least two variants, cause the disease. For instance, if one function is up-regulated and

another one down-regulated in an individual, there are no abnormalities. Deficits manifest only when all

or most functions are either up- or down-regulated. In such a combinatorial scenario, it would be difficult

to find the underlying causes since each variant itself does not lead to a deficit; only certain combinations

do. Our correlation analysis provides evidence supporting the more complex/multifactorial scenario,

where each feature makes up its own factor and manifests differently in different patients. Since our data

shows that each EEG feature is sensitive to roughly independent aspects of schizophrenia, each brain

process captured by the analysis methods might be neither necessary nor sufficient to explain important

aspects of the disease. Indeed, we found that EEG features showed very weak predictive power to key

symptoms of schizophrenia, suggesting that there is little information about individual differences in

psychopathology. Our results are an invitation to rethink the current approach in schizophrenia research

and suggest that new study designs conflating multiple features from the same and different paradigms

are more adequate, since they are more aligned with the heterogeneity of the disorder. In the next steps,

it will be important to find the right set of features, which may stem from cognitive tests but also

potentially immunological markers, of which each may contribute with a variety of features. Previous

research has shown that combining features improved classification and predictive performance in

psychosis studies [32,33].

Our study has several limitations. There are demographic differences between patients and controls,

which might affect our group comparisons. However, we attempted to minimize these demographic

effects by using education as a covariate and gender as factor in the analyses. Similarly, we cannot exclude

effects of medication in our results. Nonetheless, we find similar patterns of correlations between EEG

features, i.e., weak associations, in both patients and controls, suggesting that if there is an effect of

medication, it is small. Further, our sample size is relatively small for achieving reliable estimates of

predictive power [34–36].

Our results may explain a deep mystery in schizophrenia research. Schizophrenia has an estimated

heritability of 70 to 85% [37]. For example, the chance to also suffer from schizophrenia for monozygotic

twins is about 33% when the partner twin has the disease [38]. Furthermore, about 0.25 to 0.75% people of

a population suffer from schizophrenia and related psychotic disorders [39–41]. These values are rather stable

across cultures [42]. Given that schizophrenia patients have less offspring [43–46], this provokes the question

why schizophrenia has not been extinguished during the course of evolution [45,47]. In the above-mentioned

combinatorial scenario with many redundant functions this may simply happen because evolution

operates on the individual single-nucleotide polymorphism (SNP) level and not on the combinatorial one.

As long as most of the population shows average functioning, there will be no change of the allele

distributions. In the additive scenario, evolution may extinct harmful alleles, of which each constitutes

only a little risk, very slowly and these may be replaced by harmful de novo mutations [45]. To what extent

such considerations hold true will be shown by shallow rooting approaches using a plethora of paradigms

and a multiverse of analysis methods. In a nutshell, deep rooting will help to understand the different

aspects of the disorder, while shallow rooting will help to better diagnose schizophrenia by finding

subpopulations, leading to more personalized treatment.

304 Acknowledgements

305 We would like to thank Marc Repnow for his comments and Ben Lönnqvist for proofreading the

306 manuscript. M.H.H., E.C., A.B., and M.R. designed the research; M.R. and E.C. performed the research;

307 J.R.C., D.G., W.H.L., and O.F. analyzed data; J.R.C., D.G., O.F. A.B., P.F., and M.H.H. wrote the paper. Data

308   of 101 patients and 75 controls have already been published in different contexts [13,48–50]. The code that

309   support the findings of this study are available upon request. The authors declare no competing interests.

Funding

311   This work was partially funded by the Fundação para a Ciência e a Tecnologia under grant FCT

312   PD/BD/105785/2014 and the National Centre of Competence in Research (NCCR) Synapsy financed by the

313   Swiss National Science Foundation under grant 51NF40-185897.

References

1.   Meyer-Lindenberg A, Poline J-B, Kohn PD, et al. Evidence for Abnormal Cortical Functional
Connectivity During Working Memory in Schizophrenia. *Am J Psychiatry*. 2001;158(11):1809-1817.
doi:10.1176/appi.ajp.158.11.1809

2.   Andreasen NC, Paradiso S, O'Leary DS. "Cognitive Dysmetria" as an Integrative Theory of
Schizophrenia: A Dysfunction in Cortical-Subcortical-Cerebellar Circuitry? *Schizophr Bull*.
1998;24(2):203-218. doi:10.1093/oxfordjournals.schbul.a033321

3.   Rybakowski J, Weterle R. Niacin test in schizophrenia and affective illness. *Biol Psychiatry*.
1991;29(8):834-836. doi:10.1016/0006-3223(91)90202-W

4.   Messamore E. Niacin subsensitivity is associated with functional impairment in schizophrenia.
Schizophr Res. 2012;137(1-3):180-184. doi:10.1016/j.schres.2012.03.001

5.   Braff DL, Freedman R, Schork NJ, Gottesman II. Deconstructing Schizophrenia: An Overview of the
Use of Endophenotypes in Order to Understand a Complex Disorder. *Schizophr Bull*. 2007;33(1):21-
32. doi:10.1093/schbul/sbl049

6.   Braff DL, Light GA, Swerdlow NR. Prepulse Inhibition and P50 Suppression Are Both Deficient but not
Correlated in Schizophrenia Patients. *Biol Psychiatry*. 2007;61(10):1204-1207.
doi:10.1016/j.biopsych.2006.08.015

7.   Dickinson D, Goldberg TE, Gold JM, Elvevag B, Weinberger DR. Cognitive Factor Structure and
Invariance in People With Schizophrenia, Their Unaffected Siblings, and Controls. *Schizophr Bull*.
2011;37(6):1157-1167. doi:10.1093/schbul/sbq018

8.   Price GW, Michie PT, Johnston J, et al. A Multivariate Electrophysiological Endophenotype, from a
Unitary Cohort, Shows Greater Research Utility than Any Single Feature in the Western Australian
Family Study of Schizophrenia. *Biol Psychiatry*. 2006;60(1):1-10. doi:10.1016/j.biopsych.2005.09.010

9.   Seidman LJ, Hellemann G, Nuechterlein KH, et al. Factor structure and heritability of
endophenotypes in schizophrenia: Findings from the Consortium on the Genetics of Schizophrenia
(COGS-1). *Schizophr Res*. 2015;163(1):73-79. doi:10.1016/j.schres.2015.01.027

10.   Toomey R, Faraone SV, Seidman LJ, Kremen WS, Pepple JR, Tsuang MT. Association of
neuropsychological vulnerability markers in relatives of schizophrenic patients. *Schizophr Res*.
1998;31(2-3):89-98. doi:10.1016/S0920-9964(98)00025-5

11.   Andreou C, Leicht G, Nolte G, et al. Resting-state theta-band connectivity and verbal memory in
schizophrenia and in the high-risk state. *Schizophr Res*. 2015;161(2-3):299-307.
doi:10.1016/j.schres.2014.12.018

12.   Boutros NN, Arfken C, Galderisi S, Warrick J, Pratt G, Iacono W. The status of spectral EEG
abnormality as a diagnostic test for schizophrenia. *Schizophr Res*. 2008;99(1–3):225-237.
doi:10.1016/j.schres.2007.11.020

13.   da Cruz JR, Favrod O, Roinishvili M, et al. EEG microstates are a candidate endophenotype for
schizophrenia. *Nat Commun*. 2020;11. doi:10.1038/s41467-020-16914-1

14.   Di Lorenzo G, Daverio A, Ferrentino F, et al. Altered resting-state EEG source functional connectivity
in schizophrenia: the effect of illness duration. *Front Hum Neurosci*. 2015;9.
doi:10.3389/fnhum.2015.00234

15.   Kim D-J, Jeong J, Chae J-H, et al. An estimation of the first positive Lyapunov exponent of the EEG in
patients with schizophrenia. *Psychiatry Res Neuroimaging*. 2000;98(3):177-189. doi:10.1016/S0925-
4927(00)00052-4

16.   Nikulin VV, Jönsson EG, Brismar T. Attenuation of long-range temporal correlations in the amplitude
dynamics of alpha and beta neuronal oscillations in patients with schizophrenia. *NeuroImage*.
2012;61(1):162-169. doi:10.1016/j.neuroimage.2012.03.008

17.   Sun J, Tang Y, Lim KO, et al. Abnormal Dynamics of EEG Oscillations in Schizophrenia Patients on
Multiple Time Scales. *IEEE Trans Biomed Eng*. 2014;61(6):1756-1764.
doi:10.1109/TBME.2014.2306424

18.   Uhlhaas PJ, Singer W. Abnormal neural oscillations and synchrony in schizophrenia. *Nat Rev
Neurosci*. 2010;11(2):100-113. doi:10.1038/nrn2774

19.   Rieger K, Diaz Hernandez L, Baenninger A, Koenig T. 15 Years of Microstate Research in
Schizophrenia – Where Are We? A Meta-Analysis. *Front Psychiatry*. 2016;7.
doi:10.3389/fpsyt.2016.00022

20.   Venables NC, Bernat EM, Sponheim SR. Genetic and Disorder-Specific Aspects of Resting State EEG
Abnormalities in Schizophrenia. *Schizophr Bull*. 2009;35(4):826-839. doi:10.1093/schbul/sbn021

21.   da Cruz JR, Chicherov V, Herzog MH, Figueiredo P. An automatic pre-processing pipeline for EEG
analysis (APP) based on robust statistics. *Clin Neurophysiol*. 2018;129(7):1427-1437.
doi:10.1016/j.clinph.2018.04.600

22. McIntosh AR, Bookstein FL, Haxby JV, Grady CL. Spatial Pattern Analysis of Functional Brain Images
Using Partial Least Squares. *NeuroImage*. 1996;3(3):143-157. doi:10.1006/nimg.1996.0016

23. Tucker LR. An inter-battery method of factor analysis. *Psychometrika*. 1958;23(2):111-136.
doi:10.1007/BF02289009

24. Krishnan A, Williams LJ, McIntosh AR, Abdi H. Partial Least Squares (PLS) methods for neuroimaging:
A tutorial and review. *NeuroImage*. 2011;56(2):455-475. doi:10.1016/j.neuroimage.2010.07.034

25. Abdi H, Williams LJ. Partial Least Squares Methods: Partial Least Squares Correlation and Partial
Least Square Regression. In: Reisfeld B, Mayeno AN, eds. *Computational Toxicology*. Vol 930.
Methods in Molecular Biology. Humana Press; 2013:549-579. doi:10.1007/978-1-62703-059-5_23

26. McIntosh AR, Chau WK, Protzner AB. Spatiotemporal analysis of event-related fMRI data using
partial least squares. *NeuroImage*. 2004;23(2):764-775. doi:10.1016/j.neuroimage.2004.05.018

27. Zou H, Hastie T. Regularization and variable selection via the elastic net. *J R Stat Soc Ser B Stat
Methodol*. 2005;67(2):301-320. doi:10.1111/j.1467-9868.2005.00503.x

28. Howes OD, Kapur S. The Dopamine Hypothesis of Schizophrenia: Version III--The Final Common
Pathway. *Schizophr Bull*. 2009;35(3):549-562. doi:10.1093/schbul/sbp006

29. Burns J. The social brain hypothesis of schizophrenia. *World Psychiatry Off J World Psychiatr Assoc
WPA*. 2006;5(2):77-81.

30. Hu W, MacDonald ML, Elswick DE, Sweet RA. The glutamate hypothesis of schizophrenia: evidence
from human brain tissue studies. *Ann N Y Acad Sci*. 2015;1338:38-57. doi:10.1111/nyas.12547

31. Friston K, Brown HR, Siemerkus J, Stephan KE. The dysconnection hypothesis (2016). *Schizophr Res*.
2016;176(2-3):83-94. doi:10.1016/j.schres.2016.07.014

32. Mothi SS, Sudarshan M, Tandon N, et al. Machine learning improved classification of psychoses using
clinical and biological stratification: Update from the bipolar-schizophrenia network for
intermediate phenotypes (B-SNIP). *Schizophr Res*. 2019;214:60-69.
doi:10.1016/j.schres.2018.04.037

33. Yang H, Liu J, Sui J, Pearlson G, Calhoun VD. A Hybrid Machine Learning Method for Fusing fMRI and
Genetic Data: Combining both Improves Classification of Schizophrenia. *Front Hum Neurosci*.
2010;4. doi:10.3389/fnhum.2010.00192

34. Poldrack RA, Huckins G, Varoquaux G. Establishment of Best Practices for Evidence for Prediction: A
Review. *JAMA Psychiatry*. 2020;77(5):534. doi:10.1001/jamapsychiatry.2019.3671

35. Schnack HG, Kahn RS. Detecting Neuroimaging Biomarkers for Psychiatric Disorders: Sample Size
Matters. *Front Psychiatry*. 2016;7. doi:10.3389/fpsyt.2016.00050

36. Varoquaux G. Cross-validation failure: Small sample sizes lead to large error bars. *NeuroImage*.
2018;180:68-77. doi:10.1016/j.neuroimage.2017.06.061

37. Burmeister M, McInnis MG, Zöllner S. Psychiatric genetics: progress amid controversy. *Nat Rev*
Genet. 2008;9(7):527-540. doi:10.1038/nrg2381

38. Hilker R, Helenius D, Fagerlund B, et al. Heritability of Schizophrenia and Schizophrenia Spectrum
Based on the Nationwide Danish Twin Register. *Biol Psychiatry*. 2018;83(6):492-498.
doi:10.1016/j.biopsych.2017.08.017

39. Kessler RC, Birnbaum H, Demler O, et al. The prevalence and correlates of nonaffective psychosis in
the National Comorbidity Survey Replication (NCS-R). *Biol Psychiatry*. 2005;58(8):668-676.
doi:10.1016/j.biopsych.2005.04.034

40. Moreno-Küstner B, Martín C, Pastor L. Prevalence of psychotic disorders and its association with
methodological issues. A systematic review and meta-analyses. *PLOS ONE*. 2018;13(4):e0195687.
doi:10.1371/journal.pone.0195687

41. Saha S, Chant D, Welham J, McGrath J. A systematic review of the prevalence of schizophrenia. *PLoS*
Med. 2005;2(5):e141. doi:10.1371/journal.pmed.0020141

42. Simeone JC, Ward AJ, Rotella P, Collins J, Windisch R. An evaluation of variation in published
estimates of schizophrenia prevalence from 1990–2013: a systematic literature review. *BMC*
Psychiatry. 2015;15(1). doi:10.1186/s12888-015-0578-7

43. Avila M, Thaker G, Adami H. Genetic epidemiology and schizophrenia: a study of reproductive
fitness. *Schizophr Res*. 2001;47(2-3):233-241. doi:10.1016/S0920-9964(00)00062-1

44. Bassett AS, Bury A, Hodgkinson KA, Honer WG. Reproductive fitness in familial schizophrenia.
Schizophr Res. 1996;21(3):151-160. doi:10.1016/0920-9964(96)00018-7

45. Keller MC, Miller G. Resolving the paradox of common, harmful, heritable mental disorders: Which
evolutionary genetic models work best? *Behav Brain Sci*. 2006;29(4):385-404.
doi:10.1017/S0140525X06009095

46. MacCabe JH, Koupil I, Leon DA. Lifetime reproductive output over two generations in patients with
psychosis and their unaffected siblings: the Uppsala 1915–1929 Birth Cohort Multigenerational
Study. *Psychol Med*. 2009;39(10):1667. doi:10.1017/S0033291709005431

47. Liu C, Everall I, Pantelis C, Bousman C. Interrogating the Evolutionary Paradox of Schizophrenia: A
Novel Framework and Evidence Supporting Recent Negative Selection of Schizophrenia Risk Alleles.
Front Genet. 2019;10. doi:10.3389/fgene.2019.00389

48. da Cruz JR, Shaqiri A, Roinishvili M, et al. Neural Compensation Mechanisms of Siblings of
Schizophrenia Patients as Revealed by High-Density EEG. *Schizophr Bull*. 2020;46(4):1009-1018.
doi:10.1093/schbul/sbz133

49. Favrod O, Roinishvili M, da Cruz JR, et al. Electrophysiological correlates of visual backward masking
in patients with first episode psychosis. *Psychiatry Res Neuroimaging*. 2018;282:64-72.
doi:10.1016/j.pscychresns.2018.10.008

50.   Garobbio SA, Roinishvili M, Favrod O, et al. *Electrophysiological Correlates of Visual Backward*
Masking in Patients with Bipolar Disorder. Neuroscience; 2020. doi:10.1101/2020.05.12.090407