Large variation in the association between seasonal antibiotic use and resistance across multiple bacterial species and antibiotic classes

Daphne S. Sun[1], Stephen M. Kissler[1], Sanjat Kanjilal[2,3], Scott W. Olesen[1], Marc Lipsitch[1,4], Yonatan H. Grad[1,3]

1. Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health, Boston, MA, USA
2. Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, Boston, MA, USA
3. Division of Infectious Diseases, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA
4. Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard Chan School of Public Health, Boston, MA, USA

Correspondence: Yonatan H. Grad (ygrad@hsph.harvard.edu)

Abstract

Understanding how antibiotic use drives resistance is crucial for guiding effective strategies to

limit the spread of resistance, but the use-resistance relationship across pathogens and

antibiotics remains unclear. We applied sinusoidal models to evaluate the seasonal use-

resistance relationship across 3 species (*Staphylococcus aureus*, *Escherichia coli*, and

Klebsiella pneumoniae) and 5 antibiotic classes (penicillins, macrolides, quinolones,

tetracyclines, and nitrofurans) in Boston, Massachusetts. Use of all 5 classes and resistance in

9 of 15 species-antibiotic combinations showed statistically significant amplitudes of seasonality

(false discovery rate < 0.05). While seasonal peaks in use varied by class, resistance in all 9

species-antibiotic combinations peaked in the winter and spring. The correlations between

seasonal use and resistance thus varied widely, with resistance to all antibiotic classes being

most positively correlated with use of the winter-peaking classes (penicillins and macrolides).

These findings challenge the simple model of antibiotic use independently selecting for

resistance and suggest that stewardship strategies will not be equally effective across all

species and antibiotics. Rather, seasonal selection for resistance across multiple antibiotic

classes may be dominated by use of the most highly prescribed antibiotic classes, penicillins

and macrolides.

Introduction

Antibiotic resistance is a growing threat to society, with important public health [1] and economic

consequences [2]. Antibiotic use is considered a primary driver of resistance not only in the

pathogen targeted by the antibiotic but also in host-associated bacteria subject to 'bystander

selection' [3]. As such, stewardship programs to reduce overall antibiotic prescribing have

become a popular strategy for broadly reducing the burden of resistance. However, the efficacy

of stewardship efforts has varied widely and, in some cases, shown a limited impact on reducing

rates of resistance [4,5]. These findings reflect the complexity of the antibiotic use-resistance

relationship, underscoring the need to characterize this relationship across a wide range of

bacterial species and antibiotics and identify factors that influence the strength of this

association.

Temporal studies have shown that an association between population-level antibiotic use and

resistance can be detected on rapid timescales, where seasonal fluctuations in use have been

accompanied by seasonal fluctuations in resistance with up to a few months lag [6–9]. To

interpret the lag between seasonal use and resistance, Blanquart et al. proposed a model for

the relationship between short-term sinusoidal fluctuations in antibiotic use and resistance [10].

This model predicts that antibiotic use determines the rate of change of resistance, such that the

derivative of the prevalence of resistance should depend on the level of use. Thus, if antibiotic

use follows a sine function over a 12-month period, then peak resistance should lag peak use

by a quarter period, or 3 months. The lag can be shortened by including a 'stabilizing force' in

the model to account for forces that counteract the effect of use and drive fluctuations in

resistance towards equilibrium.

Findings from previous seasonality studies have been largely consistent with this model.

Studies that focused on antibiotics with wintertime peaks in use (e.g. penicillins, macrolides, and

quinolones) have identified positive associations with winter/spring peaks in resistance lagged

by 0-3 months in *Streptococcus pneumoniae* [6], *Escherichia coli* [7,8], *Staphylococcus aureus*

[7], and *Neisseria gonorrhoeae* [9]. In contrast, a study from the Netherlands that analyzed

antibiotics with summer/autumn peaks in use (e.g. Nitrofurantoin, Fosfomycin, Trimethoprim)

found that resistance in *E. coli* and *Klebsiella pneumoniae* still peaked in the winter/spring and

lagged use by 3-6 months [11], inconsistent with the Blanquart et al. model. The authors of this

study attribute the longer lag time to the weaker seasonal fluctuations and lower overall rates of

antibiotic use in their study population. However, it is unclear whether resistance to other

antibiotics with winter peaks in use would exhibit similarly long lag times in this population, or

whether these findings reflect a broader phenomenon where despite different seasonal patterns

of use, resistance always peaks in the winter/spring due to other ecological factors.

We aimed to characterize the seasonal relationship between antibiotic use and resistance

across antibiotic classes with winter, summer, and biannual peaks in use [6,7,9,12] in Boston,

Massachusetts. We studied three clinically relevant species—*Staphylococcus aureus*,

Escherichia coli, and *Klebsiella pneumoniae*—which represent skin/nasal and gut colonizing

bacteria [13,14] that cause a diversity of infections types and are subject to strong bystander

selection [3]. We obtained antibiotic use data from a centralized state-wide insurance claims

database and resistance data from two major Boston-area hospitals. Given the near-universal

health insurance coverage in Massachusetts [15], this analysis provided a unique opportunity to

characterize the antibiotic use-resistance relationship in a dataset that captures nearly all

antibiotic use in a population.

Results

Seasonality in antibiotic use varies across classes

The five antibiotic classes included in this study each displayed statistically significant seasonal

patterns of use (**Fig 1A**). Penicillins and macrolides were most frequently prescribed, with year-

round averages of 4.8 and 4.1 daily claims per 10,000 people, respectively. Quinolones,

tetracyclines, and nitrofurans were prescribed with year-round averages of 1.8, 1.0, and 0.5

daily claims per 10,000 people, respectively.

Penicillins had the greatest magnitude change in prescribing rate across seasons, with the

seasonal component having an amplitude of 1.1 additional daily claims per 10,000 people (peak

to mean) (95% CI, 0.96 to 1.3). This was followed by macrolides (amplitude, 0.74; 95% CI, 0.59

to 0.89), quinolones (amplitude, 0.081; 95% CI, 0.04 to 0.12), nitrofurans (amplitude, 0.04; 95%

CI, 0.02 to 0.06), and tetracyclines use (amplitude, 0.03; 95% CI, 0.01 to 0.05) (**Fig 1B**).

The timing of peak prescribing varied by antibiotic class (**Fig 1B**). Macrolide and penicillin use

peaked in the winter, around late January (phase, 1.7 months; 95% CI, 1.3 to 2.1; note that

phase is indexed to 1.0 representing January 1st) and early February (phase, 2.2 months; 95%

CI, 2.0 to 2.5), respectively. Tetracycline and nitrofuran use peaked in the summer, around mid-

June (phase, 6.5 months; 95% CI, 4.6 to 8.5) and late August (phase, 8.8 months; 95% CI, 8.2

to 9.4), respectively. Finally, quinolone use peaked twice a year in early January and early July

(phases, 1.0 (95% CI, 0.6 to 1.5) and 7.0 (95% CI, 6.6 to 7.5) months).

Seasonality in antibiotic resistance is prevalent across species and antibiotic classes

Resistance was seasonal for 9 out of 15 species-antibiotic combinations (**Fig 2, S1 Fig, S2 Fig**),

with statistically significant amplitudes of seasonality (FDR < 0.05) ranging from a peak

$\log_2$(MIC) increase of 0.028 to 0.063 above the yearly average (**Fig 3**). Ciprofloxacin resistance

and nitrofurantoin resistance were seasonal in all three species with a 12-month period (**Fig 2,**

S1 Fig, S2 Fig). Resistance to erythromycin in *S. aureus* was also seasonal with a 12-month

period (amplitude, 0.048; 95% CI, 0.012 to 0.083). Conversely, tetracycline resistance was not

seasonal in any of the three species (**Fig 2, S1 Fig, S2 Fig**). The seasonal patterns of

resistance to penicillin class antibiotics were variable across species. Oxacillin resistance in *S.*

aureus was seasonal with a 12-month period (amplitude, 0.031; 95% CI, 0.009 to 0.054), while

both penicillin resistance in *S. aureus* (amplitude, 0.010; 95% CI, -0.007 to 0.027) and

amoxicillin/clavulanate resistance in *E. coli* (amplitude, 0.010; 95% CI -0.0005 to 0.021) and *K.*

pneumoniae (amplitude, 0.034; 95% CI, 0.001 to 0.067) did not meet our criterion for

seasonality. Ampicillin resistance in *E. coli* was the only species-antibiotic combination with a

statistically significant amplitude (0.034; 95% CI, 0.002 to 0.049) that showed a 6-month period

in seasonality. However, despite having a slightly worse fit, the 12-month period model of

ampicillin resistance in *E. coli* also indicated seasonality (amplitude, 0.041; 95% CI, 0.019 to

0.062) (**S3 Fig**).

Resistance peaked in the winter to spring months in all 9 seasonal species-antibiotic

combinations, with peaks ranging from early December to mid-April (**Fig 4**). Comparing across

species, resistance in *E. coli* (median phase, 3.2; range, 2.5 to 4.5) tended to peak slightly later

in the year than resistance in *S. aureus* (median phase, 1.6; range, 0.98 to 2.1) and *K.*

pneumoniae (median phase, 1.1; range, 0.0 to 2.2). Peak resistance to macrolides and

penicillins in *S. aureus* and the first peak in resistance to ampicillin in *E. coli* occurred around

the same time of year as peak use of macrolides and penicillins, with lags of -1.2 to 2.3 months.

However, resistance to ampicillin in *E. coli* also peaked a second time during the year in

October, though use did not. Resistance to nitrofurans in all 3 species peaked between 3.2 to

5.7 months after peak nitrofuran use. Finally, resistance to quinolones in all 3 species peaked

once a year about 0.8 to 2.2 months after the first peak in quinolone use.

Since the antibiotic use dataset was restricted to outpatient prescribing for people under 65

years of age, we repeated the resistance analysis on data subset to isolates from outpatients

under 65 years old, representing 53%, 31%, and 47% of the *E. coli*, *K. pneumoniae*, and *S.*

aureus isolates, respectively. Resistance to ampicillin and nitrofurantoin in *E. coli* and

nitrofurantoin in *S. aureus* remained seasonal with statistically significant seasonal amplitudes

(FDR < 0.05) and showed the same periods and phases of seasonality as in the analyses with

the full dataset including all isolates (**S1 Table**). Resistance to ciprofloxacin in *E. coli* also

remained seasonal with a statistically significant amplitude in both the 12- and 6-month period

model; however, a 6-month period model performed marginally better than the 12-month model

(Akaike information criterion (AIC) difference, 0.2). In contrast, resistance to all antibiotics in *K.*

pneumoniae and ciprofloxacin, erythromycin, and oxacillin resistance in *S. aureus* no longer met

our criterion for seasonality (amplitude FDR < 0.05) after restricting our analysis to outpatients

under 65 years old. For ciprofloxacin resistance with a 12-month period model and erythromycin

resistance with a 6-month period model in *S. aureus*, the amplitude *p* values were < 0.05, but

did not remain significant after multiple testing correction.

To further explore whether the observed seasonality of resistance could be attributable to

seasonally varied sampling of patient demographics and sites of infection, we repeated the

resistance analysis on the full dataset with all isolates after including covariates in our model to

adjust for patient age and sex (Eqn. 3 in Materials and Methods) and patient age, sex, and site

of infection (Eqn. 4 in Materials and Methods). Resistance remained seasonal for all 9 species-

antibiotic combinations after adjusting for patient age and sex, with statistically significant

amplitudes of seasonality (FDR < 0.05), though the magnitude of the estimated amplitudes

decreased by 0-32% compared to the unadjusted model (**Table 1**). After adjusting for site of

infection in addition to age and sex, resistance to ciprofloxacin, erythromycin, and oxacillin in *S.*

aureus no longer met our criterion for seasonality (amplitude FDR > 0.05) and the estimated

amplitudes decreased by 29-62% compared to the unadjusted model, while resistance in *E. coli*

and *K. pneumoniae* remained seasonal (amplitude FDR < 0.05) with a 0-14% decrease in

amplitude compared to the unadjusted model (**Table 1**). The amplitude *p* values were < 0.05 for

ciprofloxacin and oxacillin resistance in *S. aureus* after adjusting for age, sex, and site of

infection, but did not remain significant after multiple testing correction. In this model (Eqn. 4 in

Materials and Methods), the coefficient for sex was significantly positive (FDR < 0.05; where a

positive β_s indicates that being male is associated with higher MICs) across all antibiotics in *E.*

coli (median, 0.30; range, 0.053 to 0.54) and *K. pneumoniae* (median, 0.25; range, 0.18 to 0.40)

and largely non-significant in *S. aureus* (median, -2.3e-3; range, -0.079 to 0.020) (**S2 Table**).

The coefficient for age was significantly positive (FDR < 0.05; where a positive β_a indicates that

older ages are associated with higher MICs) in *E. coli* (median, 1.9e-3; range, 7.6e-4 to 0.013)

and significantly in negative in all antibiotics except ciprofloxacin in *K. pneumoniae* (median, -

1.0e-3; range, -2.7e-3 to 8.0e-4) (**S2 Table**). In *S. aureus*, the coefficient for age was

significantly positive for ciprofloxacin, erythromycin, and oxacillin resistance, significantly

negative for penicillin resistance, and non-significant for nitrofurantoin and tetracycline

resistance (median for all antibiotics in *S. aureus*, 2.3e-3; range, -9.4e-4 to 0.016). Finally, at

least one of the coefficients for site of infection was significant (FDR < 0.05) in all 15 species-

antibiotic combinations, indicating that the site of infection was also an important determinant of

MIC (**S2 Table**).

Seasonal resistance is positively correlated with use of winter-peaking antibiotic classes

Spearman correlation coefficients between use-resistance antibiotic pairs varied widely across

antibiotics, species, and lag times, ranging from -0.91 to 0.92 (**Fig 5, S4 Fig**). The number of

statistically significant correlations between use-resistance pairs was maximized when the lag

was 0 months in *S. aureus* and *K. pneumoniae* and 1 month in *E. coli*. Resistance across

multiple antibiotics was most positively correlated with use of winter-peaking classes, penicillins

and macrolides (median Spearman's ρ across all lags, 0.45; interquartile range (IQR), 0.058 to

0.76). Resistance to most antibiotics also showed a negative correlation with use of summer-

peaking classes, tetracyclines and nitrofurans (median Spearman's ρ across all lags, -0.35;

IQR, -0.63 to -0.068). Finally, resistance was not significantly correlated with use of quinolones,

which peaked twice a year, for almost all antibiotics and species (median Spearman's ρ across

all lags, -0.21; IQR, -0.32 to -0.051).

Discussion

Under a model in which antibiotic use drives resistance, seasonal variation in antibiotic consumption is expected to be associated with variation in population-level resistance that is in phase with or lagged up to a quarter period behind use [10]. However, we found that resistance to all antibiotics, including those with summer and biannual peaks in use, best correlated temporally with use of winter-peaking antibiotics—penicillins and macrolides—at a 0-to-1-month lag.

The observed patterns of use and resistance for penicillins and macrolides were mostly consistent with previous findings [6,7,9] and with model predictions [10]. Use of penicillins and macrolides peaked in the winter, likely due to increased wintertime prescribing for respiratory infections [12]. In *S. aureus*, resistance to oxacillin and erythromycin peaked in the winter and was most correlated with penicillins and macrolides use with no lag. This was consistent with a study that compared seasonal macrolide use and resistance in methicillin-resistant *S. aureus* (MRSA) [7] and findings in other species-antibiotic combinations that have shown winter peaks in use and resistance with little to no lag [6–8]. In *E. coli*, the first peak in ampicillin resistance occurred in the spring, lagging penicillins use by about 2.3 months, but also showed a second peak in resistance 6 months later, in the absence of a second peak in penicillins use. However, we note that a 12-month period model for ampicillin resistance in *E. coli* also met our criterion for seasonality and showed a single winter peak, which is more consistent with previous findings in *E. coli* [7,8].

Antibiotic classes with different seasonal patterns of use, such as nitrofurans and quinolones, showed seasonal patterns of resistance inconsistent with model predictions and not readily explained based on their patterns of use. Nitrofurans, which are almost exclusively used to treat urinary tract infections (UTIs) [16], showed summer peaks in use during the same season as

peak UTI incidence [17–20]. However, resistance to nitrofurantoin in all 3 species peaked in the

winter and lagged use by 3.2 to 5.7 months. This was consistent with a report that nitrofurantoin

resistance lagged use by 3-6 months in *E. coli* and *K. pneumoniae* urinary tract isolates [11].

Quinolones, which are used to treat both respiratory infections and UTIs [21], showed both

winter and summer peaks in use, but only a single winter/spring peak in ciprofloxacin resistance

in all 3 species.

Several species-antibiotic combinations did not show seasonality in resistance. In each case,

this may be due to a lack of association between use and resistance, a lack of strong seasonal

variation in use of some antibiotic classes, or other factors, such as a signal too small to be

identified or dampened by the combination of inpatient and outpatient samples in our dataset.

The lack of seasonality in *S. aureus* resistance to penicillin may be explained by the high

prevalence of penicillin resistance in *S. aureus* [22] (83% in this dataset), which may limit the

observable effect of increased wintertime use of penicillins. For resistance to amoxicillin-

clavulanate in *E. coli* and *K. pneumoniae*, the estimated amplitude of seasonality missed the 5%

FDR cutoff for statistical significance, but additional data may narrow the confidence intervals

around the estimated amplitude. The absence of observed seasonality in tetracycline resistance

in all 3 species is consistent with a previous study that found no significant correlation between

tetracycline use and resistance in *E. coli* [7] and may be explained by the lack of strong

seasonal variation in tetracycline use (**Fig 1**).

Our finding that resistance to all antibiotics most correlated with use of winter-peaking antibiotic

classes suggests that the simple model in which use of a given antibiotic independently selects

for resistance is insufficient to explain the full seasonal use-resistance landscape. Below, we

discuss four factors that may contribute to this result, while acknowledging that there may be

additional unknown seasonally varying determinants of resistance.

First, the lack of association between seasonal use and resistance in some antibiotics may be

an artifact of comparing between use and resistance in overlapping but not identical

populations. Data availability limited us to comparing between antibiotic use in outpatients under

age 65 and resistance measured in inpatients and outpatients at two hospitals with patient

populations that skew towards older ages. Comparisons between population-level community

use and hospital resistance have frequently been utilized in previous ecological [23–27] and

seasonal [6–8,11] studies, due to common difficulties in both obtaining within-hospital use data

and tracking community infections that do not result in a healthcare visit. The volume of

antibiotic use in the community is much greater than in the hospital [28] and thus can have

a strong impact on resistance in both settings [29,30]. This impact appears to vary across

species-antibiotic combinations and by demographics. For quinolones, we observed biannual

peaks in use and winter peaks in resistance in all three pathogens; it may be that the

populations receiving quinolones in the summer and winter are not sampled equally in our

resistance dataset. Including only outpatients under 65 years old in our resistance analysis

resulted in the loss of statistically significant seasonality in *S. aureus* and *K. pneumoniae* for

some antibiotics. While this may be attributable to reduction in signal (this subset represents

only 30-50% of the total isolates), this result could also suggest that the observed seasonal

trends in resistance for some species-antibiotic combinations are disproportionately driven by

older populations with infections associated with hospitalization. In contrast, the seasonal

patterns of resistance among outpatients under 65 largely remained the same across antibiotics

in *E. coli* and for nitrofurantoin in *S. aureus*. Thus, the disparity in community and hospital use is

unlikely to fully explain the lack of association in seasonal patterns of use and resistance.

Second, the observed seasonal peaks in resistance could be driven by seasonal variation in the

incidence of infection with a given species, although no specific mechanism for this association

has been proposed to our knowledge. In our dataset, isolate counts peaked in the summer for

all three species considered (**S5 Fig**), while most resistance peaked in the winter. Therefore, for

seasonal variations in incidence to drive seasonality in resistance, there would need to be an

inverse association between incidence and resistance; however, since there is no proposed

mechanism for this association, we did not explore it statistically.

In contrast, a third mechanism by which resistance could vary seasonally is that certain patient

demographics, or certain sites of infection, are associated with resistance and themselves vary

seasonally. Rates of resistance have been shown to vary by age, sex, and site of infection [31–

33], and, the incidence of infections from these groups have been shown to vary by season in

our dataset (**S5 Fig**) and others [20,34]. If these factors fully explained seasonal variation in

resistance, incorporating them as covariates in our seasonal regression (assuming the

association was correctly specified) should have accounted for the seasonal signal and led to

near-zero estimates for the sinusoidal component of the regression. However, resistance

remained seasonal with amplitudes decreasing by 0-32%, but remaining statistically significant,

in all 9 species-antibiotic combinations after accounting for age and sex (**Table 1**). After also

accounting for site of infection, the seasonal amplitudes of resistance in 6 of those 9

combinations remained significant, decreasing by 0-14% compared to the unadjusted model.

However, for the 3 antibiotics in *S. aureus* where resistance was no longer seasonal, the

amplitudes decreased substantially by 29-62% compared to the unadjusted model. Therefore,

seasonally varied sampling of isolates from different demographic groups and sites of infection

likely contributes to but does not fully explain the observed seasonality in resistance.

Fourth, the winter peaks in resistance to antibiotics with different seasonal peaks in use could

be explained by co-selection, where use of one antibiotic can indirectly select for resistance to a

second antibiotic in bacteria that are co-resistant to both antibiotics [35]. Co-resistance between

penicillins/macrolides and other antibiotics is common across many bacterial species, including

those in our study [36]. Therefore, the winter peaks in resistance to other antibiotics may be

driven by co-selection by winter-peaking use of penicillins and macrolides. We might expect that

selection by use of pencillins and macrolides dominates over selection by other antibiotics

because they are prescribed at substantially higher rates and show greater seasonal variations

in use [12]. Antibiotics with higher rates of use showed stronger correlations between seasonal

use and resistance [7]. In addition, use of macrolides and penicillins have been shown to be

more strongly correlated with resistance than less frequently prescribed antibiotics [23].

There were several limitations to this study. First, we measured antibiotic use in the population

by the number of claims per capita, rather than daily doses, making the assumption that the

average dose and duration do not vary greatly within the short timescales in which we are

measuring seasonal variations in use, and that there were not major selective differences

between the effects of one prescription for different members of an antibiotic class. Second, we

were unable to link antibiotic prescriptions to specific pathogens, and thus we could not assess

the extent to which antibiotic resistance in a given species is attributable to antibiotic use for the

treatment of infections caused by that species. However, given that bystander selection has

been predicted to account for over 80% of the total antibiotic selection experienced by *S.*

aureus, *E. coli*, and *K. pneumoniae* [3], the analysis we performed comparing total antibiotic use

to resistance in each of these species may in fact yield more relevant interpretations of the use-

resistance relationship [37]. Third, the antibiotic use and resistance datasets that were available

to us for each species and antibiotic often spanned overlapping but different year ranges.

Therefore, we aggregated monthly use and resistance data across years to perform our

correlation analyses. We accounted for variability in use and resistance between years by

adjusting for annual trends in use and resistance in our model.

In conclusion, this work contributes to describing the complexity of the antibiotic use-resistance

relationship. Our finding that resistance to all antibiotics peaked in the winter/spring, regardless

of patterns of use, is consistent with studies from a range of geographic scales and regions,

including South Carolina [8], the USA overall [7,9], Israel [6], and the Netherlands [11]. This

suggests a general phenomenon in which selection or co-selection by those antibiotics with

high-volume, winter-peaking use and/or other ecological factors results in wintertime peaks of

resistance for all antibiotics and indicates that the simplest model of antibiotic use independently

driving resistance to the same antibiotic is inadequate. We show that additional factors, such as

seasonal variations in the contribution of isolates from different demographic groups, may also

contribute to the observed winter peaks in resistance. This study lays the groundwork for future

work to further identify and describe the factors that shape the use-resistance landscape across

diverse pathogens and antibiotics, with important implications for informing and monitoring the

outcome of efforts to reduce antibiotic resistance.

Materials and Methods

Antibiotic use data. Outpatient antibiotic use data was obtained from the Massachusetts All

Payer Claims Database [38], which covers >94% of outpatient prescriptions claims for

Massachusetts residents under the age of 65 [39]. Rates of use for each antibiotic class were

measured as the average daily number of antibiotic claims per 10,000 people during each

calendar month from January 2011 to May 2015. This data was subset to include only

individuals residing in 'Boston City' census tracks, as defined by the US Census Bureau [40], to

capture the antibiotic use patterns in the communities served by the hospitals in our resistance

dataset. We aggregated antibiotic use data by class according to the World Health

Organization's Anatomical Therapeutic Chemical Classification System [41] (**S3 Table**). We

included 5 antibiotic classes in our analysis, which together make up 74% of the total outpatient

antibiotic claims in Boston: penicillins, macrolides, quinolones, tetracyclines, and nitrofurans.

Given that bystander selection likely accounts for most of the antibiotic selection experienced by

S. aureus, *E. coli*, and *K. pneumoniae* [3], we included use data for all antibiotics within each

class, regardless of the target pathogen for which they were prescribed.

Antibiotic resistance data. Clinical microbiology data was obtained for *S. aureus*, *E. coli*, and

K. pneumoniae isolates collected at two tertiary care hospitals in Boston, MA: Brigham and

Women's Hospital (BWH) and Massachusetts General Hospital (MGH), from 2007-2019 and

2007-2016, respectively. Included in this analysis were all non-surveillance isolates from

inpatients and outpatients of all demographics, collected from the 5 most common sites of

infection across the 3 species: blood, skin and soft tissue, abscess/fluid, respiratory tract, and

urinary tract (**S4 Table, S5 Fig**). Isolates of the same species that were collected from the same

patient within 2 weeks were assumed to represent a single infection and thus treated as a single

isolate. Our final dataset comprised of 47,374 *S. aureus*, 130,407 *E. coli*, and 27,178 *K.*

pneumoniae isolates.

Antibiotic susceptibility testing was performed on each isolate either by automated broth

microdilution (Vitek 2, Biomerieux, Marcy l'Etoile, France) or by gradient diffusion. Resulting

minimum inhibitory concentration (MIC) values were $\log_2$-transformed. When MICs were

reported with an inequality sign, we used only the numerical value in our quantitative analyses.

Due to variations in hospital testing guidelines across the years, we excluded tests on isolates

that did not report an MIC value, either because a different test method was used (e.g., disk

diameter) or due to missing data. We excluded years/months from our analysis for each

species-antibiotic combination in each hospital where MIC values were reported for fewer than

80% of isolates or only a subset of isolate types (e.g., only testing nitrofurantoin resistance in

urinary tract isolates). **S5 Table** lists date ranges and percent resistance in each hospital,

calculated as the percentage of non-susceptible isolates out of the total number of isolates from

each hospital with a reported MIC value, for each species-antibiotic combination included in our

analysis. In S5 Table, we determined antibiotic susceptibility by applying the Clinical &

Laboratory Standards Institute 2017 breakpoints [42] to the reported MIC values of each isolate;

the determined susceptibility was then adjusted based on β-lactamase screen and cefoxitin

screen results if available for penicillin and oxacillin, respectively. For all other analyses other

than in S5 Table, we use the $\log_2$-transformed MIC as the unit of resistance. This study was

approved by the Mass General Brigham Institutional Review Board (Protocol number:

2016P001671).

Statistical methods. We quantified the extent of seasonality in antibiotic use and resistance by

fitting the use and MIC data to a pair of mathematical models, based on a previously described

method [9,10]. Both models consist of (a) a sinusoidal component to describe seasonal

deviations from average year-round use and MICs and (b) a linear component to adjust for

secular trends, such as declines in use and resistance across years [22,39]. This model makes

no assumptions about the underlying mechanism of resistance and is generalizable to any

species and antibiotic [9,10]. As in the Olesen et al. model [9], we chose to substitute MIC for

the original outcome (proportion resistance) of the Blanquart et al. model [10] to allow for

detection of seasonal variations in the quantitative level of resistance, as measured by MIC,

even if that variation occurs without crossing a defined breakpoint for antibiotic susceptibility.

To describe the seasonality of use, monthly claims data for each antibiotic class were fit to

$$u_i \sim A_{use} \cos[\omega(t_i - P_{use})] + B_{y(i)}t_i + C_{y(i)} \tag{1}$$

400   where u_i is the mean daily reported claims per 10,000 people during calendar month t_i, A_{use} is

401   the amplitude of use seasonality, ω is the frequency of seasonality where $\omega = \frac{2\pi}{period}$, P_{use} is the

402   phase of use seasonality, and $B_{y(i)}$ and $C_{y(i)}$ are the within-year slope and intercept terms. To

403   describe the seasonality of resistance, MICs for each isolate were fit to

$$y_i \sim A_{MIC} \cos[\omega(t_i - P_{MIC})] + B_{h(i)}t_i + C_{h(i)} \tag{2}$$

where y_i is the log$_2$-transformed MIC and t_i is the calendar month of collection of the i^{th} isolate,

A_{MIC} is the amplitude of resistance seasonality, ω is the frequency of seasonality where $\omega =$

$\frac{2\pi}{period}$, P_{MIC} is the phase of resistance seasonality, and $B_{h(i)}$ and $C_{h(i)}$ are the within

hospital/year slope and intercept terms.

We further fit the MIC data to two additional models to account for the effect of seasonally

varied sampling of patient demographics and sites of infection on the observed seasonality of

resistance. To describe the seasonality of resistance while adjusting for patient demographics,

age and sex, we fit the MIC data to

$$y_i \sim A_{MIC} \cos[\omega(t_i - P_{MIC})] + B_{h(i)}t_i + C_{h(i)} + \beta_a a_i + \beta_s I_{m_i} \tag{3}$$

413   where β_a and β_s are the coefficients for age and sex, respectively, a_i is the age (in years) of the

414   patient from which the i^{th} isolate is collected, and I_{m_i} is an indicator variable for whether the

415   patient is male. To describe the seasonality of resistance while adjusting for patient

416   demographics and site of infection, we fit the MIC data to

$$y_i \sim A_{MIC} \cos[\omega(t_i - P_{MIC})] + B_{h(i)}t_i + C_{h(i)} + \beta_a a_i + \beta_s I_{m_i} + \tag{4}$$

$$\beta_{bl} I_{bl_i} + \beta_{rt} I_{rt_i} + \beta_{sst} I_{sst_i} + \beta_{ab} I_{ab_i}$$

where β_{bl}, β_{rt}, β_{sst}, and β_{ab} are the coefficients for blood, respiratory tract, skin and soft tissue,

and abscess/fluid, respectively, and I_{bl_i}, I_{rt_i}, I_{sst_i}, and I_{ab_i} are the indicator variables for whether

the i^{th} isolate was collected from the blood, respiratory tract, skin and soft tissue, or

abscess/fluid, respectively.

The amplitude, phase, slope, intercept, and demographic coefficient terms in each model were

estimated by non-linear regression, using the *nls* function in R (version 3.6.2) [43]. We

examined periods of both 12 and 6 months to account for annual or biannual cycles in use and

resistance. We justified using these fixed periods by performing a wavelet analysis, using the

WaveletComp package [44] in R, on the raw antibiotic use data to show that the dominant

periods of variations in use across the included years are at 12 and 6 months (**S6 Fig**). To

determine whether to use a 12- or 6-month period for each species-antibiotic combination, we

performed model comparisons using the Akaike information criterion (AIC) and used the period

that resulted in the lower AIC (**S6 Table**, **S7 Table**). We determined that there was seasonality

in use or resistance if the amplitude was statistically significant after accounting for multiple

comparisons by applying the Benjamini-Hochberg correction with a 5% false discovery rate

(FDR).

We quantified the association between the observed seasonal patterns of use and resistance

using Spearman's rank correlations. To eliminate the impact of annual trends, we calculated

correlations between the average monthly seasonal deviates in use and resistance, aggregated

across years, rather than the raw use and MIC data. We define a 'seasonal deviate' as the

deviation in use or MIC at a given time of year from the year-round average, which we

estimated by the linear component of the models. Seasonal deviates in use for each year and

month were calculated as

$$u_i' = u_i - \widehat{B_{y(i)}}t_i - \widehat{C_{y(i)}} \tag{5}$$

442   where u_i' is the seasonal deviate of the mean reported daily claims per 10,000 people during

443   calendar month t_i, u_i is the mean reported daily claims per 10,000 people during calendar

444   month t_i, and $\widehat{B_{y(i)}}$ and $\widehat{C_{y(i)}}$ are the within-year slope and intercept terms estimated from the

445   model fit (Eqn. 1) for the corresponding year. For resistance, we calculate the seasonal deviate

446   for each isolate as

$$y_i' = y_i - \widehat{B_{h(i)}}t_i - \widehat{C_{h(i)}} \tag{6}$$

where y_i' is the seasonal deviate of the log$_2$-transformed MIC of the i^{th} isolate, y_i is the log$_2$-

transformed MIC of the i^{th} isolate, t_i is the calendar month of collection of the i^{th} isolate, and

$\widehat{B_{h(i)}}$ and $\widehat{C_{h(i)}}$ are the hospital/year slope and intercept estimated from the model fit (Eqn. 2) for

the hospital and year that the i^{th} isolate was collected in.

Since the working model for the use-resistance relationship predicts that seasonal fluctuations

in resistance can lag use by up to 3 months [10], we calculated Spearman correlations between

use and resistance seasonal deviates with no lag and lags of 1, 2, and 3 months. In addition,

because we observed some seasonal patterns of resistance that better aligned with use of non-

cognate antibiotic classes, we calculated use-resistance correlations between each pairwise

combination of target antibiotics and use classes. We only included use-resistance pairs in this

analysis for which both use and resistance met our criterion for seasonality.

All analyses were performed in R version 3.6.2 [43]. Data and code are available at:

https://github.com/gradlab/use-resistance-seasonality

Acknowledgements. The authors thank R. Monina Klevens for assistance in acquiring the

antibiotic prescribing data and Enterprise Research Infrastructure & Services at Partners

HealthCare for their computational resources and support.

Funding. This work was supported by funds from the HSPH Dean's Award (to Y.H.G.) and

Wellcome Trust (to Y.H.G.). D.S.S. is supported by the National Institutes of Health training

grant (T32AI132120).

Conflicts of interest. Y. H. G. and M. L. have received grant support from Pfizer. S. W. O. is

an employee of Biobot Analytics, Inc. The other authors declare no conflicts of interest.

References

1. U.S. Centers for Disease Control and Prevention. Antibiotic resistance threats in the United States. 2019. Available from: https://www.cdc.gov/drugresistance/pdf/threats-report/2019-ar-threats-report-508.pdf

2. Jit M, Ng DHL, Luangasanatip N, Sandmann F, Atkins KE, Robotham JV, et al. Quantifying the economic cost of antibiotic resistance and the impact of related interventions: Rapid methodological review, conceptual framework and recommendations for future studies. BMC Med. 2020;18(1):38.

3. Tedijanto C, Olesen SW, Grad YH, Lipsitch M. Estimating the proportion of bystander selection for antibiotic resistance among potentially pathogenic bacterial flora. Proc Natl Acad Sci USA. 2018;115(51):E11988–95.

4. Hennessy TW, Petersen KM, Bruden D, Parkinson AJ, Hurlburt D, Getty M, et al. Changes in antibiotic-prescribing practices and carriage of penicillin-resistant Streptococcus pneumoniae: A controlled intervention trial in rural Alaska. Clin Infect Dis. 2002;34(12):1543–50.

5. Enne VI, Livermore DM, Stephens P, Hall LMC. Persistence of sulphonamide resistance in Escherichia coli in the UK despite national prescribing restriction. Lancet. 2001;357(9265):1325–28.

6. Dagan R, Barkai G, Givon-Lavi N, Sharf AZ, Vardy D, Cohen T, et al. Seasonality of Antibiotic-Resistant Streptococcus pneumoniae That Causes Acute Otitis Media: A Clue for an Antibiotic-Restriction Policy? J Infect Dis. 2008;197(8):1094–102.

7. Sun L, Klein EY, Laxminarayan R. Seasonality and Temporal Correlation between Community Antibiotic Use and Resistance in the United States. Clin Infect Dis. 2012;55(5):687–94.

8. Ramsey EG, Royer J, Bookstaver PB, Justo JA, Kohn J, Albrecht H, et al. Seasonal variation in antimicrobial resistance rates of community-acquired Escherichia coli bloodstream isolates. Int J Antimicrob Agents. 2019;54(1):1–7.

9. Olesen SW, Torrone EA, Papp JR, Kirkcaldy RD, Lipsitch M, Grad YH. Azithromycin Susceptibility Among Neisseria gonorrhoeae Isolates and Seasonal Macrolide Use. J Infect Dis. 2019;219(4):619–23.

10. Blanquart F, Lehtinen S, Fraser C. An evolutionary model to predict the frequency of antibiotic resistance under seasonal antibiotic use, and an application to Streptococcus pneumoniae. Proc R Soc B Biol Sci. 2017;284:20170679.

11. Martínez EP, van Rosmalen J, Bustillos R, Natsch S, Mouton JW, Verbon A. Trends, seasonality and the association between outpatient antibiotic use and antimicrobial resistance among urinary bacteria in the Netherlands. J Antimicrob Chemother. 2020;75(8):2314–25.

12. Suda KJ, Hicks LA, Roberts RM, Hunkler RJ, Taylor TH. Trends and Seasonal Variation in Outpatient Antibiotic Prescription Rates in the United States, 2006 to 2010. Antimicrob Agents Chemother. 2014;58(5):2763–6.

13. Kluytmans J, Van Belkum A, Verbrugh H. Nasal Carriage of I: Epidemiology, Underlying Mechanisms, and Associated Risks. Clin Microbiol Rev. 1997;10(3):505–20.

14. Huttenhower C, Gevers D, Knight R, et al. Structure, function and diversity of the healthy human microbiome. Nature. 2012;486(7402):207–14.

15. Berchick ER, Barnett JC, Upton RD. Current population reports, P60-267 (RV), Health Insurance Coverage in the United States: 2018. Washington, DC: US Government Printing Office, 2019. Available at: https://www.census.gov/content/dam/Census/library/publications/2019/demo/p60-267.pdf

16. Guay DR. An Update on the Role of Nitrofurans in the Management of Urinary Tract
Infections. Drugs. 2001;61(3):353–64.

17. Stamm WE, McKevitt M, Roberts PL, White NJ. Natural History of Recurrent Urinary
Tract Infections in Women. Rev Infect Dis. 1991;13(1):77–84.

18. Anderson JE. Seasonality of symptomatic bacterial urinary infections in women. J
Epidemiol Community Health. 983;37:286–90.

19. Czaja CA, Scholes D, Hooton TM, Stamm WE. Population-Based Epidemiologic
Analysis of Acute Pyelonephritis. Clin Infect Dis. 2007;45(3):273–80.

20. Rosello A, Pouwels KB, Domenech De Cellès M, Van Kleef E, Hayward AC, Hopkins S,
et al. Seasonality of urinary tract infections in the United Kingdom in different age
groups: Longitudinal analysis of The Health Improvement Network (THIN). Epidemiol
Infect 2018;146(1):37–45.

21. Hooper DC. Clinical applications of quinolones. Biochim Biophys Acta. 1998;1400(1):45-
61.

22. Kanjilal S, Sater MRA, Thayer M, Lagoudas GK, Kim S, Blainey PC, et al. Trends in
Antibiotic Susceptibility in Staphylococcus Aureus in Boston, Massachusetts, from 2000
to 2014. J Clin Microbiol. 2018;56(1):e01160-17.

23. Olesen SW, Barnett ML, Macfadden DR, Brownstein JS, Hernandez-Diaz S, Lipsitch M,
et al. The distribution of antibiotic use and its association with antibiotic resistance. eLife.
2018;7:e39435.

24. Hicks LA, Chien Y-W, Taylor TH, Haber M, Klugman KP. Outpatient Antibiotic
Prescribing and Nonsusceptible Streptococcus pneumoniae in the United States, 1996–
2003. Clin Infect Dis. 2011;53(7):631–9.

25. Bronzwaer SLAM, Cars O, Buchholz U, Molstad S, Goettsch W, Veldhuijzen IK, et al.
The relationship between antimicrobial use and antimicrobial resistance in Europe.
Emerg Infect Dis. 2002;8(3):278–82.

26. Bergman M, Huikko S, Pihlajamäki M, Laippala P, Palva E, Huovinen P, et al. Effect of

macrolide consumption on erythromycin resistance in Streptococcus pyogenes in

Finland in 1997-2001. Clin Infect Dis. 2004;38(9):1251–6.

27. MacDougall C, Powell JP, Johnson CK, Edmond MB, Polk RE. Hospital and community

fluoroquinolone use and resistance in Staphylococcus aureus and Escherichia coli in 17

US hospitals. Clin Infect Dis. 20015;41(4):435–40.

28. ESPAUR Writing Committee. English Surveillance Programme for Antimicrobial

Utilisation and Resistance (ESPAUR) Report 2020-2021. Public Health England. 2021.

Available from:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment

_data/file/1033851/espaur-report-2020-to-2021-16-Nov.pdf

29. Knight GM, Costelloe C, Deeny SR, Moore LSP, Hopkins S, Johnson AP, et al.

Quantifying where human acquisition of antibiotic resistance occurs: A mathematical

modelling study. BMC Med. 2018;16(1):137.

30. Macfadden DR, Fisman DN, Hanage WP, Lipsitch M. The Relative Impact of Community

and Hospital Antibiotic Use on the Selection of Extended-spectrum Beta-lactamase-

producing Escherichia coli. Clin Infect Dis. 2019;69(1):182–8.

31. Mcgregor JC, Elman MR, Bearden DT, Smith DH. Sex- and age-specific trends in

antibiotic resistance patterns of Escherichia coli urinary isolates from outpatients. BMC

Fam Pract. 2013;14:25.

32. Garcia A, Delorme T, Nasr P. Patient age as a factor of antibiotic resistance in

methicillin-resistant Staphylococcus aureus. J Med Microbiol. 2017;66(12):1782–9.

33. Lagacé-Wiens PRS, DeCorby MR, Baudry PJ, Hoban DJ, Karlowsky JA, Zhanel GG.

Differences in antimicrobial susceptibility in Escherichia coli from Canadian intensive

care units based on regional and demographic variables. Can J Infect Dis Med Microbiol.

2008;19(4):282–6.

34. Leekha S, Diekema DJ, Perencevich EN. Seasonality of staphylococcal infections. Clin

Microbiol Infec.. 2012;18(10):927–33.

35. Pouwels KB, Muller-Pebody B, Smieszek T, Hopkins S, Robotham J V. Selection and

co-selection of antibiotic resistances among Escherichia coli by antibiotic use in primary

care: An ecological analysis. PLOS One. 2019;14(6):e0218134.

36. Chang H-H, Cohen T, Grad YH, Hanage WP, O'Brien TF, Lipsitch M. Origin and

Proliferation of Multiple-Drug Resistance in Bacterial Pathogens. Microbiol Mol Biol Rev.

2015;79(1):101–16.

37. Schechner V, Temkin E, Harbarth S, Carmeli Y, Schwaber MJ. Epidemiological

Interpretation of Studies Examining the Effect of Antibiotic Usage on Resistance. Clin

Microbiol Rev. 2013;26(2):289.

38. Center for Health Information and Analysis. Massachusetts All Payer Claims Database.

2019. Available from: https://www.chiamass.gov/ma-apcd/

39. Klevens RM, Caten E, Olesen SW, DeMaria A, Troppy S, Grad YH. Outpatient Antibiotic

Prescribing in Massachusetts, 2011–2015. Open Forum Infect Dis. 2019;6(5):ofz159.

40. US Census Bureau. American fact finder. 2020. Available from:

https://data.census.gov/cedsci/

41. World Health Organization Collaborating Centre for Drug Statistics Methodology.

Structure and Principles. 2020. Available from:

https://www.whocc.no/atc/structure_and_principles/

42. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial

Susceptibility Testing. 27th ed. CLSI M100. Wayne, PA. 2017.

43. R Core Team. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria.

44. Roesch A, Schmidbauer H. WaveletComp. 2018. Available from: https://cran.r-

project.org/web/packages/WaveletComp/WaveletComp.pdf

Figures

603
Fig 1. **Seasonal patterns of antibiotic use by class.** (A) Average daily antibiotic claims per
10,000 people by calendar month in Boston, MA from 2011 to 2015. Lines indicate LOESS
smoothing curves and shaded regions indicate 95% confidence intervals. (B) Sinusoidal model
fits for monthly prescribing rate. Points indicate monthly mean seasonal deviates in average
daily antibiotic claims per 10,000 people by calendar month and error bars indicate the standard
error of the mean. Lines indicate the point estimate for the amplitude and phase of the
sinusoidal model. Shaded regions indicate the 95% confidence intervals for the amplitude.
Asterisks indicate the amplitude of seasonality is statistically significant (FDR < 0.05).

612
Fig 2. Seasonality of antibiotic use and resistance by class in *Staphylococcus aureus*.
613
Solid lines indicate point estimates of the amplitude and phase from the best-fitting sinusoidal
model of resistance (comparing 6- and 12-month periods) to each antibiotic, colored by class.
Dashed grey lines indicate point estimates of the amplitude and phase of sinusoidal models for
use of the corresponding antibiotic class. Shaded regions indicate the 95% confidence intervals
for the amplitude. Points indicate the monthly mean seasonal deviates in resistance and error
bars indicate the standard error of the mean. Asterisks indicate the amplitude of seasonality in
resistance is statistically significant (FDR < 0.05).

Fig 3. Amplitudes of seasonality of resistance by species and antibiotic class.
Comparison of amplitudes estimated from best-fitting sinusoidal models of resistance
(comparing 6- and 12-month periods) across antibiotics in *Staphylococcus aureus*, *Escherichia coli*, and *Klebsiella pneumoniae*. Error bars indicate 95% confidence intervals of the amplitude.
Point color indicates the antibiotic class. AMC, Amoxicillin-Clavulanate; AMP, Ampicillin; CIP,
Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin; PEN, Penicillin; TET,
Tetracycline.

Fig 4. Phases of seasonality of use and resistance by species and antibiotic class. Points indicate peak month(s) of seasonal resistance estimated by the best-fitting sinusoidal model (comparing 6- and 12-month periods) for each species-antibiotic combination, and error bars indicate the 95% confidence intervals. Vertical lines indicate the peak month(s) of seasonal use estimated by the best-fitting sinusoidal model (comparing 6- and 12-month periods) for each antibiotic class, and the shaded regions indicate the 95% confidence intervals. Included are species-antibiotic combinations for which the amplitude was statistically significant (FDR < 0.05). AMC, Amoxicillin-Clavulanate; AMP, Ampicillin; CIP, Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin.

Fig 5. Spearman correlations between seasonal use and resistance with 0-3 months lag in *S. aureus*.
Spearman's rank correlation coefficients were calculated between the monthly mean seasonal deviate in resistance (in $\log_2$(MIC)) and the monthly mean seasonal deviate in use (in average daily claims per 10,000 people) with 0, 1, 2, or 3 months lag between use and resistance, for each pairwise combination of antibiotics and classes. Error bars indicate the 95% confidence intervals. Colors indicate the use antibiotic class. Mac, Macrolides; Nit, Nitrofurantoin; Pen, Penicillins; Qui, Quinolones; Tet, Tetracyclines. CIP, Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin.

Table 1. Comparison of estimated amplitudes of seasonality across three sinusoidal models for resistance.

Species	Abx	Period (months)	Amplitude (95% CI)		
			Model A (unadjusted)	**Model B** (adjusted for age and sex)	**Model C** (adjusted for age, sex, and site of infection)
E. coli	AMC	6	0.01 (-5.1e-04, 0.021)	0.011 (4.3e-04, 0.022)	0.011 (6.3e-04, 0.022)
E. coli	AMP	6	0.034 (0.019, 0.049) *	0.034 (0.019, 0.048) *	0.034 (0.019, 0.048) *
E. coli	CIP	12	0.051 (0.031, 0.072) *	0.045 (0.026, 0.065) *	0.044 (0.025, 0.064) *
E. coli	NIT	12	0.028 (0.02, 0.037) *	0.028 (0.02, 0.036) *	0.028 (0.02, 0.036) *
E. coli	TET	6	0.013 (-3.6e-03, 0.029)	0.013 (-3.4e-03, 0.029)	0.013 (-3.3e-03, 0.029)
K. pneumoniae	AMC	12	0.034 (1.2e-03, 0.067)	0.034 (1.3e-03, 0.066)	0.032 (-2.1e-04, 0.065)
K. pneumoniae	CIP	12	0.053 (0.023, 0.083) *	0.05 (0.02, 0.081) *	0.048 (0.018, 0.078) *
K. pneumoniae	NIT	12	0.035 (9.6e-03, 0.061) *	0.034 (7.9e-03, 0.06) *	0.034 (7.6e-03, 0.06) *
K. pneumoniae	TET	6	0.021 (-9.8e-03, 0.051)	0.019 (-0.011, 0.049)	0.019 (-0.012, 0.049)
S. aureus	CIP	12	0.063 (0.034, 0.093) *	0.043 (0.016, 0.07) *	0.024 (5.0e-04, 0.048)
S. aureus	ERY	12	0.048 (0.012, 0.083) *	0.042 (7.7e-03, 0.077) *	0.032 (-1.0e-03, 0.065)
S. aureus	NIT	12	0.033 (0.022, 0.043) *	0.034 (0.023, 0.044) *	0.038 (0.027, 0.048) *
S. aureus	OXA	12	0.031 (8.8e-03, 0.054) *	0.027 (5.5e-03, 0.049) *	0.023 (8.2e-04, 0.044)
S. aureus	PEN	6	0.01 (-6.6e-03, 0.027)	9.5e-03 (-7.3e-03, 0.026)	9.7e-03 (-7.1e-03, 0.027)
S. aureus	TET	12	0.013 (-7.2e-03, 0.033)	0.012 (-7.3e-03, 0.032)	0.013 (-6.9e-03, 0.033)

Model A does not adjust for the patient demographics or site of infection, Model B adjusts for patient age and sex, and Model C adjusts for patient age, sex, and site of infection. In parentheses are the 95% confidence intervals on the amplitude estimates. Asterisks indicate that the amplitude is significant after Benjamini-Hochberg multiple testing correction (false discovery rate < 0.05). AMC, Amoxicillin-Clavulanate; AMP, Ampicillin; CIP, Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin; PEN, Pencillin; TET, Tetracycline.

659
S1 Fig. Seasonality of antibiotic use and resistance by class in *Escherichia coli*. Solid
lines indicate point estimates of the amplitude and phase from the best-fitting sinusoidal model
of resistance (comparing 6- and 12-month periods) to each antibiotic, colored by class. Dashed
grey lines indicate point estimates of the amplitude and phase from sinusoidal models of use of
the corresponding antibiotic class. Shaded regions indicate the 95% confidence intervals for the
amplitude. Points indicate the monthly mean seasonal deviates in resistance and error bars
indicate the standard error of the mean. Asterisks indicate the amplitude of seasonality in
resistance is statistically significant (FDR < 0.05). Amox/Clav, Amoxicillin-Clavulanate.

668
S2 Fig. Seasonality of antibiotic use and resistance by class in *Klebsiella pneumoniae.*
Solid lines indicate point estimates of the amplitude and phase from the best-fitting sinusoidal
model (comparing 6- and 12-month periods) of resistance to each antibiotic, colored by class.
Dashed grey lines indicate point estimates of the amplitude and phase from sinusoidal models
of use of the corresponding antibiotic class. Shaded regions indicate the 95% confidence
intervals for the amplitude. Points indicate the monthly mean seasonal deviates in resistance
and error bars indicate the standard error of the mean. Asterisks indicate the amplitude of
seasonality in resistance is statistically significant (FDR < 0.05). Amox/Clav, Amoxicillin-
Clavulanate.

678
S3 Fig. Seasonality of penicillins use and ampicillin resistance in *Escherichia coli* with a
12-month period model. Solid line indicates point estimates of the amplitude and phase from a
12-month period sinusoidal model of resistance to ampicillin in *E. coli*. Dashed grey line
indicates point estimates of the amplitude and phase from a 12-month period sinusoidal model
of use of penicillin class antibiotics. Shaded regions indicate the 95% confidence intervals for
the amplitude. Points indicate the monthly mean seasonal deviates in resistance and error bars
indicate the standard error of the mean. Asterisk indicates the amplitude of seasonality in
resistance is statistically significant (FDR < 0.05).

S4 Fig. Spearman correlations between seasonal use and resistance with 0-3 months lag in (A) *E. coli* **and (B)** *K. pneumoniae.* Spearman's rank correlation coefficients were calculated between the monthly mean seasonal deviate in resistance (in $\log_2$(MIC)) and the monthly mean seasonal deviate in use (in average daily claims per 10,000 people) with 0, 1, 2, or 3 months lag between use and resistance, for each pairwise combination of antibiotics and classes. Error bars indicate the 95% confidence intervals. Colors indicate the use antibiotic class. Mac, Macrolides; Nit, Nitrofurantoin; Pen, Penicillins; Qui, Quinolones; Tet, Tetracyclines. AMP, Ampicillin; CIP, Ciprofloxacin; NIT, Nitrofurantoin.

696
697 **S5 Fig. Seasonal incidence of infection by demographic group or site of infection.** Bars
698 show the total number of isolates by month included in the resistance dataset for each species,
699 colored by (A) age group, (B) sex, and (C) site of infection. NOS, not otherwise specified.
700

701
702
703 **S6 Fig. Wavelet analysis of antibiotic use by class.** Dotted lines show 12-month (upper line)
704 and 6-month (lower line) periods. Solid lines indicate regions where the amplitude *p* value is
705 less than 0.05. Shaded areas indicate the 'cone of influence' where edge effects are important.

S1 Table. Amplitudes and phases of seasonality of resistance in outpatients under 65 years old.

Species	Antibiotic	Period	Amplitude (95% CI)	Phase (95% CI)
E. coli	AMC	6 months	0.013 (-1.7e-03, 0.027)	2.7 (1.6, 3.7)
E. coli	AMP	6 months	0.039 (0.019, 0.06) *	4.3 (3.8, 4.8)
E. coli	CIP	6 months	0.022 (5.9e-03, 0.037) *	4.6 (3.9, 5.2)
E. coli	NIT	12 months	0.033 (0.022, 0.044) *	2.6 (2.1, 3.1)
E. coli	TET	6 months	0.013 (-7.9e-03, 0.034)	3.6 (2, 5.3)
K. pneumoniae	AMC	12 months	0.028 (-0.027, 0.082)	4 (1.6, 6.4)
K. pneumoniae	CIP	6 months	0.023 (-0.014, 0.059)	4.8 (3.4, 6.2)
K. pneumoniae	NIT	6 months	0.027 (-0.013, 0.067)	5.5 (4.1, 7)
K. pneumoniae	TET	12 months	0.062 (-0.012, 0.14)	8.3 (6.2, 10)
S. aureus	CIP	12 months	0.043 (2.2e-03, 0.085)	2.2 (0.46, 3.9)
S. aureus	ERY	6 months	0.056 (8.9e-03, 0.1)	2.6 (1.8, 3.4)
S. aureus	NIT	12 months	0.041 (0.025, 0.058) *	2.4 (1.7, 3.1)
S. aureus	OXA	12 months	0.03 (-0.017, 0.078)	11 (8.2, 13)
S. aureus	PEN	12 months	0.031 (-5.7e-03, 0.069)	9.1 (7.5, 11)
S. aureus	TET	12 months	0.019 (-8.6e-03, 0.047)	11 (8.4, 13)

Amplitudes and phases were estimated from the best-fitting sinusoidal model of resistance (comparing 6- and 12-month periods) for each species-antibiotic combination, using data that was subset to include only outpatients under age 65. Asterisks indicate that the amplitude is significant after Benjamini-Hochberg multiple testing correction (false discovery rate < 0.05). AMC, Amoxicillin-Clavulanate; AMP, Ampicillin; CIP, Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin; PEN, Pencillin; TET, Tetracycline.

S2 Table. Age, sex, and site of infection coefficients for adjusted sinusoidal model of seasonal resistance.

Species	Abx	Period	β_a	β_s	β_{bl}	β_{rt}	β_{sst}	β_{ab}
E. coli	AMC	6 months	2.0e-03 (1.7e-03, 2.3e-03) *	0.25 (0.23, 0.27) *	0.23 (0.17, 0.28) *	0.35 (0.3, 0.41) *	0.045 (-2.6e-03, 0.093)	0.12 (0.059, 0.18) *
E. coli	AMP	6 months	1.9e-03 (1.5e-03, 2.3e-03) *	0.34 (0.32, 0.37) *	0.22 (0.15, 0.3) *	0.43 (0.36, 0.51) *	-4.3e-04 (-0.063, 0.063)	0.062 (-0.019, 0.14)
E. coli	CIP	12 months	0.013 (0.013, 0.014) *	0.54 (0.51, 0.56) *	0.4 (0.34, 0.47) *	0.64 (0.58, 0.7) *	0.11 (0.056, 0.17) *	0.099 (0.027, 0.17) *
E. coli	NIT	12 months	7.6e-04 (5.8e-04, 9.3e-04) *	0.053 (0.042, 0.065) *	-0.05 (-0.08, -0.02) *	-0.091 (-0.12, -0.063) *	-0.05 (-0.077, -0.023) *	-0.077 (-0.11, -0.043) *
E. coli	TET	6 months	1.1e-03 (6.0e-04, 1.5e-03) *	0.3 (0.27, 0.33) *	0.34 (0.26, 0.42) *	0.28 (0.2, 0.36) *	0.04 (-0.031, 0.11)	-0.056 (-0.15, 0.035)
K. pneumoniae	AMC	12 months	-8.2e-04 (-1.5e-03, -1.1e-04) *	0.24 (0.21, 0.27) *	0.13 (0.065, 0.2) *	0.29 (0.24, 0.34) *	0.14 (0.078, 0.21) *	0.16 (0.077, 0.25) *
K. pneumoniae	CIP	12 months	8.0e-04 (3.4e-05, 1.6e-03) *	0.4 (0.37, 0.44) *	0.079 (9.0e-03, 0.15) *	0.27 (0.22, 0.32) *	-0.041 (-0.11, 0.03)	0.044 (-0.044, 0.13)
K. pneumoniae	NIT	12 months	-1.2e-03 (-2.0e-03, -3.9e-04) *	0.18 (0.15, 0.22) *	0.08 (3.7e-03, 0.16) *	0.11 (0.056, 0.17) *	-3.7e-03 (-0.085, 0.077)	0.063 (-0.038, 0.16)
K. pneumoniae	TET	6 months	-2.7e-03 (-3.7e-03, -1.6e-03) *	0.25 (0.21, 0.3) *	5.0e-03 (-0.091, 0.1)	0.18 (0.11, 0.25) *	0.064 (-0.032, 0.16)	0.023 (-0.099, 0.14)
S. aureus	CIP	12 months	0.016 (0.015, 0.017) *	0.02 (-0.011, 0.051)	-0.79 (-0.87, -0.71) *	-0.34 (-0.4, -0.28) *	-1.1 (-1.1, -1) *	-0.9 (-0.97, -0.83) *
S. aureus	ERY	12 months	4.9e-03 (3.9e-03, 5.9e-03) *	-0.079 (-0.12, -0.035) *	-0.41 (-0.53, -0.29) *	-0.031 (-0.12, 0.056)	-0.57 (-0.66, -0.49) *	-0.049 (-0.15, 0.053)
S. aureus	NIT	12 months	-7.5e-06 (-2.4e-04, 2.2e-04)	-8.1e-04 (-0.011, 9.9e-03)	0.21 (0.18, 0.24) *	0.076 (0.056, 0.097) *	0.22 (0.2, 0.24) *	0.25 (0.22, 0.27) *
S. aureus	OXA	12 months	4.6e-03 (3.9e-03, 5.3e-03) *	-3.9e-03 (-0.034, 0.027)	-0.39 (-0.47, -0.3) *	-0.19 (-0.25, -0.13) *	-0.54 (-0.6, -0.48) *	0.026 (-0.045, 0.096)
S. aureus	PEN	6 months	-9.4e-04 (-1.4e-03, -4.5e-04) *	-2.9e-04 (-0.022, 0.022)	-0.034 (-0.093, 0.026)	0.036 (-7.9e-03, 0.08)	-0.025 (-0.066, 0.017)	0.093 (0.042, 0.14) *
S. aureus	TET	12 months	-3.1e-04 (-7.1e-04, 9.2e-05)	-0.013 (-0.03, 5.3e-03)	-0.058 (-0.11, -9.5e-03) *	-0.055 (-0.09, -0.02) *	-0.025 (-0.059, 8.0e-03)	-0.09 (-0.13, -0.048) *

Demographic and site of infection coefficients were estimated from the sinusoidal model of
resistance that was adjusted for patient age, sex, and site of infection (Eqn. 4 in Materials and
Methods). In parentheses are the 95% confidence intervals on the coefficient estimates.
Asterisks indicate that the coefficient is significant after Benjamini-Hochberg multiple testing
correction (false discovery rate < 0.05). β_a, coefficient for patient age; β_s, coefficient for patient
sex; β_{bl}, coefficient for if the isolate was a blood isolate; β_{rt}, coefficient for if the isolate was a
respiratory tract isolate; β_{sst}, coefficient for if the isolate was a skin/soft tissue isolate; β_{ab},
coefficient for if the isolate was an abscess/fluid isolate; AMC, Amoxicillin-Clavulanate; AMP,
Ampicillin; CIP, Ciprofloxacin; ERY, Erythromycin; NIT, Nitrofurantoin; OXA, Oxacillin; PEN,
Pencillin; TET, Tetracycline.
727

S3 Table. Percent of claims by individual antibiotics within each class.

Antibiotic class	Antibiotic name	ATC code	% of total use within antibiotic class
Penicillins	Amoxicillin	J01CA04	83
	Phenoxymethylpenicillin	J01CE02	16
	Dicloxacillin	J01CF01	1.1
	Ampicillin	J01CA01	0.35
	Benzylpenicillin	J01CE01	0.091
	Piperacillin	J01CA12	0.036
	Naficillin	J01CF06	0.024
	Oxacillin	J01CF04	3.0e-03
Macrolides	Azithromycin	J01FA10	51
	Clindamycin	J01FF01	28
	Erythromycin	J01FA01	17
	Clarithromycin	J01FA09	3.8
	Lincomycin	J01FF02	1.1e-03
	Telithromycin	J01FA15	7.0e-04
Quinolones	Ciprofloxacin	J01MA02	67
	Levofloxacin	J01MA12	14
	Ofloxacin	J01MA01	13
	Moxifloxacin	J01MA14	5.7
	Gatifloxacin	J01MA16	0.85
	Norofloxacin	J01MA06	0.071
	Gemifloxacin	J01MA15	9.8e-03
Tetracyclines	Doxycycline	J01AA02	79
	Minocycline	J01AA08	19
	Tetracycline	J01AA07	2
	Tigecycline	J01AA12	0.054
	Demeclocycline	J01AA01	0.053
Nitrofurans	Nitrofurantoin	J01XE01	100

Antibiotic claims data for Boston, MA residents was obtained from the Massachusetts All Payers Claims Database and subset by antibiotic class according to the following WHO ATC codes [41]: penicillins, J01C; macrolides, J01F; quinolones, J01M; tetracyclines, J01A; and nitrofurans, JO1XE. All antibiotics under each ATC header that were present in this dataset were included in the analysis. The percentage of the total claims within each antibiotic class made-up by each individual antibiotic is listed in the last column of the table.

 S4 Table. Total number of isolates by demographics.

		E. coli	*K. pneumoniae*	*S. aureus*
Total	Total	130407	27178	47374
Hospitals	BWH	53428 (41%)	11716 (43.1%)	17741 (37.4%)
	MGH	76979 (59%)	15462 (56.9%)	29633 (62.6%)
Patient type	Inpatient	23146 (17.7%)	9498 (34.9%)	16897 (35.7%)
	Outpatient	107261 (82.3%)	17680 (65.1%)	30477 (64.3%)
Site of infection	Abscess or fluid NOS	1971 (1.5%)	806 (3%)	5152 (10.9%)
	Blood	2617 (2%)	1419 (5.2%)	2802 (5.9%)
	Respiratory tract	2643 (2%)	2859 (10.5%)	11598 (24.5%)
	Skin and soft tissue	3314 (2.5%)	1324 (4.9%)	23841 (50.3%)
	Urinary tract	119862 (91.9%)	20770 (76.4%)	3981 (8.4%)
Age group	00-19	7629 (5.9%)	708 (2.6%)	4455 (9.4%)
	20-39	31247 (24%)	3003 (11%)	10496 (22.2%)
	40-64	41591 (31.9%)	8977 (33%)	17312 (36.5%)
	65+	49940 (38.3%)	14490 (53.3%)	15111 (31.9%)
Sex	Female	109948 (84.3%)	18827 (69.3%)	21565 (45.5%)
	Male	20459 (15.7%)	8351 (30.7%)	25809 (54.5%)

737
738 Table of the total number of isolates of each species within each clinical or demographic
739 category that was used in this analysis. In parentheses is the percent of the total number of
740 isolates for that species. BWH, Brigham and Women's Hospital; MGH, Massachusetts General
741 Hospital; NOS, not otherwise specified.

742 **S5 Table. Antibiotics included in analysis and percent resistance by hospital.**

Species	Antibiotic (Antibiotic class)	Dates included at BWH	Dates included at MGH	Percent resistance at BWH	Percent resistance at MGH
E. coli	Amoxicillin/ Clavulanate (Penicillins)	May 2013 – Dec 2019	Jan 2007 – Dec 2016	17.8%	17.2%
	Ampicillin (Penicillins)	Jan 2007 – Dec 2019	Jan 2007 – Dec 2016	47.8%	47.0%
	Ciprofloxacin (Quinolones)	Jan 2007 – Dec 2019	Jan 2007 – Dec 2016	22.6%	21.1%
	Nitrofurantoin (Nitrofurans)	Jan 2007 – Jun 2013	Jan 2007 – Dec 2016	6.0%	5.1%
	Tetracycline (Tetracyclines)	May 2013 – Dec 2019	Jan 2007 – Dec 2016	29.1%	29.3%
K. pneumoniae	Amoxicillin/ Clavulanate (Penicillins)	May 2013 – Dec 2019	Jan 2007 – Dec 2016	10.2%	8.2%
	Ciprofloxacin (Quinolones)	Jan 2007 – Dec 2019	Jan 2007 – Dec 2016	11.8%	11.1%
	Nitrofurantoin (Nitrofurans)	Jan 2007 – Jun 2013	Jan 2007 – Dec 2016	69.4%	68.0%
	Tetracycline (Tetracyclines)	May 2013 – Dec 2019	Jan 2007 – Dec 2016	21.3%	18.3%
S. aureus	Ciprofloxacin (Quinolones)	May 2010 – Dec 2019	Jan 2009 – Dec 2016	29.6%	30.7%
	Erythromycin (Macrolides)	May 2010 – Dec 2019	Jan 2009 – Dec 2016	51.2%	54.9%
	Nitrofurantoin (Nitrofurans)	May 2010 – Jun 2013	Jan 2009 – Dec 2016	1.1%	1.0%
	Oxacillin (Penicillins)	May 2010 – Dec 2019	Jan 2009 – Dec 2016	33.4%	35.9%
	Penicillin (Penicillins)	May 2010 – Dec 2019	Jan 2009 – Apr 2015	82.4%	84.3%
	Tetracycline (Tetracyclines)	May 2010 – Dec 2019	Jan 2009 – Dec 2016	6.6%	6.0%

744 Percent resistance was calculated as the percentage of non-susceptible isolates out of the total
745 number of isolates from each hospital with a reported minimum inhibitory concentration for that
746 antibiotic. BWH, Brigham and Women's Hospital; MGH, Massachusetts General Hospital.

747 **S6 Table. Comparison of the Akaike information criterion (AIC) values between 6- and 12-**
748 **month period models for antibiotic use.**

Antibiotic class	Antibiotic Use Regression Model	
	6-month period	12-month period
Macrolides	104.4 (+67.5)	36.9 (+0)
Nitrofurans	-201.0 (+12.1)	-213.1 (+0)
Penicillins	128.7 (+86.6)	42.0 (+0)
Quinolones	-89.5 (+0)	-79.6 (+9.9)
Tetracyclines	-155.1 (+6.1)	-161.2 (+0)

749
750 In parentheses is the difference in AIC from the model with the lower AIC.

751 S7 Table. Comparison of the AIC values between 6- and 12-month period models for
752 antibiotic resistance.

Species	Antibiotic	Antibiotic Resistance Regression Model	
		6-month period	**12-month period**
E. coli	Amoxicillin/Clavulanate	339247.4 (+0)	339249.8 (+2.4)
	Ampicillin	522468.9 (+0)	522474.6 (+5.8)
	Ciprofloxacin	498206.9 (+7.9)	498199 (+0)
	Nitrofurantoin	199978.2 (+21.3)	199956.9 (+0)
	Tetracycline	422473.8 (+0)	422474.0 (+0.2)
K. pneumoniae	Amoxicillin/Clavulanate	64714.5 (+3.8)	64710.6 (+0)
	Ciprofloxacin	89697.3 (+11.1)	89686.2 (+0)
	Nitrofurantoin	66945.1 (+7.5)	66937.6 (+0)
	Tetracycline	80426.2 (+0)	80427.8 (+1.6)
S. aureus	Ciprofloxacin	189787.4 (+21.6)	189765.8 (+0)
	Erythromycin	216426.9 (+5.1)	216421.7 (+0)
	Nitrofurantoin	52484.0 (+15.6)	52468.4 (+0)
	Oxacillin	183142.1 (+7.5)	183134.6 (+0)
	Penicillin	125212.3 (+0)	125212.5 (+0.2)
	Tetracycline	131731.2 (+1.5)	131729.7 (+0)

753
754 In parentheses is the difference in AIC from the model with the lower AIC.