Comparing Non-Communicable Disease Risk Factors in Asian Migrants and Native Koreans: An Observational Study

Short Title: Non-Communicable Disease Risk Factors

Heng Piao, M.D., Ph.D.[a,b]; Jae Moon Yun, M.D., M.P.H.[b]; Aesun Shin, M.D., Ph.D.[c,d]; Daehee Kang, M.D., Ph.D.[c,d]; Changpeng Liu, M.P.H.[a]; Xiance Tang, M.P.H.[a]; and Belong Cho, M.D., Ph.D.[b,e]

[a] Department of Medical Record, Office for DRGs (Diagnosis Related Groups), Henan Cancer Hospital, Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China.

[b] Department of Family Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Republic of Korea.

[c] Department of Preventive Medicine, Seoul National University College of Medicine, Seoul, Republic of Korea.

[d] Department of Innovative Medical Sciences, Seoul National University Graduate School, Seoul, Republic of Korea.

[e] Health Promotion Center, Seoul National University Hospital, Seoul, Republic of Korea.

Corresponding Author:

Belong Cho, M.D., Ph.D.

Department of Family Medicine, Seoul National University Hospital, Seoul National University College of Medicine

101 Daehangno, Jongno-gu, Seoul 03080, Republic of Korea.

Tel.: +82-2-2072-2195; +82-10-9865-2195

Fax: +82-2-766-3276

Email: belong@snu.ac.kr

ORCID: https://orcid.org/0000-0001-9558-689X

Word Count: 3,497 words

Abstract

Background: Regarding international migrants, the theories of healthy migration effect and sick migration effect both exist; thus, assessing the health of international migrants is crucial in the Republic of Korea, Asia, and even worldwide.

Objective: To compare non-communicable disease risk factors among Asian migrants in Korea and the Korean population.

Design: Cross-sectional (2015) and longitudinal (2009–2015) observational study using the National Health Information Database of the Korean National Health Insurance Service.

Setting: Korea

Participants: Asian migrants (n=987,214) in Korea and Korean nationals (n=1,693,281) aged ≥20 years were included.

Measurements: Regarding the age-adjusted prevalence, direct age standardization was conducted separately by sex using 10-year age bands. Multivariable logistic regression analyses were conducted and adjusted for age, sex, economic status, body mass index, smoking status, alcohol use, and physical activity.

Results: Among participants aged ≥20 years, the age-adjusted prevalence of current smokers was higher among Chinese migrant men than among Korean men (P<0.001), and among other Asian migrant women than among Korean women (P<0.001). The age-adjusted prevalence of obesity was higher in Chinese, Filipino, and other Asian migrant women than in Korean women (P<0.001, P=0.002, and P<0.001, respectively). Among participants aged 20–49 years, the age-adjusted prevalence of diabetes and hypertension was higher in Filipino migrant women than in Korean women (P=0.009 and P<0.001, respectively).

Limitations: The total numbers of diabetes and hypertension cases may have been underestimated.

Conclusion: The current status of smoking and obesity among Asian migrants of specific nationalities is worse than that among natives. Moreover, the health inequalities among Filipino migrant women in Korea, especially those aged 20–49 years, should be addressed.

Funding Source: This study was funded by Seoul National University Hospital, Seoul, Republic of Korea; Yuhan Pharmaceutical, Seoul, Republic of Korea; and Henan Cancer Hospital/Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China. The funders had no role in the design of the study; the collection, analysis, and interpretation of the data; or the decision to approve publication of the finished manuscript.

Introduction

Recent systematic reviews and meta-analyses of migrant health have supported the healthy migration hypothesis that international migrants have health advantages (1,2). This healthy migration effect suggests that international migrants, particularly new ones, are healthier than the overall population of the host country (1); this can be attributed to self-selection or immigration policies (1,3) but is inversely associated with the duration of residence in the host country (1,4). However, in Israel, a sick migration effect is shown among international migrants within the initial 20 years of residence since commencement, implying that international migrants have health disadvantages compared with the native population in the host country (5).

Compared with other ethnic groups, South Asians are more likely to have diabetes and cardiovascular disease (CVD), possibly due to influence of the socioeconomic situation, the residence region, and migration (6). The South Asian migrant population has a higher risk of diabetes than the European population, and onset is 5–10 years earlier, despite lower body mass indices (BMIs) (7). In the Republic of Korea, compared with that in Korean natives, the prevalence of obesity in Asian migrants showed increasing trends, whereas the prevalence of obesity is becoming higher in Asian migrant women than in Korean native women since 2014 (8).

Few previous studies have involved Asian migrants (those born in Asian countries and moving to another high-income East Asian country) using nationwide data (2,4,7). Therefore, we investigated differences in the prevalence of non-communicable disease (NCD) risk factors between Asian migrants of different nationalities and the Korean population to evaluate whether there is a healthy migration effect in Korea using high-quality health check-up data from the National Health Insurance Service (NHIS). Moreover, we investigated

differences in obesity, diabetes, and elevated blood pressure among Chinese, Filipino, and Vietnamese migrants in Korea compared with the general population of their home countries. Based on our previous study (8), we hypothesized that Asian migrants of different nationalities have greater health advantages regarding NCD risk factors than Koreans.

Methods

Study Population

We conducted an observational retrospective population-based analysis of data collected during a longitudinal study. The study population constituted Asian migrants and the general Korean population aged ≥20 years who underwent health check-ups between 2009 and 2015; data were obtained from the National Health Information Database (NHID) established by the Korean NHIS (8,9). The NHID included eligibility data, national health check-up data, healthcare utilization, long-term care insurance, and health care provider information (9). Information on income-based insurance contributions, demographic variables, and date of death was included (9). The NHID is suitable for population-based studies aimed at primary and secondary prevention of diseases (9,10). The health check-up data for Asian migrants could be extracted from customized NHID data; this data represented most Asian migrants who underwent annual voluntary health check-ups in Korea (9,11). The health check-up data of the Korean population were obtained from the 1 million-sample cohort of the NHID, which is a random sample representing 2.2% of the Korean-born population, representative of the nationwide Korean population (12). Considering that the NHID is more extensive than it seems, the NHIS conducted systematic stratified random sampling with 2.2% within each stratum for the construction of the 1 million-sample cohort, of which, the target variable for sampling was the individual's total annual medical expenses (12).

Asian migrants originated from China, Japan, Vietnam, the Philippines, Indonesia, Thailand, Uzbekistan, Sri Lanka, Mongolia, Bangladesh, Pakistan, and India. Asian migrants were divided into Chinese, Japanese, Filipino, Vietnamese, and other Asian migrants to ensure sufficient samples after age standardization in 2015 (Appendix Methods).

Measurements

Age, sex, and monthly insurance premiums are included in the data collected by the NHIS; insurance premiums are determined by income level in Korea (8,13). Prevalence was defined as the number of participants with cases in a given year per number of participants in the population who underwent a health check-up during this period. Incidence was defined as the number of participants with new cases during the specified period per number of participants in the population who underwent health check-ups more than once during this period (Appendix Methods).

The primary outcome was the prevalence of NCD risk factors, including current smoking, physical inactivity, obesity, diabetes, elevated blood pressure, hypertension, and hypercholesterolemia, in 2015 (14-16). The secondary outcomes were the incidence of type 2 diabetes (T2D) and hypertension during 2009–2015. To ascertain incident T2D and hypertension, we excluded participants with diabetes and hypertension at the first health check-up between 2009 and 2014. Incident T2D and hypertension were determined for each participant between the first and last health check-ups (11).

Statistics

We conducted several sets of analyses. The research design used and statistical analyses performed for this and previous studies are shown in Appendix Figure 1. First, to calculate the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors among

Chinese, Japanese, Filipino, Vietnamese, and other Asian migrants compared with Koreans aged ≥20 years, direct age standardization was conducted by sex using 10-year age bands, wherein the World Standard Population was used as the standard population (17). For Asian migrants compared with Koreans aged 20–29, 30–39, and 40–49 years, we examined age-adjusted prevalence ratios (aPRs) and 95% log-normal confidence intervals (CIs) (18). Moreover, we conducted direct age standardization in Asian migrants compared with Koreans aged 20–49 years, wherein the World Standard Population was used as the standard population (17).

Second, to examine the differences in the development of T2D and hypertension between Asian migrants according to nationality and the native Korean population, multivariable logistic regression analyses were conducted and adjusted for the following covariates: age (continuous, years), sex, economic status, BMI (continuous, kg/m^2), smoking status, alcohol use, and physical activity. Adjusted odds ratios (ORs) and 95% CIs of incident T2D and hypertension determinants were examined between the first and last health check-ups. Further, we conducted multivariable logistic regression analyses stratified by age (20–39 and ≥40 years) and sex (11).

Finally, we examined differences in the age-adjusted prevalence of obesity, elevated blood pressure, and diabetes among Chinese, Filipino, and Vietnamese migrants compared with the general population of their respective home countries (11). Age standardization was conducted using 10-year age bands, wherein the World Standard Population was used as the standard population (17). Health data among general populations in China, the Philippines, and Vietnam were obtained from the Global Health Observatory Data Repository of the World Health Organization (available from:

http://apps.who.int/gho/data/node.main.A867?lang=en/) (Appendix Methods).

All analyses were performed using SAS version 9.3 (SAS Institute Inc., Cary, NC, USA). P-values <0.05 were considered statistically significant.

IRB Approval

The Institutional Review Board (IRB) of Seoul National University Hospital approved the study on August 21, 2018 (IRB number: E-1808-093-966). According to the IRB criteria, the need of participant informed consent was waived.

Role of the Funding Source

This study was funded by Seoul National University Hospital, Seoul, Republic of Korea; Yuhan Pharmaceutical, Seoul, Republic of Korea; and Henan Cancer Hospital/Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China. The funders had no role in the design of the study; the collection, analysis, and interpretation of the data; or the decision to approve publication of the finished manuscript.

Results

Table 1 shows the characteristics of Asian migrants of different nationalities and Koreans in 2015. In 2015, the age-adjusted prevalence of current smoking was higher in Chinese migrant men than in Korean men (P<0.001; Table 2) and higher in other Asian migrant women than in Korean women (P<0.001; Table 3). The age-adjusted prevalence of obesity was higher in Chinese, Filipino, and other Asian migrant women than in Korean women (P<0.001, P=0.002, and P<0.001, respectively; Table 3). The age-adjusted prevalence of hypercholesterolemia was higher in Japanese and Vietnamese migrant men than in Korean

men (P=0.004 and P=0.016, respectively; Table 2) and higher in Japanese and Vietnamese migrant women than in Korean women (P<0.001 and P<0.001, respectively; Table 3).

In 2015, among participants aged 30–39 years, the prevalence of diabetes was higher in Chinese migrant men than in Korean men (prevalence ratio [PR], 1.20; 95% CI: 1.07–1.33). Among participants aged 40–49 years, the prevalence of obesity and hypertension was higher in Chinese migrant women than in Korean women (PR, 1.17; 95% CI: 1.13–1.22; and PR, 1.10; 95% CI: 1.04–1.17, respectively) (Appendix Tables 1 and 2).

In 2015, among participants aged 20–29 and 30–39 years, the prevalence of hypercholesterolemia was higher in Filipino migrant men than in Korean men (PR, 1.37; 95% CI: 1.18–1.61; and PR, 1.09; 95% CI: 1.00–1.18, respectively). Among participants aged 30–39 years, the prevalence of diabetes was higher in Filipino migrant women than in Korean women (PR, 2.08; 95% CI: 1.47–2.94) (Appendix Tables 3 and 4).

In 2015, among participants aged 20–49 years, the age-adjusted prevalence of hypercholesterolemia was higher in Filipino migrant men than in Korean men (P=0.010), that of diabetes was higher in Filipino migrant women than in Korean women (aPR, 1.49; 95% CI: 1.11–2.02; P=0.009), and that of hypertension was higher in Filipino migrant women than in Korean women (aPR, 1.81; 95% CI: 1.48–2.21; P<0.001) (Figure 1 and Appendix Table 5).

During 2009–2015, the number of participants who underwent health check-ups more than once and for whom data regarding diabetes were available was 505,342. The number of participants with incident T2D between the first and last health check-ups was 22,284. In multivariate analyses, compared with that among native Koreans, the OR for developing T2D after adjusting for covariates was 0.82 (95% CI: 0.78–0.86) among Asian migrants. However,

among Vietnamese migrant men aged 20–39 years, the OR for developing T2D was 1.32 (95% CI: 1.11–1.57) compared with that among Korean men (Appendix Figure 2).

During 2009–2015, the number of participants who underwent health check-ups more than once and for whom data regarding hypertension were available was 431,433. The number of participants with hypertension between the first and last health check-ups was 48,007. In multivariate analyses, compared with that among native Koreans, the OR for developing hypertension after adjusting for covariates was 0.77 (95% CI: 0.75–0.79) among Asian migrants. However, compared with that among Korean women, the ORs for developing hypertension were 1.49 (95% CI: 1.05–2.11) and 2.22 (95% CI: 1.17–4.19) among Filipino migrant women aged 20–39 and >40 years, respectively (Appendix Figure 3).

Compared with the general population in their home countries, most Asian migrants in Korea were healthier in terms of the prevalence of obesity, diabetes, and elevated blood pressure. However, the age-adjusted prevalence of obesity was higher in Filipino migrant men than in the general male population in the Philippines in 2015. Moreover, in 2014, the age-adjusted prevalence of diabetes was higher in Filipino migrant men than in Philippine men (Figure 2).

Discussion

The age-adjusted prevalence of smoking was higher in Chinese migrant men than in Korean men and higher in other Asian migrant women than in Korean women aged ≥20 years in 2015. The prevalence of smoking among migrants is changing according to the situation at home and in the host countries (19,20). Smoking is harmful to the health of both the migrant and native populations. Risk factors attributed to social inequality may contribute to more than half of the major NCDs, especially CVD and lung cancer (14,21,22).

In 2015, the age-adjusted prevalence of obesity was higher in Chinese, Filipino, and other Asian migrant women than in Korean women aged ≥20 years. In our previous study analyzing the characteristics of migrants, the number of women who underwent health check-ups was less than the number of men; however, the number of women increased with each year during 2009–2015 (8). Obesity is a major health problem, particularly in migrant women (8,11,23). Unhealthy lifestyles, socioeconomic and cultural factors, genetics, and gene-environment interactions may lead to obesity, diabetes, and hypertension (11,24,25). Compared with Koreans, we could not demonstrate concurrence between obesity and diabetes/hypertension among Asian migrants, consistent with previous findings showing that there was no concurrence between obesity and T2D among international migrants from the Western Pacific to Europe (23). Nonetheless, there was concurrence between obesity and hypertension in Chinese migrant women aged 40–49 years. Socioeconomic inequalities, cultural stress, and obesity may be the main factors contributing to hypertension among migrants (4,14,26). Among Filipino migrant women aged 20–49 years, we found concurrence between obesity and diabetes/hypertension compared with Korean women, probably because of unhealthy diets, lack of exercise, early life factors, poor adherence to medication, low socioeconomic status, genetics, and gene-environment interactions (11,24,25). The stress generated by adapting to a new culture may contribute to a greater prevalence of hypertension among Filipino migrant women aged 20–49 years; this appears to be more important than unhealthy diets or lack of exercise (26,27).

Here, Japanese and Vietnamese migrants had a higher prevalence of hypercholesterolemia than Koreans. Among participants aged 20–49 years, Filipino migrant men had a higher prevalence of hypercholesterolemia than Korean men, especially those aged 20–29 and 30–39 years. Previous studies have shown that because of Americanized diets, which are higher in fat than traditional Japanese diets, total cholesterol levels became higher in Japanese

Americans than in Japanese natives (28,29). Michael Marmot's doctoral thesis on coronary heart disease (CHD) in Japanese Americans illustrated that participants who adhered to Japanese culture and had closer social networks were less likely to develop CHD compared to those who were more accustomed to Western culture and lifestyles (30). Although diet and hypercholesterolemia may increase the prevalence of CHD, the protective effect of Japanese culture on CHD is important, irreplaceable, and even independent of unhealthy diets and smoking (28,30,31). Therefore, the protective effect of social and cultural factors on cardiovascular health should be addressed while advocating healthy lifestyles.

In 2015, Chinese migrant men had a greater prevalence of diabetes than Korean men, whereas Filipino migrant women had a greater prevalence of diabetes than Korean women, aged 30–39 years. Migrants aged 30–39 years compared with Koreans may be more susceptible to diabetes after several migrant-related lifestyle changes (7,11,25). However, in multivariate analyses, during 2009–2015, among participants aged 20–39 years, Vietnamese migrant men were more likely to have T2D than Korean men. This finding is consistent with previous studies showing that international migrants developed diabetes approximately 10–20 years earlier than the local population of their host countries (7,11,24). Vietnamese migrant men aged 20–39 years may be more susceptible to T2D because of greater vulnerability to stress from Korean society and culture, under the combined effect of unhealthy diets, genetics, and gene-environment interactions (14,32).

Considering the important relationship between the pressure of cultural changes and elevated blood pressure among migrants, the reason why most Asian migrants are less likely to develop hypertension than Koreans may be due to the similar cultures and relatively close proximity to their home country (11,26,27). Asian migrants may be more likely to adapt to Korean society and maintain large social networks, ensuring sufficient social support. These

factors may help Asian migrants in Korea reduce stress and deal with conflicts more easily than when adapting to a different culture. Nevertheless, we cannot exclude the possibility that Asian migrants may return to their home countries either because of poor health or financial difficulties in Korea (19,33). Therefore, while the overall incidence of diabetes and hypertension was lower among Asian migrants in this and previous studies, the true incidence may be higher. However, Filipino migrant women were more likely to develop hypertension than Korean women, possibly due to social and cultural challenges encountered by Filipino women during the process of acculturation to Korea (26,27).

Compared with the general population in their home countries, most Asian migrants in Korea were healthier regarding the prevalence of obesity, diabetes, and elevated blood pressure. Nevertheless, the prevalence of obesity and diabetes was higher among Filipino migrant men than among the general Philippine population men, potentially due to unhealthy lifestyles, low socioeconomic status, genetics, and gene–environment interactions (11,24,25). During 2009–2015, the age-adjusted prevalence of diabetes, hypertension, and hypercholesterolemia was lower among most Asian migrants—regardless of sex—than among Koreans (8). Based on previous studies, the pursuit for better economic, educational, or social and living environment needs, combined with the self-selection of migrants and the immigration policy of the Korean government, may explain the healthy migration advantage in some NCD risk factors among most Asian migrants in Korea (1,2,34). However, regarding Israel, the reason for the sick migration effect may be attributed to the Jewish population's international migration to Israel for ideological and not economic motivations compared with the healthy migration effect in European countries (5).

Our study has several strengths. This is a large-scale observational study comparing the prevalence of NCD risk factors among Asian migrants of different nationalities and the host

population in Korea. The design was cross-sectional in 2015 and longitudinal during 2009–2015. Few similar studies have investigated the healthy migration effect among international migrant populations of various ethnic groups within the host populations of high-income countries in Asia using big data from the same source. Second, longitudinal data were collected to examine the differences in cases of newly diagnosed T2D and hypertension between Asian migrants according to nationality and Koreans, after adjusting for lifestyle factors. Regarding longitudinal analysis, a recent review has addressed the importance of answering research questions on migrant health using longitudinal data (19). However, due to the lack of complete standardized worldwide migrant data, even a global-scale study that verified the healthy migration effects in high-income countries could only be conducted by pooling mortality data and considering heterogeneity (high-quality data from different sources or sectors) and did not include international migrants in Asia (2). Therefore, the innovative point here is that, while comparing Asian migrants with Korean natives using health check-up data from the Korean NHIS, we conducted age standardization using the World Standard Population and compared the health data of Asian migrants with the health data in their home countries, targeting the general populations of China, the Philippines, and Vietnam, published by the Global Health Observatory Data Repository of the World Health Organization (17).

Our study has several limitations. First, the total number of diabetes and hypertension cases was underestimated because we only selected individuals who had National Health Insurance and underwent health check-ups. Further, the study population could have been healthier than the entire population if the individuals were more attentive to their health and fully used medical resources. However, a previous study using the same data (during 2009–2015) showed that the prevalence of diabetes in Korean men and women aged ≥20 years showed an increasing trend, in line with the results of a recent study on Korean adults aged ≥30 years

based on the Korean National Health and Nutrition Examination Survey (8,35). Moreover, the prevalence of diabetes and hypertension is more likely to have been underestimated in Asian migrants than in Koreans (11). For example, in Singapore, although international migrant workers have National Health Insurance when employed, they may not utilize necessary health services or even know that they have been covered (36). Second, we did not adjust for family history, nutritional factors, stress, and depression in the statistical analysis of relationships among Asian migrants according to nationality and newly diagnosed T2D or hypertension compared with Koreans. Third, because of the limited use of data on the type of international migrants (migrant workers, international students, and marriage migrants), migrant status factors were not considered; however, we used economic status instead of migrant status to compensate (8). Fourth, we could not distinguish between essential and secondary hypertensive participants. However, secondary hypertension is only expected in 5–10% of participants with hypertension, which may not affect our conclusions (37,38). Fifth, there are insufficient samples among Japanese, Filipino, and Vietnamese migrants, regarding several NCD risk factors of a specific age group. Nevertheless, we overcame these limitations by using cross-sectional and longitudinal designs based on age, sex, and socioeconomic status of Asian migrants of different nationalities in Korea. Finally, we were unaware of their residence duration and could not calculate the exact follow-up time of incident T2D and hypertension. However, even using the Kaplan–Meier method with adequate follow-up time, it was impossible to accurately compare changes in diabetes and hypertension over time in multiple populations of different nationalities during 2009–2015. This results from differences in countries of birth, early and lifestyle factors, and cultural backgrounds between Asian migrants and Korean natives. However, we performed an annual trend analysis of lifestyle, socioeconomic, and health-related factors among Asian migrants and Koreans during 2009–2015 in our previous study (8). These trend analyses provide insight into

changes in the overall prevalence of obesity, diabetes, and hypertension among Asian migrants compared with Koreans over time.

In conclusion, considering the adverse effects of socioeconomic and cultural factors on public health, strategies targeting current smoking and obesity among Asian migrants of specific nationalities in Korea are needed to address primary and secondary NCD prevention, especially targeting CVD and lung cancer. Moreover, health inequalities exist among Filipino migrants, especially women aged 20–49 years.

Acknowledgments

We thank Carol Brayne and Caroline Lee (University of Cambridge), Jae-Heon Kang (Sungkyunkwan University), Jian Li (University of California), Jong-Koo Lee and Young-Ho Khang (Seoul National University), and Shu-Zheng Liu (Zhengzhou University) for their constructive suggestions and comments. We also thank Editage (www.editage.cn), Luyao Zhang (Zhengzhou University), and Mohamed S. Bangura (Dalian Medical University) for their writing support.

Disclosures

Belong Cho reports grants and non-financial support from Yuhan Pharmaceutical during the conduct of the study. Heng Piao reports grants and non-financial support from Yuhan Pharmaceutical during the conduct of the study. The remaining authors have no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; there are no other relationships or activities that could influence the submitted work. The authors had access to all the study data, take responsibility for the accuracy of the analysis, and had authority over manuscript preparation and the decision to submit the manuscript for publication. All authors approve the manuscript and agree to adhere to all terms outlined in *Annals of Internal Medicine* Information for Authors, including terms for copyright.

References

1. Vang ZM, Sigouin J, Flenon A, et al. Are immigrants healthier than native-born Canadians? A systematic review of the healthy immigrant effect in Canada. Ethn Health 2017;22(3):209–41.

2. Aldridge RW, Nellums LB, Bartlett S, et al. Global patterns of mortality in international migrants: a systematic review and meta-analysis. Lancet 2018;392(10164):2553–66.

3. Marmot MG, Adelstein AM, Bulusu L. Lessons from the study of immigrant mortality. Lancet 1984;1(8392):1455–7.

4. Rechel B, Mladovsky P, Ingleby D, et al. Migration and health in an increasingly diverse Europe. Lancet 2013;381(9873):1235–45.

5. Constant AF, García-Muñoz T, Neuman S, et al. A "healthy immigrant effect" or a "sick immigrant effect"? Selection and policies matter. Eur J Health Econ 2018;19(1):103–21.

6. Misra A, Tandon N, Ebrahim S, et al. Diabetes, cardiovascular disease, and chronic kidney disease in South Asia: current status and future directions. BMJ 2017;357:j1420.

7. Sattar N, Gill JM. Type 2 diabetes in migrant South Asians: mechanisms, mitigation, and management. Lancet Diabetes Endocrinol 2015;3(12):1004–16.

8. Piao H, Yun JM, Shin A, et al. Longitudinal study of diabetic differences between international migrants and natives among the Asian population. Biomol Ther (Seoul) 2020;28(1):110–8.

9.   Seong SC, Kim YY, Khang YH, et al. Data resource profile: the national health

        information database of the National Health Insurance Service in South Korea. Int

        J Epidemiol 2016;46(3):799–800.

10.   Song SO, Jung CH, Song YD, et al. Background and data configuration process of

        a nationwide population-based study using the Korean national health insurance

        system. Diabetes Metab J 2014;38(5):395–403.

11.   Heng P. A comparison of cardiovascular risk factors among Asian migrants and

        the native population in Korea. The Graduate School of Seoul National

        University; 2020.

12.   Lee J, Lee JS, Park SH, et al. Cohort profile: the national health insurance

        service–national sample cohort (NHIS-NSC), South Korea. Int J Epidemiol

        2017;46(2):e15.

13.   Lee H, Cho J, Shin DW, et al. Association of cardiovascular health screening with

        mortality, clinical outcomes, and health care cost: a nationwide cohort study. Prev

        Med 2015;70:19–25.

14.   Di Cesare M, Khang YH, Asaria P, et al. Inequalities in non-communicable

        diseases and effective responses. Lancet 2013;381(9866):585–97.

15.   O'Donnell CJ, Elosua R. Cardiovascular risk factors. Insights from Framingham

        heart study. Rev Esp Cardiol 2008;61(3):299–310.

16.   Grundy SM, Cleeman JI, Daniels SR, et al. Diagnosis and management of the

        metabolic syndrome: an American Heart Association/National Heart, Lung, and

        Blood Institute scientific statement. Circulation 2005;112(17):2735–52.

17. Ahmad OB, Boschi-Pinto C, Lopez AD, et al. Age standardization of rates: a new WHO standard. J Geneva. World Health Organization 2001;9.

18. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. BMC Med Res Methodol 2003;3(1):21.

19. Abubakar I, Aldridge RW, Devakumar D, et al. The UCL-Lancet Commission on Migration and Health: the health of a world on the move. Lancet 2018;392(10164):2606–54.

20. Weber MF, Banks E, Sitas F. Smoking in migrants in New South Wales, Australia: report on data from over 100 000 participants in the 45 and up study. Drug Alcohol Rev 2011;30(6):597–605.

21. Blakely T, Fawcett J, Hunt D, et al. What is the contribution of smoking and socioeconomic position to ethnic inequalities in mortality in New Zealand? Lancet 2006;368(9529):44–52.

22. Roth GA, Mensah GA, Johnson CO, et al. Global burden of cardiovascular diseases and risk factors, 1990–2019: update from the GBD 2019 study. J Am Coll Cardiol 2020;76(25):2982–3021.

23. Meeks KA, Freitas-Da-Silva D, Adeyemo A, et al. Disparities in type 2 diabetes prevalence among ethnic minority groups resident in Europe: a systematic review and meta-analysis. Intern Emerg Med 2016;11(3):327–40.

24. Agyemang C, van den Born BJ. Non-communicable diseases in migrants: an expert review. J Travel Med 2019;26(2):tay107.

25. Misra A, Ganda OP. Migration and its impact on adiposity and type 2 diabetes. Nutrition 2007;23(9):696–708.

26. Gibson J, Stillman S, McKenzie D, et al. Natural experiment evidence on the effect of migration on blood pressure and hypertension. Health Econ 2013;22(6):655–72.

27. Steffen PR, Smith TB, Larson M, et al. Acculturation to Western society as a risk factor for high blood pressure: a meta-analytic review. Psychosom Med 2006;68(3):386–97.

28. Marmot MG, Syme SL, Kagan A, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: prevalence of coronary and hypertensive heart disease and associated risk factors. Am J Epidemiol 1975;102(6):514–25.

29. Yano K, Rhoads GG, Kagan A, et al. Dietary intake and the risk of coronary heart disease in Japanese men living in Hawaii. Am J Clin Nutr 1978;31(7):1270–9.

30. Marmot MG, Syme SL. Acculturation and coronary heart disease in Japanese-Americans. Am J Epidemiol 1976;104(3):225–47.

31. Matsumoto YS. Social stress and coronary heart disease in Japan: a hypothesis. Milbank Mem Fund Q 1970;48(1):9–36.

32. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. Nat Rev Endocrinol 2018;14(2):88–98.

33. Pablos-Méndez A. Mortality among Hispanics. JAMA 1994;271(16):1237.

34. Kennedy S, Mcdonald J, Biddle N. The healthy immigrant effect and immigrant selection: evidence from four countries. Social and economic dimensions of an aging population research papers. McMaster University; 2006;164.

35. Lee JW, Kang HT, Lim HJ, et al. Trends in diabetes prevalence among Korean adults based on Korean National Health and Nutrition Examination Surveys III–VI. Diabetes Res Clin Pract 2018;138:57–65.

36. Ang JW, Chia C, Koh CJ, et al. Healthcare-seeking behaviour, barriers and mental health of non-domestic migrant workers in Singapore. BMJ Glob Health 2017;2(2):e000213.

37. Tziomalos K. Secondary hypertension: novel insights. Curr Hypertens Rev 2020;16(1):11.

38. Lee HY, Shin J, Kim GH, et al. Korean Society of Hypertension guidelines for the management of hypertension: part II-diagnosis and treatment of hypertension. Clin Hypertens 2018;25(1):1–24.

Tables

Table 1. Characteristics of Asian migrants and Koreans aged ≥20 years who underwent health check-ups, according to nationality, in 2015*

Variable	Native	Migrant				
	Korean	Chinese	Japanese	Filipino	Vietnamese	Other Asian
N	266,162	101,223	4,445	11,888	19,224	63,765
Age (years)	49.4 ± 14.0	46.4 ± 11.4	47.4 ± 9.2	32.5 ± 6.0	30.3 ± 9.2	30.3 ± 6.3
Sex (%)						
Men	53.1	52.8	28.7	80.3	70.9	92.7
Women	46.9	47.2	71.3	19.7	29.1	7.3
Economic status (%)						
First quartile (low)	15.5	23.4	24.5	34.9	27.4	28.0
Second quartile	25.0	45.1	19.2	53.2	53.1	57.5
Third quartile	21.3	27.9	25.3	10.6	18.0	12.9
Fourth quartile (high)	38.2	3.6	31.0	1.3	1.5	1.6
Body mass index (kg/m^2)	24.0 ± 3.4	23.7 ± 3.3	23.2 ± 3.6	24.1 ± 3.1	21.8 ± 2.4	23.3 ± 3.2
Systolic blood pressure (mm	122.1 ± 14.5	121.6 ± 14.8	117.6 ± 14.3	121.2 ± 12.8	115.7 ± 12.0	118.8 ± 12.0

Hg)

Fasting serum glucose (mg/dL)	99.8 ± 24.2	97.7 ± 21.9	95.4 ± 19.1	94.5 ± 18.6	90.3 ± 14.9	91.4 ± 16.8
Total serum cholesterol (mg/dL)	195.3 ± 38.9	190.4 ± 38.0	205.4 ± 38.2	192.0 ± 37.6	182.8 ± 37.5	185.0 ± 37.5
Smoking status (%)						
Ex-smoker	18.0	10.7	10.6	4.2	4.1	4.1
Current smoker	21.5	28.7	9.5	19.8	20.1	30.9
Any alcohol use (%)	50.1	46.2	30.5	46.4	39.9	37.9
Physical activity (%)	70.2	51.6	64.4	49.9	49.1	45.4

* Data are expressed as the means ± standard deviation, or %.

Table 2. Comparison of the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors between Asian migrant men of

different nationalities and Korean men aged ≥20 years who underwent health check-ups in 2015

Variable Men	Age-adjusted prevalence (95% CI)*							
	Current smoking	Any alcohol use	Physical inactivity	Low-income level	Obesity	Hypertension	Diabetes	Hypercholesterolemia
Korean	39.9 (39.5–40.3)	70.3 (69.8–70.9)	24.8 (24.5–25.1)	10.9 (10.7–11.1)	41.7 (41.3–42.1)	22.4 (22.2–22.7)	9.6 (9.5–9.8)	10.2 (10.0–10.4)
Asian migrants								
Chinese	52.8 (52.1–53.6)	68.6 (67.8–69.4)	45.8 (45.1–46.4)	19.6 (19.2–20.1)	35.6 (35.0–36.2)	20.4 (20.0–20.8)	7.4 (7.2–7.7)	7.6 (7.4–7.9)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Japanese‡	26.8 (22.1–31.5)	70.4 (63.3–77.5)	27.8 (23.4–32.3)	10.4 (6.7–14.0)	37.9 (32.5–43.3)	18.3 (15.7–20.9)	8.0 (6.3–9.7)	14.4 (11.6–17.1)

P-value for difference†	<0.001	0.990	0.161	0.778	0.187	0.006	<0.001	0.004
Filipino§	23.0 (20.0–26.0)	49.6 (40.1–59.1)	52.9 (49.4–56.5)	28.6 (20.9–36.2)	41.4 (31.9–51.0)	20.8 (10.3–31.3)	8.9 (3.2–14.5)	11.2 (5.9–16.5)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	0.958	0.772	0.804	0.695
Vietnamese§	33.2 (30.8–35.5)	51.0 (47.0–54.9)	51.5 (47.1–55.9)	21.7 (18.8–24.7)	13.5 (11.3–15.6)	9.2 (6.7–11.6)	3.4 (2.1–4.8)	12.9 (10.4–15.4)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.016
Other Asian	35.8 (31.6–39.2)	45.4 (41.4–49.4)	52.9 (49.4–56.5)	24.0 (21.7–26.3)	38.3 (34.6–42.1)	18.4 (14.9–22.0)	5.1 (3.8–6.5)	9.9 (8.4–11.4)
P-value for difference†	0.026	<0.001	<0.001	<0.001	0.092	0.046	<0.001	0.709

* Data are expressed as %; age standardization with 10-year age bands; the World Standard Population was used as the standard population. CI, confidence interval.

† P-value for the difference was calculated among Asian migrant men compared with Korean men, according to nationality.

‡ Age-adjusted prevalence of diabetes (95% CI) between Japanese migrants and Korean men aged ≥30 years: 8.0 (6.3–9.7) and 12.3 (12.1–12.5), respectively.

§ Age-adjusted prevalence of current smoking (95% CI) among Filipino migrants, Vietnamese migrants, and Korean men aged 20–59 years: 23.0 (20.0–26.0), 33.2 (30.8–35.5), and 44.2 (43.7–44.7), respectively.

Table 3. Comparison of the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors between Asian migrant women of

different nationalities and Korean women aged ≥20 years who underwent health check-ups in 2015

Variable Women	Age-adjusted prevalence (95% CI)*							
	Current smoking	Any alcohol use	Physical inactivity	Low-income level	Obesity	Hypertension	Diabetes	Hypercholesterolemia
Korean	3.8 (3.7–4.0)	37.7 (37.3–38.2)	32.0 (31.6–32.3)	19.2 (19.0–19.5)	23.2 (22.9–23.4)	16.0 (15.8–16.2)	5.9 (5.8–6.0)	9.5 (9.3–9.7)
Asian migrants								
Chinese	2.9 (2.7–3.1)	23.4 (22.8–24.0)	46.5 (45.8–47.3)	29.6 (29.0–30.2)	25.6 (25.1–26.1)	14.6 (14.2–14.9)	4.4 (4.2–4.6)	8.3 (8.1–8.6)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Japanese‡	4.7 (3.2–6.2)	28.9 (24.6–33.3)	34.5 (31.1–37.9)	22.8 (20.4–25.2)	18.3 (16.0–20.6)	13.2 (11.2–15.2)	5.6 (4.0–7.2)	13.8 (12.0–15.7)

P-value for difference†	0.207	<0.001	0.140	0.002	<0.001	0.012	0.040	<0.001
Filipino§	2.0 (1.3–2.7)	11.9 (8.8–15.0)	51.6 (44.9–58.4)	42.0 (36.9–47.0)	30.9 (25.2–36.6)	19.1 (13.2–24.9)	5.2 (2.2–8.2)	11.2 (6.9–15.6)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	0.002	0.266	0.672	0.398
Vietnamese§	2.2 (1.1–3.2)	5.0 (4.2–5.7)	56.4 (52.9–59.8)	38.5 (35.9–41.1)	12.7 (10.8–14.6)	10.7 (8.4–13.0)	5.6 (3.9–7.3)	12.9 (10.6–15.2)
P-value for difference†	0.004	<0.001	<0.001	<0.001	<0.001	<0.001	0.764	<0.001
Other Asian	6.4 (4.8–7.9)	12.8 (11.1–14.5)	53.2 (49.4–57.1)	36.3 (33.3–39.3)	35.5 (31.4–39.5)	16.1 (12.7–19.6)	6.0 (3.8–8.1)	11.4 (9.0–13.9)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	0.949	0.961	0.094

* Data are expressed as %; age standardization with 10-year age bands; the World Standard Population was used as the standard population. CI, confidence interval.

† P-value for difference was calculated among Asian migrant women compared with Korean women, according to nationality.

‡ Age-adjusted prevalence of diabetes (95% CI) between Japanese migrants and Korean women aged ≥30 years: 5.6 (4.0–7.2) and 7.6 (7.5–7.8), respectively.

§ Age-adjusted prevalence of current smoking (95% CI) among Filipino migrants, Vietnamese migrants, and Korean women aged 20–59 years: 2.0 (1.3–2.7), 2.2 (1.1–3.2), and 4.4 (4.2–4.5), respectively.

Figure Legends

Figure 1. Comparison of age-adjusted prevalence ratios of diabetes and hypertension between Asian migrants of different nationalities and Koreans aged 20–49 years who underwent health check-ups in 2015*. * Age-adjusted prevalence ratios and 95% confidence intervals (CIs) of diabetes and hypertension were examined, with age standardization using 10-year age bands. The World Standard Population was used as the standard population.

Figure 2. Age-adjusted prevalence of obesity, elevated blood pressure, and diabetes among Chinese, Filipino, and Vietnamese migrants, compared with the general population in their home countries*. * Data are expressed as %. Age standardization was conducted using 10-year age bands. The World Standard Population was used as the standard population. Obesity was defined as a body mass index $\geq$25 kg/m^2 for the Asian population. Elevated blood pressure was defined as blood pressure $\geq$140/90 mmHg. Diabetes was defined as fasting blood glucose $\geq$126 mg/dL, a previous diabetes diagnosis, or the prescription of antidiabetic drugs. Regarding obesity and elevated blood pressure, 2015 data were used; for diabetes, 2014 data were used.

Age (20-49 years)	Adjusted prevalence ratio (95% CI)
Diabetes	
Chinese migrant men	1.02 (0.96–1.08)
Chinese migrant women	0.95 (0.85–1.06)
Filipino migrant men	0.89 (0.75–1.07)
Filipino migrant women	1.49 (1.11–2.02)
Vietnamese migrant men	0.63 (0.49–0.80)
Vietnamese migrant women	0.89 (0.66–1.21)
Other Asian migrant men	0.79 (0.72–0.86)
Other Asian migrant women	0.92 (0.70–1.22)
Hypertension	
Chinese migrant men	0.92 (0.89–0.96)
Chinese migrant women	1.06 (1.00–1.13)
Filipino migrant men	0.83 (0.74–0.92)
Filipino migrant women	1.81 (1.48–2.21)
Vietnamese migrant men	0.42 (0.34–0.51)
Vietnamese migrant women	0.70 (0.56–0.89)
Other Asian migrant men	0.54 (0.51–0.58)
Other Asian migrant women	1.06 (0.90–1.26)

*Age–adjusted prevalence ratios and 95% confidence intervals (CIs) of diabetes and hypertension were examined; age standardization with 10-year age bands; the world standard population was used as the standard population.

*Data are expressed as %; age standardization with 10-year age bands; the world standard population was used as the standard population; obesity was defined as a BMI≥25 kg/m² for the Asian population; elevated blood pressure was defined as blood pressure ≥ 140/90 mm Hg; diabetes was defined as a fasting blood glucose ≥126 mg/dL, previous diabetes diagnosis, or prescription of antidiabetic drugs; regarding obesity and elevated blood pressure, we used the data in 2015; regarding diabetes, we used the data in 2014.

† Source: data from the global health observatory data repository of World Health Organization. Available from: http://apps.who.int/gho/data/node.main.A867?lang=en/.