Comparing Non-Communicable Disease Risk Factors in Asian Migrants and Native Koreans: An Observational Study

Brief title: Comparison of Non-Communicable Disease Risk Factors

Heng Piao, MD, PhD[a,b]*; Jae Moon Yun, MSc MD[b]; Aesun Shin, MD, PhD[c,d]; Daehee Kang, MD, PhD[a,c,d]; Belong Cho, MD, PhD[b,e]*

[a] Department of Cancer Epidemiology and Prevention, Henan Cancer Hospital, Affiliate Cancer Hospital of Zhengzhou University, Zhengzhou 450003, China
[b] Department of Family Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul 03080, Republic of Korea
[c] Department of Preventive Medicine, Seoul National University College of Medicine, Seoul 03080, Republic of Korea
[d] Department of Innovative Medical Sciences, Seoul National University Graduate School, Seoul 03080, Republic of Korea
[e] Health Promotion Center, Seoul National University Hospital, Seoul 03080, Republic of Korea

*Heng Piao and Belong Cho are co-corresponding authors.

Address for correspondence
Belong Cho, MD, PhD
Department of Family Medicine, Seoul National University Hospital, Seoul National University College of Medicine
101 Daehangno, Jongno-gu, Seoul 03080, Republic of Korea
Tel: 82-2-2072-2195, 82-10-9865-2195
Fax: 82-2-766-3276
E-mail: belong@snu.ac.kr

Heng Piao, MD, PhD
Department of Cancer Epidemiology and Prevention, Henan Cancer Hospital, Affiliate Cancer Hospital of Zhengzhou University
127 Dongming Road, Jinshui District, Zhengzhou 450008, China
Phone: 86-133-1433-6468, 82-10-8712-6579
Fax: 82-2-766-3276
E-mail: jlpmh505@snu.ac.kr

Total word count: 4679 words

SHORT TWEET
A comparison of non-communicable disease risk factors between Asian migrants and native Koreans

FUNDING
This study was funded by the Seoul National University Hospital, Seoul, Republic of Korea; Yuhan Pharmaceutical, Seoul, Republic of Korea; and Henan Cancer Hospital/Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

DISCLOSURES

ACKNOWLEDGMENTS
We thank Carol Brayne and Caroline Lee (University of Cambridge), Jae-Heon Kang (Sungkyunkwan University), Jian Li (University of California), Jong-Koo Lee and Young-Ho Khang (Seoul National University), and Shu-Zheng Liu (Zhengzhou University) for their constructive suggestions and comments. We also thank Editage (www.editage.cn), Luyao Zhang (Zhengzhou University), and Mohamed S. Bangura (Dalian Medical University) for English language editing.

ABSTRACT
Background: Regarding international migrants, the theories of healthy migration effect and sick migration effect both exist; thus, assessing the health of international migrants is crucial in Korea, Asia, and even worldwide.
Objectives: To compare non-communicable disease risk factors among Asian migrants in Korea and the Korean population.
Methods: We conducted a cross-sectional (2015) and longitudinal (2009–2015) observational study using the National Health Information Database of the Korean National Health Insurance Service. Asian migrants (n=987,214) in Korea and Korean nationals (n=1,693,281) aged ≥20 years were included. Regarding the age-adjusted prevalence, direct age standardization was conducted separately by sex using 10-year age bands. Multivariable logistic regression analyses were conducted and adjusted for age, sex, economic status, BMI, smoking status, alcohol use, and physical activity.
Results: Among participants aged ≥20 years, the age-adjusted prevalence of current smokers was higher among Chinese migrant men than among Korean men (P<0.001) and among other Asian migrant women than among Korean women (P<0.001). The age-adjusted prevalence of obesity was higher in Chinese, Filipino, and other Asian migrant women than in Korean women (P<0.001; P=0.002; P<0.001, respectively). Among participants aged 20–49 years, the age-adjusted prevalence of diabetes and hypertension was higher in Filipino migrant women than in Korean women (P=0.009 and P<0.001, respectively).
Conclusions: The current status of smoking and obesity among Asian migrants of specific nationalities is worse than that among natives. Moreover, the health inequalities among Filipino migrant women in Korea, especially those aged 20–49 years, should be addressed.

CONDENSED ABSTRACT
Whether a health advantage exists in terms of non-communicable disease risk factors in Korea is unclear. We aimed to investigate the healthy migration effect using health check-up data from the National Health Insurance Service. Among participants aged ≥20 years, the problem of current smoking among Chinese migrant men and other Asian migrant women and that of obesity among Chinese, Filipino, and other Asian migrant women in Korea needs to be addressed. The prevalence of diabetes and hypertension in Filipino migrant women was worse than that in Korean women among the age group of 20–49 years.

Keywords: healthy migration effect, Filipino migrant women, obesity, diabetes, hypertension, smoking

ABBREVIATIONS
aPR: adjusted prevalence ratio
BMI: body mass index
CI: confidence interval
CVD: cardiovascular disease
NCD: non-communicable disease
NHID: National Health Information Database
NHIS: National Health Insurance Service
OR: odds ratio
PR: prevalence ratio
T2D: type 2 diabetes

INTRODUCTION

Recent systematic reviews and meta-analyses of migrant health have supported the healthy migration hypothesis that international migrants have health advantages (1,2). This healthy migration effect suggests that international migrants, particularly new, are healthier than the overall population of the host country (1). The healthy migration effect can be attributed to self-selection or immigration policies (1,3); nevertheless, the healthy migration effect is inversely associated with the duration of residence in the host country (1,4). However, in Israel, a sick migration effect is shown among international migrants within the initial 20 years of residence since commencement, implying that international migrants have health disadvantages compared with the native population in the host country (5).

South Asians are more likely to have diabetes and cardiovascular disease (CVD) than other ethnic groups, influenced by the socioeconomic situation, residence region, and migration (6). Moreover, the South Asian migrant population has a higher risk of developing diabetes compared with the European population, and onset is still 5–10 years earlier despite lower BMIs (7). In the Republic of Korea, compared with Korean natives, the prevalence of obesity in Asian migrants showed increasing trends; compared to Korean native women, the prevalence of obesity is becoming higher in Asian migrant women since 2014 (8).

So far, few previous studies have involved Asian migrants, those born in Asian countries and moving to another high-income East Asian country, using nationwide data. Therefore, we aimed to investigate differences in the prevalence of non-communicable disease (NCD) risk factors, including obesity, diabetes, and hypertension, between Asian migrants of different nationalities and the Korean population to evaluate whether there is a healthy migration effect in Korea, using high-quality health check-up data from the National Health Insurance Service (NHIS). In addition, we investigated differences in obesity, diabetes, and elevated blood pressure among Chinese, Filipino, and Vietnamese migrants in

Korea compared with the general population of their home countries. Based on our previous study, we hypothesized that Asian migrants of different nationalities have greater health advantages regarding NCD risk factors than Koreans (8).

METHODS

Study population

We conducted a retrospective population-based analysis of data collected during a longitudinal study. The study population constituted Asian migrants and the general Korean population aged ≥20 years who underwent health check-ups between 2009 and 2015; data were obtained from the National Health Information Database (NHID) established by the Korean NHIS (8,9). The NHID included eligibility data, National Health check-up data, healthcare utilization, long-term care insurance, and health care provider information (9). Namely, concerning the eligibility database, the information on income-based insurance contributions, demographic variables, and date of death was included (9). Moreover, the NHID is suitable for population-based studies aimed at primary and secondary prevention of diseases, including diabetes and hypertension (9,10). The health check-up data for Asian migrants could be extracted from customized NHID data and represented most Asian migrants who underwent annual voluntary health check-ups in Korea (9,11). Moreover, the health check-up data of the Korean population were obtained from the 1 million-sample cohort of the NHID, which is a random sample representing 2.2% of the Korean-born population, representative of the nationwide Korean population (12). Notably, considering that the NHID is more extensive than it seems, the NHIS conducted systematic stratified random sampling with 2.2% within each stratum for the construction of the 1 million-sample cohort, of which the target variable for sampling was the individual's total annual medical expenses (12). More details of the study population are shown in the Supplementary Text.

Here, Asian migrants originated from China, Japan, Vietnam, the Philippines, Indonesia, Thailand, Uzbekistan, Sri Lanka, Mongolia, Bangladesh, Pakistan, and India. For the study, Asian migrants were divided into Chinese, Japanese, Filipino, Vietnamese, and other Asian migrants to ensure sufficient samples after age standardization in 2015. Additional details regarding the classification of Asian migrants are shown in the Supplementary Text. The Institutional Review Board (IRB) of Seoul National University Hospital approved the study on August 21, 2018 (IRB number: E-1808-093-966). According to the IRB criteria, the need for obtaining informed participant consent was waived.

Measurements

Age, sex, and monthly insurance premiums are included in the data collected by the NHIS, and insurance premiums are determined by income level in Korea (8,13). Prevalence was defined as the number of participants with cases in a given year per number of participants in the population who underwent a health check-up during this period. Incidence was defined as the number of participants with new cases during the specified period per number of participants in the population who underwent health check-ups more than once during this period. Additional details regarding the definition of variables are shown in the Supplementary Text.

The primary study outcome was the prevalence of NCD risk factors, including current smoking, physical inactivity, obesity, diabetes, elevated blood pressure, hypertension, and hypercholesterolemia, in 2015 (14-16). The secondary study outcomes were the incidence of T2D and hypertension during 2009–2015. To ascertain incident T2D and hypertension, we excluded participants with diabetes and hypertension at the first health check-up between 2009 and 2014. Incident T2D and hypertension were determined for each participant between the first and last health check-ups (11).

Statistical analysis

We conducted several sets of analyses. The research design used and statistical analyses performed for this and previous studies are shown in Supplementary Figure 1. First, to calculate the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors among Chinese, Japanese, Filipino, Vietnamese, and other Asian migrants compared with Koreans aged ≥20 years, direct age standardization was conducted by sex using 10-year age bands, in which the world standard population was used as the standard population (17). For Asian migrants compared with Koreans aged 20–29, 30–39, and 40–49 years, we also examined age-adjusted prevalence ratios (aPRs) and 95% log-normal confidence intervals (CIs) (18). Moreover, we conducted direct age standardization in Asian migrants compared with Koreans aged 20–49 years, in which the world standard population was used as the standard population (17).

Second, to examine the differences in the development of T2D and hypertension between Asian migrants according to nationality and the native Korean population, multivariable logistic regression analyses were conducted and adjusted for the following covariates: age (continuous, years), sex, economic status, BMI (continuous, kg/m^2), smoking status, alcohol use, and physical activity. For these analyses, adjusted odds ratios (ORs) and 95% CIs of incident T2D and hypertension determinants were examined between the first and last health check-ups. Further, we conducted multivariable logistic regression analyses stratified by age (20–39 and ≥40 years) and sex (11).

Finally, we examined differences in the age-adjusted prevalence of obesity, elevated blood pressure, and diabetes among Chinese, Filipino, and Vietnamese migrants, compared with the general population of their respective home countries (11). Age standardization was conducted using 10-year age bands, and the world standard population was used as the

standard population (17). Health data among general populations in China, the Philippines, and Vietnam were obtained from the Global Health Observatory data repository of the World Health Organization. Data are available from http://apps.who.int/gho/data/node.main.A867?lang=en/, which is of the same year for comparison with Asian migrants, according to nationalities. More details regarding direct age standardization are shown in the Supplementary Text.

All analyses were performed using SAS version 9.3 (SAS Institute Inc., Cary, NC, USA), and *P*-values <0.05 were considered statistically significant.

RESULTS

Table 1 shows the characteristics of Asian migrants of different nationalities and Koreans in 2015. In 2015, the age-adjusted prevalence of current smoking was higher in Chinese migrant men than in Korean men (P<0.001; Table 2). Moreover, the age-adjusted prevalence of current smoking was higher in other Asian migrant women than in Korean women (P<0.001; Table 3). The age-adjusted prevalence of obesity was higher in Chinese migrant women, Filipino migrant women, and other Asian migrant women than in Korean women (P<0.001, P=0.002, P<0.001, respectively; Table 3). Additionally, the age-adjusted prevalence of hypercholesterolemia was higher in Japanese and Vietnamese migrant men than in Korean men (P=0.004, P=0.016, respectively; Table 2). Furthermore, the age-adjusted prevalence of hypercholesterolemia was higher in Japanese migrant women and Vietnamese migrant women than in Korean women (P<0.001, P<0.001, respectively; Table 3).

In 2015, among participants aged 30–39 years, the prevalence of diabetes was higher in Chinese migrant men than in Korean men (prevalence ratio [PR], 1.20; 95% CI, 1.07−1.33). Among participants aged 40–49 years, the prevalence of obesity and hypertension was higher in Chinese migrant women than in Korean women (PR, 1.17; 95%

CI, 1.13−1.22) and (PR, 1.10; 95% CI, 1.04−1.17), respectively (Supplementary Tables A1 and A2).

In 2015, among participants aged 20–29 years and 30–39 years, the prevalence of hypercholesterolemia was higher in Filipino migrant men than in Korean men (PR, 1.37; 95% CI, 1.18−1.61) and (PR, 1.09; 95% CI, 1.00−1.18), respectively. Moreover, among participants aged 30–39 years, the prevalence of diabetes was higher in Filipino migrant women than in Korean women (PR, 2.08; 95% CI, 1.47−2.94) (Supplementary Tables A3 and A4).

In 2015, among participants aged 20–49 years, the age-adjusted prevalence of hypercholesterolemia was higher in Filipino migrant men than in Korean men (P=0.010). The age-adjusted prevalence of diabetes was higher in Filipino migrant women than in Korean women (aPR, 1.49; 95% CI, 1.11−2.02; P=0.009). The age-adjusted prevalence of hypertension was higher in Filipino migrant women than in Korean women (aPR, 1.81; 95% CI, 1.48−2.21; P<0.001) (Central illustration and Supplementary Table B).

During 2009–2015, the number of participants who underwent health check-ups more than once and had information on diabetes was 505,342. The number of participants with incident T2D between the first and last health check-ups was 22,284. In multivariate analyses, compared with the native Korean population, the OR for developing T2D after adjusting for covariates was 0.82 (95% CI, 0.78−0.86) among Asian migrants. However, among Vietnamese migrant men aged 20–39 years, the OR for developing T2D was 1.32 (95% CI, 1.11−1.57), compared with Korean men (Figure 1).

During 2009–2015, the number of participants who underwent health check-ups more than once and for whom data regarding hypertension was available was 431,433; the number of participants with hypertension between the first and last health check-ups was 48,007. In multivariate analyses, compared with the native Korean population, the OR for developing

hypertension after adjusting for covariates was 0.77 (95% CI, 0.75−0.79) among Asian migrants. However, compared with Korean women, the ORs of developing hypertension were 1.49 (95% CI, 1.05−2.11) and 2.22 (95% CI, 1.17−4.19) among Filipino migrant women aged 20–39 years and those aged >40 years, respectively (Figure 2).

Compared to the general population in their home countries, we found that most Asian migrants in Korea were healthier in terms of the prevalence of obesity, diabetes, and elevated blood pressure. However, the age-adjusted prevalence of obesity was higher in Filipino migrant men than in men from the general population in the Philippines in 2015. Moreover, in 2014, the age-adjusted prevalence of diabetes was higher among Filipino migrant men than among men in the Philippines (Figure 3B).

DISCUSSION

The age-adjusted prevalence rates of current smoking were higher in Chinese migrant men than in Korean men; the age-adjusted prevalence of current smoking was higher in other Asian migrant women than in Korean women among participants aged ≥20 years in 2015. Moreover, the prevalence of smoking among migrants is changing according to the situation of the home and host countries (19,20). Furthermore, smoking is harmful to the health of both the migrant and native populations, and risk factors attributed to social inequality can contribute to more than half of the major NCDs, especially CVD and lung cancer (14,21,22).

In 2015, among participants aged ≥20 years, the age-adjusted prevalence of obesity was higher in Chinese, Filipino, and other Asian migrant women than in Korean women. Additionally, in our previous study, analyzing the characteristics of migrants, the number of women who underwent health check-ups was less than men, but the number of women increased with each year during 2009–2015 (8). Obesity is a major health problem of migrants, particularly migrant women (8,11,23). In migrants, unhealthy lifestyles,

socioeconomic and cultural factors, genetics, and interaction of genetics with the environment may lead to obesity, diabetes, and hypertension (11,24,25). Compared with Koreans, we could not find a clear overall pattern of concurrence between obesity and diabetes/hypertension among Asian migrants. This is consistent with previous research findings, showing that there was no concurrence pattern between obesity and T2D among international migrants from the Western Pacific to Europe (including Chinese and Vietnamese migrants) (23). Nonetheless, there was a clear pattern of concurrence between obesity and hypertension in Chinese migrant women aged 40–49 years. Therefore, socioeconomic inequalities, cultural stress, and obesity may be the main factors contributing to hypertension among migrants (4,14,26). Among Filipino migrant women aged 20–49 years, we found a distinct pattern of concurrence between obesity and diabetes/hypertension compared with Korean women, probably because of unhealthy diets, lack of exercise, early life factors, poor adherence to medication, low socioeconomic status, and both genetic and genetic–environmental interactions (11,24,25). Moreover, the stress generated by adapting to a new cultural process may contribute to a greater prevalence of hypertension among Filipino migrant women aged 20–49 years; this appears to be more important than unhealthy diets or lack of exercise (26,27).

Here, Japanese and Vietnamese migrants have a higher prevalence of hypercholesterolemia than Koreans. Among participants aged 20–49 years, Filipino migrant men have a higher prevalence of hypercholesterolemia than Korean men, especially targeting the age groups of 20–29 and 30–39 years. Previous studies have shown that, because of Americanized diets, which are higher in fat than traditional Japanese diets, total cholesterol levels became higher in Japanese Americans than in Japanese natives (28,29). Furthermore, Michael Marmot's doctoral thesis on coronary heart disease (CHD) in Japanese Americans illustrated that participants who were of adherence to deep Japanese culture and have closer

social networks were less likely to develop CHD, compared with participants who were more acculturated into western culture and lifestyles (30). Although diet and hypercholesterolemia may increase the prevalence of CHD, the protective effect of Japanese culture on CHD is important, irreplaceable, and even independent of unhealthy diets and smoking (28,30,31). Therefore, the protective effect of social and cultural factors on cardiovascular health should be addressed while advocating healthy lifestyles.

In 2015, among participants aged 30–39 years, Chinese migrant men had a greater prevalence of diabetes than Korean men, whereas Filipino migrant women had a greater prevalence of diabetes than Korean women. The migrant age group of 30–39 years compared with Koreans might be more susceptible to diabetes after several migrant-related lifestyle changes (7,11,25). However, in multivariate analyses, during 2009–2015, among participants aged 20–39 years, Vietnamese migrant men were more likely to have T2D than Korean men. This finding is consistent with previous studies, showing that international migrants developed diabetes about 10–20 years earlier than the local population of their host countries (7,11,24). Moreover, Vietnamese migrant men aged 20–39 years might be more susceptible to T2D because of greater vulnerability to the stress from the Korean society and culture, under the combined effect of unhealthy diet, genetic, and genetic–environmental interactions (14,32).

Considering the important relationship between the pressure of cultural changes and elevated blood pressure among migrants, the reason that most Asian migrants were less likely to develop hypertension than Koreans might be the similar cultures and the relatively close geographic location to their home country (11,26,27). Asian migrants might be more likely to adapt to the Korean social environment and maintain large social networks, ensuring sufficient social support. These factors might help Asian migrants in Korea reduce stress and deal with conflicts more easily than when adapting to a different society. Nevertheless, we

cannot exclude the possibility that Asian migrants may return to their home countries either because of poor health or financial difficulties in Korea (19,33). Therefore, while the detectable overall incidence of diabetes and hypertension was lower in this and previous studies among Asian migrants, the true incidence might be higher. However, here, Filipino migrant women had a higher probability of developing hypertension than Korean women. This might be caused by the social and cultural challenges encountered by Filipino women during the process of acculturation to Korea (26,27).

Compared with the general population in their home countries, we found that most Asian migrants in Korea were healthier regarding the prevalence of obesity, diabetes, and elevated blood pressure. Nevertheless, the prevalence of obesity and diabetes among Filipino migrant men was greater than among men from the general Philippine population, potentially due to unhealthy lifestyles, low socioeconomic status, genetics, and interaction between genetics and the environment (11,24,25). During 2009–2015, the age-adjusted prevalence of diabetes, hypertension, and hypercholesterolemia was lower among most Asian migrants— regardless of sex—than among Koreans (8). Based on previous studies, the pursuit for better economic, educational, or social and living environmental needs, combined with the self-selection of migrants and the immigration policy of the Korean government, may explain the healthy migration advantage in some NCD risk factors, especially targeting diabetes and hypertension, among most Asian migrants in Korea (1,2,34). However, regarding Israel, the reason for the sick migration effect could be attributed to the Jewish population's international migration to Israel for ideological and not economic motivations, compared with the healthy migration effect in the 16 European countries (5).

Our study had several strengths. This was a large-scale observational study comparing the prevalence of NCD risk factors among Asian migrants of different nationalities and the host population in Korea, of which the design was cross-sectional in

2015 and longitudinal during 2009–2015. Thus far, few similar previous studies investigated the healthy migration effect among international migrant populations of various ethnic groups within the host populations of high-income countries in Asia, using big data coming from the same source. Second, longitudinal data were collected to examine differences in cases of newly diagnosed T2D and hypertension between Asian migrants according to nationality and Koreans, adjusting for lifestyle factors. Regarding longitudinal analysis, a recent review study has addressed the importance of answering research questions on migrant health using longitudinal data (19). However, due to lack of complete and standardized worldwide migrant data, even a global-scale study that verifies the healthy migration effects in high-income countries could only be conducted by pooling mortality data and considering heterogeneity—of which the high-quality data were from different sources or sectors—and did not include international migrants in Asia (2). Therefore, the innovative point here is that, while comparing Asian migrants with Korean natives regarding health, using health check-up data of the Korean NHIS, we conducted age standardization by the world standard population and compared the health of Asian migrants with the health data in their home countries, targeting the general populations in China, Philippines, and Vietnam, published by the Global Health Observatory data repository of the World Health Organization (17).

Our study had several limitations. First, the total numbers of diabetes and hypertension cases were underestimated because we only selected individuals who had national health insurance and underwent health check-ups. Further, the study population could have been healthier than the population as a whole if the individuals were more attentive to their health and fully used medical resources. However, a previous study using the same data (during 2009–2015) showed that the prevalence of diabetes in Korean men and women aged ≥20 years showed an increasing trend, in line with the results of a recent study on Korean adults aged ≥30 years based on the Korean National Health and Nutrition

Examination Survey (8,35). Moreover, the prevalence of diabetes and hypertension is more likely to have been underestimated in Asian migrants than in Koreans (11). For example, in Singapore, although international migrant workers have national health insurance when employed, they may not utilize necessary health services or even know that they have been covered (36). Second, we did not adjust for family history, nutritional factors, stress, and depression in the statistical analysis of the relationship between Asian migrants according to nationality and newly diagnosed T2D or hypertension compared with Koreans. Third, because of the limited use of data on the type of international migrant—namely, migrant workers, international students, and marriage migrants—migrant status factors were not considered; however, we used economic status instead of migrant status to compensate (8). Fourth, we could not distinguish between essential and secondary hypertensive participants here. However, secondary hypertension is only expected in 5%–10% of participants with hypertension, which may not affect our research conclusion (37,38). Fifth, there are insufficient samples among Japanese, Filipino, and Vietnamese migrants, regarding several non-communicable risk factors of a specific age group. Nevertheless, we overcame these limitations by using cross-sectional and longitudinal study designs based on age, sex, and socioeconomic status of Asian migrants of different nationalities in Korea. Finally, we were unaware of their residence duration and could not calculate the exact follow-up time of incident T2D and hypertension. However, even using the Kaplan-Meier method with adequate follow-up time, it was impossible to accurately compare changes in diabetes and hypertension over time in multiple populations of different nationalities during 2009–2015. This results from differences in countries of birth, early and lifestyle factors, and cultural backgrounds between Asian migrants and Korean natives. However, we performed an annual trend analysis for lifestyle, socioeconomic, and health-related factors among Asian migrants and Koreans during 2009–2015 in our previous study (8). These trend analyses provide

insight into changes in the overall prevalence of obesity, diabetes, and hypertension among Asian migrants compared with Koreans over time.

CONCLUSIONS

In conclusion, considering the adverse effects of socioeconomic and cultural factors on public health, strategies targeting current smoking and obesity among Asian migrants of specific nationalities in Korea are needed to address primary and secondary NCD prevention, especially targeting CVD and lung cancer. Moreover, health inequalities exist among Filipino migrants, especially Filipino migrant women aged 20–49 years.

CLINICAL PERSPECTIVES

Systems-based practice: In clinical practice and public health policy development, the perspective of general and preventative medicine should be considered to investigate the healthy migration effect over time, fully considering socioeconomic and cultural factors. The identified relationships between Asian migrant nationality and NCD risk factors provide basic evidence for targeting high-risk groups and improving policy development in Korea.

Translational outlook: To distinguish between genetics, environment, and their interaction, more rigorous and innovative study designs should be used to conduct research on the health effects of culture, social environment, and ethnicity/race on the occurrence of NCDs, including dyslipidemia, CVD, and cancer.

REFERENCES

1. Vang ZM, Sigouin J, Flenon A, Gagnon A. Are immigrants healthier than native-born Canadians? A systematic review of the healthy immigrant effect in Canada. Ethn Health 2017;22:209-241.

2. Aldridge RW, Nellums LB, Bartlett S, et al. Global patterns of mortality in international migrants: a systematic review and meta-analysis. The Lancet 2018;392:2553-2566.

3. Marmot MG, Adelstein AM, Bulusu L. Lessons from the study of immigrant mortality. Lancet 1984;1:1455-1457.

4. Rechel B, Mladovsky P, Ingleby D, Mackenbach JP, McKee M. Migration and health in an increasingly diverse Europe. Lancet 2013;381:1235-1245.

5. Constant AF, García-Muñoz T, Neuman S, Neuman T. A "healthy immigrant effect" or a "sick immigrant effect"? Selection and policies matter. Eur J Health Econ 2018;19:103-121.

6. Misra A, Tandon N, Ebrahim S et al. Diabetes, cardiovascular disease, and chronic kidney disease in South Asia: current status and future directions. Bmj 2017;357:j1420.

7. Sattar N, Gill JM. Type 2 diabetes in migrant south Asians: mechanisms, mitigation, and management. Lancet Diabetes Endocrinol 2015;3:1004-1016.

8. Piao H, Yun JM, Shin A, Cho B. Longitudinal Study of Diabetic Differences between International Migrants and Natives among the Asian Population. Biomol Ther (Seoul) 2020;28:110-118.

9. Seong SC, Kim Y-Y, Khang Y-H et al. Data resource profile: the national health information database of the National Health Insurance Service in South Korea. Int J Epidemiol 2016;46:799-800.

10. Song SO, Jung CH, Song YD et al. Background and data configuration process of a nationwide population-based study using the korean national health insurance system. Diabetes Metab J 2014;38:395-403.

11. Heng P. A comparison of cardiovascular risk factors among Asian migrants and the native population in Korea. The Graduate School of Seoul National University, 2020. Available at: http://dcollection.snu.ac.kr/common/orgView/000000163370. Accessed 13 October, 2020

12. Lee J, Lee JS, Park S-H, Shin SA, Kim K. Cohort profile: the national health insurance service–national sample cohort (NHIS-NSC), South Korea. Int J Epidemiol 2017;46:e15-e15.

13. Lee H, Cho J, Shin DW et al. Association of cardiovascular health screening with mortality, clinical outcomes, and health care cost: a nationwide cohort study. Preventive medicine 2015;70:19-25.

14. Di Cesare M, Khang YH, Asaria P et al. Inequalities in non-communicable diseases and effective responses. Lancet 2013;381:585-597.

15. O'Donnell CJ, Elosua R. Cardiovascular risk factors. Insights from Framingham Heart Study. Rev Esp Cardiol 2008;61:299-310.

16. Grundy SM, Cleeman JI, Daniels SR et al. Diagnosis and management of the metabolic syndrome: an American Heart Association/National Heart, Lung, and Blood Institute scientific statement. Circulation 2005;112:2735-2752.

17. Ahmad OB, Boschi-Pinto C, Lopez AD, Murray CJ, Lozano R, Inoue M. Age standardization of rates: a new WHO standard. J Geneva: World Health Organization 2001;9.

18. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. BMC Med Res Methodol 2003;3:21.

19. Abubakar I, Aldridge RW, Devakumar D et al. The UCL-Lancet Commission on Migration and Health: the health of a world on the move. Lancet 2018;392:2606-2654.

20. Weber MF, Banks E, Sitas F. Smoking in migrants in New South Wales, Australia: report on data from over 100 000 participants in The 45 and Up Study. Drug Alcohol Rev 2011;30:597-605.

21. Blakely T, Fawcett J, Hunt D, Wilson N. What is the contribution of smoking and socioeconomic position to ethnic inequalities in mortality in New Zealand? Lancet 2006;368:44-52.

22. Roth GA, Mensah GA, Johnson CO et al. Global burden of cardiovascular diseases and risk factors, 1990-2019: Update from the GBD 2019 study. J Am Coll Cardiol 2020;76:2982-3021.

23. Meeks KA, Freitas-Da-Silva D, Adeyemo A et al. Disparities in type 2 diabetes prevalence among ethnic minority groups resident in Europe: a systematic review and meta-analysis. Intern Emerg Med 2016;11:327-340.

24. Agyemang C, Born B-Jvd. Non-communicable diseases in migrants: an expert review. J Travel Med 2018;26:tay107.

25. Misra A, Ganda OP. Migration and its impact on adiposity and type 2 diabetes. Nutrition 2007;23:696-708.

26. Gibson J, Stillman S, McKenzie D, Rohorua H. Natural experiment evidence on the effect of migration on blood pressure and hypertension. Health Econ 2013;22:655-672.

27. Steffen PR, Smith TB, Larson M, Butler L. Acculturation to Western society as a risk factor for high blood pressure: a meta-analytic review. Psychosom Med 2006;68:386-397.

28. Marmot MG, Syme SL, Kagan A, Kato H, Cohen JB, Belsky J. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: prevalence of coronary and hypertensive heart disease and associated risk factors. Am J Epidemiol 1975;102:514-525.

29. Yano K, Rhoads GG, Kagan A, Tillotson J. Dietary intake and the risk of coronary heart disease in Japanese men living in Hawaii. Am J Clin Nutr 1978;31:1270-1279.

30. Marmot MG, Syme SL. Acculturation and coronary heart disease in Japanese-Americans. Am J Epidemiol 1976;104:225-247.

31. Matsumoto YS. Social stress and coronary heart disease in Japan: a hypothesis. Milbank Mem Fund Q 1970;48:9-36.

32. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. Nat Rev Endocrinol 2018;14:88.

33. Pablos-Méndez A. Mortality among Hispanics. JAMA 1994;271:1237.

34. Kennedy S, Mcdonald J, Biddle N. The Healthy immigrant effect and immigrant selection: evidence from four Countries. Social and economic dimensions of an aging population research papers. McMaster University 2006:164.

35. Lee J-w, Kang H-T, Lim H-J, Park B. Trends in diabetes prevalence among Korean adults based on Korean National Health and Nutrition Examination Surveys III–VI. Diabetes Res Clin Pract 2018;138:57-65.

36. Ang JW, Chia C, Koh CJ, et al. Healthcare-seeking behaviour, barriers and mental health of non-domestic migrant workers in Singapore. BMJ global health 2017;2:e000213.

37. Tziomalos K. Secondary hypertension: novel insights. Curr Hypertens Rev 2020;16:11.

38. Lee HY, Shin J, Kim GH et al. 2018 Korean Society of Hypertension Guidelines for the management of hypertension: part II-diagnosis and treatment of hypertension. Clin Hypertens 2019;25:20.

FIGURE LEGENDS

Figure 1. Multivariable adjusted odds ratios of developing type 2 diabetes between the first and last health check-ups according to nationality during 2009–2015*

Figure 2. Multivariable adjusted odds ratios of developing hypertension between the first and last health check-ups, according to nationality during 2009–2015*

Figure 3. Age-adjusted prevalence of obesity, elevated blood pressure, and diabetes among Chinese, Filipino, and Vietnamese migrants, compared with the general population in their home countries*

Central illustration. Comparison of age-adjusted prevalence ratios of diabetes and hypertension between Asian migrants of different nationalities and Koreans aged 20–49 years who underwent health check-ups in 2015*

TABLES

Table 1. Characteristics of Asian migrant and Koreans aged 20 years and older who underwent health check-ups, according to nationalities in 2015*

Variable	Native	Migrant				
	Korean	Chinese	Japanese	Filipino	Vietnamese	Other Asian
N	266,162	101,223	4,445	11,888	19,224	63,765
Age (years)	49.4 ±14.0	46.4 ±11.4	47.4 ±9.2	32.5 ±6.0	30.3 ±9.2	30.3 ±6.3
Sex (%)						
Men	53.1	52.8	28.7	80.3	70.9	92.7
Women	46.9	47.2	71.3	19.7	29.1	7.3
Economic status (%)						
First quartile (low)	15.5	23.4	24.5	34.9	27.4	28.0
Second quartile	25.0	45.1	19.2	53.2	53.1	57.5
Third quartile	21.3	27.9	25.3	10.6	18.0	12.9
Fourth quartile (high)	38.2	3.6	31.0	1.3	1.5	1.6
Body mass index (kg/m^2)	24.0 ±3.4	23.7 ±3.3	23.2 ±3.6	24.1 ±3.1	21.8 ±2.4	23.3 ±3.2

Systolic blood pressure (mm Hg)	122.1 ±14.5	121.6 ±14.8	117.6 ±14.3	121.2 ±12.8	115.7 ±12.0	118.8 ±12.0
Fasting serum glucose (mg/dL)	99.8 ±24.2	97.7 ±21.9	95.4 ±19.1	94.5 ±18.6	90.3 ±14.9	91.4 ±16.8
Total serum cholesterol (mg/dL)	195.3 ±38.9	190.4 ±38.0	205.4 ±38.2	192.0 ±37.6	182.8 ±37.5	185.0 ±37.5
Smoking status (%)						
Ex-smoking	18.0	10.7	10.6	4.2	4.1	4.1
Current smoking	21.5	28.7	9.5	19.8	20.1	30.9
Any alcohol use (%)	50.1	46.2	30.5	46.4	39.9	37.9
Physical activity (%)	70.2	51.6	64.4	49.9	49.1	45.4

*Data are expressed as the means ±standard deviation, or %.

Table 2. Comparison of the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors between Asian migrant men of different nationalities and Korean men aged 20 years and older who underwent health check-ups in 2015

Variable	Age-adjusted prevalence (95% CI)*							
Men	Current smoking	Any alcohol use	Physical inactivity	Low-income level	Obesity	Hypertension	Diabetes	Hypercholesterolemia
Korean	39.9 (39.5–40.3)	70.3 (69.8–70.9)	24.8 (24.5–25.1)	10.9 (10.7–11.1)	41.7 (41.3–42.1)	22.4 (22.2–22.7)	9.6 (9.5–9.8)	10.2 (10.0–10.4)
Asian migrant								
Chinese	52.8 (52.1–53.6)	68.6 (67.8–69.4)	45.8 (45.1–46.4)	19.6 (19.2–20.1)	35.6 (35.0–36.2)	20.4 (20.0–20.8)	7.4 (7.2–7.7)	7.6 (7.4–7.9)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Japanese‡	26.8 (22.1–31.5)	70.4 (63.3–77.5)	27.8 (23.4–32.3)	10.4 (6.7–14.0)	37.9 (32.5–43.3)	18.3 (15.7–20.9)	8.0 (6.3–9.7)	14.4 (11.6–17.1)

P-value for difference†	<0.001	0.990	0.161	0.778	0.187	0.006	<0.001	0.004
Filipino §	23.0 (20.0–26.0)	49.6 (40.1–59.1)	52.9 (49.4–56.5)	28.6 (20.9–36.2)	41.4 (31.9–51.0)	20.8 (10.3–31.3)	8.9 (3.2–14.5)	11.2 (5.9–16.5)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	0.958	0.772	0.804	0.695
Vietnamese §	33.2 (30.8–35.5)	51.0 (47.0–54.9)	51.5 (47.1–55.9)	21.7 (18.8–24.7)	13.5 (11.3–15.6)	9.2 (6.7–11.6)	3.4 (2.1–4.8)	12.9 (10.4–15.4)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	0.016
Other Asian	35.8 (31.6–39.2)	45.4 (41.4–49.4)	52.9 (49.4–56.5)	24.0 (21.7–26.3)	38.3 (34.6–42.1)	18.4 (14.9–22.0)	5.1 (3.8–6.5)	9.9 (8.4–11.4)
P-value for difference†	0.026	<0.001	<0.001	<0.001	0.092	0.046	<0.001	0.709

*Data are expressed as %; age standardization with 10-year age bands; the world standard population was used as the standard population.

CI, confidence interval.

†*P*-value for the difference was calculated among Asian migrant men compared with Korean men, according to nationality.

‡Age-adjusted prevalences of diabetes (95% CI) between Japanese migrants and Korean men aged ≥30 years were 8.0 (6.3–9.7) and 12.3 (12.1–12.5), respectively.

§Age-adjusted prevalences of current smoking (95% CI) among Filipino migrants, Vietnamese migrants, and Korean men aged 20–59 years were 23.0 (20.0–26.0), 33.2 (30.8–35.5), and 44.2 (43.7–44.7), respectively

Table 3. Comparison of the age-adjusted prevalence of lifestyle, socioeconomic, and health-related factors between Asian migrant women of different nationalities and Korean women aged 20 years and older who underwent health check-ups in 2015

Variable	Age-adjusted prevalence (95% CI)*							
Women	Current smoking	Any alcohol use	Physical inactivity	Low-income level	Obesity	Hypertension	Diabetes	Hypercholesterolemia
Korean	3.8 (3.7–4.0)	37.7 (37.3–38.2)	32.0 (31.6–32.3)	19.2 (19.0–19.5)	23.2 (22.9–23.4)	16.0 (15.8–16.2)	5.9 (5.8–6.0)	9.5 (9.3–9.7)
Asian migrant								
Chinese	2.9 (2.7–3.1)	23.4 (22.8–24.0)	46.5 (45.8–47.3)	29.6 (29.0–30.2)	25.6 (25.1–26.1)	14.6 (14.2–14.9)	4.4 (4.2–4.6)	8.3 (8.1–8.6)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
Japanese‡	4.7 (3.2–6.2)	28.9 (24.6–33.3)	34.5 (31.1–37.9)	22.8 (20.4–25.2)	18.3 (16.0–20.6)	13.2 (11.2–15.2)	5.6 (4.0–7.2)	13.8 (12.0–15.7)

P-value for difference†	0.207	<0.001	0.140	0.002	<0.001	0.012	0.040	<0.001
Filipino §	2.0 (1.3–2.7)	11.9 (8.8–15.0)	51.6 (44.9–58.4)	42.0 (36.9–47.0)	30.9 (25.2–36.6)	19.1 (13.2–24.9)	5.2 (2.2–8.2)	11.2 (6.9–15.6)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	0.002	0.266	0.672	0.398
Vietnamese §	2.2 (1.1–3.2)	5.0 (4.2–5.7)	56.4 (52.9–59.8)	38.5 (35.9–41.1)	12.7 (10.8–14.6)	10.7 (8.4–13.0)	5.6 (3.9–7.3)	12.9 (10.6–15.2)
P-value for difference†	0.004	<0.001	<0.001	<0.001	<0.001	<0.001	0.764	<0.001
Other Asian	6.4 (4.8–7.9)	12.8 (11.1–14.5)	53.2 (49.4–57.1)	36.3 (33.3–39.3)	35.5 (31.4–39.5)	16.1 (12.7–19.6)	6.0 (3.8–8.1)	11.4 (9.0–13.9)
P-value for difference†	<0.001	<0.001	<0.001	<0.001	<0.001	0.949	0.961	0.094

*Data are expressed as %; age standardization with 10-year age bands; the world standard population was used as the standard population.

CI, confidence interval.

†*P*-value for difference was calculated among Asian migrant women compared with Korean women, according to nationality.

‡Age-adjusted prevalences of diabetes (95% CI) between Japanese migrant and Korean women aged ≥30 years were 5.6 (4.0–7.2) and 7.6 (7.5–7.8), respectively.

§Age-adjusted prevalences of current smoking (95% CI) among Filipino migrant, Vietnamese migrant, and Korean women aged 20–59 years were 2.0 (1.3–2.7), 2.2 (1.1–3.2), and 4.4 (4.2–4.5), respectively.

Type 2 diabetes	Adjusted odds Ratio (95% CI)
Age (20–39 years)	
Chinese migrant men	0.95 (0.84–1.07)
Chinese migrant women	0.78 (0.57–1.06)
Japanese migrant men	0.26 (0.07–1.06)
Japanese migrant women	0.91 (0.22–3.72)
Filipino migrant men	1.05 (0.89–1.24)
Filipino migrant women	1.01 (0.57–1.79)
Vietnamese migrant men	1.32 (1.11–1.57)
Vietnamese migrant women	1.26 (0.68–2.35)
Other Asian migrant men	0.97 (0.88–1.08)
Other Asian migrant women	0.65 (0.38–1.10)
Age (≥40 years)	
Chinese migrant men	0.75 (0.69–0.81)
Chinese migrant women	0.64 (0.57–0.72)
Japanese migrant men	0.64 (0.39–1.04)
Japanese migrant women	0.75 (0.47–1.20)
Filipino migrant men	0.82 (0.51–1.31)
Filipino migrant women	1.53 (0.56–4.21)
Vietnamese migrant men	1.03 (0.41–2.56)
Vietnamese migrant women	1.56 (0.73–3.35)
Other Asian migrant men	0.61 (0.45–0.81)
Other Asian migrant women	0.73 (0.37–1.42)
Overall	0.82 (0.78–0.86)

* Adjusted odds rartios and 95% confidence intervals (CIs) of incident type 2 diabetes determinants were determined using the multivariable logistic regression model; adjusted for age (continuous, years), economic status, BMI (cintinuous, kg/m^2), smoking status, any alcohol use, and physical activity.

Hypertension	Adjusted odds Ratio (95% CI)
Age (20-39 years)	
Chinese migrant men	0.70 (0.65–0.77)
Chinese migrant women	0.72 (0.59–0.88)
Japanese migrant men	0.58 (0.32–1.04)
Japanese migrant women	0.83 (0.34–2.05)
Filipino migrant men	0.76 (0.68–0.85)
Filipino migrant women	1.49 (1.05–2.11)
Vietnamese migrant men	0.55 (0.48–0.63)
Vietnamese migrant women	0.79 (0.45–1.39)
Other Asian migrant men	0.55 (0.51–0.59)
Other Asian migrant women	1.04 (0.76–1.42)
Age (≥40 years)	
Chinese migrant men	0.85 (0.80–0.90)
Chinese migrant women	0.78 (0.72–0.84)
Japanese migrant men	0.73 (0.53–1.01)
Japanese migrant women	1.16 (0.90–1.49)
Filipino migrant men	0.82 (0.58–1.15)
Filipino migrant women	2.22 (1.17–4.19)
Vietnamese migrant men	0.42 (0.17–1.04)
Vietnamese migrant women	0.78 (0.42–1.46)
Other Asian migrant men	0.70 (0.57–0.86)
Other Asian migrant women	0.97 (0.65–1.44)
Overall	0.77 (0.75–0.79)

* Adjusted odds rartios and 95% confidence intervals (CIs) of incident hypertension determinants were determined using the multivariable logistic regression model; adjusted for age (continuous, years), economic status, BMI (continuous, kg/m^2), smoking status, any alcohol use, and physical activity.

A

Men — Women

Prevalence (%)

Obesity | Raised blood pressure | Diabetes

■ Chinese migrants in Korea　□ The general population in China†

B

Men — Women

Prevalence (%)

Obesity | Raised blood pressure | Diabetes

■ Filipino migrants in Korea　□ The general population in Philippines†

C

Men — Women

Prevalence (%)

Obesity | Raised blood pressure | Diabetes

■ Vietnamese migrants in Korea　□ The general population in Vietnam†

* Data are expressed as %; age standardization with 10-year age bands; the world standard population was used as the standard population; obesity was defined as a BMI ≥25 kg/m² for the Asian population; elevated blood pressure was defined as blood pressure ≥ 140/90 mm Hg; diabetes was defined as a fasting blood glucose ≥126 mg/dL, previous diabetes diagnosis, or prescription of antidiabetic drugs; regarding obesity and elevated blood pressure, we used the data in 2015; regarding diabetes, we used the data in 2014.

† Source: data from the global health observatory data repository of World Health Organization. Available from: http://apps.who.int/gho/data/node.main.A867?lang=en/.

Age (20-49 years)	Adjusted prevalence ratio (95% CI)

Diabetes

Chinese migrant men	1.02 (0.96–1.08)
Chinese migrant women	0.95 (0.85–1.06)
Filipino migrant men	0.89 (0.75–1.07)
Filipino migrant women	1.49 (1.11–2.02)
Vietnamese migrant men	0.63 (0.49–0.80)
Vietnamese migrant women	0.89 (0.66–1.21)
Other Asian migrant men	0.79 (0.72–0.86)
Other Asian migrant women	0.92 (0.70–1.22)

Hypertension

Chinese migrant men	0.92 (0.89–0.96)
Chinese migrant women	1.06 (1.00–1.13)
Filipino migrant men	0.83 (0.74–0.92)
Filipino migrant women	1.81 (1.48–2.21)
Vietnamese migrant men	0.42 (0.34–0.51)
Vietnamese migrant women	0.70 (0.56–0.89)
Other Asian migrant men	0.54 (0.51–0.58)
Other Asian migrant women	1.06 (0.90–1.26)

* Age–adjusted prevalence ratios and 95% confidence intervals (CIs) of diabetes and hypertension were examined; age standardization with 10-year age bands; the world standard population was used as the standard population.

Hypothesis: Asian migrants of different nationalities have a greater health advantage in terms of non-communicable disease risk factors than Koreans.

Data: The National Health Information Database established by the Korean National Health Insurance Service
Subject: Adults aged 20 years and older (987,214 Asian migrants and 1,693,281 Koreans) who underwent health check-ups (2009–2015)
Asian migrants: Chinese, Japanese, Filipino, Vietnamese, and other Asian migrants

Objective 1: The prevalence of non-communicable disease risk factors in Asian migrants of different nationalities versus Koreans in 2015

Objective 2: The incidence of type 2 diabetes and hypertension in Asian migrants of different nationalities versus Koreans (2010–2015)

Objective 3: The age-adjusted prevalence of obesity, elevated blood pressure, and diabetes in Asian migrants of different nationalities versus the general population of their home countries

Direct age standardization with 10-year age bands; the world standard population (the standard population)

Age-specific prevalence among participants aged 20–29, 30–39, and 40–49 years

Multivariable logistic regression analyses stratified by age (20–39 and ≥40 years) and sex after adjusting for covariates

Each participant who developed type 2 diabetes and hypertension between the first and last health check-ups was considered a new incident case.

Direct age standardization with 10-year age bands; the world standard population (the standard population)