Modeling the Potential Role of Media Campaigns on the Control of Listeriosis

C.W. Chukwu[1,a], and F. Nyabadza[1,b]

[a,b]Mathematics and Applied Mathematics, University of Johannesburg,
Auckland Park Kingsway Campus, PO Box 524, 2006,
Johannesburg, South Africa

[a]cwochukwu@uj.ac.za, [b]fnyabadza@uj.ac.za

Keywords: Media campaigns. Equilibria. Listeria. Simulations. Food contamination constant.

Abstract. The human Listeriosis infection is a food-borne disease caused by the consumption of food products contaminated by the bacterial pathogen, Listeria. In this paper, we propose a mathematical model to analyse the impact of media awareness campaigns on the spread and control of Listeriosis. The model equilibria are determined and the model has three equilibria namely; the disease-free, the Listeria-free, and the endemic equilibria. The food contamination constant ($\mathcal{R}_f$) is determined and the stability analysis shows that the disease-free steady state is locally asymptotically stable if $\mathcal{R}_f < 1$, the Listeria-free and endemic steady states are locally asymptotically stable whenever $\mathcal{R}_f > 1$. Sensitivity analysis is done to determine the model parameters that most affect the severity of the disease. Numerical simulations are carried out to determine the role of media awareness campaigns on the spread of the Listeriosis. The results show that an increase in the intensity of the media awareness campaigns and removal of contaminated food products, a decrease in the contact rate, results in fewer humans getting infected leading to disease eradication, while an increase in the depletion of media awareness campaigns results in more humans been infected with Listeriosis. These findings may significantly impact policy and decision making in the control of Listeriosis disease.

Introduction

Listeriosis is a serious and severe food-borne disease that affects the human population globally. The disease is caused by a bacteria called *Listeria monocytogenes* which exists in the soil, water, and the environment as its primary host [1]. The bacteria can also been found in ready-to-eat (RTE) foods [2]. Human beings contract Listeriosis through the ingestion of contaminated ready-to-eat food products such as cantaloupes, meat, Ricotta Salata cheese, vegetables, polony, bean sprouts, ham, and or directly from the environment [3].

Before the 2017 outbreak in South Africa, an average of 60 to 80 confirmed Listeriosis disease cases were recorded annually (i.e. approximately 1 case per week). The recent outbreak in South Africa which occurred from 1 January 2017 to 17 July 2018 had 1 060 confirmed cases, with a total of 216 (26.8%) deaths. This was the world largest ever documented Listeriosis outbreak [4,5]. The source of the disease was traced to be from contaminated RTE processed meat products. Also, Listeriosis outbreaks resulting from human consumption of different kinds of contaminated RTE food products occur commonly in the USA, Canada and Europe [3].

Media campaigns and media-driven awareness programs such as print media, social media, internet, television, radio and advertisements play an important role in the dissemination of information about the spread of infectious disease outbreaks [6–8]. In particular, they influence the behavior of individuals

and the public health care systems in the implementation of control strategies to contain the spread of diseases. Dissemination of information educates people and helps them to take precautionary measures, practice better hygiene, factory workers wearing of clean cloves to avoid crosscontamination of food products during food productions to reduce the possibility of humans getting infected with Listeriosis.

Researchers have used mathematical models to describe the effect and impact of media campaign or several awareness programs on the dynamics of infectious diseases such as Ebola [9], Human Immunodeficiency Virus [10], human influenza [11], Cholera [12] e.t.c. Misra et al. [13], used a non-linear mathematical model which assumed that due to awareness programs by media, once the population becomes aware of the disease spread, they avoid contacts with the infectives and therefore form a new class of individuals called the aware class, who become susceptible again if their awareness wanes over time. The model analysis revealed that the number of infectives decrease with an increase in media campaigns. Kaur [14], extended the work by Misra by assuming that aware susceptible do not lose awareness but can also interact with infected individuals and get infected albeit at a lower rate. Their study suggests that with the increase in the rate of implementation of awareness programs via media, there is a subsequent decline in the number of the infected in any targeted population under consideration. Authors in [9] used a mathematical model to describe the transmission dynamics of Ebola in the presence of asymptomatic cases and the impact of media campaigns on the disease transmission was represented by a linearly decreasing function. Their results show that messages sent through media have a more significant effect on the reduction of Ebola cases if they are more effective and spaced out. An $SIRS$ model was proposed in [15] to investigate the impact of awareness programs by considering private and public awareness, which reduces the contact rate between unaware and aware population, and effect/impact of public information campaigns on disease prevalence. It was shown that both private and public awareness can reduce the size of epidemic outbreaks. A smoking cessation model with media campaigns was presented [16]. The results show that the reproduction number is suppressed when media campaigns that focus on smoking cessation is increased. Thus, spreading information to encourage smokers to quit smoking was an effective intervention. According to [17], an $SIRS$ model was used to analyze the role of information and limited optimal treatment on disease prevalence. The model considered the growth rate of information proportional to a saturated function of infected individuals. The result from the analysis shows that the combined effects of information and treatment are more effective and economical during an infection. Exponential functions have also been used to model the impact of media awareness campaigns on peoples behaviour which affects the evolution of infectious diseases [18–20]. In particular, the effects of Twitter messages on reducing the transmission rate of the influenza virus was studied in [20]. The result reveals that Twitter messages had a substantial influence on the dynamics of influenza disease spread.

To the best of our knowledge, there are very few mathematical models on the dynamics of Listeriosis, let alone those that investigate the potential role of media awareness camapaigns on the dynamics of Listeriosis. This work is motivated by the work done in [13]. We formulate a mathematical model to study the impact and effects of media campaigns on the dynamics of Listeriosis disease in the human population. We decribe the model in detail in the following section

Model Description

The human population is divided into four sub-classes, viz: Susceptibles $S_h(t)$, aware susceptibles $S_a(t)$, the infected $I_h(t)$, and the recovered $R_h(t)$. Individuals are recruited at a rate proportional to the size of

the human population $N(t)$, $\mu_h N(t)$, where

$$N(t) = S_h(t) + S_a(t) + I_h(t) + R_h(t),$$

and μ_h is the natural birth/mortality rate. Upon infection with Listeria from contaminated food, the susceptibles move into the infectious class $I_h(t)$ at a rate $\lambda_h(t)$, where $\lambda_h(t) = \beta_{f_1} F_c$ with β_{f_1} being the rate at which humans gets infected and $F_c(t)$ the contaminated food products. Here, $\lambda_h(t)$ describes the force of infection by the consumption of contaminated food products. The susceptible individuals can also move to the awareness class at a rate ρ as a result of the interaction with the media campaigns. We assume that media campaigns wane over time and the aware individuals can revert to being susceptible again at a rate ω_1. The infected individuals recover at a rate γ with immunity after treatment. These individuals who recover after some time can also lose their immunity and become susceptible again at a rate δ_h. We assume a constant human population $N(t)$ which consist of those individuals who do not work in the factory over the modelling time. Further, we assume that aware individuals can not be infected as their awareness protects them from contracting the disease. Given that, the bacteria survive even at $4^0 C$, it can die or grow in its host or the environment at significantly low temperatures. Let r_l denote the net growth rate of Listeria, $L(t)$. Our model assumes a logistic growth of Listeria with carrying capacity K_L. The non-contaminated food products $F_n(t)$ can be contaminated as a result of interacting with the bacteria from the environment that comes, via the workers, exchange of gloves or utensils during food manufacturing and also through the contact with contaminated food $F_c(t)$ at a rate $\lambda_f(t)$, where $\lambda_f(t) = \beta_L L(t) + \beta_{f_2} F_c(t)$. The parameters β_L and β_{f_2} are contact rate of Listeria and the contamination rate of non-contaminated food by contaminated food products respectively. The contaminated food products are then responsible for the transmission of Listeriosis disease to the human population through ingestion of the contaminated food products. The total amount of food products, $F(t)$, at any given time is given by

$$F(t) = F_n(t) + F_c(t),$$

where θ_f is the rate of food processing and μ_f the rate of removal of food products through consumption. Let M_a be the cumulative density of media campaigns with maximum intensity, M, at which media awareness campaigns are implemented, π_0 the rate of implementation of the media awareness campaigns and μ_0 the rate of depletion of media awareness campaigns.

Fig. 1: Model flow diagram. The solid arrows represents the transitions between the compartments, while the dotted lines represents the influences on the solid arrows and the compartments.

The above model descriptions and Figure 1 gives the following systems of non-linear ordinary differential equations:

$$
\begin{aligned}
\frac{dS_h}{dt} &= \mu_h N + \delta_h R_h + \omega_1 S_a - \lambda_h S_h - \mu_h S_h - \rho M_a S_h, \\[2mm]
\frac{dS_a}{dt} &= \rho M_a S_h - (\omega_1 + \mu_h) S_a, \\[2mm]
\frac{dI_h}{dt} &= \lambda_h S_h - (\mu_h + \gamma) I_h, \\[2mm]
\frac{dR_h}{dt} &= \gamma I_h - (\mu_h + \delta_h) R_h, \\[2mm]
\frac{dL}{dt} &= r_l L \left(1 - \frac{L}{K_L} \right), \\[2mm]
\frac{dM_a}{dt} &= \pi_0 I_h - \mu_0 M_a, \\[2mm]
\frac{dF_n}{dt} &= \theta_f - \lambda_f F_n - \mu_f F_n, \\[2mm]
\frac{dF_c}{dt} &= \lambda_f F_n - \mu_f F_c.
\end{aligned}
\right\} \qquad (1)
$$

All parameters for the model system (1) except for r_l, are assumed to be non-negative for all time $t > 0$. By setting

$$s_h = \frac{S_h}{N}, \quad s_a = \frac{S_a}{N}, \quad i_h = \frac{I_h}{N}, \quad l = \frac{L}{K_L}, \quad m_a = \frac{M_a}{M}, \quad f_n = \frac{F_n}{F}, \quad f_c = \frac{F_c}{F},$$

and given that $R_h(t) = N(t) - S_h(t) - S_a(t) - I_h(t)$ we have

$$\left.\begin{aligned}
\frac{ds_h}{dt} &= \mu_h + \delta_h(1 - s_h - s_a - i_h) + \omega_1 s_a - (\tilde{\lambda}_h + \mu_h + \tilde{\rho} m_a)s_h, \\[2mm]
\frac{ds_a}{dt} &= \tilde{\rho} m_a s_h - (\omega_1 + \mu_h)s_a, \\[2mm]
\frac{di_h}{dt} &= \tilde{\lambda}_h s_h - (\mu_h + \gamma)i_h, \\[2mm]
\frac{dl}{dt} &= r_l l(1 - l), \\[2mm]
\frac{dm_a}{dt} &= \pi i_h - \mu_0 m_a, \\[2mm]
\frac{df_n}{dt} &= \tilde{\theta}_f - (\tilde{\lambda}_f + \mu_f)f_n, \\[2mm]
\frac{df_c}{dt} &= \tilde{\lambda}_f f_n - \mu_f f_c,
\end{aligned}\right\} \tag{2}$$

where

$$\tilde{\lambda}_h = \beta_1 f_c, \quad \tilde{\lambda}_f = \beta_2 l + \beta_3 f_c, \quad \tilde{\theta}_f = \frac{\theta_f}{F}, \quad \alpha = \frac{N}{M} \quad \tilde{\rho} = \rho M, \quad \pi = \pi_0 \alpha,$$

with $\beta_1 = \beta_{f_1} F, \beta_2 = \beta_l K_L, \beta_3 = \beta_{f_2} F$, and initial conditions

$$\begin{aligned}
s_h(0) &= s_{h0} > 0, s_a(0) = s_{a0} > 0, i_h(0) = i_{h0} \geq 0, l(0) = l_0 \geq 0, \\
m_a(0) &= m_{a0} \geq 0, f_n(0) = f_{n0} \geq 0, f_c(0) = f_{c0} \geq 0.
\end{aligned} \tag{3}$$

Model Properties and Analysis

Existence and Uniqueness of Solutions

We now show that the solutions of systems (2) are bounded. We thus have the following result.

Theorem 1. *The solutions of model system (2) are contained in the region $\Omega \in \mathbb{R}^7_+$, which is given by*

$$\Omega = \left\{ (s_h, s_a, i_h, l, m_a, f_n, f_c) \in \mathbb{R}^7_+ : 0 \leq s_h + s_a + i_h \leq 1, 0 \leq l \leq 1, 0 \leq m_a \leq \frac{\pi}{\mu_0}, 0 \leq f_n + f_c \leq \frac{\tilde{\theta}_f}{\mu_f} \right\},$$

for the initial conditions (3) in Ω.

Proof. Considering the total change in the human population from the model system (2) given by

$$\frac{dn}{dt} = \mu_h(1 - s_h - s_a - i_h) + \delta_h(1 - s_h - s_a - i_h) - \gamma i_h, \tag{4}$$

for $n = s_h + s_a + i_h \leq 1$. Setting $\Theta_1 = (s_h + s_a + i_h)$ we obtain

$$\begin{aligned}\frac{d\Theta_1}{dt} &= \mu_h(1 - \Theta_1) + \delta_h(1 - \Theta_1) - \gamma i_h, \\ &\leq (\mu_h + \delta_h)(1 - \Theta_1),\end{aligned}$$

whose solution is

$$\Theta_1 \leq 1 - \Theta_1(0) \exp[-(\mu_h + \delta_h)t],$$

where $\Theta_1(0) = s_h(0) + s_a(0) + i_h(0)$ is the initial condition. We note that $0 \leq \Theta_1 \leq 1 - \Theta_1(0)e^{-(\mu_h + \delta_h)t}$, so that $n(t)$ is bounded provided that $\Theta_1(0) \geq 0$.

The equation

$$\frac{dl}{dt} = r_l l(1 - l)$$

for the Listeria compartment has a standard solution for a logistic equation

$$l(t) = \frac{1}{1 + \Theta_2 \exp[-r_l t]},$$

which is bounded with $\Theta_2 = \frac{\exp[r_l c]}{\mu_f}$, where c is a constant.

On the other hand, the solutions of $f_n(t)$ yields

$$f_n(t) = \frac{\tilde{\theta}_f}{\mu_f} - \Theta_3 \exp[-\mu_f t],$$

where $\Theta_3 = \frac{\exp[-\mu_f c]}{\mu_f}$, c constant of integration. We have that $f_n(t) \leq \frac{\tilde{\theta}_f}{\mu_f}$ as $t \to \infty$, thus, $f_n(t)$ is also bounded.

From the last equation of the system (2) we have that

$$\begin{aligned}\frac{df_c}{dt} &= (\beta_2 l + \beta_3 f_c)\frac{\tilde{\theta}_f}{\mu_f} - \mu_f f_c, \\ &\leq (\beta_2 + \beta_3 f_c)\frac{\tilde{\theta}_f}{\mu_f} - \mu_f f_c.\end{aligned} \tag{5}$$

If we set $\omega = (\beta_2 + \beta_3 f_c)\frac{\tilde{\theta}_f}{\mu_f}$, then equation (5) becomes

$$\frac{df_c}{dt} \leq \omega - \mu_f f_c,$$

whose solutions yields

$$f_c(t) \geq \frac{\omega}{\mu_f} - \Theta_4 \exp[-\mu_f t],$$

where $\Theta_4 = \Theta_3$. Here, $f_c(t) \leq \frac{\omega}{\mu_f}$, hence bounded. We thus conclude that all the solutions of system (2) are bounded, biologically feasible, and remains in Ω for all $t \in [0, \infty)$. This completes the proof.

Positivity of Solutions

We prove the positivity of the solutions of model system (2) with initial conditions (3). First, we state the following Lemma as given in [21].

Lemma 2. *Suppose* $\Omega \subset \mathbb{R} \times C^n$ *is open,* $f_i \in C(\Omega, \mathbb{R}), i = 1, 2, \ldots, n.$ *If*

$$f_i|_{x_i(t)=0, X_t \in C_{0+}^n} \geq 0, \quad X_t = (x_{1t}, \cdots, x_{nt})^T \quad i = 1, 2, \cdots, n,$$

then C_{+0}^n *is the invariant domain of the following equations*

$$\dot{x}_i(t) = f_i(t, X_t), \quad t \geq 0, \ i = 1, 2, \cdots, n. \tag{6}$$

If $f_i|_{x_i(t)=0, X_t \in C_{-0}^n} \leq 0, \quad X_t = (x_{1t}, \cdots, x_{nt})^T \quad i = 1, 2, \cdots, n,$ *then* C_{-0}^n *is the invariant domain of equations (6).*

We have the following Theorem on the invariance of system (2).

Theorem 3. *Each solution* $(s_h(t), s_a(t), I_h(t), l(t), m_a(t), f_n(t), f_c(t))$ *of the model system (2) with the non-negative initial conditions (3) is non-negative for all* $t > 0$.

Proof. Let $X = (s_h, s_a, I_h, l, m_a, f_n, f_c)^T$ and

$$g(X) = (g_1(X), g_2(X), g_3(X), g_4(X), g_5(X), g_6(X), g_7(X))^T,$$

then we can re-write the model system (2) as follows:

$$\dot{X} = g(X)$$

where

$$g(X) = \begin{pmatrix} g_1(X) \\ g_2(X) \\ g_3(X) \\ g_4(X) \\ g_5(X) \\ g_6(X) \\ g_7(X) \end{pmatrix} = \begin{pmatrix} \mu_h + \delta_h(1 - s_h - s_a - i_h) + \omega_1 s_a - (\tilde{\lambda}_h + \mu_h + \tilde{\rho}m_a)s_h \\ \tilde{\rho}m_a s_h - (\omega_1 + \mu_h)s_a \\ \tilde{\lambda}_h s_h - (\mu_h + \gamma)i_h \\ r_l l(1 - l) \\ \pi_0 \alpha i_h - \mu_0 m_a \\ \tilde{\theta}_f - (\tilde{\lambda}_f + \mu_f)f_n \\ \tilde{\lambda}_f f_n - \mu_f f_c \end{pmatrix}. \tag{7}$$

From (7), setting all the classes to zero, we have that

$$\begin{aligned} \frac{ds_h(t)}{dt}\Big|_{s_h=0} &= (\mu_h + \omega_1 s_a) > 0, \\ \frac{ds_a(t)}{dt}\Big|_{s_a=0} &= \rho M m_a s_h > 0, \quad \frac{di_h(t)}{dt}\Big|_{i_h=0} = \tilde{\lambda}_h s_h > 0, \quad \frac{dl(t)}{dt}\Big|_{l=0} = 0, \\ \frac{dm_a(t)}{dt}\Big|_{m_a=0} &= \pi_0 \alpha i_h > 0, \quad \frac{df_n(t)}{dt}\Big|_{f_n=0} = \tilde{\theta}_f > 0, \quad \frac{df_c(t)}{dt}\Big|_{f_c=0} = \tilde{\lambda}_f f_n > 0. \end{aligned}$$

Thus, it follows that from Lemma 2 that $\mathbb{R}_+^7$ is an invariant set and positive.

Steady States Analysis

The steady states of the model system (2) are obtained by equating the right side of equations (2) to zero, so that

$$\mu_h + \delta_h(1 - s_h^* - s_a^* - i_h^*) + \omega_1 s_a^* - (\beta_1 f_c^* + \mu_h + \tilde{\rho} m_a^*) s_h^* = 0,$$

$$\tilde{\rho} m_a^* s_h^* - (\omega_1 + \mu_h) s_a^* = 0,$$

$$\beta_1 f_c^* s_h^* - (\mu_h + \gamma) i_h^* = 0,$$

$$r_l l^* (1 - l^*) = 0, \qquad (8)$$

$$\pi i_h^* - \mu_0 m_a^* = 0,$$

$$\tilde{\theta}_f - (\beta_2 l^* + \beta_3 f_c^* + \mu_f) f_n^* = 0,$$

$$(\beta_2 l^* + \beta_3 f_c^*) f_n^* - \mu_f f_c^* = 0,$$

From the fourth equation of system (8), we have $l^* = 0$ or $l^* = 1$. We consider the two cases separately.

CASE A: If $l^* = 0$, (i.e. if there is no Listeria in the environment) then from the second last equation of (8) we have that

$$f_n^* = \frac{\tilde{\theta}_f}{\beta_3 f_c^* + \mu_f}. \qquad (9)$$

Substituting (9) into the last equation of (8) we obtain

$$\beta_3 \tilde{\theta}_f f_c^* - \mu_f f_c^* (\beta_3 f_c^* + \mu_f) = 0,$$

and upon simplification we have

$$f_c^* = 0 \quad \text{or} \quad f_c^* = \frac{\beta_3 \tilde{\theta}_f - \mu_f^2}{\beta_3 \mu_f}. \qquad (10)$$

Thus, if $l^* = 0$ then $f_c^* = 0, \tilde{\lambda}_h = \tilde{\lambda}_f = 0, f_n^* = \frac{\tilde{\theta}_f}{\mu_f}$ and $i_h^* = 0, m_a^* = 0, s_a^* = 0$. Also, from the first equation of (8)

$$s_h^* = 1$$

This results in the disease free steady states (DFS) given by

$$E_0^* = \left(s_h^*, s_a^*, i_h^*, l_0^*, m_a^*, f_n^*, f_c^*\right) = \left(1, 0, 0, 0, 0, \frac{\tilde{\theta}_f}{\mu_f}, 0\right).$$

On the other hand, from (10), we have

$$f_c^* = \frac{\mu_f}{\beta_3}\left(\frac{\beta_3 \tilde{\theta}_f}{\mu_f^2} - 1\right) = \frac{\mu_f}{\beta_3}\left(\mathcal{R}_f - 1\right).$$

where

$$\mathcal{R}_f = \frac{\beta_3 \tilde{\theta}_f}{\mu_f^2}.$$

We thus have the following result on the existence of f_c^*.

Lemma 4. *The existence of f_c^* is subject to $\mathcal{R}_f > 1$.*

However, we note that $\frac{\beta_3 \tilde{\theta}_f}{\mu_f^2} = \frac{\beta_3}{\mu_f} \cdot \frac{\tilde{\theta}_f}{\mu_f}$, where $\frac{\tilde{\theta}_f}{\mu_f}$ is the equilibrium size of the non-contaminated food products with $\frac{1}{\mu_f}$ as the duration of food contamination. So, $\mathcal{R}_f$ can be defined as the "food contamination constant" that measures the growth of contaminated food due to the contamination of uncontaminated food by contaminated food products. This is equivalent to the basic reproduction number $(\mathcal{R}_0)$ in disease modelling, see [22].

If $f_c^* = \frac{\mu_f}{\beta_3}(\mathcal{R}_f - 1)$, then expressing the second, third, and fifth equations of (8) in terms of i_h^* we obtain the following equation

$$m_a^* = \phi_0 i_h^*,$$

$$s_h^* = \phi_1 i_h^*, \qquad (11)$$

$$s_a^* = \phi_2 i_h^{*2},$$

where $\phi_0 = \frac{\pi}{\mu_0}$, $\phi_1 = \frac{\beta_3(\gamma + \mu_h)}{\beta_1 \mu_f (\mathcal{R}_f - 1)}$, and $\phi_2 = \frac{\tilde{\rho} \phi_0 \phi_1}{(\mu_h + \omega_1)}$. Substituting all the expressions in equation (11) into the first equation of (8), and after some algebraic simplifications we obtain the following quadratic equation

$$\xi_2 i_h^{*2} + \xi_1 i_h^* + \xi_0 = 0, \qquad (12)$$

where

$$\xi_0 = -\beta_3 (\mu_h + \delta_h) < 0,$$

$$\xi_1 = \gamma + \mu_h + \delta_h + \frac{\beta_3(\mu_h + \gamma)(\mu_h + \delta_h)}{\beta_1 \mu_f (\mathcal{R}_f - 1)} > 0 \text{ if } \mathcal{R}_f > 1,$$

$$\xi_2 = \frac{\pi \tilde{\rho} \beta_3 (\mu_h + \gamma)(\mu_h + \delta_h)}{\beta_1 \mu_0 \mu_f (\mu_h + \omega_1)(\mathcal{R}_f - 1)} > 0 \text{ if } \mathcal{R}_f > 1.$$

However, we note that the solutions of the quadratic equation (12) are given by

$$i_h^* = \frac{-\xi_1 \pm \sqrt{\xi_1^2 - 4\xi_2 \xi_0}}{2\xi_2}.$$

The solutions to (12) has one positive root when $\mathcal{R}_f > 1$. Biologically, this implies that the disease will persist and eventually invade the human population. This results in the Listeria steady states (LFS)

$$E_1^* = \left(s_h^*, s_a^*, l_0^*, m_a^*, f_n^*, f_c^* \right) = \left(\phi_1 i_h^*, \ \phi_2 i_h^*, \ 0, \ \phi_0 i_h^*, \ \frac{\tilde{\theta}_f}{\mu_f \left((\mathcal{R}_f - 1) + 1 \right)}, \ \frac{\mu_f}{\beta_3}(\mathcal{R}_f - 1) \right).$$

We note that at Listeria disease free steady state, there are contaminated food products which may result in Listeriosis infection in the human population.

Remark 1 *We note that, when $l^* = 0$, we have two steady states E_0^* and E_1^*. The existence of E_1^* is subject to the contaminated food generation number $(\mathcal{R}_f)$ been greater than 1. As long as $\mathcal{R}_f > 1$, even without Listeria in the environment, we will have the disease in the human population.*

CASE B: If $l^* = 1$, then from last equation of (8) solving for new f_n^*, we have that

$$f_n^{*+} = \frac{\mu_f f_c^*}{\beta_3 f_c^* + \beta_2}. \tag{13}$$

Substituting (13), into second last equation of (8) we obtain the following expression in terms of f_c^* after some algebraic simplifications

$$\nu_2 f_c^{*2} + \nu_1 f_c^* + \nu_0 = 0, \tag{14}$$

where

$$\nu_0 = -\beta_3 \tilde{\theta}_f < 0,$$

$$\nu_1 = \mu_f(\beta_2 + \mu_f)\left(1 - \frac{\beta_3 \tilde{\theta}_f}{\mu_f(\beta_2 + \mu_f)}\right),$$

$$\nu_2 = \mu_f \beta_3 > 0.$$

The solutions of the quadratic equation (14) given by

$$f_c^* = \frac{-\nu_1 \pm \sqrt{\nu_1^2 - 4\nu_2\nu_0}}{2\nu_2},$$

has one positive root irrespective of the signs of ν_1. The solutions of f_c^* say f_c^{*+} exists, but cannot be determined due to its intractability. Hence, as long as $l^* = 1$, we have a positive f_c^{*+}.

Now, we express the second, third, and fifth equation of (8) in terms of i_h^*, and obtain the following expressions

$$m_a^* = \Psi_0 i_h^*,$$

$$s_h^* = \Psi_1 i_h^*, \tag{15}$$

$$s_a^* = \Psi_2 i_h^{*2},$$

respectively, where $\Psi_0 = \frac{\pi}{\mu_0}$, $\Psi_1 = \frac{(\gamma + \mu_h)}{\beta_1 f_c^{*+}}$, and $\Psi_2 = \frac{\tilde{\rho}\Psi_0\Psi_1}{\mu_h + \omega_1}$. Similarly, substituting all the expressions from equations (15) into the first equation of (8), and after some algebraic manipulations we obtain the following quadratic equation in terms of i_h^*

$$\xi_5 i_h^{*2} + \xi_4 i_h^* + \xi_3 = 0, \tag{16}$$

where

$$\xi_3 = -(\mu_h + \delta_h) < 0,$$

$$\xi_4 = \frac{(\mu_h + \gamma)(\mu_h + \delta_h) + \beta_1 f_c^{*+}(\delta_h + \mu_h + \gamma)}{\beta_1 f_c^{*+}} > 0,$$

$$\xi_5 = \frac{\pi\rho(\gamma + \mu_h)(\delta_h + \mu_h)}{\beta_1 \mu_0 f_c^{*+}(\mu_h + \omega_1)} > 0.$$

The solutions (i_h^*) of the quadratic equation (16) given by

$$i_h^* = \frac{-\xi_4 \pm \sqrt{\xi_4^2 - 4\xi_5\xi_3}}{2\xi_5},$$

exists and has one positive root. We thus have the following result on the existence of new i_h^* say i_h^{*+}.

Lemma 5. *The steady state i_h^{*+} exists whenever f_c^{*+} exists.*

This results in the endemic steady states (ESS) given by

$$E_2^* = \left(s_h^{*+}, s_a^{*+}, l_1^*, m_a^{*+}, f_n^{*+}\right) = \left(\Psi_1 i_h^{*+}, \Psi_2 i_h^{*+}, 1, \Psi_0 i_h^{*+}, \frac{\mu_f f_c^{*+}}{\beta_3 f_c^{*+} + \beta_2}\right).$$

Hence, at endemic steady state, there are contaminated food products which result in the persistence of the Listeria infections in the human population.

Local Stability of the Disease Free Steady State

To analyse local stability of the disease free steady state (DFS), we show that the eigenvalues of the Jacobian matrix at DFS have negative real parts. We now state the following theorem for the DFS.

Theorem 6. *The disease free steady state (E_0^*), is locally asymptotically stable if $r_l < 0$ and $\mathcal{R}_f < 1$, otherwise it is unstable.*

Proof. The Jacobian of system (2) is given by the block matrix

$$J = \left(\begin{array}{c|c} A_1 & A_2 \\ \hline A_3 & A_4 \end{array}\right), \tag{17}$$

where

$$A_1 = \begin{pmatrix} -(\delta_h + \tilde{\rho}m_a^*) & -\delta_h + \omega_1 & -\delta_h & 0 \\ \tilde{\rho}m_a^* & -(\omega_1 + \mu_h) & 0 & 0 \\ \beta_1 f_c^* & 0 & -(\mu_h + \gamma) & 0 \\ 0 & 0 & 0 & r_l - 2r_l l^* \end{pmatrix}, \quad A_3 = \begin{pmatrix} 0 & 0 & \pi & 0 \\ 0 & 0 & 0 & -\beta_2 f_n^* \\ 0 & 0 & 0 & \beta_2 f_n^* \end{pmatrix},$$

$$A_2 = \begin{pmatrix} -\tilde{\rho}s_h^* & 0 & -\beta_1 s_h^* \\ \tilde{\rho}s_h^* & 0 & 0 \\ 0 & 0 & \beta_1 s_h^* \\ 0 & 0 & 0 \end{pmatrix} \quad \text{and } A_4 = \begin{pmatrix} -\mu_0 & 0 & 0 \\ 0 & -(\beta_2 l^* + \beta_3 f_c^* + \mu_f) & -\beta_3 f_n^* \\ 0 & \beta_2 l^* + \beta_3 f_c^* & \beta_3 f_n^* - \mu_f \end{pmatrix}.$$

Evaluating (17) at DFS we have that

$$J(E_0^*) = \left(\begin{array}{c|c} J_1(E_0^*) & J_2(E_0^*) \\ \hline J_3(E_0^*) & J_4(E_0^*) \end{array}\right),$$

where

$$J_1(E_0^*) = \begin{pmatrix} -\delta_h & \omega_1 - \delta_h) & -\delta_h & 0 \\ 0 & -(\omega_1 + \mu_h) & 0 & 0 \\ 0 & 0 & -(\mu_h + \gamma) & 0 \\ 0 & 0 & 0 & r_l \end{pmatrix}, \quad J_2(E_0^*) = \begin{pmatrix} -\tilde{\rho} & 0 & -\beta_1 \\ \tilde{\rho} & 0 & 0 \\ 0 & 0 & \beta_1 \\ 0 & 0 & 0 \end{pmatrix},$$

$$J_3(E_0^*) = \begin{pmatrix} 0 & \pi_0\alpha & 0 \\ 0 & 0 & -\beta_2\frac{\tilde{\theta}_f}{\mu_f} \\ 0 & 0 & \beta_2\frac{\tilde{\theta}_f}{\mu_f} \end{pmatrix} \quad \text{and } J_4(E_0^*) = \begin{pmatrix} -\mu_0 & 0 & 0 \\ 0 & -\mu_f & -\beta_3\frac{\tilde{\theta}_f}{\mu_f} \\ 0 & 0 & \beta_3\frac{\tilde{\theta}_f}{\mu_f} - \mu_f \end{pmatrix}.$$

The eigenvalues of $J(E_0^*)$ are; $\lambda_1 = -\delta_h$, $\lambda_2 = -(\omega_1 + \mu_h)$, $\lambda_3 = -(\mu_h + \gamma)$, and $\lambda_4 = r_l$, $\lambda_5 = -\mu_0$, $\lambda_6 = -\mu_f$, and $\lambda_7 = \mu_f(\mathcal{R}_f - 1) < 0$ provided that $\mathcal{R}_f < 1$. We note that if r_l is negative, all the eigenvalues are negative, hence E_0^* is locally asymptotically stable. However, if r_l is positive, E_0^* unstable. Note that if $r_l < 0$ then Listeria in the environment dies out, resulting in a locally asymptotically stable equilibrium point while it establishes itself when $r_l > 0$ resulting in an unstable steady state.

Local Stability of the Listeria Free Steady State

We state the following theorem for the local stability of Listeria free steady state.

Theorem 7. *The Listeria free steady state (E_1^*), is locally asymptotically stable if $r_l < 0$ and $\mathcal{R}_f > 1$, otherwise it is unstable.*

Proof. Evaluating (17) at Listeria free steady state, we have that

$$J(E_1^*) = \left(\begin{array}{c|c} J_1(E_1^*) & J_2(E_1^*) \\ \hline J_3(E_1^*) & J_4(E_1^*) \end{array} \right),$$

where

$$J_1(E_1^*) = \begin{pmatrix} -(\delta_h + \tilde{\rho}\phi_0 i_h^*) & -\delta_h + \omega_1 & -\delta_h & 0 \\ \tilde{\rho}\phi_0 i_h^* & -(\omega_1 + \mu_h) & 0 & 0 \\ \frac{\beta_1\mu_f}{\beta_3}(\mathcal{R}_f - 1) & 0 & -(\mu_h + \gamma) & 0 \\ 0 & 0 & 0 & r_l \end{pmatrix}, \quad J_2(E_1^*) = \begin{pmatrix} -\tilde{\rho}\phi_1 i_h^* & 0 & -\beta_1\phi_1 i_h^* \\ \tilde{\rho}\phi_1 i_h^* & 0 & 0 \\ 0 & 0 & \beta_1\phi_1 i_h^* \\ 0 & 0 & 0 \end{pmatrix},$$

$$J_3(E_1^*) = \begin{pmatrix} 0 & \pi_0\alpha & 0 \\ 0 & 0 & \frac{-\beta_2\tilde{\theta}_f}{\mu_f\mathcal{R}_f} \\ 0 & 0 & \frac{\beta_2\tilde{\theta}_f}{\mu_f\mathcal{R}_f} \end{pmatrix}, \quad \text{and } J_4(E_1^*) = \begin{pmatrix} -\mu_0 & 0 & 0 \\ 0 & -\mu_f\mathcal{R}_f & \frac{-\beta_3\tilde{\theta}_f}{\mu_f\mathcal{R}_f} \\ 0 & \mu_f(\mathcal{R}_f - 1) & \frac{\beta_3\tilde{\theta}_f}{\mu_f\mathcal{R}_f} - \mu_f \end{pmatrix}.$$

The eigenvalues from $J_1(E_1^*)$ are; $\lambda_1 = r_l$ and the solutions to the cubic equation

$$\lambda^3 + a_2\lambda^2 + a_1\lambda + a_0 = 0,$$

where

$$a_0 = \frac{1}{\beta_3\mu_0}\left(\beta_3(\gamma+\mu_h)[\pi\tilde{\rho}\mu_h i_h^* + \delta_h(\pi\tilde{\rho}i_h^* + \mu_0(\mu_h+\omega_1))] + (\beta_1\mu_0\mu_f\delta_h)(\mu_h+\omega_1)(\mathcal{R}_f-1)\right),$$

$$a_1 = \frac{1}{\beta_3\mu_0}\left(\beta_3\left(\pi\tilde{\rho}i_h^*(2\mu_h+\delta_h+\gamma)+\mu_0\left((\mu_h+\gamma)(\mu_h+\omega_1)+\delta_h(2\mu_h+\omega_1+\gamma)\right)\right)\right.$$
$$\left. +(\beta_1\mu_0\mu_f\delta_h)(\mathcal{R}_f-1)\right),$$

$$a_2 = 2\mu_h + \delta_h + \omega_1 + \gamma + \frac{\pi\tilde{\rho}i_h^*}{\mu_0}.$$

Given that a_0, a_1 and a_2 are positive provided that $\mathcal{R}_f > 1$, we note that

$$a_2 a_1 - a_0 = \frac{1}{\beta_3\mu_0^2}\left(\Upsilon_0 + \beta_1\delta_h\mu_0\mu_f(\pi\tilde{\rho}i_h^* + \mu_0(\gamma+\delta_h+\mu_h))(\mathcal{R}_f-1)\right).$$

for

$$\Upsilon_0 = \beta_3(\pi\tilde{\rho}i_h^* + \mu_0(\delta_h+\mu_h+\omega_1))(\pi\tilde{\rho}i_h^*(\gamma+\delta_h+2\mu_h)+\mu_0(\gamma+\delta_h+\mu_h)(2\mu_h+\omega_1+\gamma)).$$

We also, note that $a_2 a_1 - a_0 > 0$ if $\mathcal{R}_f > 1$. Hence, by the Routh-Hurwitz criterion the eigenvalues of $J(E_1^*)$ have negative real parts. The rest of the eigenvalues are determined from $J_4(E_1^*)$, which are; $\lambda_5 = -\mu_0$ and the solutions to the quadratic equation

$$\lambda^2 + a_4\lambda + a_3 = 0,$$

where

$$a_3 = \mu_f^2(\mathcal{R}_f-1) \text{ and } a_4 = \mu_f\mathcal{R}_f.$$

We thus have $a_3 > 0$ if $\mathcal{R}_f > 1$ and a_4 is always positive. Also, the eigenvalues have negative real part by Routh-Hurwitz criteria. Therefore, all the eigenvalues of $J(E_1^*)$ are negative if $r_l < 0$. Thus, E_1^* is locally asymptotically stable, otherwise unstable if $r_l > 0$.

Local Stability of the Endemic Steady State

We now state the following theorem on the local stability of endemic steady state.

Theorem 8. *The endemic steady state (E_2^*), is locally asymptotically stable if $\mathcal{R}_f > 1$, otherwise it is unstable.*

Proof. Evaluating (17) at endemic steady state, we have that

$$J(E_2^*) = \left(\begin{array}{c|c} J_1(E_2^*) & J_2(E_2^*) \\ \hline J_3(E_2^*) & J_4(E_2^*) \end{array}\right),$$

where

$$J_1(E_2^*) = \begin{pmatrix} -(\delta_h + \tilde\rho\Psi_0 i_h^{*+}) & -\delta_h + \omega_1 & -\delta_h & 0 \\ \tilde\rho\Psi_0 i_h^{*+} & -(\omega_1 + \mu_h) & 0 & 0 \\ \beta_1 f_n^{*+} & 0 & -(\mu_h + \gamma) & 0 \\ 0 & 0 & 0 & -r_l \end{pmatrix},$$

$$J_2(E_2^*) = \begin{pmatrix} -\tilde\rho\Psi_1 i_h^{*+} & 0 & -\beta_1\Psi_1 i_h^{*+} \\ \tilde\rho\Psi_1 i_h^{*+} & 0 & 0 \\ 0 & 0 & \beta_1\Psi_1 i_h^{*+} \\ 0 & 0 & 0 \end{pmatrix},$$

$$J_3(E_2^*) = \begin{pmatrix} 0 & \pi_0\alpha & 0 \\ 0 & 0 & -\beta_2 f_n^{*+} \\ 0 & 0 & \beta_2 f_n^{*+} \end{pmatrix} \text{ and } J_4(E_2^*) = \begin{pmatrix} -\mu_0 & 0 & 0 \\ 0 & -(\beta_2 + \beta_3 f_c^{*+} + \mu_f) & -\beta_3 f_n^{*+} \\ 0 & \beta_2 + \beta_3 f_c^{*+} & \beta_3 f_n^{*+} - \mu_f \end{pmatrix}.$$

The eigenvalues from $J_1(E_2^*)$ are; $\lambda_1 = -r_l$ and the solutions to the cubic equation

$$\lambda^3 + b_2\lambda^2 + b_1\lambda + b_0 = 0,$$

where

$$b_0 = \frac{1}{\mu_0}\left(\pi\tilde\rho i_h^{*+}\mu_h(\gamma + \mu_h) + \delta_h\left(\pi\tilde\rho i_h^{*+}(\gamma + \mu_h) + \mu_0(\gamma + \beta_1 f_c^{*+} + \mu_h)(\mu_h + \omega_1)\right)\right),$$

$$b_1 = \frac{1}{\mu_0}\left(\pi\tilde\rho i_h^{*+}(\gamma + \delta_h + 2\mu_h) + \mu_0\left((\gamma + \mu_h)(\mu_h + \omega_1) + \delta_h(\gamma + \beta_1 f_c^{*+} + 2\mu_h + \omega_1)\right)\right),$$

$$b_2 = 2\mu_h + \delta_h + \omega_1 + \gamma + \frac{\pi\tilde\rho i_h^{*+}}{\mu_0}.$$

Note that b_0, b_1 and b_2 are positive and

$$b_2 b_1 - b_0 = \frac{1}{\mu_0^2}\left(\Upsilon_1 + \Upsilon_2\right) > 0,$$

for

$$\Upsilon_1 = \pi^2\tilde\rho^2 i_h^{*+2}(\gamma + \delta_h + 2\mu_h) + \mu_0^2(\gamma + \delta_h + \mu_h)\Big((\mu_h + \omega_1)(\gamma + 2\mu_h + \omega_1)$$
$$+ \delta_h(\gamma + \beta_1 f_c^{*+} + 2\mu_h + \omega_1)\Big) \text{ and }$$

$$\Upsilon_2 = \pi\tilde\rho\mu_0 i_h^{*+}(\delta_h^2 + (\gamma + 2\mu_h)^2 + (2\gamma + 3\mu_h)\omega_1 + \delta_h[\beta_1 f_c^{*+} + 5\mu_h + 2(\gamma + \omega_1)]).$$

Thus, by the Routh-Hurwitz criteria, the eigenvalues have negative real parts. The remaining eigenvalues are determined from $J_4(E_2^*)$, which are; $\lambda_5 = -\mu_0$ and the solutions to the quadratic equation

$$\lambda^2 + b_4\lambda + b_3 = 0,$$

where

$$b_3 = \frac{\mu_f\left((\beta_2 + \beta_3 f_c^{*+})^2 + \beta_2\mu_f\right)}{\beta_2 + \beta_3 f_c^{*+}} > 0,$$

$$b_4 = \beta_2 + \beta_3 f_c^{*+} + \frac{(2\beta_2 + \beta_3 f_c^{*+})\mu_f}{\beta_2 + \beta_3 f_c^{*+}} > 0.$$

Hence, the eigenvalues have negative real parts by Routh-Hurwitz criterion. Therefore, all the eigenvalues of $J(E_2^*)$ have negative real parts. Thus, E_2^* is locally asymptotically stable, otherwise unstable.

Numerical Results

Parameter Estimation

Numerical simulations of the model system (2) were done using the set of parameters values given in Table 1. Most of the parameter values were estimated since there are very few mathematical models in literature, that have been done on Listeria Monocytogenes disease dynamics and hence the parameter values are elusive. We used a fourth order Runge-Kutta numerical scheme implemented in Matlab to perform the simulations with the initial conditions: $s_h(0) = 0.42, s_a(0) = 0.53, i_h(0) = 0.05, l(0) = 0.1, m_a(0) = 0.25, f_n = 0.2$, and $f_c = 0.8$. The initial conditions were hypothetically chosen for the numerical simulations presented in section, and are thus only for illustrative purpose and do not represent any observed scenario.

Table 1: Parameter values used for numerical simulations.

Parameter	Symbol	Value/day	Source
Mortality rate of humans	μ_h	0.02/365	[24]
Recovery rate of humans	γ	0.034	[24]
Rate of lost of immunity for humans	δ_h	0.2	[24]
Waining rate of aware susceptibles	ω_1	0.25	[13]
Rate at which non-aware susceptibles becomes aware	$\tilde{\rho}$	0.9	Assumed
Net growth rate of Listeria	r_l	0.1	Assumed
Depletion rate of media campaigns	μ_0	0.03	[14]
Rate of implementation of media campaign	π	0.001	[14]
Contact rate between Listeria and humans	β_1	0.0056	Assumed
Food contamination rate by Listeria	β_2	0.09	Assumed
The food contamination rate	β_3	0.0052	Assumed
Removal rate of food products	μ_f	0.009	Assumed
Rate of food processing	$\tilde{\theta}_f$	0.055	Assumed

(a)

(b)

Fig. 2: Scatter plots of parameters with the negative ($\tilde{\rho}$) and positive (ω_1) PRCCs.

Sensitivity Analysis

Sobol sensitivity analysis [23], were used to determine the model parameters that are sensitive to changes in some variable of the model system (2). We performed simulations for some chosen parameters π, μ_0, $\tilde{\rho}$, ω_1, β_1, β_2, β_3, and μ_f versus some of the model state variables s_a, i_h and f_c to show their respective partial rank correlation coefficients (PRCC) over time. These parameters and state variables were selected because they are the most significant in the Listeria disease transmission and control according to our model formulation. The simulations were done using Matlab with 1000 runs over 800 days as depicted in Figure 3, and the scatter plots for parameters with positive and negative PRCC shown in Figure 2. Thus, an increase in the rate at which susceptible individuals move into the aware susceptible class results in a fewer number of humans been infected with the Listeriosis and parameter ω_1 with negative correlation signifies that if aware individuals revert to being susceptible, then they are prone to contracting the disease as depicted in Figure 2 respectively. This highlights the importance of media campaigns in disease control.

Fig. 3: Partial correlation coefficients of parameters π, μ_0, $\tilde{\rho}$, ω_1, β_1, β_2, β_3, and μ_f over time. Note that, ρ in the the legend represents $\tilde{\rho}$ due to Matlab do not compile tilda.

Effects of Varying π and μ_0 on Listeriosis Spread

This subsection is devoted to a numerical simulations that show the effects of increasing and decreasing media campaigns over time. Figure 4(a) reveals that an increase in the awareness campaigns result in

a decrease in the number of infected humans, while a reduction of media campaigns results in more humans been infected with Listeriosis as shown in Figure 4(b).

Fig. 4: (a) Effects of implementing of media campaigns (π) on the fraction of infected humans. (b) Effects of depletion of media campaign (μ_0) on the fraction of infected humans. (c) Effects of implementation of media campaign (π) on the fraction of aware susceptible humans. (d) Effects of depletion of media campaign (μ_0) on the fraction of aware susceptible humans.

Effects of β_3 and μ_f on $\mathcal{R}_f$

In Figure 5, we present a contour of the food contamination rate β_3 and rate of food product removal, μ_f, versus the food contamination constant, $\mathcal{R}_f$. An increase in the contamination of non-contaminated food by contaminated food products results in an increase in the value of $\mathcal{R}_f$. Hence, more humans get infected with Listeriosis. Also, an increase in the removal of contaminated food products results in a decrease in the values of $\mathcal{R}_f$ which implies that fewer humans contract the disease. Note that, the changes in the value of μ_f do not significantly impact $\mathcal{R}_f$ when compared to β_3.

Fig. 5: Contour plot of contaminated food generation number $(\mathcal{R}_f)$ as a function of β_3 and μ_f.

Effects of $\tilde{\theta}_f$ and μ_f on $\mathcal{R}_f$
Figure 6, illustrates a mesh plot showing the variation of $\tilde{\theta}_f$, μ_f and the contaminated food generation number $\mathcal{R}_f$. It is seen that an increase in the rate of food processing increases $\mathcal{R}_f$, while an increase in the removal of contaminated food results in a decrease in $\mathcal{R}_f$. Note that at the intersection of the plane $\mathcal{R}_f = 1$ and the mesh plot, we obtain all values of $\tilde{\theta}_f$ and μ_f necessary for the eradication of the disease.

Fig. 6: Effects of β_3 and μ_f on contaminated food generation number $(\mathcal{R}_f)$.

Impacts of Varying $\tilde{\rho}$ and μ_f on Infected Humans
Figure 7 depicts the effects of varying parameters $\tilde{\rho}$ and μ_f. In Figure 7(a), we observe that increasing

$\tilde{\rho}$ decreases the number of infected humans. This results show that media campaigns have the potential to reduce the number of infectives. Also, we note that an increase in $\tilde{\rho}$ does not impact the value of the contaminated food generation number, $\mathcal{R}_f$. On the other hand, increasing μ_f also reduces the number of infected individuals and $\mathcal{R}_f$ (see Figure 7(b)). So the removal of contaminated food products during an outbreak is an important intervention in the control of Listeriosis.

Fig. 7: (a) Effects varying the rate of non-aware susceptible becomes aware susceptible ($\tilde{\rho}$) on fraction of infected humans. The value for the $\mathcal{R}_f$ is 3.5309. (b) Effects varying removal of contaminated food products (μ_f) on the fraction of infected humans.

Conclusion

A deterministic model on the potential role of media campaigns on Listeriosis disease transmissions was developed and analysed. Stability analysis of the model was done in terms of the food contamination constant $\mathcal{R}_f$. The model exhibited three different steady states which are: the disease-free, Listeria free and endemic equilibria. The disease-free and the Listeria free steady states are locally asymptotically stable if the net growth rate $r_l < 0$ and $\mathcal{R}_f < 1$ respectively, while the endemic equilibrium state is locally asymptotically stable if $\mathcal{R}_f > 1$.

On the other hand from our numerical results, it was established that an increase in the removal of contaminated food products and increase in the rate at which the susceptible individuals become aware susceptible individuals leads to a decrease in the number of infected humans (see Figure 7). Thus to effectively control Listeria disease spread, policymakers, public health, governments, and global stakeholders are advised to implement media campaigns that do not wane as time progresses. This means that media campaigns need to be effective. Also, these interventions from the campaigns should target mainly susceptible individuals on the danger of Listeriosis. As people adhere to the media campaign effectively, it helps reduce and control the number of infected humans which lead to less disease trans-

mission. Furthermore, practising proper hygiene, decontamination of contaminated food products such as; thermal processing, removal of contamination by excision, irradiation by ionizing, high-pressure processing and the application of antimicrobial chemicals are also other measures to control Listeriosis.

The model presented in this paper is not without fallibility. First, the model was not fitted to epidemiological data. This would have been useful in the validation of key epidemiological parameters. Second, the contribution of infected humans and contaminated food products to the gowth of the bacteria is assumed to be negligible. However, one can argue that Listeria in the environment could also result from contaminated food products and infected humans. Last, we assumed that the aware susceptibles can not be infected bi Listeria. This assumption means that the awareness programs are not imperfect, which may be contested. Despite these shortcomings, the results obtained in this paper is still important in the control Listeriosis disease transmission dynamics in an event of an outbreak.

References

[1] H. Hof, History and epidemiology of Listeriosis, FEMS Immunology and Medical Microbiology, 35(3), (2003), pp. 199–202.

[2] EFSA Panel on Biological Hazards (BIOHAZ) and Ricci, et. al, Listeria monocytogenes contamination of ready-to-eat foods and the risk for human health in the EU, EFSA Journal, 16(1),(2018), pp.e05134.

[3] E. Tambo, C.S. Yah, and G. Madjou, Deadly Listeriosis outbreaks in South Africa and Australia: re-inforcing food safety surveillance and emergency response actions, Journal of Advances in Virology Research, 1(1), (2018), pp.1–9.

[4] National Listeria Incident Management Team, `http://www.health.gov.za/index.php/component/phocadownload/category/439`, Accessed 20 August 2020.

[5] EFSA, European Food safty Authority,`https://ecdc.europa.eu/sites/portal/files/documents/2018_ECDC-EFSA_ROA_Update1_UI-444_Listeria.pdf`, Accessed 3 July 2020.

[6] M. Ghosh, P. Chandra, P. Sinha, and J.B. Shukla, Modeling and analysis of the spread of carrier dependent infectious diseases with environmental effects, Journal of Biological Systems 11(3), (2003), pp. 325–-335.

[7] R. Naresh, S. Pandey, and A.K. Misra, Analysis of a vaccination model for carrier dependent infectious diseases with environmental effects, Nonlinear Analysis. Modelling and Control 13(3), (2008), pp. 331–350.

[8] J. Li, Effects of behavior change on the spread of AIDS epidemic, Mathematical and Computer Modelling 16(6–7), (1992), pp. 103–-111.

[9] D. Njankou, S. Diane and F. Nyabadza, Modelling the potential role of media campaigns in ebola transmission dynamics, International Journal of Differential Equations, 2017, Article ID 3758269, (2017), 13 pp.

[10] F. Nyabadza, C. Chiyaka, Z. Mukandavire, and S. Musekwa, Analysis of an HIV/AIDS model with public-health information campaigns and individual withdrawal, Journal of Biological Systems, 18(02), (2010), pp. 357–375, `https://doi:10.1142/S0218339010003329`.

[11] J.M. Tchuenche, N. Dube, C.P. Bhunu, R.J. Smith, and C.T. Bauch, The impact of media coverage on the transmission dynamics of human influenza. BMC Public Health, 11(1), (2011), p. S5.

[12] C. Yang, X. Wang, D. Gao, and J. Wang, Impact of awareness programs on cholera dynamics: two modeling approaches. Bulletin of Mathematical Biology, 79(9), (2017), pp. 2109–2131.

[13] A.K. Misra, A. Sharma and J.B. Shukla, Modeling and analysis of effects of awareness programs by media on the spread of infectious diseases, Mathematical and Computer Modelling, 53(5-6), (2011), (2011), pp. 1221–1228.

[14] N. Kaur, M. Ghosh, and S.S. Bhatia, Modeling and analysis of an SIRS epidemic model with effect of awareness programs by media, International Journal Mathematical Computational National Physics Engineering, 8(1),(2014), pp. 233–239.

[15] G.O. Agaba, Y.N. Kyrychko and K.B. Blyuss, Mathematical model for the impact of awareness on the dynamics of infectious diseases, Mathematical Biosciences, 286, (2017), pp. 22–30.

[16] A. Sharma and A.K. Misra, Backward bifurcation in a smoking cessation model with media campaigns, Applied Mathematical Modelling, 39(3-4), (2015), pp. 1087–-1098.

[17] A. Kumar, P.K. Srivastava and Y. Takeuchi, Modeling the role of information and limited optimal treatment on disease prevalence, Journal of Theoretical Biology, 414, (2017), pp. 103–119.

[18] Y. Xiao, T. Zhao, and S. Tang, Dynamics of an infectious diseases with media/psychology induced non-smooth incidence, Mathematical Biosciences and Engineering, 10(2), (2013), pp. 445–461.

[19] L. Yiping and C. Jing-an, The impact of media coverage on the dynamics of infectious disease, International Journal of Biomathematics, 1(01), (2008), pp. 65–74.

[20] K.A. Pawelek, A. Oeldorf-Hirsch and L. Rong, Modeling the impact of Twitter on influenza epidemics, Mathematical Biosciences and Engineering, 11(6), (2014), pp. 1337–-1356.

[21] X. Yang, L. Chen and J. Chen, Permanence and positive periodic solution for the single-species nonautonomous delay diffusive models, Computers and Mathematics with Applications, 32(4), (1996), pp. 109–116.

[22] O. Diekmann, J.A.P. Heesterbeek and J.A.J. Metz, On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations, Journal of Mathematical Biology, 28(4), (1990), pp. 365–382.

[23] X.Y. Zhang, M.N. Trame, L.J. Lesko, and S. Schmidt, Sobol sensitivity analysis: a tool to guide the development and evaluation of systems pharmacology models, CPT: Pharmacometrics & Systems Pharmacology, 4(2), (2015), pp.69–-79.

[24] C.W. Chukwu, and F. Nyabadza, A theoretical model of listeriosis driven by cross contamination of ready-to-eat food products, International Journal of Mathematics and Mathematical Sciences, 09 March 2020, (2020), 14 pp.