Liver immune abnormalities persist after cure of Hepatitis C Virus by anti-

viral therapy

Xia Wu[1, *], Jessica B. Roberto[1], Allison Knupp[1], Alexander L. Greninger[1,3], Camtu D. Truong[1],

Nicole Hollingshead[1], Heidi L. Kenerson[2], Marianne Tuefferd[4], Antony Chen[4], Helen Horton[4],

Keith R. Jerome[1,3], Stephen J. Polyak[1], Raymond S. Yeung[2] and Ian N. Crispe[1]

[1]Department of Laboratory Medicine and Pathology, and [2]Department of Surgery, University of

Washington, Seattle, WA, USA 98195

[3]Fred Hutchinson Cancer Research Institute, Seattle, Washington, USA.

[4]Infectious Diseases and Vaccines, Janssen Research and Development, B-2340 Beerse,

Belgium.

*To whom correspondence should be addressed:

Xia Wu, Ph.D.

Department of Medicine, University of Washington,

750 Republican St., E630, Seattle, WA 98109

E-mail: xiawu2@uw.edu Tel: 1-206-221-3343

Keywords: Hepatitis C virus, direct-acting antiviral treatment, liver slice culture, innate

immunity, inflammation, fibrosis

Short Title: Immune abnormalities in human liver after HCV cure

Electronic word count

5926 words

Number of figures and tables

6 figures, 1 table

Financial support

This work was supported by NIH NIAID R56 grant AI143683; US Department of Defense grant CA150370; Janssen Research and Development; the Seattle Foundation; and the University of Washington.

Authors' contributions

X.W., J.B.R., A.C., H.H., R.S.Y. and I.N.C. designed research; X.W., J.B.R., A.K., A.L.G., N.H., H.L.K, M.T. and A.C. performed research; K.R.J., S.J.P. and R.S.Y. contributed reagents/analytic tools; X.W., A.L.G., C.D.T. and I.N.C. analyzed data; X.W., S.J.P. and I.N.C. wrote the manuscript; all authors reviewed the manuscript and provided suggestions.

Abstract

Direct-Acting Antiviral (DAA) drugs are highly effective in eliminating HCV in most

chronically infected subjects, but it is unclear whether the multiple mechanisms employed by

HCV to disable both innate and adaptive immunity cease to function as soon as HCV is

eliminated. To test this, we evaluated the capacity of human liver tissue to respond to TLR3 and

TLR4 ligands using non-infected liver tissue, tissue with active HCV infection, and tissue from

which HCV had been eliminated by DAA. We found that DAA-treated, formerly HCV-infected

liver tissue manifested ongoing abnormalities of innate immunity that mapped to liver non-

parenchymal cells (NPC). Hepatic innate immunity was not suppressed but enhanced in HCV-

infected tissue, and these abnormalities were not corrected in the successfully DAA-treated liver

tissue. In conclusion, ongoing immune activation persists in formerly HCV-infected but now

DAA-cured liver.

(136 words)

Introduction

HCV is a positive-strand RNA virus classified in its own genus, *Hepacivirus*, within the

Flaviviridae family. Approximately 71 million individuals are infected with HCV worldwide

[1]. The acute phase of infection is often subclinical, but 55–85% of infected individuals

develop a chronic infection leading to progressive liver pathology [2]. The World Health

Organization (WHO) estimated that approximately 399,000 people died from HCV- related

diseases in 2016, mostly from cirrhosis and hepatocellular carcinoma (HCC).

The recent introduction of highly effective direct-acting antiviral (DAA) drugs has presented a

window of opportunity to compare innate immunity in virus-free human subjects, chronically

HCV-infected but untreated individuals, and individuals successfully treated for HCV with

DAA. This window is closing as most HCV patients in resource-rich countries are now being

treated with DAA, with substantial benefit [3]. However, the extent to which such drugs revert

the liver to normal is controversial, with some studies suggesting that HCV clearance with DAA

therapy leaves residual abnormalities in innate immunity in peripheral monocytes and NK cells

[4, 5], adaptive immunity in peripheral CD4+ and CD8+ T cells [6-9], in γδ-T cells [10], in

mucosal associated invariant T cells [11], and in the persisting risk to hepatocellular cancer [12-

15]. Other studies also found abnormal serum lipids [16], persistent epigenetic modifications

[17], and sustained hepatic inflammation [18] in liver biopsies after DAA treatment. To our

knowledge, no studies have evaluated hepatic immunity of untreated versus DAA-treated HCV

infection in the context of intact liver tissue.

We developed a liver slice culture model in which human liver tissue can be sustained for up to 2

weeks [19], allowing the evaluation of innate immune responses in liver slices collected from

non-infected subjects, chronic HCV-infected patients, and patients whose HCV was cured by

DAA treatment. Because of the emerging concept that the liver is not normal after HCV

eradication, the objective of the study was to investigate which aspects of immune pathology are

not corrected by DAA cure of chronic HCV infection. Our data indicate that, in formerly HCV-

infected but now DAA-cured liver, ongoing immune activation co-exists with fibrosis.

Materials and Methods

Liver slice culture

Fresh liver tissues were obtained from patients undergoing liver resection at the University of

Washington Medical Center (Seattle, WA, USA).   Liver tissue was obtained under Institutional

Review Board protocol #00001852. Liver slices were cultured and assayed for immune gene

function with *ex vivo* stimulation assay with polyinosinic–polycytidylic acid (poly-I:C) or

lipopolysaccharide endotoxin (LPS) as described previously [19]. *Gene expression data*

included 8 HCV-infected, 10 DAA-cured, and 9 non-infected control patients. Liver slices of

four of these HCV-infected and eight DAA-cured patients were further analyzed with fibrosis

staining. Additional details are provided in supplementary materials.

RNA isolation and qRT-PCR analysis

The RNA of liver slices or the purified liver cells was isolated with TRIzol and the Direct-zol

RNA MiniPrep Kit (Zymo Research, Irvine, CA, USA). Gene expression was analyzed with

fluidigm platform as previously described [19], with normalization to the arithmetic mean of Ct

values of ACTB, HPRT, and GAPDH. Additional details are provided in supplementary

materials.

HCV RNA measurement

HCV RNA was measured with an Food and Drug Administration-approved Abbott Real-Time

HCV assay (Abbott Molecular, Des Plaines, IL, USA) as previously described [18, 20]. Fifty ng

of total RNA for each sample was input in the assay, and HCV IU/ng total RNA was calculated

from the calibration curve of positive controls. The reliable detection limit for HCV with this

assay was 1.2 IU/ 50 ng liver total RNA.

Results

Ex vivo **detection of HCV RNA with liver slices**

We first examined the HCV status of liver slices that were collected from three groups of

patients, including untreated patients with chronic HCV infection, patients previously infected

with HCV but treated and cured with DAA therapy, and patients who had no history of HCV

infection (**Fig. 1A**).   Liver slices collected from all eight chronic HCV patients contained

detectable HCV RNA. None of the non-infected or DAA-treated subjects were positive for HCV

RNA *ex vivo* (**Fig. 2A**). Chronic HCV-infected liver slices continued to be HCV positive after 7

days of *ex vivo* culture (**Fig. 2B**), and the viral load on day 7 did not significantly differ from day

0 (Wilcoxon matched-pairs signed rank test, two-tailed, P=0.4688). The detection of HCV RNA

in *ex vivo* cultured liver slices is consistent with a previous report that human liver slices could

be infected *ex vivo* with HCV [21].

Persistent fibrosis in untreated HCV versus DAA-treated livers

Previous reports on the impact of liver fibrosis with DAA treatment relied on the indirect

measurements with transient elastography and noninvasive fibrosis indices [22, 23]. We

examined the liver fibrosis directly using picrosirius red and trichrome staining (**Fig. S1**). Six

out of eight DAA-cured patients met the definition of cirrhosis (i.e. fibrosis score of 4,

Scheuer/Batts-Ludwig method) in livers despite the absence of HCV RNA (**Fig. 2C**). Likewise,

the untreated chronic HCV livers were fibrotic, as previously reported [24]. None of the non-

infected subjects had liver fibrosis. Thus, liver fibrosis was not corrected by DAA treatment,

despite the elimination of HCV RNA, when we evaluated livers for fibrosis out to 20 months

from the completion of DAA treatment (**Table S1**).

Ongoing immune activation persists in liver slices of chronic HCV-infected and DAA-cured

subjects

We used multiplex qRT-PCR to examine 140 genes functioning in immune activation and

suppression, and tissue repair.   Forty immune genes differed in expression in chronic HCV-

infected livers versus non-infected patients (Mann Whitney test, two-tailed, $P < 0.05$) (**Fig. 3A**).

This included antiviral interferon (IFN)-stimulated genes (ISGs) including *IFIT1/2/3, ISG15,*

RSAD2, MX1, CXCL9, CXCL10, consistent with previous reports [25-28]. In addition, immune

activation genes including MHC genes *HLA-A, HLA-DRA*, *CIITA*, *CD80, CD86*, tumor necrosis

factor (TNF) superfamily genes *OX40, 4-1BB, TNFSF9, TNFRSF18* and inflammation genes

including *CASP1, IL-12A, IL-12B, TNF, IL-18*, and *NFKB* were also significantly up-regulated

in HCV-infected livers (**Fig. S3**).

DAA treatment restored the antiviral genes to baseline, as reported previously [29], but some

immune genes were remained elevated after cure of HCV infection (**Figs. 3BC, S2, S3**). These

included immune activation genes (*HLA-A, CIITA, 4-1BB, CCL5, CCR2*), inflammatory genes

(*CASP1, IL-12A*), and immune suppression genes (*CTLA4, TIGIT, IDO1, ARG1*).

A recent paper [17] reported HCV-induced epigenetic alterations persisted in liver tissue in

DAA-cured patients after SVR. The phosphoinositol-3-kinase (PI3K) pathway-linked genes

SPHK1 and *SOX9* were among those genes harboring epigenetic modification insensitive to

DAA treatment. We therefore analyzed a more extended set of PI3K related genes in different

liver tissue, and found that *SPHK1, BTG2, SOX9, SNAP25*, and *GPRC5B* were significantly up-

regulated in chronic HCV infected livers (**Fig. S3E**). Furthermore, *SPHK1, BTG2*, and *SOX9*

remained significantly up-regulated after HCV clearance with DAA treatment (**Fig. S3E**). The

up-regulation of PI3K genes may be associated with persistent liver fibrosis [30, 31].

The dysregulation of immunoregulatory genes in DAA-cured liver tissue maps to hepatic

non-parenchymal immune cells

To identify the cellular origin of the dysregulated immune genes, we mapped gene expression to

specific liver cell types, using FACS purified liver cells [19] (**Figure 1B**). Six liver wedge

samples of non-infected liver tissue were large enough in size and anatomically ideal for

perfusion through the vasculature, which allowed them to be dissociated and analyzed for gene

expression. We could not do this with samples of DAA-cured patients due to limitations in the

amount of liver tissue obtained. Compared with the total liver and hepatocyte enriched fraction,

the majority of the antiviral genes, immune activation, inflammation genes were significantly

enriched in the NPC (**Table 1**). Even though NPC may not be the cell type targeted by HCV

infection, previous studies demonstrate that NPC can be activated through paracrine signaling by

the neighboring HCV-infected hepatocytes [27, 32, 33].

Innate immune responses were altered in HCV infected and DAA-cured liver slices

We stimulated liver slices with poly-I:C (TLR3 agonist) and LPS (TLR4 agonist) for varying

times using the method established in a previous study [19]. We chose day 7 as the time to add

novel immune stimuli because the immune induction by tissue slicing was broadly stabilized

after 4 days of *ex vivo* culture [19]. The immune genes were recorded at the day 7 baseline prior

to the treatment. A number of the immune genes remained at elevated baseline in HCV-infected

or DAA-cured samples compared with non-infected control (**Fig. 4**). The PBS mock stimulation

did not significantly induce IFNs and ISGs in liver slices [19]. Expression of TLR3, TLR4 and

NF-κB were not significantly different at day 7 among chronic HCV-infected, DAA-treated and

non-infected liver slices (**Fig. S3A and S3F**).

Against this baseline, we describe three major findings. First, robust induction of IFNs and ISGs

by both TLR3 and TLR4 agonists was observed in liver slices from all three groups of liver

specimens (*i.e.*, non-infected, HCV-infected, and DAA-cured). *IFNB1* and *IFNL3* (IL28B)

peaked at 2-4 h, prior to the maximum abundance of ISGs (*IFIT 1/2/3, RSAD2* and *MX1*) which

occurred at 4-8 h, consistent with the sequential signaling cascade induced by TLR3 and TLR4

pathways.

Second, for chronic HCV-infected livers, PAMP sensing appeared robust and enhanced

compared to non-infected liver tissue. For example, antiviral genes *MX1, RSAD2, IFNG*,

chemokine genes *CCL3, CX3CL1, CXCL9*, inflammatory genes *TNF, CASP1* and *IL-10* were not

suppressed but induced more strongly by TLR3 agonist in liver slices of chronic HCV-infected

liver at least at one time point (Mann-Whitney test, two tailed, P < 0.05) (**Figs 5 and 6**).   Other

common activation markers for TLR3 signaling, including *IFNB1, IL28B, IFIT 1/2/3, ISG15,*

CXCL10, IL-1B and *CCL-5*, were also responsive in chronic HCV-infected slices with elevation

of greater than 10-fold, as strong as changes in the non-infected liver slices.   Thus, TLR3

sensing and signaling did not appear to be impaired in chronic HCV-infected liver slices.

Notably, *TNF, IL-10* and *RSAD2* remained abnormally induced at 12 h and 24 h in chronic HCV

liver slices, indicating persistent inflammation.

TLR4 sensing and signaling were similarly robust in chronic HCV-infected livers. *IFIT1, IFIT3,*

RSAD2, MX1, CCL3, CASP1 and *IL-10* were again identified along with *IL-1A, CCL7* and

TGFB1 at greater abundance post stimulation in chronic HCV-infected liver slices on at least one

time point (Mann-Whitney test, two tailed, $P < 0.05$) (**Figs 5 and 6**). *IFIT1, IFIT3, IL-1A,*

CCL3, IL-10, CASP1, and *TGFB1* were more strongly elevated at 12 h and 24 h in chronic HCV

liver slices, again revealing that innate immune signaling was not disabled in these liver tissue.

Third, most of the hyper-induction of the genes in HCV-infected tissue (7/8 genes in TLR3

response, 10/12 genes in TLR4 response) was not reversed in the tissue from formerly HCV+,

but now successfully treated donors (**Figs 5 and 6**). *IFNG, CX3CL1*, and *IL-10* induction by the

TLR3 agonist, and *TNF, MX1* and *CCL5* induction by the TLR4 agonist exacerbated differences

at 12 h and 24 h in tissue from the DAA-treated livers, consistent with the sustained

inflammation in these tissues (**Fig. S7**). However, *TGFB1, CCL3* and *IL-18* were examples of

genes whose expression was restored to normal by DAA treatment (**Fig. 6B**). The shared

abnormality in TLR3 and TLR4 signaling pathway in HCV-infected and DAA-treated liver

tissue is in line with epigenetic changes in TNF and IFNG signaling and inflammatory response

identified for such liver tissue [17]. Importantly, since NPCs were the main cellular

compartment expressing TLR3 and TLR4 (**Fig. S4G**) as well as the majority of the immune

genes, the hyper-activation signals during TLR3 and TLR4 response were likely contributed by

the abnormal NPC. Analysis of fold-change (delta-delta Ct) also identified the greater induction

of inflammatory-related genes (*TNF, IL-1B, IL-8, IL-10, CX3CL1* and *CTGF*) in chronic HCV-

infected or DAA-cured liver tissues during poly-I:C or LPS responses (**Fig. S8**).

Discussion

Here we showed that immune abnormalities persist after the elimination of HCV by anti-viral

therapy, and the persisting inflammatory and immune signatures mapped to hepatic non-

parenchymal cells. An improved method for culture of human liver slices [19] enabled us to

further investigate the immunopathology of liver slices *ex vivo*. Innate immunity was not

suppressed but enhanced in HCV-infected tissue, and these abnormalities were not corrected

after effective DAA treatment.

In human hepatocyte cultures [34, 35] HCV infection triggered cleavage of MAVS and TRIF

host proteins by HCV NS3/4A protease, disrupting TLR3-antiviral signaling. Reduction of

protein levels of MAVS and TRIF was also observed previously in chronic HCV-infected livers

[36, 37]. Nevertheless, we observed a robust response to TLR3 signaling in the HCV-infected

liver, which supports the idea that ISG induction is weak in HCV infected cells, but strong in

HCV infected liver due to paracrine activation of NPC. The data on the direct effect of HCV on

the TLR3 response in liver tissue is novel and is distinct from previous work focused on the IFN-

induced gene expression in chronically HCV-infected liver by pegylated IFN therapy [38, 39],

since such direct activation of IFN signaling bypasses MAVS and TRIF.

In DAA-cured, formerly HCV+ liver slices, a number of antiviral genes were hyper-activated by

poly-I:C and LPS. The gene induction pattern, coupled with our ability to assign these genes to

hepatic non-parenchymal cells, indicates excessive immune activation in NPC, with risk of both

increased inflammation, and impaired immunity [40-42]. Histological analysis of DAA-cured,

formerly HCV+ liver biopsy samples also reported sustained inflammation and cirrhosis despite

the clearance of HCV [18] (and this study). Such ongoing liver inflammation that was not cured

by DAA treatment, could result from the properties of liver-resident immune cells, including NK

cells and Kupffer cells, which were pre-activated during HCV infection; or from the abnormal

fibrotic liver environment in DAA-treated livers.

The immune abnormalities observed in HCV-infected liver tissue were distinct from those

identified in a recent study of TLR immune response with chronic HBV-infected liver biopsy

tissues [43]. Here we used several approaches to improve the TLR ligand treatment method to

enable enhanced detection of differences from non-infected, control liver tissue. However, the

logistics of our approach made it impractical to study a wide variety of TLR ligands, so we

concentrated on TLR3 and TLR4 firstly to interrogate a major pathway against RNA viruses, and

second to engage both the TRIF and the MyD88 signaling pathways. We would not expect the

defects induced by HBV and HCV to be identical, but some common features emerge: other

studies with chronic HBV-infected hepatic tissue have reported immune interference by HBV

[44].

From the days of IFN-based therapy, we know that HCV-induced liver disease can regress when

the virus is eradicated from the liver [45]. However, halting or reversing liver disease upon IFN

cure of HCV has not been universal; some patients with advanced disease still go on to develop

HCC [46-48]. DAA drugs are now able to cure the majority of HCV infections, even in subjects

with advanced liver disease. Nevertheless, clearance of HCV with DAA treatment only partially

restores immune cell function. In this study, we have identified immune genes that were restored

to normal by DAA treatment, and more importantly immune genes that were not restored *ex vivo*

(*i.e.*, at day 0 time point), including a subset of immune activation genes, inflammation genes,

immune suppression, and PI3K pathway genes.

The era during which some, but not all HCV patients were treated with DAA was brief and has

passed. Within this narrow time window, we worked with tissue from patients who were treated

for surgical resection of a liver lesion. Accordingly, there are inherent limitations to this study.

First, due to the small sample size, we did not further classify patient subgroups based on HCV

genotypes. Likewise, other subject characteristics including gender, age, genetic polymorphisms

(such as IL28B), and liver fibrosis status which may play a role in immunity and responses to

HCV infection [13, 49, 50] were not used as factors to subdivide patient groups in the

downstream analysis. Surprisingly, we observed significant effects that cut across these hidden

variables. Among these, the identification of aberrant immune gene expression in NPC, and

alteration of signaling transduction in TLR3 and TLR4 pathways in chronic HCV-infected and

formerly HCV+ but now DAA-cured liver tissues are novel. Second, we cannot determine how

far the sustained abnormality is due to the host response to hepatic fibrosis or HCC rather than

HCV-mediated. Future study may extend the conclusion through analysis of a more diverse

group of clinical situations, including chronic HCV patients and DAAs-SVR cases with or

without fibrosis or HCC, as well as non-HCV related liver tissues with fibrosis or HCC.

Acknowledgements

We thank Pathology Research Services Laboratory, and Histology and Imaging Core (HIC) at

UW for their help in data and image acquisition.

Conflict of Interest

The authors have no conflict of interest to declare.

References

1.   Polaris Observatory HCVC. Global prevalence and genotype distribution of hepatitis C
virus infection in 2015: a modelling study. Lancet Gastroenterol Hepatol. 2017;2(3):161-76.
Epub 2017/04/14. doi: 10.1016/S2468-1253(16)30181-9. PubMed PMID: 28404132.

2.   Ringelhan M, McKeating JA, Protzer U. Viral hepatitis and liver cancer. Philos Trans R
Soc Lond B Biol Sci. 2017;372(1732). Epub 2017/09/13. doi: 10.1098/rstb.2016.0274. PubMed
PMID: 28893941; PubMed Central PMCID: PMCPMC5597741.

3.   Kim D, Li AA, Gadiparthi C, Khan MA, Cholankeril G, Glenn JS, et al. Changing Trends
in Etiology-Based Annual Mortality From Chronic Liver Disease, From 2007 Through 2016.
Gastroenterology. 2018;155(4):1154-63 e3. Epub 2018/07/17. doi: 10.1053/j.gastro.2018.07.008.
PubMed PMID: 30009816; PubMed Central PMCID: PMCPMC6467699.

4.   Zhang Q, Wang Y, Zhai N, Song H, Li H, Yang Y, et al. HCV core protein inhibits
polarization and activity of both M1 and M2 macrophages through the TLR2 signaling pathway.
Sci Rep. 2016;6:36160. Epub 2016/10/28. doi: 10.1038/srep36160. PubMed PMID: 27786268;
PubMed Central PMCID: PMCPMC5082373.

5.   Spaan M, van Oord G, Kreefft K, Hou J, Hansen BE, Janssen HL, et al. Immunological
Analysis During Interferon-Free Therapy for Chronic Hepatitis C Virus Infection Reveals
Modulation of the Natural Killer Cell Compartment. J Infect Dis. 2016;213(2):216-23. Epub
2015/08/01. doi: 10.1093/infdis/jiv391. PubMed PMID: 26223768.

6.   Aregay A, Sekyere SO, Deterding K, Port K, Dietz J, Berkowski C, et al. Elimination of
Hepatitis C Virus has limited impact on the functional and mitochondrial impairment of HCV-

specific CD8+ T cell responses. Journal of hepatology. 2019. Epub 2019/07/12. doi:
10.1016/j.jhep.2019.06.025. PubMed PMID: 31295532.

7.   Najafi Fard S, Schietroma I, Corano Scheri G, Giustini N, Serafino S, Cavallari EN, et al.
Direct-acting antiviral therapy enhances total CD4+ and CD8+ T-cells responses, but does not
alter T-cells activation among HCV mono-infected, and HCV/HIV-1 co-infected patients. Clin
Res Hepatol Gastroenterol. 2018;42(4):319-29. Epub 2017/12/28. doi:
10.1016/j.clinre.2017.11.006. PubMed PMID: 29279268.

8.   Telatin V, Nicoli F, Frasson C, Menegotto N, Barbaro F, Castelli E, et al. In Chronic
Hepatitis C Infection, Myeloid-Derived Suppressor Cell Accumulation and T Cell Dysfunctions
Revert Partially and Late After Successful Direct-Acting Antiviral Treatment. Front Cell Infect
Microbiol. 2019;9:190. Epub 2019/07/02. doi: 10.3389/fcimb.2019.00190. PubMed PMID:
31259160; PubMed Central PMCID: PMCPMC6588015.

9.   Vranjkovic A, Deonarine F, Kaka S, Angel JB, Cooper CL, Crawley AM. Direct-Acting
Antiviral Treatment of HCV Infection Does Not Resolve the Dysfunction of Circulating CD8(+)
T-Cells in Advanced Liver Disease. Front Immunol. 2019;10:1926. Epub 2019/08/29. doi:
10.3389/fimmu.2019.01926. PubMed PMID: 31456810; PubMed Central PMCID:
PMCPMC6700371.

10.   Ghosh A, Mondal RK, Romani S, Bagchi S, Cairo C, Pauza CD, et al. Persistent gamma
delta T-cell dysfunction in chronic HCV infection despite direct-acting antiviral therapy induced
cure. Journal of viral hepatitis. 2019;26(9):1105-16. Epub 2019/05/11. doi: 10.1111/jvh.13121.
PubMed PMID: 31074195.

11.   Hengst J, Strunz B, Deterding K, Ljunggren HG, Leeansyah E, Manns MP, et al.
Nonreversible MAIT cell-dysfunction in chronic hepatitis C virus infection despite successful
interferon-free therapy. Eur J Immunol. 2016;46(9):2204-10. Epub 2016/06/15. doi:
10.1002/eji.201646447. PubMed PMID: 27296288.

12.   Reig M, Marino Z, Perello C, Inarrairaegui M, Ribeiro A, Lens S, et al. Unexpected high
rate of early tumor recurrence in patients with HCV-related HCC undergoing interferon-free
therapy. Journal of hepatology. 2016;65(4):719-26. Epub 2016/04/17. doi:
10.1016/j.jhep.2016.04.008. PubMed PMID: 27084592.

13.   Conti F, Buonfiglioli F, Scuteri A, Crespi C, Bolondi L, Caraceni P, et al. Early
occurrence and recurrence of hepatocellular carcinoma in HCV-related cirrhosis treated with
direct-acting antivirals. Journal of hepatology. 2016;65(4):727-33. Epub 2016/06/29. doi:
10.1016/j.jhep.2016.06.015. PubMed PMID: 27349488.

14.   Kuftinec G, Loehfelm T, Corwin M, Durbin-Johnson B, Candido M, Hluhanich R, et al.
De novo hepatocellular carcinoma occurrence in hepatitis C cirrhotics treated with direct-acting
antiviral agents. Hepat Oncol. 2018;5(1):HEP06. Epub 2018/10/12. doi: 10.2217/hep-2018-0003.
PubMed PMID: 30302197; PubMed Central PMCID: PMCPMC6168046.

15.   Dong Ji ea. Increase in Incidence of Hepatocellular Carcinoma in Chronic Hepatitis C
Chinese After Sustained Virologic Response with Direct Acting Antiviral Agents as Compared

to Pegylated Interferon Plus Ribavirin Therapy: A Prospective Open-Label Study. Preprint with
THE LANCET. 2019;preprint.

16.   Gitto S, Cicero AFG, Loggi E, Giovannini M, Conti F, Grandini E, et al. Worsening of
Serum Lipid Profile after Direct Acting Antiviral Treatment. Ann Hepatol. 2018;17(1):64-75.
Epub 2018/01/10. doi: 10.5604/01.3001.0010.7536. PubMed PMID: 29311405.

17.   Hamdane N, Juhling F, Crouchet E, El Saghire H, Thumann C, Oudot MA, et al. HCV-
Induced Epigenetic Changes Associated With Liver Cancer Risk Persist After Sustained
Virologic Response. Gastroenterology. 2019;156(8):2313-29 e7. Epub 2019/03/06. doi:
10.1053/j.gastro.2019.02.038. PubMed PMID: 30836093.

18.   Whitcomb E, Choi WT, Jerome KR, Cook L, Landis C, Ahn J, et al. Biopsy Specimens
From Allograft Liver Contain Histologic Features of Hepatitis C Virus Infection After Virus
Eradication. Clin Gastroenterol Hepatol. 2017;15(8):1279-85. Epub 2017/05/16. doi:
10.1016/j.cgh.2017.04.041. PubMed PMID: 28501538.

19.   Wu X, Roberto JB, Knupp A, Kenerson HL, Truong CD, Yuen SY, et al. Precision-cut
human liver slice cultures as an immunological platform. Journal of immunological methods.
2018;455:71-9. Epub 2018/02/07. doi: 10.1016/j.jim.2018.01.012. PubMed PMID: 29408707;
PubMed Central PMCID: PMCPMC6689534.

20.   Elmasry S, Wadhwa S, Bang BR, Cook L, Chopra S, Kanel G, et al. Detection of Occult
Hepatitis C Virus Infection in Patients Who Achieved a Sustained Virologic Response to Direct-
Acting Antiviral Agents for Recurrent Infection After Liver Transplantation. Gastroenterology.
2017;152(3):550-3 e8. Epub 2016/11/14. doi: 10.1053/j.gastro.2016.11.002. PubMed PMID:
27838287; PubMed Central PMCID: PMCPMC5285320.

21.   Lagaye S, Shen H, Saunier B, Nascimbeni M, Gaston J, Bourdoncle P, et al. Efficient
replication of primary or culture hepatitis C virus isolates in human liver slices: A relevant ex
vivo model of liver infection. Hepatology. 2012;56(3):861-72. doi: 10.1002/hep.25738.

22.   Dolmazashvili E, Abutidze A, Chkhartishvili N, Karchava M, Sharvadze L, Tsertsvadze
T. Regression of liver fibrosis over a 24-week period after completing direct-acting antiviral
therapy in patients with chronic hepatitis C receiving care within the national hepatitis C
elimination program in Georgia: results of hepatology clinic HEPA experience. European journal
of gastroenterology & hepatology. 2017;29(11):1223-30. doi:
10.1097/MEG.0000000000000964. PubMed PMID: 28857900.

23.   Hsu WF, Lai HC, Su WP, Lin CH, Chuang PH, Chen SH, et al. Rapid decline of
noninvasive fibrosis index values in patients with hepatitis C receiving treatment with direct-
acting antiviral agents. BMC gastroenterology. 2019;19(1):63. doi: 10.1186/s12876-019-0973-5.
PubMed PMID: 31029101; PubMed Central PMCID: PMC6486982.

24.   Mittal S, El-Serag HB, Sada YH, Kanwal F, Duan Z, Temple S, et al. Hepatocellular
Carcinoma in the Absence of Cirrhosis in United States Veterans is Associated With
Nonalcoholic Fatty Liver Disease. Clin Gastroenterol Hepatol. 2016;14(1):124-31 e1. Epub

2015/07/22. doi: 10.1016/j.cgh.2015.07.019. PubMed PMID: 26196445; PubMed Central
PMCID: PMCPMC4690789.

25.   Negash AA, Ramos HJ, Crochet N, Lau DT, Doehle B, Papic N, et al. IL-1beta
production through the NLRP3 inflammasome by hepatic macrophages links hepatitis C virus
infection with liver inflammation and disease. PLoS Pathog. 2013;9(4):e1003330. Epub
2013/05/02. doi: 10.1371/journal.ppat.1003330. PubMed PMID: 23633957; PubMed Central
PMCID: PMCPMC3635973.

26.   Lau DT, Fish PM, Sinha M, Owen DM, Lemon SM, Gale M, Jr. Interferon regulatory
factor-3 activation, hepatic interferon-stimulated gene expression, and immune cell infiltration in
hepatitis C virus patients. Hepatology. 2008;47(3):799-809. Epub 2008/01/19. doi:
10.1002/hep.22076. PubMed PMID: 18203148.

27.   Brownell J, Wagoner J, Lovelace ES, Thirstrup D, Mohar I, Smith W, et al. Independent,
parallel pathways to CXCL10 induction in HCV-infected hepatocytes. Journal of hepatology.
2013;59(4):701-8. doi: 10.1016/j.jhep.2013.06.001. PubMed PMID: 23770038; PubMed Central
PMCID: PMC3779522.

28.   Brownell J, Bruckner J, Wagoner J, Thomas E, Loo YM, Gale M, Jr., et al. Direct,
interferon-independent activation of the CXCL10 promoter by NF-kappaB and interferon
regulatory factor 3 during hepatitis C virus infection. J Virol. 2014;88(3):1582-90. doi:
10.1128/JVI.02007-13. PubMed PMID: 24257594; PubMed Central PMCID:
PMCPMC3911583.

29.   Meissner EG, Wu D, Osinusi A, Bon D, Virtaneva K, Sturdevant D, et al. Endogenous
intrahepatic IFNs and association with IFN-free HCV treatment outcome. The Journal of clinical
investigation. 2014;124(8):3352-63. Epub 2014/07/02. doi: 10.1172/JCI75938. PubMed PMID:
24983321; PubMed Central PMCID: PMCPMC4109554.

30.   Wang L, Feng Y, Xie X, Wu H, Su XN, Qi J, et al. Neuropilin-1 aggravates liver
cirrhosis by promoting angiogenesis via VEGFR2-dependent PI3K/Akt pathway in hepatic
sinusoidal endothelial cells. EBioMedicine. 2019;43:525-36. doi: 10.1016/j.ebiom.2019.04.050.
PubMed PMID: 31060904; PubMed Central PMCID: PMC6558257.

31.   Cai CX, Buddha H, Castelino-Prabhu S, Zhang Z, Britton RS, Bacon BR, et al.
Activation of Insulin-PI3K/Akt-p70S6K Pathway in Hepatic Stellate Cells Contributes to
Fibrosis in Nonalcoholic Steatohepatitis. Digestive diseases and sciences. 2017;62(4):968-78.
doi: 10.1007/s10620-017-4470-9. PubMed PMID: 28194671.

32.   Dreux M, Garaigorta U, Boyd B, Decembre E, Chung J, Whitten-Bauer C, et al. Short-
range exosomal transfer of viral RNA from infected cells to plasmacytoid dendritic cells triggers
innate immunity. Cell Host Microbe. 2012;12(4):558-70. Epub 2012/10/23. doi:
10.1016/j.chom.2012.08.010. PubMed PMID: 23084922; PubMed Central PMCID:
PMCPMC3479672.

33.   Wieland S, Makowska Z, Campana B, Calabrese D, Dill MT, Chung J, et al.
Simultaneous detection of hepatitis C virus and interferon stimulated gene expression in infected

human liver. Hepatology. 2014;59(6):2121-30. Epub 2013/10/15. doi: 10.1002/hep.26770.
PubMed PMID: 24122862; PubMed Central PMCID: PMCPMC3975814.

34.   Loo YM, Owen DM, Li K, Erickson AK, Johnson CL, Fish PM, et al. Viral and
therapeutic control of IFN-beta promoter stimulator 1 during hepatitis C virus infection. Proc
Natl Acad Sci U S A. 2006;103(15):6001-6. Epub 2006/04/06. doi: 10.1073/pnas.0601523103.
PubMed PMID: 16585524; PubMed Central PMCID: PMCPMC1458687.

35.   Li K, Foy E, Ferreon JC, Nakamura M, Ferreon AC, Ikeda M, et al. Immune evasion by
hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein
TRIF. Proc Natl Acad Sci U S A. 2005;102(8):2992-7. Epub 2005/02/16. doi:
10.1073/pnas.0408824102. PubMed PMID: 15710891; PubMed Central PMCID:
PMCPMC548795.

36.   Bellecave P, Sarasin-Filipowicz M, Donze O, Kennel A, Gouttenoire J, Meylan E, et al.
Cleavage of mitochondrial antiviral signaling protein in the liver of patients with chronic
hepatitis C correlates with a reduced activation of the endogenous interferon system. Hepatology.
2010;51(4):1127-36. Epub 2010/01/02. doi: 10.1002/hep.23426. PubMed PMID: 20044805.

37.   Kar P, Kumar D, Gumma PK, Chowdhury SJ, Karra VK. Down regulation of TRIF,
TLR3, and MAVS in HCV infected liver correlates with the outcome of infection. J Med Virol.
2017;89(12):2165-72. Epub 2017/05/10. doi: 10.1002/jmv.24849. PubMed PMID: 28480979.

38.   Sarasin-Filipowicz M, Oakeley EJ, Duong FH, Christen V, Terracciano L, Filipowicz W,
et al. Interferon signaling and treatment outcome in chronic hepatitis C. Proc Natl Acad Sci U S
A. 2008;105(19):7034-9. doi: 10.1073/pnas.0707882105. PubMed PMID: 18467494; PubMed
Central PMCID: PMC2383932.

39.   McGilvray I, Feld JJ, Chen L, Pattullo V, Guindi M, Fischer S, et al. Hepatic cell-type
specific gene expression better predicts HCV treatment outcome than IL28B genotype.
Gastroenterology. 2012;142(5):1122-31 e1. doi: 10.1053/j.gastro.2012.01.028. PubMed PMID:
22285807.

40.   Zhang X, Starnbach MN. An Excess of the Proinflammatory Cytokines IFN-gamma and
IL-12 Impairs the Development of the Memory CD8+ T Cell Response to Chlamydia
trachomatis. Journal of immunology. 2015;195(4):1665-75. Epub 2015/07/17. doi:
10.4049/jimmunol.1500457. PubMed PMID: 26179901; PubMed Central PMCID:
PMCPMC4530050.

41.   Benci JL, Xu B, Qiu Y, Wu TJ, Dada H, Twyman-Saint Victor C, et al. Tumor Interferon
Signaling Regulates a Multigenic Resistance Program to Immune Checkpoint Blockade. Cell.
2016;167(6):1540-54 e12. Epub 2016/12/03. doi: 10.1016/j.cell.2016.11.022. PubMed PMID:
27912061; PubMed Central PMCID: PMCPMC5385895.

42.   McAlpine W, Sun L, Wang KW, Liu A, Jain R, San Miguel M, et al. Excessive
endosomal TLR signaling causes inflammatory disease in mice with defective SMCR8-WDR41-
C9ORF72 complex function. Proc Natl Acad Sci U S A. 2018;115(49):E11523-E31. Epub

2018/11/18. doi: 10.1073/pnas.1814753115. PubMed PMID: 30442666; PubMed Central
PMCID: PMCPMC6298088.

43.   Suslov A, Boldanova T, Wang X, Wieland S, Heim MH. Hepatitis B Virus Does Not
Interfere With Innate Immune Responses in the Human Liver. Gastroenterology.
2018;154(6):1778-90. Epub 2018/02/07. doi: 10.1053/j.gastro.2018.01.034. PubMed PMID:
29408639.

44.   Li TY, Yang Y, Zhou G, Tu ZK. Immune suppression in chronic hepatitis B infection
associated liver disease: A review. World J Gastroenterol. 2019;25(27):3527-37. Epub
2019/08/02. doi: 10.3748/wjg.v25.i27.3527. PubMed PMID: 31367154; PubMed Central
PMCID: PMCPMC6658392.

45.   D'Ambrosio R, Colombo M. Should surveillance for liver cancer be modified in hepatitis
C patients after treatment-related cirrhosis regression? Liver international : official journal of the
International Association for the Study of the Liver. 2016;36(6):783-90. Epub 2016/03/05. doi:
10.1111/liv.13106. PubMed PMID: 26936383.

46.   Makiyama A, Itoh Y, Kasahara A, Imai Y, Kawata S, Yoshioka K, et al. Characteristics
of patients with chronic hepatitis C who develop hepatocellular carcinoma after a sustained
response to interferon therapy. Cancer. 2004;101(7):1616-22. Epub 2004/09/21. doi:
10.1002/cncr.20537. PubMed PMID: 15378504.

47.   van der Meer AJ, Feld JJ, Hofer H, Almasio PL, Calvaruso V, Fernandez-Rodriguez CM,
et al. Risk of cirrhosis-related complications in patients with advanced fibrosis following
hepatitis C virus eradication. Journal of hepatology. 2017;66(3):485-93. Epub 2016/10/27. doi:
10.1016/j.jhep.2016.10.017. PubMed PMID: 27780714.

48.   van der Meer AJ, Veldt BJ, Feld JJ, Wedemeyer H, Dufour JF, Lammert F, et al.
Association between sustained virological response and all-cause mortality among patients with
chronic hepatitis C and advanced hepatic fibrosis. JAMA. 2012;308(24):2584-93. Epub
2012/12/27. doi: 10.1001/jama.2012.144878. PubMed PMID: 23268517.

49.   Schott E, Witt H, Hinrichsen H, Neumann K, Weich V, Bergk A, et al. Gender-dependent
association of CTLA4 polymorphisms with resolution of hepatitis C virus infection. Journal of
hepatology. 2007;46(3):372-80. Epub 2006/12/08. doi: 10.1016/j.jhep.2006.09.011. PubMed
PMID: 17150279.

50.   Yan Z, Wang Y. Viral and host factors associated with outcomes of hepatitis C virus
infection (Review). Mol Med Rep. 2017;15(5):2909-24. Epub 2017/03/25. doi:
10.3892/mmr.2017.6351. PubMed PMID: 28339063.

Figure legends

Figure 1. Experimental Overview. (A) Liver slices were collected from resected livers from three groups of patients including non-HCV-infected patients (controls), chronic HCV-infected patients, and patients with previous history of HCV infection who were cured by DAA treatment (DAA-cured patients). Day 0 liver slices were harvested immediately after slicing, which revealed the *ex vivo* immunopathology of liver cells. Day 7 baseline differences were compared with liver slices collected after seven days of *ex vivo* culture and before the addition of TLR agonist. Innate immune responses were analyzed with day 7 liver slices which were stimulated with poly-I:C (TLR3 agonist) or LPS (TLR4 agonist) with time points of 2, 4, 8, 12, 24 hours of treatment. (B) Specific liver cell types were purified from fresh resected liver wedges from non-HCV-infected individuals with perfusion, differential centrifugation and fluorescence-activated cell sorting (FACS) techniques. Immune gene expression was analyzed with liver cell samples enriched with hepatocytes, total non-parenchymal cells (NPC), Kupffer cells (KC), liver sinusoidal endothelial cells (LSEC), or hepatic stellate cells (HSC).

Figure 2. Analysis of HCV RNA and liver fibrosis with liver slices. (A) All eight chronic HCV samples (blue color) were confirmed with positive HCV RNA detection. None of the DAA cured subjects (red color) or the non-infected controls (black color) was positive for HCV RNA. n.d., meaning non-detected. The limit of reliable detection (LoD) was 1.2 IU/ 50 ng liver total RNA. (B) HCV RNA remained robustly detected in the day 7 liver slices cultured from chronically HCV-infected patients. The viral load between day 0 and day 7 liver slices was not statistically significant (P=0.4688, Wilcoxon matched-pairs signed rank test). (C) Fibrosis analysis of the day 0 *ex vivo* liver specimens with trichrome staining and picrosirius red staining indicated that in DAA-treated, and now HCV-negative patients there was persistent fibrosis, similar to untreated HCV and different from tissue without a history of HCV

infection. The scoring system was based on the Scheuer/Batts-Ludwig method. The box-and-whisker plot

included data of 10 non-infected patients, 4 chronic HCV patients, and 8 DAA-treated patients, with the

minimum and maximum data points for each subgroup being shown. Statistical significance was based

on Mann-Whitney test. **, P < 0.01. ††, P < 0.01. For additional details, see Fig S1.

Figure 3. Differential expression of immune genes at the day 0 baseline in chronic HCV-infected

and DAA-cured patients versus non-infected controls. (A) Day 0 differentially expressed genes in

chronic HCV-infected versus non-infected patients were highlighted with red markers (increased genes in

HCV) and green markers (decreased genes in HCV).   Gene subsets persistent to DAA treatment were

highlighted. In comparison, HLA-DR, MX1 and CXCL10 were examples of genes that were restored to

normal by DAA therapy at the day 0 baseline. (B) Day 0 differential expressed immune genes in DAA-

cured versus non-infected liver slices. (C) Venn diagram showing commonality of immune gene changes

in different patient groups. Comparison of D0 baseline for DAA-cured versus HCV-infected livers is

provided in Fig S2.

Figure 4. Immune gene changes at the day 7 baseline in chronic HCV-infected and DAA-cured

livers. (A) Hierarchical clustering of immune genes with statistically significant elevation in HCV or

DAA-cured subjects at Day 0 or Day 7 time points. Genes were colored according to their group-level

fold changes, which were calculated with the mean Ct value of each gene within each patient group.

Individual gene plots are provided in Fig. S3. (B) Summary diagram for genes that were statistically

significant at day 7 time point. Genes were categorized based on their statistical significance in chronic

HCV-infected and DAA-treated samples. The cell-type-specific expression was mapped to their cell

types analyzed with the day 0 perfused liver wedge samples. Many of the immune genes were enriched

in expression in the NPC compartment (Fig. S4 and S5). Genes that were also statistically significantly

evaluated at the day 0 time point are indicated with bold fonts.

Figure 5. Poly-I:C and LPS induced TLR3 and TLR4 response with chronic HCV-infected liver

slices and DAA-cured liver slices. The relative RNA abundance was normalized to ACTB, GAPDH and

HPRT1. Statistical significance was based on two-tailed Mann-Whitney test. *****, statistical

significantly different between non-infected versus chronic HCV liver slices. **†**, statistical

significantly different between non-infected versus DAA-cured liver slices. **‡**, statistical

significantly different between chronic HCV versus DAA-cured liver slices. Levels of statistical

significance was *, P <0.05 **, P<0.01, ***, P<0.001. TNF, IFNG, MX1 and IL-10 are

examples of genes that were hyper-activated in TLR3 and TLR4 response in chronic HCV-

infected liver slices and in DAA-cured liver slices.

Figure 6. Altered TLR3 and TLR4 response in chronic HCV-infected liver slices not

restored in DAA-cured liver slices. (A) Hierarchical clustering of immune responsive genes

with 2-fold or greater induction during LPS stimulation. Genes and time points were colored

based on the statistical significance P value of HCV-infected versus control, and DAA-cured

versus control according to two-tailed Mann-Whitney test. The poly-I:C induced heat map is in

Fig. S6. The gene clusters that were altered by chronic HCV infection and not restored to

normal by DAA therapy were highlighted.   Data of individual genes are shown in Fig. S7. (B)

Summary of statistically significant genes (P < 0.05) that were altered by HCV infection and not

restored by DAA treatment with the mapping the enriched liver cell types.

Table

Table 1 Immune gene changes *ex vivo* (Day 0)

	Fold change			Statistical significance[2]			Cell-type-specific information[3]
	[1]HCV/ Con	DAA/ Con	DAA/ HCV	HCV/ Con	DAA/ Con	DAA/ HCV	
Antiviral genes							
Cured by DAA at D0 baseline[4]							
CXCL10	54.39	2.93	0.05	0.0003	0.4598	0.0004	NPC, KC
RSAD2	7.68	1.05	0.14	0.0007	0.7197	0.0007	NPC
IFIT1	7.16	1.51	0.21	0.0007	0.211	0.002	NPC
IFIT3	7.02	1.61	0.23	0.0002	0.0653	0.0004	NPC
CXCL9	6.80	1.45	0.21	0.0016	0.7197	0.0017	NPC, HSC
IFIT2	5.39	1.82	0.34	0.0007	0.0535	0.033	NPC, HSC
ISG15	6.16	1.04	0.17	0.0115	0.8421	0.0012	likely NPC, HSC
MX1	7.05	1.01	0.14	0.0004	0.9682	0.0002	likely NPC
Immune activation genes							
NOT cured by DAA at D0 baseline[5]							
HLA-A	3.84	1.61	0.42	0.0002	0.022	0.0012	NPC, LSEC
CCL5	3.30	1.92	0.58	0.0033	0.022	0.0878	NPC
CCR2	9.93	4.45	0.45	0.0007	0.0104	0.02	likely NPC
4-1BB (CD137, TNFRSF9)	6.35	3.17	0.50	0.0007	0.0279	0.0702	likely NPC
CIITA	5.03	4.64	0.92	0.0007	0.0003	0.6691	likely KC
Cured by DAA at D0 baseline							
HLA-DRA	2.53	1.43	0.56	0.0052	0.2775	0.1088	NPC
CD80	3.54	1.91	0.54	0.0012	0.1823	0.0553	NPC
CD86	2.32	1.25	0.54	0.0418	0.447	0.1613	NPC
OX40 (TNFRSF4)	3.93	2.46	0.63	0.0033	0.0789	0.3638	LSEC
TNFRSF18	5.62	2.97	0.53	0.0006	0.0745	0.351	HSC, LSEC
TNFSF9	2.97	2.08	0.70	0.036	0.2224	0.2721	likely KC
CXCL1	2.73	1.01	0.37	0.1447	0.9048	0.003	HSC
IFNG	2.69	0.83	0.31	0.1738	>0.9999	0.0431	NPC
Inflammation genes							
NOT cured by DAA at D0 baseline							
CASP1	3.13	2.71	0.87	0.0007	0.0435	0.0136	NPC
IL-12A	3.71	1.91	0.51	0.0037	0.0434	0.0185	LSEC

Cured by DAA at D0 baseline							
IL-12B	2.86	1.29	0.45	0.0262	0.7577	0.1738	HSC
TNF	2.77	1.06	0.38	0.0115	0.7802	0.0046	NPC, KC, HSC
IL-18	2.40	0.90	0.37	0.0164	0.6607	0.025	NPC, HSC
NFKB	1.46	1.01	0.69	0.0283	>0.9999	0.0317	N/A
Immune Suppression genes							
NOT cured by DAA at D0 baseline							
CTLA4	6.52	2.70	0.41	0.0003	0.0172	0.033	likely NPC
TIGIT	4.77	3.20	0.67	0.0003	0.0172	0.033	likely NPC
IDO1	4.35	2.52	0.58	0.0052	0.0435	0.1613	NPC
ARG1	0.45	0.39	0.89	0.0229	0.0056	0.6806	Hep, uncommon in KC, LSEC
Cured by DAA at D0 baseline							
B7-H4	4.49	3.70	0.83	0.0059	0.0541	0.7104	HSC
VISTA (VSIR)	2.10	1.40	0.67	0.0021	0.1333	0.3638	KC, LSEC, HSC
CD47	1.43	1.24	0.87	0.0418	0.3562	0.3148	NPC, KC, LSEC, HSC
PDCD1	5.16	2.30	0.44	0.0205	0.0592	0.1416	not cell-type-specific
LAG3	1.88	1.19	0.64	0.0164	>0.9999	0.0712	not cell-type-specific
TIM3	2.18	0.99	0.46	0.0079	0.8633	0.0164	NPC
FOXP3	2.94	0.70	0.24	0.1416	0.8633	0.0079	not cell-type-specific
PI3K pathway genes							
NOT cured by DAA at D0 baseline							
SPHK1	3.50	2.13	0.61	0.0052	0.04	0.2523	HSC
BTG2	2.19	2.10	0.96	0.0907	0.0188	0.8371	N/A
SOX9	2.62	2.30	0.88	0.0549	0.04	>0.9999	Not cell-type-specific
SNAP25	3.03	1.88	0.62	0.0164	0.0503	0.351	N/A
GPRC5B	2.30	1.91	0.83	0.0311	0.1615	0.351	N/A
Tissue repair genes							
Cured by DAA at D0 baseline							
TIMP1	1.98	1.29	0.65	0.0311	0.447	0.2698	LSEC
MMP9	2.04	1.13	0.56	0.0164	0.7802	0.1331	Not cell-type-specific
PDGFB	1.62	0.93	0.58	0.2523	0.8633	0.0164	LSEC
TGFB1	2.18	1.27	0.58	0.0712	0.447	0.0431	KC, LSEC, HSC

Liver tissue was perfused, and total livers, hepatocytes (Hep) and NPC populations (NPC) were obtained with differential centrifugation. Kupffer cells (KC), liver sinusoidal endothelial cells (LSEC), and hepatic stellate cells (HSC) enriched populations were obtained with FACS from NPC (Fig. 1B). RNA was extracted from each cell type, followed by qRT-PCR gene expression analysis using the fluidigm platform.

[1]The comparisons were HCV/ Con, chronic HCV-infected versus non-infected controls; DAA/ Con, DAA-cured versus non-infected controls; DAA/HCV, DAA-cured versus chronic HCV-infected.

[2]Statistical significance was based on non-parametric two-tailed Mann-Whitney test.
[3]Cell-type-specific information was based on analysis of the non-HCV-infected samples. N/A means cell-type-specific gene information not analyzed. This was due to the limit of sample availability and the assay through-put with fluidigm assays.

[4]Cured by DAA at D0 baseline categorized the gene subsets that were normalized to the same level as non-infected controls.

[5]Not cured by DAA at D0 baseline categorized gene subsets that were not normalized to the correct levels as the non-infected controls.

Figure 1

A

Resected liver cores

(**Non-infected** vs. **Chronic HCV-infected** vs. **DAA-cured**)

B

Figure 2

A

B

C

Figure 3

A **Non-infected vs. Chronic HCV**

Non-infected livers (log2 scale) vs Chronic HCV livers (log2 scale)

Up-regulated in chronic HCV livers

Labeled genes: ARG1, HLA-A, HLA-DRA, MX1, CASP1, CCL5, CXCL10, CCR2, CIITA, IDO1, SPHK1, TIGIT, CTLA4, 4-1BB, IL12A

C **Day 0 genes**

Non-infected vs. HCV | Non-infected vs. DAA-cured
28 | 12 | 2

Non-infected vs. HCV | HCV vs. DAA-cured
22 | 18 | 5

B **Non-infected vs. DAA-cured**

Non-infected livers (log2 scale) vs DAA-cured livers (log2 scale)

Up-regulated in DAA-cured livers

Labeled genes: ARG1, HLA-A, CASP1, CCL5, BTG2, SOX9, CCR2, CIITA, IDO1, SPHK1, TIGIT, CTLA4, 4-1BB, IL12A

Figure 4

Figure 5

Figure 6