

Title Page

COVID-19 Vaccine Acceptance Among Health Care Workers in the United States

Rahul Shekhar, MD¹; Abu Baker Sheikh, MD¹; Shubhra Upadhyay, MD¹; Mriganka Singh, MD³;

Saket Kottewar, MD⁴; Hamza Mir², Eileen Barrett, MD¹; Suman Pal, MD¹

1 - Department of Internal Medicine, University of New Mexico Health Sciences Center,
Albuquerque, New Mexico USA

2 – Data Analyst, University of New Mexico Health Sciences Center, Albuquerque, New Mexico
US

3 - Department of Geriatrics and Palliative Care, University Hospitals of School of Medicine Case
Western University, Cleveland, Ohio USA

4- Department of Hospital Medicine. University of Texas Health San Antonio, San Antonio,
Texas USA

Corresponding author:

Abu Baker Sheikh

absheikh@salud.unm.edu,

2211 Lomas BLVD, NE,

Albuquerque, NM 87106

Phone – 6316334125

Funding: This is a non-funded study

IRB approval: University of New Mexico Health Sciences Center

Key Words: Covid-19, vaccine, healthcare workers, United States

Conflicts of interest: All authors do not report any conflict of interest

ABSTRACT:

Title: COVID-19 Vaccine Acceptance Among Health Care Workers in the United States

Background:

Acceptance of any COVID-19 vaccine is an important challenge to address and will play a major role in combating the pandemic. Healthcare workers (HCWs) are amongst the first group to receive vaccination, so it is important to consider their attitudes about COVID-19 vaccination to better address barriers to widespread vaccination acceptance.

Methods:

We conducted a cross sectional study to assess the attitude of HCWs toward COVID-19 vaccination. Data was collected between October 7th and November 9th, 2020. We received 4080 responses out of which 3479 were complete responses and were included in final analysis.

Results:

36% of respondents were willing to take the vaccine as soon as it became available while 56% were not sure or would wait to review more data. Vaccine acceptance increased with increasing age, education, and income level. Lower acceptance was noted in females (31%), Black (10%), Hispanic (30%) and conservative/republican (21%) HCWs, and those working in a rural setting (26%). In the multinomial regression model, direct medical care providers and those with heart disease and cancer had lower likelihood of waiting to review than accepting vaccines. Safety (69%), effectiveness (69%) and rapidity of development/approval (74%) were noted as most common concerns regarding COVID-19 vaccine in our survey.

Conclusion:

Acceptance of COVID-19 vaccine is low, with the majority of healthcare workers choosing to wait to review more data before deciding. Overall attitude toward vaccination was positive but specific concerns regarding COVID-19 vaccine are prevalent. Differences in vaccine acceptance were noted along lines of social determinants of health which should be addressed to avoid magnifying health disparities.

Key Words:

Covid-19, vaccine, healthcare workers, United States

Introduction:

COVID-19 has rapidly become a major global public health crisis, affecting 79.7 million individuals, and causing 1.7 million deaths. The USA has reported more than 18.5 million cases and 320,000 deaths. To curb this pandemic, apart from effective public health measures such as social distancing, wearing face masks, hand washing and avoidance of crowded indoor spaces, educating the general population, efficacious vaccination is emerging as essential to mitigating disease and death.²⁻⁵

Uptake of any COVID-19 vaccine is an important challenge to address. In a recent survey, more than one-third of lay respondents were unsure or did not intend to take the vaccine⁷. Clinicians are an important source of information for vaccines and physician communication can improve adherence to vaccination recommendations^{8,9,10}. Thus, the role of healthcare workers becomes particularly important in advising patients and communities, and as well as through role modeling behavior. Healthcare workers are being prioritized among the high-risk groups who will be considered as initial candidates for early vaccination. As such, it is important to consider healthcare worker attitudes about COVID-19 vaccination to better address barriers to widespread vaccination.

Methods:

Design: We conducted a cross sectional study to assess the attitude of healthcare workers (HCWs) toward COVID-19 vaccination. An online questionnaire was created using REDCap electronic data capture tools hosted at University of New Mexico. The survey was modified from the general population survey by Fisher et al⁷ to capture more information pertinent to healthcare workers. The data was collected anonymously, and no personal identifying information was collected. This study was approved by University of New Mexico Hospitals (UNMH) Institutional review board.

Sampling: A snowball sampling was utilized. The survey tool was distributed via links posted on social media platforms in various HCW groups and distributed to administrative leaders at five major hospital systems to disseminate among their employees. Data was collected between October 7th and November 9th, 2020. We received 4080 responses out of which 3479 were complete responses and were included in final analysis.

Participants: All adults (>18 years of age) working in a healthcare setting regardless of patient care contact and role in health care settings were eligible to participate in the study. Informed consent was obtained prior to enrollment in the study.

Measures:

Demographic information collected included age, gender, ethnicity, race, state of primary residence in majority of last six months, occupation, marital status, the number of household members excluding participant, annual household income, location of healthcare setting (rural, suburban or urban), education level and political orientation (conservative/ republican, unaffiliated, Democrat/Liberal, or do not wish to answer).

Self-perceived risk of COVID-19 was gauged by the question “Do you think you are at risk of getting COVID-19 in the next 1 year?” The responses allowed for graded self-perceived risk (No I am confident I won't get infected, Yes I am concerned that I will get mild symptoms which will probably not require hospitalization, Yes I am concerned that I will get moderate symptoms which will probably need hospitalization, Yes I am concerned that I will get severe symptom which will probably require admission to the Intensive care unit, I believe I already have the disease and I am immune to it (Not diagnosed by a test), No, I already have recovered and won't get re-infected (Diagnosed by a test)).

Exposure to COVID-19 was assessed by questions “Have you directly or indirectly taken care of the COVID-19 patients?” and “Have you, your family member or someone you know been diagnosed with COVID-19 (Excluding your patients)?”

Acceptance of COVID-19 vaccine was assessed by question - “When COVID-19 vaccination becomes available would you take it?” Participant could choose response from among the options - Yes, as soon as I can get it; Yes, only if it is required by employer; No, I will wait for 3 months to review safety profile; No, I will wait for 6 months to review safety profile; No, I will wait for at least 1 year to review safety profile; I will not get the vaccine; Not sure.

Attitude toward vaccination was assessed by agreement with perception/concern statements. General perception about vaccines were assessed by statements - “I do not believe vaccines work”; “I do not believe vaccines are safe”; “I do not get vaccinated for religious reasons”; “I do not get vaccinated for reasons of personal freedom/choice”; “I do not get vaccinated for a fear of needles/doctors/hospitals”. Concerns regarding COVID-19 vaccine were evaluated by statements - “I am worried about the safety / adverse effects of COVID-19 vaccine”; “I am worried about effectiveness of COVID-19 vaccine”; “I am worried about the out of pocket cost/Insurance coverage of the vaccine”; “I am concerned about adverse effect of vaccine on my pre-existing conditions”; “I am worried about the rapidity of the development and approval of COVID-19 vaccine”; “I do not need the vaccine for my risk level”; “I am worried about the rapidity of the development and approval of COVID-19 vaccine”. Agreement was measured on Likert scale from strongly disagree to strongly agree.

Outcome:

We have divided HCW into four major categories, Direct medical provider (DMP) which includes If the participant is a Physician/ Resident/Medical students and advanced practice provider including Nurse Practitioner (NP) and / Physician's assistant (PA), Direct patient care provider (DPCP) including Registered Nurse / Patient care tech/ Paramedics/ Rehab services - Respiratory, physical, occupational speech therapist, nutritionist, social workers, case managers care coordinators, administrative staff and others without direct patient care providers. If the participant is DMP their primary specialty was also recorded and is divided into primary medical, primary surgical diagnostic and others.

The primary outcome of the survey was whether HCW are willing to take the potential upcoming COVID-19 vaccine or not. Responses were collected against willingness to take it as soon as it becomes available, yes but only if it is required by employer (grouped into one as yes responses), wait for safety data review for 3 months, 6 months or a year or not sure (grouped into one response wait for review) and not willing to take it.

We performed a multinomial logistic regression due to the overall description of the data. For doing this we split the sample into 3 groups according to the outcome variable, would a participant take the COVID-19 vaccination or not or would wait to review safety data. Due to the variables being categorical and the model being a multinomial logistic regression we used the likelihood-ratio test leading to the derived chi-square and p-values for analysis. We used the R programming language (R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>) to develop the model and analyze its results. The multinomial regression evaluated the prediction of membership into the Covid-19 vaccination acceptance categories (Yes, No, Wait for Review). The reference group here was 'Yes' that the participant would take the vaccination as soon as possible. Statistical significance was assumed for $p < 0.05$.

Results:

3,479 HCWs completed survey. Most participants were younger than 40 years of age (1877, 54%), female (2598, 75%), White (2803, 81%), completed a bachelor's degree or higher (2788, 80%) identified as Democrat or liberal (1521, 44%) and had no chronic medical conditions (2039, 59%). Most reported working in an urban area (2229, 64%), in primary medical and medical subspecialties (54%) and provided direct patient care (79%). Majority of the participants perceived themselves to be at risk for acquiring COVID-19 (3043, 87%) and 21% think they will acquire serious disease requiring admission to hospital (747, 21%) but less than 80% were confident that they would not get the disease (267, 8%). Roughly Half of the HCW have directly taken care of COVID-19 positive patients (1570, 45%). Most of the participants believed that the COVID-19 vaccination should be voluntary (1665, 48%). Additional sample characteristics are shown in Table 1.

Only about one-third (1247, 36%) of respondents were willing to take COVID-19 vaccine as soon as it became available at the time of the survey. Majority of the HCW were not sure or would

wait to review safety data before getting vaccinated (1953, 56%). Among the respondents who want to wait, 11% will like to wait for 3 months, 10% will like to wait for six months and 20% will like to wait at least one year. Only 8% (279) of respondents were not willing to take the vaccination at all.

Chi square indicated a significant association was noted between the choice that participants make about receiving the COVID-19 vaccination and multiple predictor variables (Table 2). We note that acceptance for COVID-19 vaccine increased with increasing age. In the 18-30 age group only 34% of respondents were willing to take COVID-19 vaccine as soon as it became available which increased to 47% in the >70 age group. A similar trend was noted with education and income level. Education and income levels represent a higher proportion of HCWs willing to take the vaccine as soon as it becomes available. Differences in vaccine acceptance were noted by gender and racial identity. Female HCWs had lower vaccine acceptance at 31% compared to male HCWs (49%) and trans/non-binary HCWs (43%). Black HCWs had lower acceptance (19%) with majority choosing to wait to review (65%) whereas Asian HCWs had high vaccine acceptance (44%). A majority of Native American HCWs (80%) and all Native Hawaiian/Other pacific islander HCWs (100%) chose to wait to review data for COVID vaccine. Vaccine acceptance was lower among those identifying as Hispanic or Latino (30%). Geographical variation in vaccine acceptance was also noted in our survey with West having the lowest (33%) and South having the highest (48%) vaccine acceptance. HCWs employed in a rural setting had lower acceptance (26%) of the vaccine. Those identifying as democrat/liberal had higher vaccine acceptance (42%). HCWs who believe themselves to be immune to COVID-19 and those who feel confident they won't get infected had the highest rates of refusing COVID-19 vaccine at 22% and 27% respectively. HCWs who had not taken care of COVID-19 patients had higher rates of vaccine refusal (9.2%).

Acceptance of COVID-19 vaccine was also associated with a plan to recommend vaccination to friends and family, and with higher likelihood of wanting COVID-19 vaccine for HCWs to be mandatory. Of note, a large majority of our respondents who plan to not get COVID-19 vaccine would also not recommend the vaccine to friends or family and want vaccination to be voluntary.

The multinomial logistic regression evaluated the prediction of membership into the Covid-19 vaccination acceptance categories (Yes, No, Wait for Review). The reference group here was 'Yes' that the participant would take the vaccination as soon as possible. Analysis revealed that the model as a whole was good for predicting the categories of our outcome variable as the accuracy on the training and testing data predictions was above 80% and the p-value was less than 0.05 for the whole model. Regarding the relative risk comparison in our model regarding variables/categories we have found trends which would be important for future research. DMPs had 53% less likely to wait to review more data than accepting the vaccine as soon as it is made available as compared to HCWs without direct patient contact (RRR=0.47). Participants who had heart disease had 77% lower relative likelihood of waiting to review more data for the vaccine in comparison to taking the vaccine right now (RRR= 0.23). A participant who had cancer had 62% lower relative likelihood of responding "wait for review" versus "yes" to the

vaccination (RRR = 0.38). Participants who were diagnosed with COVID-19 themselves had nearly 3-fold (Have you, your family member or someone you know been diagnosed with COVID-19 (Excluding your patients) = I was Diagnosed = RRR = 3.4) higher relative likelihood of responding wait for review rather than yes for the vaccination for COVID-19. Participants who thought they had COVID-19 already and were immune to it had a 64% lower relative likelihood of responding “Wait for Review” versus “Yes”. (RRR = 0.36).

While overall concerns regarding vaccination were low, concerns regarding COVID-19 vaccines were prevalent (Figure 1/Supplementary Table 4). Most HCWs believe in general vaccination works (90%), are safe (86%) and did not mention personal (87%) or religious belief (95%) as a reason for not vaccinating. Most participants endorsed concerns (agree or strongly agree) about vaccine safety/adverse effects (69%), effectiveness (69%), and rapidity of development/approval (74%). Majority of HCWs trust their Doctors and Health care professionals recommending the COVID-19 vaccine (73%) but nearly half of the respondents do not trust information provided by the government about COVID-19 and its severity (46%) and one-third do not trust regulatory authorities like CDC or FDA overseeing the vaccine development and safety (34%).

Discussion:

Since the announcement of efforts to develop a COVID-19 vaccine, several surveys have been conducted to gauge public perception and acceptance of the vaccine through 2020. Most of the surveys have focused on the general population. However, the rollout of the vaccination is to be tiered to various subgroups of the population based on limited availability in the initial phases. HCWs are likely to be among the first subgroups of the population to have access to the vaccine. HCWs are also likely to be an important source of information about the vaccine for the general population. As such, it is crucial to assess predictors of vaccine uptake among HCWs which will help institutions and policy makers target resources to maximize uptake. To the best of our knowledge, this is the largest survey of HCWs in the USA about COVID-19 vaccination.

In our survey, only one-third of respondents were amenable to COVID-19 vaccination immediately, while more than half of respondents preferred to defer their decision until reviewing more data. This contrasts with the findings of a general population survey in April 2020⁷ that reported half the participants were willing to take the vaccine. Another study in May 2020 reported 67% acceptance of COVID-19 vaccine among US adults¹¹. The high percentage of respondents waiting to review more data in our study is expected as HCWs are more likely to base healthcare decisions on published scientific literature of efficacy and safety which was largely lacking during the time of our study. This also highlights the importance of publication and dissemination of scientific data regarding the vaccine which will be a crucial factor to determine eventual uptake of the vaccine among HCWs. Overall, only 8% respondents said they would refuse the COVID-19 vaccine which shows a potential for high uptake of the vaccine

among HCWs. Increasing vaccination acceptance has substantial benefits. To eventually slow down the spread of COVID-19 and its mortality it is imperative to achieve herd immunity by vaccination before immunity by natural infection. With R_0 of 3, the threshold herd immunity will be achieved by immunizing at least 70 % of the population assuming the vaccine is 100 % effective⁽¹⁷⁾. With recent EUA of 2 vaccines in the USA claiming to be almost 95 % effective in phase III clinical trials, this number could be even higher. Thus, it is imperative to vaccinate a maximum number of HCWs to prevent infection among HCWs and loss of critical workforce when there is already a critical shortage. Studies have shown that vaccination of HCWs with influenza vaccine decreases mortality and absenteeism¹⁹⁻²². It would be reasonable to expect a similar trend with COVID-19 vaccine. It is important to note that 97% of HCWs in our survey had received influenza vaccine in the previous year indicating a generally favorable perception of vaccination.

The low initial acceptance of COVID-19 vaccine among HCWs could also have broader consequences. Studies have shown that HCWs who are vaccinated are more likely to recommend vaccines to friends, family, and their patients¹²⁻¹⁴. This has also been borne out in our study where we see a strong association among HCWs who plan to be vaccinated and plan to recommend the vaccine to friends and family.

COVID-19 vaccine acceptance increased with increasing age, income, and education level. Black race also had lower vaccine acceptance while Asian race had higher vaccine acceptance. Vaccine acceptance was also lower in HCWs identifying as Hispanic or Latino. This mirrors the trends seen with general population surveys^{7,11}. The toll of COVID-19 pandemic has disproportionately affected low-income communities and BIPOC⁽¹⁸⁾. Lower vaccine uptake could magnify the health disparities among these communities. Targeting messaging and outreach would be required to achieve higher vaccination rates. It will also be important to understand the factors driving low vaccine uptake which would be instrumental in addressing key concerns and directing resources to increase uptake.

HCWs identifying as conservative or Republican in their political leanings had lower acceptance of the vaccine whereas those identifying as liberal or Democrat had higher vaccine acceptance. This is in line with other general US population survey^{7,15,16}. This difference could be in part due to differential messaging regarding vaccines in news and social media targeting these populations and in part due to very divergent responses to the pandemic by political leaders of major parties²³. A unified and consistent messaging from political leaders will be essential to bridge this gap.

Vaccine acceptance was higher among HCWs involved in direct patient care and in HCWs with chronic medical conditions. This could mirror trends in perceived risk of COVID-19 infection. More HCWs who perceived themselves to be at risk for COVID-19 infection were willing to accept than refuse the vaccine. However, even among HCWs directly involved in patient care, vaccine acceptance was also lower among HCWs identified as DPCP than among DMP. This is concerning since DPCP (such as RNs, RTs, etc) often have more direct and prolonged patient contact. They are, therefore, at high risk of acquiring the disease. DPCP are also one of the key

resources of the healthcare system where critical shortages have been noted during the pandemic. They therefore represent a key subgroup whose health is essential to continue our response to the pandemic.

To understand the factors driving vaccine uptake, we assessed HCWs attitude toward vaccination and toward COVID-19 vaccine. Concerns regarding vaccination in general were low in our study, this is in keeping with other studies that show generally positive attitudes of healthcare workers toward vaccination. However, concerns specific to COVID-19 vaccination were prevalent. We noted a high prevalence of concerns regarding vaccine efficacy, adverse effects, and rapidity of development. This was particularly noted among HCWs who do not plan to take the COVID-19 vaccine. Of note, while HCWs who did not want to be vaccinated reported poor trust in regulatory authorities and government, the trust in medical professionals prescribing the vaccine was somewhat higher. This could suggest an important role for dissemination of information through medical agencies and professional societies to increase uptake among HCWs.

A major strength of our study is the large sample of HCWs surveyed. Our survey population is also diverse with representation from different gender, age groups, ethnic and racial backgrounds, and roles in healthcare.

We recognize limitations of our study. Due to sampling methods, our study population may not be representative of all US HCWs. Social desirability bias may also affect the interpretation of our study results, though the responses were anonymized to minimize this. Most importantly, our study was conducted when information regarding COVID-19 vaccines under development were limited and findings of the clinical trials had not been made public. As such, it is possible that with this information now being publicly available, both vaccine acceptance and attitude toward COVID-19 vaccines have changed.

References

1. COVID-19 Map - Johns Hopkins Coronavirus Resource Center. Johns Hopkins Coronavirus Resource Center. <https://coronavirus.jhu.edu/map.html>. Published 2020. Accessed December 24, 2020.
2. Adaptive Phase IB-II Randomized Clinical Trial Of Preventive Vaccine Consisting Of Autologous Dendritic Cells Loaded With Antigens From Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), With Or Without GM-CSF, In Subjects Negative For COVID-19 Infection And Anti-SARS-CoV-2 Antibodies. *ClinicalTrials.gov identifier (NCT number): NCT04386252*. <https://clinicaltrials.gov/ct2/show/NCT04386252>
3. A Randomized, Double-blind, Placebo-controlled Phase 3 Study to Assess the Efficacy and Safety of Ad26.COV2.S for the Prevention of SARS-CoV-2-mediated COVID-19 in Adults Aged 18 Years and Older. *ClinicalTrials.gov Identifier: NCT04505722*. <https://clinicaltrials.gov/ct2/show/NCT04505722>
4. A Phase 2a, Randomized, Observer-Blind, Placebo Controlled, Dose-Confirmation Study to Evaluate the Safety, Reactogenicity, and Immunogenicity of mRNA-1273 SARS-COV-2 Vaccine in Adults Aged 18 Years and Older. *ClinicalTrials.gov Identifier: NCT04405076*. <https://clinicaltrials.gov/ct2/show/NCT04405076>
5. A Phase 1/Phase 2, Randomized, Double-Blind, Placebo-Controlled, Dose-Ranging Trial to Evaluate the Safety, Tolerability and Immunogenicity of V591 (COVID-19 Vaccine) in Healthy Younger and Older Participants. *ClinicalTrials.gov Identifier: NCT04498247*. <https://www.clinicaltrials.gov/ct2/show/NCT04498247>
6. Phadke VK, Bednarczyk RA, Salmon DA, Omer SB. Association Between Vaccine Refusal and Vaccine-Preventable Diseases in the United States: A Review of Measles and Pertussis [published correction appears in *JAMA*. 2016 May 17;315(19):2125] [published correction appears in *JAMA*. 2016 May 17;315 (19):2125]. *JAMA*. 2016;315(11):1149-1158. doi:10.1001/jama.2016.1353
7. Fisher KA, Bloomstone SJ, Walder J, Crawford S, Fouayzi H, Mazor KM. Attitudes Toward a Potential SARS-CoV-2 Vaccine: A Survey of U.S. Adults [published online ahead of print, 2020 Sep 4]. *Ann Intern Med*. 2020;M20-3569. doi:10.7326/M20-3569
8. Wheeler M, Bутtenheim AM. Parental vaccine concerns, information source, and choice of alternative immunization schedules. *Hum Vaccin Immunother*. 2013;9(8):1782-1789. doi:10.4161/hv.25959
9. Salmon DA, Moulton LH, Omer SB, DeHart MP, Stokley S, Halsey NA. Factors associated with refusal of childhood vaccines among parents of school-aged children: a case-control study. *Arch Pediatr Adolesc Med*. 2005;159(5):470-476. doi:10.1001/archpedi.159.5.470
10. Omer SB, Salmon DA, Orenstein WA, deHart MP, Halsey N. Vaccine refusal, mandatory immunization, and the risks of vaccine-preventable diseases. *N Engl J Med*. 2009;360(19):1981-1988. doi:10.1056/NEJMsa0806477
11. Malik, A. A., Mcfadden, S. M., Elharake, J., & Omer, S. B. (2020). Determinants of COVID-19 vaccine acceptance in the US. *EClinicalMedicine*, 26, 100495. doi:10.1016/j.eclinm.2020.100495
12. Paterson P, Meurice F, Stanberry LR, Glismann S, Rosenthal SL, Larson HJ. Vaccine hesitancy and healthcare providers. *Vaccine*. 2016 Dec 20;34(52):6700–6.
13. Zhang J, While AE, Norman IJ. Nurses' knowledge and risk perception towards seasonal influenza and vaccination and their vaccination behaviours: A cross-sectional survey. *Int J Nurs Stud*. 2011 Oct 1;48(10):1281–9.
14. LaVela SL, Smith B, Weaver FM, Legro MW, Goldstein B, Nichol K. Attitudes and Practices Regarding Influenza Vaccination Among Healthcare Workers Providing Services to Individuals

- With Spinal Cord Injuries and Disorders. *Infect Control Hosp Epidemiol* [Internet]. 2004 Nov [cited 2020 Oct 28];25(11):933–40. Available from: </core/journals/infection-control-and-hospital-epidemiology/article/attitudes-and-practices-regarding-influenza-vaccination-among-healthcare-workers-providing-services-to-individuals-with-spinal-cord-injuries-and-disorders/C672BE66C1D0649B3E0FA423C040AC91>
15. Reiter PL, Pennell ML, Katz ML. Acceptability of a COVID-19 vaccine among adults in the United States: How many people would get vaccinated? *Vaccine*. 2020 Sep 29;38(42):6500–7.
 16. Pogue K, Jensen JL, Stancil CK, Ferguson DG, Hughes SJ, Mello EJ, et al. Influences on Attitudes Regarding Potential COVID-19 Vaccination in the United States. *Vaccines* [Internet]. 2020 Oct 3 [cited 2020 Dec 15];8(4):582. Available from: <https://www.mdpi.com/2076-393X/8/4/582>
 17. Fontanet A, Cauchemez S. COVID-19 herd immunity: where are we? [Internet]. Vol. 20, *Nature Reviews Immunology*. Nature Research; 2020 [cited 2020 Dec 18]. p. 583–4. Available from: <https://www.defense.gouv.fr/content/download/583466/9938746/>
 18. Webb Hooper M, Nápoles AM, Pérez-Stable EJ. COVID-19 and Racial/Ethnic Disparities [Internet]. Vol. 323, *JAMA - Journal of the American Medical Association*. American Medical Association; 2020 [cited 2020 Dec 24]. p. 2466–7. Available from: <https://www.chicago.gov/city/en/sites/covid->
 19. Lemaitre M, Meret T, Rothan-Tondeur M, Belmin J, Lejonc JL, Luquel L, et al. Effect of influenza vaccination of nursing home staff on mortality of residents: A cluster-randomized trial. *J Am Geriatr Soc* [Internet]. 2009 Sep [cited 2020 Dec 24];57(9):1580–6. Available from: <https://pubmed-ncbi-nlm-nih-gov.libproxy.unm.edu/19682118/>
 20. Zaffina S, Gilardi F, Rizzo C, Sannino S, Brugaletta R, Santoro A, et al. Seasonal influenza vaccination and absenteeism in health-care workers in two subsequent influenza seasons (2016/17 and 2017/18) in an Italian pediatric hospital. *Expert Rev Vaccines* [Internet]. 2019 Apr 3 [cited 2020 Dec 24];18(4):411–8. Available from: <https://www-tandfonline-com.libproxy.unm.edu/doi/abs/10.1080/14760584.2019.1586541>
 21. Murti M, Otterstatter M, Orth A, Balshaw R, Halani K, Brown PD, et al. Measuring the impact of influenza vaccination on healthcare worker absenteeism in the context of a province-wide mandatory vaccinate-or-mask policy. *Vaccine*. 2019 Jul 9;37(30):4001–7.
 22. Imai C, Toizumi M, Hall L, Lambert S, Halton K, Merollini K. A systematic review and meta-analysis of the direct epidemiological and economic effects of seasonal influenza vaccination on healthcare workers. Ortiz JR, editor. *PLoS One* [Internet]. 2018 Jun 7 [cited 2020 Dec 24];13(6):e0198685. Available from: <https://dx.plos.org/10.1371/journal.pone.0198685>
 23. Romer D, Jamieson KH. Conspiracy theories as barriers to controlling the spread of COVID-19 in the U.S. *Soc Sci Med*. 2020 Oct 1;263:113356.




FIGURE 1: Attitude of Healthcare Workers toward Vaccines

Table 1: Participant characteristics

Variable	N = 3,479 ^a
Age	
18-30 years	816 (23%)
31-40 years	1,061 (30%)
41-50 years	686 (20%)
51-60 years	571 (16%)
61-70 years	326 (9.4%)
> 70 years	19 (0.5%)
Gender	
Female	2,598 (75%)
Male	864 (25%)
Trans/ Gender non-binary/ not specified above	7 (0.2%)
Do not wish to reply	10 (0.3%)
Ethnicity	
Hispanic or Latino	560 (16%)
NOT Hispanic or Latino	2,763 (79%)
Unknown / Not Reported	36 (1.0%)
Do not wish to answer	120 (3.4%)
Race	
White or Caucasian	2,803 (81%)
Asian	218 (6.3%)
Black or African American	74 (2.1%)
Native Americans/Alaska Native	30 (0.9%)

Native Hawaiian or Other Pacific Islander	6 (0.2%)
More Than One Race	126 (3.6%)
Unknown / Other	70 (2.0%)
Do not wish to answer	152 (4.4%)
State of Residence	
Midwest	1,433 (41%)
North East	81 (2.3%)
South	314 (9.0%)
West	1,651 (47%)
Occupation	
Direct Patient Care Providers (DPCP)	1,573 (45%)
Direct Medical Providers (DMP)	1,207 (35%)
Administrative staff working in hospital without direct patient contact	295 (8.5%)
Others without direct patient contact	404 (12%)
Primary Area of Work	
Primary medical and medical subspecialty	1,882 (54%)
Primary surgical and surgical subspecialty	363 (10%)
Diagnostic subspecialty	246 (7.1%)
Others	988 (28%)
Annual Income	
< \$30,000	117 (3.4%)
\$30,001 - \$70,000	741 (21%)
\$70,001 - \$100,00	759 (22%)
\$100,001 - \$150,000	796 (23%)
>\$150,001	1,066 (31%)
Health Care facility in	

Rural area	293 (8.4%)
Suburban area	957 (28%)
Urban area	2,229 (64%)
Education	
No formal education	1 (<0.1%)
High school graduate, diploma or the equivalent (for example: GED)	46 (1.3%)
Some college credit, no degree	169 (4.9%)
Trade/technical/vocational training	111 (3.2%)
Associate degree	364 (10%)
Bachelor's degree	1,046 (30%)
Master's degree	606 (17%)
Professional degree	297 (8.5%)
Doctorate degree	839 (24%)
Political Identification	
Conservative-Republican	746 (21%)
Democrat-Liberal	1,519 (44%)
Unaffiliated	640 (18%)
Do not wish to answer	574 (16%)
Medical comorbidities	
No Chronic Condition	2,039 (59%)
Diabetes Mellitus Type-1/Type-2	136 (3.9%)
Heart Disease	36 (1.0%)
Hypertension	449 (13%)
COPD Asthma - Lung Disease	318 (9.1%)
Obesity BMI>30	461 (13%)

Cancer	125 (3.6%)
Immuno-compromised/on immunosuppressants	110 (3.2%)
Smoking	288 (8.3%)
Other Medical Conditions	505 (15%)
Have you, your family member or someone you know been diagnosed with COVID-19 (Excluding your patients)	
I was Diagnosed with COVID-19	90 (2.6%)
Family Member was Diagnosed with COVID-19	447 (13%)
Someone Personal was Diagnosed with COVID-19	1,793 (52%)
No one Personal was Diagnosed with COVID-19	1,400 (40%)
Do you think you are at risk of getting COVID-19 in the next 1 year?	
I believe I already have the disease and I am immune to it (Not diagnosed by a test)	138 (4.0%)
No, I am confident I won't get infected	266 (7.6%)
No, I already have recovered and won't get re-infected (Diagnosed by a test)	34 (1.0%)
Yes, I am concerned that I will get mild symptoms which will probably not require hospitalization	2,294 (66%)
Yes, I am concerned that I will get moderate symptoms which will probably need hospitalization	572 (16%)
Yes, I am concerned that I will get severe symptom which will probably require admission to the Intensive care unit	175 (5.0%)
Have you directly or indirectly taken care of the COVID-19 patients?	
No	1,263 (36%)
Yes, but no direct patient contact	646 (19%)
Yes, I have direct patient contact	1,570 (45%)
Would you take the COVID-19 Vaccine	
No	279 (8.0%)
Wait for Review	1,953 (56%)

Yes	1,247 (36%)
Would you advise friends and family to get vaccinated for COVID-19?	
No	519 (15%)
Not sure	1,376 (40%)
Yes	1,584 (46%)
COVID-19 Vaccine for health care workers should be:	
Mandated by the employer, like Influenza vaccine	792 (23%)
Mandated by the Federal government for all health care workers	338 (9.7%)
Mandated by the State government for all health care workers	99 (2.8%)
Not sure	585 (17%)
Voluntary	1,665 (48%)
Statistics presented: n (%)	
Direct Patient Care Provider: Registered Nurse / Patient care technician/ Paramedics/ Rehab services – Respiratory therapist/Physical therapist/Occupation therapist/Speech/ Nutritionist/ social workers/ care coordinators	
Direct Medical Provider: Physician/ Resident/Medical students and Advanced practice provider (Nurse Practitioner / Physician's assistant	
Diagnostic Subspecialty: Radiology / Cath/ Endoscopy/ Pulmonary/ Echo/ Sleep Laboratory Technicians	

Table 2: Participant characteristics by vaccine acceptance

Variable	No, N = 279 ^a	Wait for Review, N = 1,953 ^a	Yes, N = 1,247 ^a	p-value ^b
Age				<0.001*
18-30 years	72 (8.8%)	467 (57%)	277 (34%)	
31-40 years	83 (7.8%)	615 (58%)	363 (34%)	
41-50 years	70 (10%)	389 (57%)	227 (33%)	
51-60 years	43 (7.5%)	306 (54%)	222 (39%)	
61-70 years	10 (3.1%)	167 (51%)	149 (46%)	
> 70 years	1 (5.3%)	9 (47%)	9 (47%)	
Gender				<0.001*
Female	240 (9.2%)	1,540 (59%)	818 (31%)	
Male	37 (4.3%)	402 (47%)	425 (49%)	
Trans/ Gender non-binary/ not specified above	0 (0%)	4 (57%)	3 (43%)	
Do not wish to reply	2 (20%)	7 (70%)	1 (10%)	
Ethnicity				<0.001*
Hispanic or Latino	55 (9.8%)	337 (60%)	168 (30%)	
NOT Hispanic or Latino	191 (6.9%)	1,536 (56%)	1,036 (37%)	
Unknown / Not Reported	8 (22%)	16 (44%)	12 (33%)	
Do not wish to answer	25 (21%)	64 (53%)	31 (26%)	
Race				<0.001*
White or Caucasian	221 (7.9%)	1,545 (55%)	1,037 (37%)	
Asian	1 (0.5%)	122 (56%)	95 (44%)	

Black or African American	12 (16%)	48 (65%)	14 (19%)	
Native Americans/Alaska Native	3 (10%)	24 (80%)	3 (10%)	
Native Hawaiian or Other Pacific Islander	0 (0%)	6 (100%)	0 (0%)	
More Than One Race	4 (3.2%)	82 (65%)	40 (32%)	
Unknown / Other	9 (13%)	41 (59%)	20 (29%)	
Do not wish to answer	29 (19%)	85 (56%)	38 (25%)	
State of Residence				0.003*
Midwest	146 (10%)	776 (54%)	511 (36%)	
North East	2 (2.5%)	45 (56%)	34 (42%)	
South	11 (3.5%)	151 (48%)	152 (48%)	
West	120 (7.3%)	981 (59%)	550 (33%)	
Occupation				<0.001*
Direct patient care providers (DPCP)	187 (12%)	969 (62%)	417 (27%)	
Direct Medical Provider (DMP)	30 (2.5%)	582 (48%)	595 (49%)	
Administrative staff working in hospital without direct patient contact	25 (8.5%)	170 (58%)	100 (34%)	
Others without direct patient contact	37 (9.2%)	232 (57%)	135 (33%)	
Annual Income				<0.001*
< \$30,000	19 (16%)	53 (45%)	45 (38%)	
\$30,001 - \$70,000	81 (11%)	420 (57%)	240 (32%)	
\$70,001 - \$100,00	80 (11%)	463 (61%)	216 (28%)	
\$100,001 - \$150,000	60 (7.5%)	472 (59%)	264 (33%)	
>\$150,001	39 (3.7%)	545 (51%)	482 (45%)	

Health Care facility in				<0.001*
Rural area	52 (18%)	164 (56%)	77 (26%)	
Suburban area	98 (10%)	523 (55%)	336 (35%)	
Urban area	129 (5.8%)	1,266 (57%)	834 (37%)	
Education				<0.001*
No formal education	0 (0%)	1 (100%)	0 (0%)	
High school graduate, diploma, or the equivalent (for example: GED)	9 (20%)	26 (57%)	11 (24%)	
Some college credit, no degree	24 (14%)	94 (56%)	51 (30%)	
Trade/technical/vocational training	18 (16%)	69 (62%)	24 (22%)	
Associate degree	56 (15%)	220 (60%)	88 (24%)	
Bachelor's degree	109 (10%)	627 (60%)	310 (30%)	
Master's degree	40 (6.6%)	362 (60%)	204 (34%)	
Professional degree	6 (2.0%)	139 (47%)	152 (51%)	
Doctorate degree	17 (2.0%)	415 (49%)	407 (49%)	
Political Identification				<0.001*
Conservative-Republican	93 (12%)	390 (52%)	263 (35%)	
Democrat-Liberal	42 (2.8%)	833 (55%)	644 (42%)	
Unaffiliated	48 (7.5%)	373 (58%)	219 (34%)	
Do not wish to answer	96 (17%)	357 (62%)	121 (21%)	
Did you get the Influenza vaccine Last Year?				<0.001*
No	50 (43%)	56 (48%)	10 (8.6%)	
Yes	229 (6.8%)	1,897 (56%)	1,237 (37%)	
If you have Children under 18 years old, Have they been vaccinated for other diseases				<0.001*
No	38 (26%)	67 (46%)	41 (28%)	
Not Applicable	120 (6.0%)	1,132 (56%)	754 (38%)	

Yes	121 (9.1%)	754 (57%)	452 (34%)	
Medical Condition = Chronic Condition				0.046
No	107 (7.4%)	819 (57%)	514 (36%)	
Yes	172 (8.4%)	1,134 (56%)	733 (36%)	
Diabetes Mellitus Type-1/Type-2				0.076
No	272 (8.1%)	1,876 (56%)	1,195 (36%)	
Yes	7 (5.1%)	77 (57%)	52 (38%)	
Heart Disease				0.067
No	276 (8.0%)	1,935 (56%)	1,232 (36%)	
Yes	3 (8.3%)	18 (50%)	15 (42%)	
Hypertension				0.8
No	253 (8.3%)	1,701 (56%)	1,076 (36%)	
Yes	26 (5.8%)	252 (56%)	171 (38%)	
COPD Asthma - Lung Disease				0.701
No	258 (8.2%)	1,778 (56%)	1,125 (36%)	
Yes	21 (6.6%)	175 (55%)	122 (38%)	
Obesity BMI>30				0.022*
No	250 (8.3%)	1,671 (55%)	1,097 (36%)	
Yes	29 (6.3%)	282 (61%)	150 (33%)	
Cancer				0.399
No	274 (8.2%)	1,882 (56%)	1,198 (36%)	
Yes	5 (4.0%)	71 (57%)	49 (39%)	
Immuno-compromised/on immunosuppressants				0.354
No	269 (8.0%)	1,889 (56%)	1,211 (36%)	
Yes	10 (9.1%)	64 (58%)	36 (33%)	
Smoking				0.013*

No	262 (8.2%)	1,778 (56%)	1,151 (36%)	
Yes	17 (5.9%)	175 (61%)	96 (33%)	
Other Medical Conditions				0.629
No	227 (7.6%)	1,663 (56%)	1,084 (36%)	
Yes	52 (10%)	290 (57%)	163 (32%)	
I was Diagnosed with COVID-19				0.009*
No	272 (8.0%)	1,901 (56%)	1,216 (36%)	
Yes	7 (7.8%)	52 (58%)	31 (34%)	
Family Member was Diagnosed with COVID-19				0.695
No	244 (8.0%)	1,712 (56%)	1,076 (35%)	
Yes	35 (7.8%)	241 (54%)	171 (38%)	
Someone Personal was Diagnosed with COVID-19				0.003*
No	168 (10.0%)	924 (55%)	594 (35%)	
Yes	111 (6.2%)	1,029 (57%)	653 (36%)	
No one Personal was Diagnosed with COVID-19				0.143
No	134 (6.4%)	1,181 (57%)	764 (37%)	
Yes	145 (10%)	772 (55%)	483 (34%)	
Do you think you are at risk of getting COVID-19 in next 1 year?				<0.001*
I believe I already have the disease and I am immune to it (Not diagnosed by a test)	30 (22%)	68 (49%)	40 (29%)	
No, I am confident I won't get infected	71 (27%)	128 (48%)	67 (25%)	
No, I already have recovered and won't get re-infected (Diagnosed by a test)	4 (12%)	18 (53%)	12 (35%)	

Yes, I am concerned that I will get mild symptoms which will probably not require hospitalization	153 (6.7%)	1,283 (56%)	858 (37%)	
Yes, I am concerned that I will get moderate symptoms which will probably need hospitalization	15 (2.6%)	350 (61%)	207 (36%)	
Yes, I am concerned that I will get severe symptom which will probably require admission to the Intensive care unit	6 (3.4%)	106 (61%)	63 (36%)	
Have you directly or indirectly taken care of the COVID-19 patients?				<0.001*
No	116 (9.2%)	714 (57%)	433 (34%)	
Yes, but no direct patient contact	45 (7.0%)	358 (55%)	243 (38%)	
Yes, I have direct patient contact	118 (7.5%)	881 (56%)	571 (36%)	
I would get the vaccine to prevent COVID-19 in myself				<0.001*
No	277 (20%)	816 (58%)	324 (23%)	
Yes	2 (<0.1%)	1,137 (55%)	923 (45%)	
I would get the vaccine to prevent COVID-19 in friends and family members				<0.001*
No	276 (21%)	723 (56%)	286 (22%)	
Yes	3 (0.1%)	1,230 (56%)	961 (44%)	
I would get the vaccine to prevent COVID-19 in community				<0.001*
No	277 (26%)	560 (52%)	237 (22%)	
Yes	2 (<0.1%)	1,393 (58%)	1,010 (42%)	
I would not get the vaccine				<0.001*
No	3 (<0.1%)	1,787 (59%)	1,235 (41%)	
Yes	276 (61%)	166 (37%)	12 (2.6%)	
Would you advise friends and family to get vaccinated for COVID-19?				<0.001*
No	230 (44%)	272 (52%)	17 (3.3%)	

Not sure	49 (3.6%)	1,162 (84%)	165 (12%)	
Yes	0 (0%)	519 (33%)	1,065 (67%)	
COVID-19 Vaccine for health care workers should be:				<0.001*
Mandated by the employer, like Influenza vaccine	0 (0%)	245 (31%)	547 (69%)	
Mandated by the Federal government for all health care workers	0 (0%)	90 (27%)	248 (73%)	
Mandated by the State government for all health care workers	0 (0%)	33 (33%)	66 (67%)	
Not sure	5 (0.9%)	457 (78%)	123 (21%)	
Voluntary	274 (16%)	1,128 (68%)	263 (16%)	
*Statistics presented: n (%)				
*Statistical tests performed: Likelihood-Ratio Test, Chi-Square Test				
Direct Patient Care Provider: Registered Nurse / Patient care technician/ Paramedics/ Rehab services – Respiratory therapist/Physical therapist/Occupation therapist/Speech/ Nutritionist/ social workers/ care coordinators				
Direct Medical Provider: Physician/ Resident/Medical students and Advanced practice provider (Nurse Practitioner / Physician’s assistant)				
Diagnostic Subspecialty: Radiology / Cath/ Endoscopy/ Pulmonary/ Echo/ Sleep Laboratory Technicians				

Table 3: Relative Risk Ratio multinomial regression model

Variable	RRR = Relative Risk Ratio			
	Opinion on Vaccination: No Vs. Yes		Opinion on Vaccination: Wait for Review Vs. Yes	
	RRR ¹	p-value ²	RRR ¹	p-value ²
Age				
18-30 years	0.000	0.563	0.595	0.565
31-40 years	0.000	0.689	0.355	0.244
41-50 years	0.000	0.670	0.295	0.170
51-60 years	0.000	0.295	0.303	0.171
61-70 years	0.000	0.586	0.223	0.087
> 70 years	Reference	Reference	Reference	Reference
Gender				
Female	51.297	0.687	0.793	0.155
Trans/ Gender non-binary/ not specified above	0.000	0.000	0.000	0.000
Do not wish to reply	28.238	0.844	3.18E+06	0.379
Male	Reference	Reference	Reference	Reference
Ethnicity				
Hispanic or Latino	1.42E+03	0.610	1.404	0.119
Unknown / Not Reported	0.147	0.946	1.141	0.878
Do not wish to answer	1.80E+07	0.388	1.834	0.332
NOT Hispanic or Latino	Reference	Reference	Reference	Reference
Race				
Asian	0.000	0.638	1.217	0.500
Black or African American	0.002	0.754	0.755	0.621
Native Americans/Alaska Native	0.000	0.000	0.587	0.524
Native Hawaiian or Other Pacific Islander	0.024	0.000	7.88E+13	0.000
More Than One Race	0.000	0.652	1.207	0.583
Unknown / Other	0.000	0.333	0.748	0.600
Do not wish to answer	1.80E+07	0.388	1.834	0.332
White or Caucasian	Reference	Reference	Reference	Reference
State of Residence				
North East	1.92E+05	0.633	1.742	0.185
South	0.000	0.542	0.853	0.544
West	0.418	0.916	1.183	0.308
Midwest	Reference	Reference	Reference	Reference
Occupation				
Direct Patient Care Provider	48.841	0.733	0.596	0.053
Direct Medical Provider	260.209	0.708	0.472	0.015*
Administrative staff working in hospital without direct patient contact	2.380	0.954	0.845	0.599

Others without direct patient contact	Reference	Reference	Reference	Reference
Annual Income				
< \$30,000	2.50E+04	0.602	0.609	0.294
\$30,001 - \$70,000	83.987	0.620	1.063	0.789
\$70,001 - \$100,00	6.509	0.790	0.876	0.535
>\$150,001	4.808	0.869	1.419	0.079
\$100,001 - \$150,000	Reference	Reference	Reference	Reference
Health Care facility in				
Rural area	2.432	0.932	0.975	0.925
Suburban area	9.203	0.737	0.785	0.152
Urban area	Reference	Reference	Reference	Reference
Education				
Some college credit, no degree	0.000	0.428	0.752	0.762
Trade/technical/vocational training	0.000	0.344	0.340	0.264
Associate degree	0.000	0.259	0.666	0.658
Bachelor's degree	0.000	0.428	0.283	0.159
Master's degree	0.000	0.353	0.362	0.261
Professional degree	0.000	0.461	0.308	0.205
Doctorate degree	0.000	0.270	0.298	0.188
High school graduate, diploma or the equivalent (for example: GED)	Reference	Reference	Reference	Reference
Political Identification				
Democrat-Liberal	7.233	0.868	0.821	0.325
Unaffiliated	23.926	0.700	0.885	0.591
Do not wish to answer	0.295	0.891	1.253	0.372
Conservative-Republican	Reference	Reference	Reference	Reference
Did you get the Influenza vaccine Last Year?	0.000	0.290	0.295	0.091
If you have Children under 18 years old, Have they been vaccinated for other diseases				
Yes	0.688	0.975	3.198	0.012*
Not Applicable	0.022	0.781	1.957	0.129
No	Reference	Reference	Reference	Reference
Medical co-morbidities				
No Chronic Condition	0.007	0.633	0.821	0.411
Diabetes Mellitus Type-1/Type-2	0.000	0.710	1.177	0.656
Heart Disease	3.07E+09	0.603	0.233	0.047*
Hypertension	0.514	0.950	0.929	0.764
COPD Asthma - Lung Disease	486.848	0.677	0.947	0.849
Obesity BMI>30	0.008	0.691	0.706	0.138
Cancer	0.000	0.697	0.384	0.018*
Immuno-compromised/on immunosuppressants	0.000	0.690	1.046	0.912
Smoking	0.000	0.669	1.611	0.096

Other Medical Conditions	0.003	0.682	0.896	0.638
Have you, your family member or some one you know been diagnosed with COVID-19(Excluding your patients) ?				
I was Diagnosed with COVID-19	6.46E+06	0.641	3.407	0.019*
Family Member was Diagnosed with COVID-19	20.875	0.799	0.597	0.052
Someone Personal was Diagnosed COVID-19	54.754	0.844	1.122	0.723
No one Personal was Diagnosed COVID-19	7.714	0.913	1.007	0.985
Do you think you are at risk of getting COVID-19 in the next 1 year?				
I believe I already have the disease and I am immune to it (Not diagnosed by a test)	0.002	0.608	0.357	0.024*
No, I already have recovered and won't get re-infected (Diagnosed by a test)	2.84E+03	0.852	1.226	0.833
Yes I am concerned that I will get mild symptoms which will probably not require hospitalization	0.587	0.966	0.697	0.260
Yes I am concerned that I will get moderate symptoms which will probably need hospitalization	5.474	0.904	1.099	0.787
Yes I am concerned that I will get severe symptom which will probably require admission to the Intensive care unit	0.078	0.927	1.032	0.948
No I am confident I won't get infected	Reference	Reference	Reference	Reference
Have you directly or indirectly taken care of the COVID-19 patients?				
Yes, but no direct patient contact	0.729	0.970	1.055	0.784
Yes, I have direct patient contact	0.633	0.959	1.143	0.464
No	Reference	Reference	Reference	Reference
Would you advise friends and family to get vaccinated for COVID-19?				
Yes	0.000	0.679	0.082	0.000*
Not sure	0.000	0.278	0.483	0.099
No	Reference	Reference	Reference	Reference
COVID-19 Vaccine for health care workers should be:				
Mandated by the employer, like Influenza vaccine	31.144	0.907	0.463	0.000*
Mandated by the Federal government for all health care workers	6.69E+03	0.735	0.355	0.000*
Mandated by the State government for all health care workers	2.21E+04	0.793	0.374	0.007*

Not sure	0.004	0.709	1.322	0.179
Voluntary	Reference	Reference	Reference	Reference
¹ Statistics presented: Likelihood Ratio Test, RRR (Value)				
² Statistical tests performed: P-Value - All Options (*<0.05 Sig.)				
Direct Patient Care Provider: Registered Nurse/ Patient care technician/ Paramedics/ Rehab service - Respiratory therapist/ Physical therapist/ Occupational therapist/ Speech/ Nutritionist/Social Workers/ Care Coordinators				
Direct Medical Provider: Physician/ Residents/ Medical Students and Advanced Practice Providers (Nurse Practitioners/ Physician's Assistant)				

Table 4: Attitude of Healthcare workers toward vaccine

I would not get the vaccine because I do not believe they work	
Strongly disagree	2,493 (72%)
Disagree	85 (2.4%)
Indifferent	630 (18%)
Agree	190 (5.5%)
Strongly agree	81 (2.3%)
I would not get the vaccine because I believe they are not safe	
Strongly disagree	2,120 (61%)
Disagree	140 (4.0%)
Indifferent	889 (26%)
Agree	251 (7.2%)
Strongly agree	79 (2.3%)
I would not get the vaccine due to religious reasons	
Strongly disagree	2,974 (85%)
Disagree	28 (0.8%)
Indifferent	315 (9.1%)
Agree	129 (3.7%)
Strongly agree	33 (0.9%)
I would not get the vaccine as its my personal choice/freedom	
Strongly disagree	2,629 (76%)
Disagree	187 (5.4%)
Indifferent	398 (11%)
Agree	171 (4.9%)
Strongly agree	94 (2.7%)
I would not get the vaccine as I fear needles	
Strongly disagree	3,001 (86%)
Disagree	23 (0.7%)
Indifferent	372 (11%)
Agree	70 (2.0%)
Strongly agree	13 (0.4%)
I would not get the vaccine as I am worried about the effects of the COVID-19 Vaccine	
Strongly disagree	293 (8.4%)
Disagree	344 (9.9%)
Indifferent	427 (12%)
Agree	1,441 (41%)
Strongly agree	974 (28%)
I would not get the vaccine as I am worried about the effectiveness of the Vaccine	
Strongly disagree	215 (6.2%)
Disagree	324 (9.3%)
Indifferent	540 (16%)

Agree	1,607 (46%)
Strongly agree	793 (23%)
I would not get the vaccine as I am worried about the out of pocket cost coverage of the vaccine	
Strongly disagree	482 (14%)
Disagree	734 (21%)
Indifferent	1,340 (39%)
Agree	678 (19%)
Strongly agree	245 (7.0%)
I would not get the vaccine due to the adverse effects of the vaccine on pre existing conditions	
Strongly disagree	1,511 (43%)
Disagree	682 (20%)
Indifferent	596 (17%)
Agree	437 (13%)
Strongly agree	253 (7.3%)
I would not get the vaccine as there is no need for the vaccine for my risk level	
Strongly disagree	1,823 (52%)
Disagree	939 (27%)
Indifferent	452 (13%)
Agree	167 (4.8%)
Strongly agree	98 (2.8%)
I would not get the vaccine as I have had prior reactions to other vaccinations	
Strongly disagree	2,316 (67%)
Disagree	736 (21%)
Indifferent	226 (6.5%)
Agree	146 (4.2%)
Strongly agree	55 (1.6%)
I would not get the vaccine as I am worried about the rapidity of the COVID vaccine	
Strongly disagree	316 (9.1%)
Disagree	288 (8.3%)
Indifferent	319 (9.2%)
Agree	1,238 (36%)
Strongly agree	1,318 (38%)
I would not get the vaccine as I do not trust pharmaceutical companies	
Strongly disagree	617 (18%)
Disagree	861 (25%)
Indifferent	964 (28%)
Agree	658 (19%)
Strongly agree	379 (11%)
I would not get the vaccine as I do not trust the FDA/CDC	

Strongly disagree	586 (17%)
Disagree	727 (21%)
Indifferent	631 (18%)
Agree	903 (26%)
Strongly agree	632 (18%)
I would not get the vaccine as I do not trust the doctors recommending the vaccine	
Strongly disagree	1,481 (43%)
Disagree	1,046 (30%)
Indifferent	650 (19%)
Agree	201 (5.8%)
Strongly agree	101 (2.9%)
I would not get the vaccine as I do not trust the government about the COVID-19 severity	
Strongly disagree	666 (19%)
Disagree	594 (17%)
Indifferent	621 (18%)
Agree	792 (23%)
Strongly agree	806 (23%)

Table 5: Attitude of US Healthcare workers towards vaccine breakdown by vaccine acceptance

	No, N = 279	Wait for Review, N = 1,953	Yes, N = 1,247	p-value
I would not get the vaccine because I do not believe they work				<0.001 *
Strongly disagree	81 (29%)	1,351 (69%)	1,061 (85%)	
Disagree	86 (31%)	425 (22%)	119 (9.5%)	
Indifferent	71 (25%)	99 (5.1%)	20 (1.6%)	
Agree	23 (8.2%)	45 (2.3%)	17 (1.4%)	
Strongly agree	18 (6.5%)	33 (1.7%)	30 (2.4%)	
I would not get the vaccine because I believe they are not safe				<0.001 *
Strongly disagree	61 (22%)	1,078 (55%)	981 (79%)	
Disagree	80 (29%)	610 (31%)	199 (16%)	
Indifferent	63 (23%)	160 (8.2%)	28 (2.2%)	
Agree	40 (14%)	82 (4.2%)	18 (1.4%)	
Strongly agree	35 (13%)	23 (1.2%)	21 (1.7%)	
I would not get the vaccine due to religious reasons				0.161
Strongly disagree	139 (50%)	1,668 (85%)	1,167 (94%)	
Disagree	62 (22%)	204 (10%)	49 (3.9%)	
Indifferent	53 (19%)	56 (2.9%)	20 (1.6%)	
Agree	13 (4.7%)	9 (0.5%)	6 (0.5%)	
Strongly agree	12 (4.3%)	16 (0.8%)	5 (0.4%)	
I would not get the vaccine as its my personal choice/freedom				<0.001 *
Strongly disagree	63 (23%)	1,452 (74%)	1,114 (89%)	
Disagree	37 (13%)	280 (14%)	81 (6.5%)	
Indifferent	45 (16%)	98 (5.0%)	28 (2.2%)	
Agree	71 (25%)	100 (5.1%)	16 (1.3%)	
Strongly agree	63 (23%)	23 (1.2%)	8 (0.6%)	
I would not get the vaccine as I fear needles				0.749
Strongly disagree	185 (66%)	1,654 (85%)	1,162 (93%)	
Disagree	60 (22%)	243 (12%)	69 (5.5%)	
Indifferent	24 (8.6%)	35 (1.8%)	11 (0.9%)	

Agree	8 (2.9%)	14 (0.7%)	1 (<0.1%)	
Strongly agree	2 (0.7%)	7 (0.4%)	4 (0.3%)	
I would not get the vaccine as I am worried about the effects of the COVID-19 Vaccine				<0.001 *
Strongly disagree	2 (0.7%)	47 (2.4%)	244 (20%)	
Disagree	0 (0%)	56 (2.9%)	288 (23%)	
Indifferent	13 (4.7%)	175 (9.0%)	239 (19%)	
Agree	40 (14%)	980 (50%)	421 (34%)	
Strongly agree	224 (80%)	695 (36%)	55 (4.4%)	
I would not get the vaccine as I am worried about the effectiveness of the Vaccine				0.11
Strongly disagree	2 (0.7%)	49 (2.5%)	164 (13%)	
Disagree	3 (1.1%)	103 (5.3%)	218 (17%)	
Indifferent	31 (11%)	283 (14%)	226 (18%)	
Agree	58 (21%)	996 (51%)	553 (44%)	
Strongly agree	185 (66%)	522 (27%)	86 (6.9%)	
I would not get the vaccine as I am worried about the out of pocket cost coverage of the vaccine				<0.001 *
Strongly disagree	112 (40%)	617 (32%)	611 (49%)	
Disagree	42 (15%)	418 (21%)	274 (22%)	
Indifferent	79 (28%)	437 (22%)	162 (13%)	
Agree	16 (5.7%)	318 (16%)	148 (12%)	
Strongly agree	30 (11%)	163 (8.3%)	52 (4.2%)	
I would not get the vaccine due to the adverse effects of the vaccine on pre existing conditions				0.155
Strongly disagree	71 (25%)	707 (36%)	733 (59%)	
Disagree	44 (16%)	387 (20%)	251 (20%)	
Indifferent	75 (27%)	388 (20%)	133 (11%)	
Agree	34 (12%)	297 (15%)	106 (8.5%)	
Strongly agree	55 (20%)	174 (8.9%)	24 (1.9%)	
I would not get the vaccine as there is no need for the vaccine for my risk level				<0.001 *
Strongly disagree	24 (8.6%)	904 (46%)	895 (72%)	
Disagree	37 (13%)	649 (33%)	253 (20%)	
Indifferent	93 (33%)	288 (15%)	71 (5.7%)	
Agree	66 (24%)	81 (4.1%)	20 (1.6%)	
Strongly agree	59 (21%)	31 (1.6%)	8 (0.6%)	
I would not get the vaccine as I have had prior reactions to other vaccinations				0.005*

Strongly disagree	99 (35%)	1,202 (62%)	1,015 (81%)	
Disagree	73 (26%)	482 (25%)	181 (15%)	
Indifferent	58 (21%)	140 (7.2%)	28 (2.2%)	
Agree	30 (11%)	98 (5.0%)	18 (1.4%)	
Strongly agree	19 (6.8%)	31 (1.6%)	5 (0.4%)	
I would not get the vaccine as I am worried about the rapidity of the COVID vaccine				<0.001 *
Strongly disagree	7 (2.5%)	69 (3.5%)	240 (19%)	
Disagree	7 (2.5%)	65 (3.3%)	216 (17%)	
Indifferent	24 (8.6%)	143 (7.3%)	152 (12%)	
Agree	37 (13%)	714 (37%)	487 (39%)	
Strongly agree	204 (73%)	962 (49%)	152 (12%)	
I would not get the vaccine as I do not trust pharmaceutical companies				0.029*
Strongly disagree	5 (1.8%)	185 (9.5%)	427 (34%)	
Disagree	15 (5.4%)	424 (22%)	422 (34%)	
Indifferent	67 (24%)	636 (33%)	261 (21%)	
Agree	66 (24%)	480 (25%)	112 (9.0%)	
Strongly agree	126 (45%)	228 (12%)	25 (2.0%)	
I would not get the vaccine as I do not trust the FDA/CDC				0.019*
Strongly disagree	3 (1.1%)	161 (8.2%)	422 (34%)	
Disagree	9 (3.2%)	367 (19%)	351 (28%)	
Indifferent	52 (19%)	397 (20%)	182 (15%)	
Agree	67 (24%)	609 (31%)	227 (18%)	
Strongly agree	148 (53%)	419 (21%)	65 (5.2%)	
I would not get the vaccine as I do not trust the doctors recommending the vaccine				<0.001 *
Strongly disagree	22 (7.9%)	604 (31%)	855 (69%)	
Disagree	46 (16%)	695 (36%)	305 (24%)	
Indifferent	96 (34%)	487 (25%)	67 (5.4%)	
Agree	53 (19%)	132 (6.8%)	16 (1.3%)	
Strongly agree	62 (22%)	35 (1.8%)	4 (0.3%)	
I would not get the vaccine as I do not trust the government about the COVID-19 severity				0.59
Strongly disagree	7 (2.5%)	255 (13%)	404 (32%)	
Disagree	16 (5.7%)	334 (17%)	244 (20%)	
Indifferent	55 (20%)	374 (19%)	192 (15%)	

Agree	55 (20%)	503 (26%)	234 (19%)	
Strongly agree	146 (52%)	487 (25%)	173 (14%)	