Thinner cortex is associated with psychosis onset in individuals at Clinical High Risk for Developing Psychosis: An ENIGMA Working Group mega-analysis

Short Title: Cortical Thickness in CHR

Authors

Jalbrzikowski, Maria; Hayes, Rebecca A.; Wood, Stephen J.; Nordholm, Dorte; Zhou, Juan H.; Fusar-Poli, Paolo; Uhlhaas, Peter J.; Takahashi, Tsutomu; Sugranyes, Gisela; Kwak, Yoo Bin; Mathalon, Daniel H.; Katagiri, Naoyuki; Hooker, Christine I.; Smigielski, Lukasz; Colibazzi, Tiziano; Via, Esther; Tang, Jinsong; Koike, Shinsuke; Rasser, Paul E.; Michel, Chantal; Lebedeva, Irina; ten Velden Hegelstad, Wenche; de la Fuente-Sandoval, Camilo; Waltz, James A.; Mizrahi, Romina R.M.; Corcoran, Cheryl; Resch, Franz; Tamnes, Christian K.; Haas, Shalaila S.; Lemmers-Jansen, Imke L.J.; Agartz, Ingrid; Allen, Paul; Andreassen, Ole A.; Atkinson, Kimberley; Bachman, Peter; Baeza, Inmaculada; Baldwin, Helen; Bartholomeusz, Cali F.; Brønnick, Kolbjørn S.; Catalano, Sabrina; Chee, Michael W.L.; Chen, Xiaogang; Cho, Kang Ik K.; Cooper, Rebecca E.; Cropley, Vanessa L.; Ebdrup, Bjørn H.; Fortea, Adriana; Glenthøj, Louise B.; Glenthøj, Birte Y.; de Haan, Lieuwe; Hamilton, Holly K.; Harris, Mathew A.; Haut, Kristen M.; He, Ying; Heekeren, Karsten; Heinz, Andreas; Hooker, Christine I.; Hubl, Daniela; Hwang, Wu Jeong; Kaess, Michael; Kasai, Kiyoto; Kim, Minah; Kindler, Jochen; Klaunig, Mallory, J.; Kristensen, Tina D.; Kwon, Jun Soo; Lawrie, Stephen M.; Lee, Jimmy; León-Ortiz, Pablo; Lin, Ashleigh; Loewy, Rachel L.; Ma, Xiaoqian; Mathalon, Daniel H.; McGorry, Patrick; McGuire, Philip; Mizuno, Masafumi; Møller, Paul; Moncada-Habib, Tomas; Nelson, Barnaby; Nemoto, Takahiro; Nordentoft, Merete; Omelchenko, Maria A.; Oppedal, Ketil; Ouyang, Lijun; Pantelis, Christos; Pariente, Jose C.; Raghava, Jayachandra; Reyes-Madrigal, Francisco; Roach, Brian J.; Røssberg, Jan I.; Rössler, Wulf; Salisbury, Dean F.; Sasabayashi, Daiki; Schall, Ulrich; Schiffman, Jason; Schlagenhauf, Florian; Schmidt, Andre; Sørensen, Mikkel E.; Suzuki, Michio; Theodoridou, Anastasia; Tomyshev, Alexander S.; Værnes, Tor G.; Velakoulis, Dennis; Venegoni, Gloria D.; Vinogradov, Sophia; Wenneberg, Christina; Westlye, Lars T.; Yamasue, Hidenori; Yuan, Liu; Yung, Alison R.; van Amelsvoort, Thérèse A.M.J.; Turner, Jessica A.; van Erp, Theo G.M.; Thompson, Paul M.; Hernaus, Dennis.

Author Affiliations

Amsterdam UMC, Amsterdam, The Netherlands (de Haan); Arkin, Amsterdam, The Netherlands (de Haan); Center for Clinical Intervention and Neuropsychiatric Schizophrenia Research, CINS, Mental Health Centre Glostrup, University of Copenhagen, Glostrup, Denmark (Ebdrup, Glenthøj, Sørensen, Kristensen); Center for Evolutionary Cognitive Sciences, Graduate School of Art and Sciences, The University of Tokyo (Koike); Center for Sleep and Cognition, Yong Loo Lin School of Medicine, National University of Singapore, Singapore (Zhou); Center for the Neurobiology of Learning and Memory, 309 Qureshey Research Lab, Irvine, CA, USA (Van Erp); Center for Translational Magnetic Resonance Research, Yong Loo Lin School of Medicine, National University of Singapore, Singapore (Zhou); Centre for Clinical Intervention and Neuropsychiatric Schizophrenia Research (CINS), Glostrup, Denmark (Raghava); Centre for Mental Health, Faculty of Health, Arts and Design, School of Health Sciences, Swinburne

University, Melbourne, Australia (Pantelis, Cropley); Centre for Neuropsychiatric Schizophrenia Research, CNSR, Glostrup, Denmark (Raghava); Centre for Sleep and Cognition, Yong Loo Lin School of Medicine, National University of Singapore, Singapore (Chee); Centre for Youth Mental Health, University of Melbourne, Australia (Wood, Bartholomeusz, Nelson); Child and Adolescent Mental Health Research Group, Institut de Recerca Sant Joan de Déu, Barcelona, Spain (Via, Dolz Abadia, Muñoz-Samons, Tomyshev, Tor); Child and Adolescent Psychiatry and Psychology Department, Hospital Sant Joan de Déu, Barcelona, Spain (Via, Dolz Abadia, Muñoz-Samons, Tor); Clinic for Child and Adolescent Psychiatry, University Hospital of Heidelberg, Heidelberg, Germany (Resch); Clinical Translational Neuroscience Laboratory, Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, CA, USA (Van Erp); Columbia University Department of Psychiatry, New York, USA (Colibazzi); Copenhagen Research Center for Mental Health - CORE, Mental Health Center Copenhagen, Copenhagen University Hospital, Denmark (Nordholm, Glenthøj, Kristensen, Nordentoft, Wenneberg); Department for Mental Health Research and Development, Division of Mental Health and Addiction, Vestre Viken Hospital Trust, Norway (Møller); Department of Brain and Behavioral Sciences, University of Pavia, Italy (Fusar-Poli); Department of Brain and Cognitive Sciences, Seoul National University College of Natural Sciences, Republic of Korea (Kwak, Hwang); Department of Child and Adolescent Psychiatry and Psychology, Institute of Neuroscience, Hospital Clinic Barcelona, Fundació Clínic Recerca Biomèdica, Universitat de Barcelona, Spain (Fortea); Department of Child and Adolescent Psychiatry and Psychology, Institute of Neuroscience, Hospital Clinic Barcelona, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), CIBERSAM, Universitat de Barcelona, Spain (Sugranyes, Baeza); Department of Child and Adolescent Psychiatry, Psychiatric University Hospital Zurich, University of Zurich, Switzerland (Smigielski); Department of Child and Adolescent Psychiatry, Center of Psychosocial Medicine, University of Heidelberg, Heidelberg, Germany (Kaess); Department of Child and Adolescent Psychiatry, Charité Universitätsmedizin, Berlin, Germany (Uhlhaas); Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Denmark (Ebdrup, Glenthøj); Department of Clinical Physiology, Nuclear Medicine and PET, Functional Imaging Unit, Glostrup, Denmark (Raghava); Department of Neuropsychiatry, Graduate School of Medicine, The University of Tokyo, Japan (Kasai); Department of Neuropsychiatry, Seoul National University Hospital, Republic of Korea (Kim, Kwon); Department of Neuropsychiatry, Toho University School of Medicine, Japan (Katagiri, Mizuno, Nemoto); Department of Neuropsychiatry, University of Toyama Graduate School of Medicine and Pharmaceutical Sciences, Toyama, Japan (Takahashi, Sasabayashi, Suzuki); Department of Pharmacology and Toxicology, University of Toronto, Canada (Koppel); Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway (Agartz); Department of Psychiatry & Behavioral Sciences, University of California, San Francisco, CA, USA (Hamilton); Department of Psychiatry & Neuropsychology, School for Mental Health and Neuroscience, Maastricht University, The Netherlands (van Amelsvoort, Hernaus); Department of Psychiatry (UPK), University of Basel, Switzerland (Borgwardt, Schmidt); Department of Psychiatry and Behavioral Sciences, Rush University Medical Center, Chicago IL (Hooker); Department of Psychiatry and Psychotherapy I, LVR-Hospital Cologne, Cologne, Germany (Heekeren); Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Germany (Heinz, Schlagenhauf); Department of Psychiatry and Psychotherapy, Charité Universitätsmedizin, Berlin, Germany (Rössler); Department of Psychiatry and Psychotherapy, University of Lübeck, Germany (Borgwardt); Department of Psychiatry, Hamamatsu University School of Medicine, Hamamatsu City, Japan (Yamasue); Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA (Corcoran, Haas); Department of Psychiatry, Psychiatry Neuroimaging Laboratory, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA (Cho); Department of Psychiatry,

Psychotherapy and Psychosomatics, Psychiatric University Hospital Zurich, University of Zurich, Switzerland (Smigielski, Heekeren, Rössler, Theodoridou); Department of Psychiatry, Rush University Medical Center, Chicago, IL, USA (Haut); Department of Psychiatry, Seoul National University College of Medicine, Republic of Korea (Kim, Kwon); Department of Psychiatry, Sir Run Run Shaw Hospital, School of Medicine, Zhejiang University, Hangzhou, Zhejiang, China (Tang); Department of Psychiatry, the Second Xiangya Hospital, Central South University, Changsha, Hunan, China (Chen, He, Ma, Ouyang, Yuan); Department of Psychiatry, University of California, San Francisco, CA, USA (Mathalon, Loewy, Roach); Department of Psychiatry, University of Edinburgh, UK (Atkinson, Harris); Department of Psychiatry, University of Minnesota, MN, USA (Vinogradov); Department of Psychiatry, University of Pittsburgh, PA, USA (Jalbrzikowski, Hayes, Bachman, Catalano, Salisbury); Department of Psychological Science, University of California, Irvine, CA, USA (Schiffman); Department of Psychology, University of Maryland, Baltimore County, MD, USA (Klaunig, Schiffman); Department of Psychology, University of Oslo, Norway (Westlye); Department of Psychology, University of Roehampton, London, UK (Allen); Department of Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK (Lemmers-Jansen, Baldwin); Department of Psychosis, Institute of Mental Health, Singapore (Lee); Division of Psychiatry, University of Edinburgh, Edinburgh, UK (Lawrie, Venegoni); Douglas Research Center, Montreal, Canada (Mizrahi); Early Intervention in Psychosis Advisory Unit for South-East Norway, TIPS Sør-Øst, Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway (Værnes); EPIC Lab, Department of Psychosis Studies, King's College London, UK (Fusar-Poli); Faculty of Behavioural and Movement sciences, Department of Clinical, Neuro and Developmental Psychology, Vrije Universiteit Amsterdam, The Netherlands (Lemmers-Jansen); Faculty of Social Sciences,University of Stavanger, Norway (Hegelstad); Florey Institute of Neuroscience and Mental Health, Australia (Pantelis); Georgia State University, Atlanta, GA, USA (Turner); Hunan Key Laboratory of Psychiatry and Mental Health, Changsha, Hunan, China (Ouyang, Yuan, Yung); Imaging Genetics Center, Mark & Mary Stevens Institute for Neuroimaging & Informatics, Keck USC School of Medicine, Los Angeles, CA, USA (Thompson); Institute of Neuroscience and Psychology, University of Glasgow, Glasgow, UK (Uhlhaas); James J Peters VA, Bronx, NY, USA (MIRECC) (Corcoran); Key Laboratory of Medical Neurobiology of Zhejiang Province, Zhejiang, China (Tang); King's College London, UK (McGuire); Laboratory of Experimental Psychiatry, Instituto Nacional de Neurología y Neurocirugía, Mexico City, Mexico (de la Fuente-Sandoval, León-Ortiz, Moncada-Habib, Reyes-Madrigal); Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore (Lee); Maastricht University, The Netherlands (Venegoni); Magnetic Resonance Imaging core facility, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain (Pariente); Maryland Psychiatric Research Center, University of Maryland School of Medicine, MD, USA (Waltz); McGill University, Montreal, Canada (Mizrahi); Melbourne Neuropsychiatry Centre, Department of Psychiatry, University of Melbourne & Melbourne Health, Melbourne, Australia (Cooper, Cropley, Velakoulis); Mental Health Research Center, Moscow, Russia (Lebedeva, Omelchenko); National Clinical Research Center for Geriatric Disorders, Xiangya Hospital, Central South University, Changsha, China (Chen, Ouyang, Yuan, Yung); Neuropsychiatry, The Royal Melbourne Hospital, Australia (Velakoulis); New York State Psychiatric Institute, NY, USA (Colibazzi); NIHR Maudsley Biomedical Research Centre (South London and Maudsley NHS Foundation Trust & King's College London), UK (Baldwin); NORMENT, Division of Mental Health and Addiction, Oslo University Hospital & Institute of Clinical Medicine, University of Oslo, Oslo, Norway (Tamnes, Agartz, Andreassen, Røssberg, Værnes, Westlye); Orygen, The University of Melbourne, Melbourne, Australia (Wood, Bartholomeusz, McGorry, Nelson, Yung); Priority Centre for Brain & Mental Health Research, The University of Newcastle, Australia (Rasser, Schall); Priority Research Centre for Stroke and Brain Injury, The University of Newcastle, Australia (Rasser); Priority Research Centre Grow Up Well, The University of

Newcastle, Australia (Schall); PROMENTA Research Center, Department of Psychology, University of Oslo, Norway (Tamnes); Psychosis Studies, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK (Allen); Research Center for Idling Brain Science, University of Toyama, Toyama, Japan (Takahashi, Sasabayashi, Suzuki); San Francisco Veterans Affairs Health Care System, San Francisco, CA (Hamilton); School of Psychology, University of Birmingham, UK (Wood); Stavanger Medical Imaging Laboratory (SMIL), Stavanger University Hospital, Norway (Oppedal); Telethon Kids Institute, The University of Western Australia, Perth, Australia (Lin); The University of Tokyo Institute for Diversity and Adaptation of Human Mind (UTIDAHM), Japan (Koike, Kasai); TIPS Network for Clinical Research in Psychosis, Stavanger University Hospital, Norway (Hegelstad); Translational Research Center, University Hospital of Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland (Hubl); University Hospital of Child and Adolescent Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland (Michel, Kaess, Kindler)

Address for Correspondence
3501 Forbes Ave, Suite 430
Pittsburgh, PA, United States of America, 15213
jalbrzikowskime@upmc.edu/dennis.hernaus@maastrichtuniversity.nl
Word Count (Total): 3413 (including Key Points, Abstract, and Main Text)
Word Count (Abstract): 332 (max 350)
Word Count (Body): 2961 (max 3000)
Tables: 1 main text/20 supplemental
Figures: 3 main text/5 supplemental
Citation no.: 123
179

Key Points

Question: How do baseline brain morphometric features relate to later psychosis conversion in individuals at clinical high risk (CHR)?

Findings: In the largest coordinated international analysis to date, reduced baseline cortical thickness, but not cortical surface area or subcortical volume, was more pronounced in CHR, in a manner highly consistent with thinner cortex in established psychosis. Regions that displayed greater cortical thinning in future psychosis converters additionally displayed abnormal associations with age.

Meaning: CHR status and later transition to psychosis is robustly associated with reduced cortical thickness. Abnormal age associations and specificity to cortical thickness may point to aberrant postnatal brain development in CHR, including pruning and myelination.

Abstract

Importance: The ENIGMA clinical high risk for psychosis (CHR) initiative, the largest pooled CHR-neuroimaging sample to date, aims to discover robust neurobiological markers of psychosis risk in a sample with known heterogeneous outcomes.

Objective: We investigated baseline structural neuroimaging differences between CHR subjects and healthy controls (HC), and between CHR participants who later developed a psychotic disorder (CHR-PS+) and those who did *not* (CHR-PS-). We assessed associations with age by group and conversion status, and similarities between the patterns of effect size maps for psychosis conversion and those found in other large-scale psychosis studies.

Design, Setting, and Participants. Baseline T1-weighted MRI data were pooled from 31 international sites participating in the ENIGMA CHR Working Group. MRI scans were processed using harmonized protocols and analyzed within a mega- and meta-analysis framework from January-October 2020.

Main Outcome(s) and Measure(s): Measures of regional cortical thickness (CT), surface area (SA), and subcortical volumes were extracted from T1-weighted MRI scans. Independent variables were group (CHR, HC) and conversion status (CHR-PS+, CHR-PS-, HC).

Results: The final dataset consisted of 3,169 participants (CHR=1,792, HC=1,377, age range: 9.5 to 39.8 years, 45% female). Using longitudinal clinical information, we identified CHR-PS+ (N=253) and CHR-PS- (N=1,234). CHR exhibited widespread thinner cortex compared to HC (average d=-0.125, range: -0.09 to -0.17), but not SA or subcortical volume. Thinner cortex in the fusiform, superior temporal, and paracentral regions was associated with psychosis conversion (average d=-0.22). Age showed a stronger negative association with left fusiform and left paracentral CT in HC, compared to CHR-PS+. Regional CT psychosis conversion effect sizes resembled patterns of CT alterations observed in other ENIGMA studies of psychosis.

Conclusions and Relevance: We provide evidence for widespread subtle CT reductions in CHR. The pattern of regions displaying greater CT alterations in CHR-PS+ were similar to those reported in other large-scale investigations of psychosis. Additionally, a subset of these regions displayed abnormal age associations. Widespread CT disruptions coupled with abnormal age associations in CHR may point to disruptions in postnatal brain developmental processes.

1. INTRODUCTION

The clinical high-risk paradigm is a widely used framework to investigate mechanisms underlying psychosis vulnerability. Help-seeking individuals who do not meet diagnostic criteria for a psychotic disorder, but typically present with subthreshold psychotic symptoms and accumulating risk factors, are considered at clinical high-risk (CHR) for developing psychosis[1–3]. An estimated 18-20% of CHR individuals develop a psychotic disorder within 2 years of identification[4], although conversion rates vary[5–11], likely due to heterogeneous recruitment and sampling strategies, and interventions applied[2,13]. However, despite decades of research, the nature of morphometrical alterations associated with psychosis conversion remains largely unknown. Here, we aim to address this question by combining all available structural neuroimaging data in CHR to date, in an attempt to better understand brain mechanisms associated with psychosis risk and conversion in this population.

A large body of work has used structural magnetic resonance imaging (sMRI) to investigate morphometric brain alterations in CHR individuals[14–37]. However, the extent to which characteristic baseline structural neuroimaging differences exist between those at CHR who later develop a psychotic disorder (CHR-PS+) compared to those who do *not* (CHR-PS-) is debated. Many studies failed to find baseline differences between CHR-PS+ and CHR-PS-[15,35,38,39], though a meta-analysis and multi-center study found lower prefrontal and temporal volumes or cortical thickness (CT) in CHR-PS+[40,41]. High attrition rates in CHR samples[42–44], coupled with low psychosis conversion rates[4,45], often yield insufficient power to detect between-group structural brain differences. Moreover, small sample sizes can be associated with inflated effect sizes[46,47], so effect sizes of prior studies that found structural brain alterations in CHR may be overestimated. Although multi-site consortia aim to address these challenges[34,48–50], the largest published sMRI studies to date included fewer than 50 CHR-PS+[38,41]. Furthermore, it is currently unknown whether group differences are robust enough to predict outcomes.

Importantly, many CHR participants are adolescents or young adults, a timeframe associated with psychosis onset[51,52]. Several brain regions implicated in psychosis show protracted developmental courses continuing through adolescence[53–63], suggesting that morphometric alterations associated with psychosis risk vary with age. Indeed, there are developmental influences on psychotic symptom presentation[64–68], perhaps driven by differences in regional brain changes. It is not fully understood how age-related patterns in brain morphometry in CHR differ from normal development. Thus, using a developmental framework to examine whether morphometric alterations in CHR are influenced by age may provide important insights into mechanisms associated with psychosis risk, and the stability of neuroimaging measures associated with psychosis risk across development.

Finally, it is unknown whether baseline brain alterations associated with future conversion to psychosis resemble those observed in other large-scale psychosis studies. Understanding whether morphometric alterations in CHR overlap with those observed in individuals who have schizophrenia[69–71] and individuals with a genetic subtype of psychosis[72,73] will provide insights into convergent or distinct alterations across the psychosis spectrum.

To address these questions, we founded the Enhancing Neuro Imaging Genetics through Meta-Analysis (ENIGMA) Clinical High Risk Working Group in 2018. Using baseline

sMRI data and longitudinal clinical information from 31 sites, this study addressed the following questions:

1. Do CHR and healthy control (HC) participants differ in CT, surface area (SA), and/or subcortical volumes?
2. Is there a neuroanatomic signature associated with future transition to a psychotic disorder (CHR-PS+ versus CHR-PS- versus HC)?
3. Do structural neuroimaging measures identified in Aims 1 and 2 display group differences in age associations, suggestive of abnormal developmental trajectories?
4. Is the pattern of morphometric alteration associated with psychosis conversion similar to that observed in ENIGMA studies of psychosis?

2. METHODS

Participants

We included 1,792 CHR and 1,377 HC from 31 sites participating in the ENIGMA CHR Working Group (Table 1). CHR data consisted of CHR-PS+ (N=253), CHR-PS- (N=1,234), and CHR individuals without follow-up data (CHR-UNK, N=305). CHR participants met CAARMS (N=821) or SIPS (N=971) CHR criteria (see details in Supplement). Site-specific inclusion/exclusion criteria are detailed in **eTable 1**. All sites obtained Institutional Review Board approval prior to data collection. Informed written consent was obtained from every participant or the participant's guardian (for participants <18 years). All studies were conducted in accordance with the Declaration of Helsinki[74].

Image Acquisition and Processing

Thirty-one sites contributed T1-weighted MRI brain scans from 50 MRI scanners (**eTable 2** for MRI acquisition details). We conducted power calculations to estimate the sample size necessary to detect statistically significant effects (see **Supplement** for details).

We extract structural gray matter measures of 68 CT, 68 SA measures, and 16 subcortical volume measures using the FreeSurfer analysis software[75–78]. We implemented the ENIGMA consortium quality assessment pipeline[70,73,79–82] (**Supplement** for further details).

Prior to conducting the mega-analysis, all neuroimaging data were adjusted for scanner protocol effects using neuroComBat[83], a modified version of ComBat[84] that increases the statistical power and precision of neuroimaging effect size estimates[83].

Statistical Analyses

Group and conversion-related differences in structural neuroimaging metrics

We assessed group differences using general linear models (GLMs) within a mega-analysis framework, with each sMRI measure (i.e., regional CT, SA or subcortical volume) as the dependent variable and group (HC/CHR) or conversion status (CHR-PS+/CHR-PS-/HC) as

the independent variable. We included age, age^2, sex, and estimated total intracranial volume
(ICV) as covariates in all models, and corrected for multiple comparisons (N=155) using the
False Discovery Rate (FDR[85]) method. We considered q-values<0.05 statistically significant.
For all structural neuroimaging measures, we conducted pairwise comparisons and
calculated Cohen's d effect sizes between two (CHR vs. HC) or three groups of interest (CHR-
PS+ vs. HC/CHR-PS+ vs. CHR-PS-/CHR-PS- vs. HC). We also conducted analyses within a
meta-analysis framework and fitted linear mixed models of sMRI data (not adjusted with
neuroCombat), with scanner as a random effect. We investigated sMRI differences associated
with the specific psychosis-risk syndromes (e.g., Attenuated Positive Symptom Syndrome;
details in **Supplement**).
To evaluate the stability of group and conversion status differences, we performed
analyses statistically controlling for psychotropic medication exposure at baseline. To assess
effects of site, we conducted jackknife resampling analyses, i.e., iteratively removing one site's
data and re-running respective analyses[86,87]. sMRI measures that failed to show a group or
conversion status effect at q<0.05 in >10% of jackknife iterations (i.e., 4/31 sites) were
considered "unstable".
To assess the meaningfulness of obtained effect sizes we used two analytic
approaches: equivalence testing (to assess whether observed differences fell within the upper
and lower bounds of a predefined smallest effect size of interest, providing support for the
absence of a meaningful effect) and minimal-effects testing (to assess whether observed effects
were greater than the same pre-defined effect size[88]). Upper/lower bounds (representing the
positive/negative predefined smallest effect size of interest) were set to d = +/- 0.15 (further
details and effect size rationale, see **Supplement**).
336
Group and conversion-related differences in sMRI age-associations
We used general additive models[89,90] to model group and conversion status differences
in the relationship between age and sMRI measures. General additive models extend GLMs by
allowing for a non-linear relationship between independent and dependent variables. Here, age
is modeled as a smooth function (continuous derivatives), which is akin to the "age-related"
slope in a GLM. Analyses were confined to sMRI measures that displayed statistically significant
effects of I) HC vs. CHR (N=56 measures) and II) CHR-PS+ vs. HC/CHR-PS+ vs. CHR-PS-
/CHR-PS- vs. HC (N=4). We examined the effect of baseline age, group/conversion status, and
the interaction between the two variables. Sex and estimated ICV were included as covariates.
Similar to prior work examining age effects during adolescent development[91,92], we restricted
our sample's age range to 12-25 years (**eTable 3**). Details on post-hoc analyses for significant
interaction effects are provided in the **Supplement**.
349
Comparison of psychosis conversion-related effects to other ENIGMA findings
We computed Spearman's rank correlations to assess the extent to which the pattern of
observed effect sizes (Cohen's d's for CHR-PS+ vs. HC and CHR-PS-) correlated with the
pattern found in prior psychosis studies, specifically the ENIGMA Schizophrenia(SZ vs. HC[69,70])
and ENIGMA 22q11.2 Deletion Syndrome (22q11DS with psychosis vs. 22q11DS without
psychosis[72,73]) Working Groups. As a control, we compared CHR-PS+/CHR-PS- vs. HC effect
sizes to effect sizes published by the Major Depressive Disorder Working Group (MDD vs.
HC[93]). Additional details of correlation analyses are reported in the **Supplement**.

3. RESULTS

Sample Characteristics

Site demographics are reported in **Table 1**. See **eTables 4-6** for IQ and symptom comparisons. Few CHR participants reported typical (<1%) and/or atypical antipsychotic (12.4%) medication use (**eTable 7**).

Widespread lower CT in CHR versus HC

Compared to HC, the CHR group had smaller global neuroimaging measures: estimated ICV (d=-0.13, CI=-0.2 to -0.06), mean CT (d=-0.18, CI=-0.25 to -0.11) and total SA (d=-0.15, CI=-0.22 to -0.08). Fifty-three additional regions exhibited a significant effect of group (q<0.05, **eTable 8**). The largest group effects were observed for widespread lower CT in CHR vs. HC (42/68 comparisons, d range=-0.09 to -0.17; **Figure 1A** for overview; **eTable 8** and **eFigure 1** for details). Few subcortical (3/16) and SA (8/68) group differences were observed (d range=-0.09 to -0.16). No group-by-sex interactions were detected (all q>0.05).

Group difference magnitudes remained similar when controlling for antipsychotic and psychotropic medication use (**eTable 9**). 91% of group differences remained statistically significant (q<0.05) in >90% of jackknife resampling iterations (eFigure 2), providing evidence that site effects were not driving results. Equivalence testing revealed that effect sizes for group differences were likely not meaningfully greater than our *a priori*-defined smallest effect size of interest in 3/56 measures (**eFigure 1, eTable 10**).

Mega- and meta-analyses of neuroimaging data prior to neuroCombat harmonization provided similar results (Spearman's ρ>0.93; **eTable 11**). No sMRI measures were uniquely sensitive to psychosis-risk syndrome (see **Supplement** and **eTables 12-15** forresults).

Thinner paracentral, fusiform, and superior temporal CT are associated with psychosis conversion

Forty-eight structural neuroimaging measures (N_{CT}=37) exhibited a significant overall effect of psychosis conversion status (q<0.05, **Figure 1B**; **eFigure 3** and **eTable 16**). The three groups differed from each other on four structural neuroimaging measures: in comparison to HC and CHR-PS-, CHR-PS+ exhibited thinner cortex in bilateral paracentral, right superior temporal, and left fusiform regions (**Figure 1C**, average d of four sMRI measures=-0.22). CHR-PS+ and CHR-PS- (vs. HC) exhibited thinner cortex in the left superior temporal and right fusiform regions;similar trends were observed for CHR-PS+ vs. CHR-PS- differences (p<0.08; Figure 1C). Results remained stable when length of follow-up period was included as a covariate.

Using minimal-effects testing, we observed that effect sizes for bilateral paracentral (L Z=-2.43, p=0.02; R Z=-1.86 p=0.06), right superior temporal (Z=-2.29, p=0.02) and left fusiform (Z=-2.00, p=0.05) in CHR-PS+ vs. HC were all significantly *greater* than 0.15, at least at trend level, underscoring the presence of notable group differences.

In the remaining follow-up comparisons, CHR-PS+ and CHR-PS- differed from HC at p<0.05, but there were no significant differences between CHR-PS+ and CHR-PS- (29/48 comparisons; **eTable 16**). We observed no conversion status-by-sex interactions (all q>0.05).

When we controlled for antipsychotic or psychotropic medication exposure, most overall psychosis conversion effects remained statistically significant (**eTable 17**). sMRI measures that differentiated the three groups showed an overall conversion effect at $q<0.05$ in all jackknife iterations, ruling out site effects as a potential confound (**eFigure 2**). No pairwise comparisons for structural neuroimaging measures that showed an overall conversion effect at $q<0.05$ were statistically equivalent to our predefined smallest effect size of interest (**eTable 18**). There were no statistically significant psychosis-risk syndrome-by-conversion status interactions (**eTable 19**).

CHR-PS+ fail to show expected age-associations with fusiform and paracentral CT

There were no significant group-by-age interactions in the 56 neuroimaging measures that differed between CHR and HC ($q>0.05$, **eTable 20**). In contrast, 2/4 sMRI measures in which all groups differed (HC vs. CHR-PS+ vs. CHR-PS-) displayed a statistically significant conversion status-by-age interaction.

In left fusiform, age-CT associations differed between CHR-PS+ vs. HC (F=9.8, p=4.9e-05, q=5.9e-04) and CHR-PS- vs. HC (F=8.7, p=1.5e-04, q=9.1e-04, Figure 2A, left). Between ages 12-16, HC showed a stronger negative association between age and CT, compared to CHR-PS+ and CHR-PS-, suggesting a reduced rate of CT change in CHR-PS+ and CHR-PS-. Though the age-by-conversion status interaction was not statistically significant, a similar pattern emerged for the right fusiform CT (Figure 2A, right).

Age-effects in the left paracentral CT differed between CHR-PS+ vs. HC (F=5.9, p=4.9e-03, q=0.02, Figure 2B, left). Between 12-15.8 years of age, HC showed a stronger negative association between age and CT in comparison to CHR-PS+. Age-CT associations did not differ between CHR-PS- vs. HC (F=0.2, p=0.69). This pattern of results was not observed for the right paracentral CT (Figure 2B, right).

We found no significant age-by-conversion status interactions for the superior temporal CT (Figure 2C); all three groups showed negative CT-age associations.

CT aberrations in CHR-PS+ resemble the pattern observed in SZ and 22q11DS with a psychotic disorder diagnosis, but not MDD

Figure 3A provides a visual overview of CT alterations in CHR-PS+ relative to SZ and individuals with 22q11DS and a psychotic disorder. The pattern of baseline CT alterations in CHR-PS+ (CHR-PS+ vs. HC effect sizes) correlated significantly with that observed in SZ (ρ=0.35, $p_{permute}$=0.004, **Figure 3B** top) and individuals with 22q11DS and psychosis (ρ=0.43, $p_{permute}$=0.001, **Figure 3B** bottom). CT alterations in CHR did *not* correlate with those observed in MDD (CHR-PS+ ρ=-0.03) and slopes for CHR-PS+/SZ and CHR-PS+/MDD correlations significantly differed (Steiger's Z=2.06, $p_{permute}$=0.008). No significant correlations were observed for SA (SZ ρ=-0.03; 22q.11DS ρ=-0.06, eFigure 5). Subcortical volume alterations in CHR-PS+ correlated with those observed in SZ (ρ=0.54, p=0.03) and a similar non-significant trend was observed for 22q11DS and psychosis (ρ=0.46, p=0.07). Associations for CHR-PS- (vs. HC) effect sizes were similar to those reported here (see **Supplement**).

4. DISCUSSION

We conducted the largest multisite neuroimaging investigation to date in CHR participants, examining baseline structural neuroimaging measures associated with later transition to psychosis. We found widespread thinner cortex in CHR, consistent with previously-reported CT alterations in individuals with an established psychotic disorder. Compared to CHR-PS- and HC, at baseline, CHR-PS+ exhibited thinner cortex in bilateral paracentral, right fusiform and left superior temporal regions, with effect sizes significantly greater than what we considered to be meaningful *a priori*. Our results were robust to effects of medication exposure, sex, site, and length of follow-up period. Findings from this international effort suggest that conversion to psychosis amongst those at high risk is associated with thinner cortex at baseline.

We identified widespread regional thinner cortex in CHR compared to HC. Thinner cortex has been observed in SZ, as well as other psychiatric disorders[69,79,93–95]. Importantly, the overall pattern of thinner cortex in CHR resembled that observed in SZ and individuals with 22q11DS and a psychotic disorder, but not in MDD. For CHR-PS+, correlations with SZ CT alterations were significantly greater than the relationship observed with MDD CT alterations. Taken together, our results suggest that the overall constellation of reported CT alterations resembles the regional pattern of alterations observed in SZ and genetic disorders associated with psychosis, and thus argues in favour of psychosis-specific thinner cortex in CHR

We also observed thinner cortex specific to psychosis conversion in paracentral, superior temporal and fusiform CT; CHR-PS+ exhibit significantly lower CT than CHR-PS- and HC in these regions. Lower baseline CT and/or volume in these regions has previously been reported in CHR-PS+[96,97] (data not used here). Furthermore, longitudinal CT decreases in these regions have been associated with transition to psychosis in CHR[17,98,99]. The magnitude of altered CT in CHR-PS+ in paracentral, superior temporal and fusiform regions was highly consistent with findings in SZ[70,100–102] and lower fusiform and paracentral CT has been observed in non-clinical voice hearers[103]. Thus, the observed pattern of CT alterations in CHR-PS+ is consilient with morphometric disruptions across the wider psychosis spectrum.

Consistent with previous CHR studies examining baseline neuroimaging associations with later conversion to psychosis[96,104], we did not observe widespread subcortical volume or SA alterations associated with later psychosis transition. Taken together, these results suggest that CT reductions may be among the most widespread, robust, and specific morphometric changes associated with psychosis risk and conversion, compared to SA or subcortical volume.

The neuroanatomic pattern of group differences and age-associated disruptions observed in CHR may provide important insights into mechanisms underlying increased risk for psychosis. Pre-clinical models [105,106] and recent genome-wide association studies[107] suggest that genetic variants associated with SZ are linked to the regulation of neural progenitor cells during fetal development, while genetic markers associated with CT were associated with regulatory processes in adulthood. Thus, CT alterations may be the end result of maladaptive maturation-related mechanisms that occur during post-fetal development, including proliferation, synaptic pruning and/or myelination[108–111]. Thinner CT, particularly in early adolescence[112,113] (**Figure 2**), could reflect abnormal synaptic plasticity or pruning, which have both been implicated in *in vitro* SZ models[114]. Although excessive synaptic pruning is one plausible explanation for thinner cortex associated with psychosis transition, recent evidence suggests

that intracortical myelination and/or expression of myelin-related genes may be mechanisms
driving cortical thinning[115,116]. To better understand neurobiological mechanisms underlying
psychosis transition in CHR, investigations of concomitant measures of cortical thickness,
macroscale white matter tracts, and intracortical myelination are necessary.
Even if CT reductions in CHR were robust, effect sizes for between-group differences
were nevertheless small-to-moderate and accounted for ~1% of the variance in CHR-PS+ vs.
CHR-PS- comparisons. The subtle nature of these morphometric alterations underscores the
importance of adequate statistical power, achievable only through large-scale multi-site
collaborations. Consistent with recent work showing that SZ polygenic risk scores only improved
differentiation of CHR-PS+ and HC (not CHR-PS+ from CHR-PS-)[117], we anticipate that
baseline, univariate sMRI metrics will to have a similar impact on psychosis risk prediction
algorithms. Given the logistic and financial challenges that MRI brings, the use of MRI metrics in
isolation may not be feasible or useful for psychosis risk prediction. A viable solution may be to
adopt sequential assessment frameworks, as recently implemented[34]. Alternatively, sMRI
alterations may be a better predictor of general psychopathology, and would be better suited for
transdiagnostic risk prediction models[118,119].
503
Limitations
One limitation common to multi-site studies is that data were collected from multiple
scanners, although leave-one-out analyses suggest that site effects were not prominent.
Secondly, this initial study focused on baseline cross-sectional data, and did not investigate
progressive sMRI changes (trajectories) associated with psychosis conversion, as identified in
prior work[17,19,38,97–99,120–123]. This is a future goal of the ENIGMA CHR Working Group, now that
feasibility of this collaboration has been established.
511
Conclusions and Future Directions
In the largest study of brain abnormalities in CHR to date, we found robust evidence for a
subtle, widespread pattern of CT alterations, consistent with observations in psychosis. The
specificity of these alterations to CT - as well as age-associated deviations in regions sensitive
to psychosis conversion - may point to abnormal development processes. These findings also
point to age ranges (i.e., early adolescence) when morphometric abnormalities in CHR might be
greatest.
519

Acknowledgements
BHE (Bjørn H. Ebdrup) is part of the Advisory Board of Eli Lilly Denmark A/S, Janssen-
Cilag, Lundbeck Pharma A/S, and Takeda Pharmaceutical Company Ltd; and has
received lecture fees from Bristol-Myers Squibb, Otsuka Pharma Scandinavia AB, Eli
Lilly Company, Boehringer Ingelheim Danmark A/S, and Lundbeck Pharma A/S. DH
(Dennis Hernaus) has received financial compensation as a consultant for P1vital
Produicts Ltd. F-RM (Francisco Reyes-Madrigal) has received compensation as a
speaker for Janssen (Johnson & Johnson). All reported activities were unrelated to the
work presented in this manuscript.

References

1. Miller TJ, McGlashan TH, Rosen JL, et al. Prodromal assessment with the structured interview for prodromal syndromes and the scale of prodromal symptoms: predictive validity, interrater reliability, and training to reliability. *Schizophr Bull*. 2003;29(4):703.

2. Yung AR, McGorry PD, McFarlane CA, Jackson HJ, Patton GC, Rakkar A. Monitoring and care of young people at incipient risk of psychosis. *Schizophr Bull*. 1996;22(2):283-303. doi:10.1093/schbul/22.2.283

3. Yung AR, Phillips LJ, McGorry PD, et al. Prediction of psychosis. A step towards indicated prevention of schizophrenia. *Br J Psychiatry Suppl*. 1998;172(33):14-20.

4. Fusar-Poli P, Bonoldi I, Yung AR, et al. Predicting psychosis: meta-analysis of transition outcomes in individuals at high clinical risk. *Arch Gen Psychiatry*. 2012;69(3):220-229. doi:10.1001/archgenpsychiatry.2011.1472

5. Simon AE, Umbricht D. High remission rates from an initial ultra-high risk state for psychosis. *Schizophr Res*. 2010;116(2-3):168-172. doi:10.1016/j.schres.2009.10.001

6. Ruhrmann S, Schultze-Lutter F, Salokangas RKR, et al. Prediction of psychosis in adolescents and young adults at high risk: results from the prospective European prediction of psychosis study. *Arch Gen Psychiatry*. 2010;67(3):241-251. doi:10.1001/archgenpsychiatry.2009.206

7. Cannon TD, Cadenhead K, Cornblatt B, et al. Prediction of Psychosis in Youth at High Clinical Risk: A Multisite Longitudinal Study in North America. *Arch Gen Psychiatry*. 2008;65(1):28. doi:10.1001/archgenpsychiatry.2007.3

8. Yung AR, Phillips LJ, Yuen HP, et al. Psychosis prediction: 12-month follow up of a high-risk ("prodromal") group. *Schizophr Res*. 2003;60(1):21-32. doi:10.1016/s0920-9964(02)00167-6

9. Cornblatt BA, Lencz T, Smith CW, Correll CU, Auther AM, Nakayama E. The schizophrenia prodrome revisited: a neurodevelopmental perspective. *Schizophr Bull*. 2003;29(4):633-651. doi:10.1093/oxfordjournals.schbul.a007036

10. Zhang T, Li H, Tang Y, et al. Validating the Predictive Accuracy of the NAPLS-2 Psychosis Risk Calculator in a Clinical High-Risk Sample from the SHARP (Shanghai At Risk for Psychosis) Program. *Am J Psychiatry*. 2018;175(9):906-908. doi:10.1176/appi.ajp.2018.18010036

11. Tor J, Dolz M, Sintes A, et al. Clinical high risk for psychosis in children and adolescents: a systematic review. *Eur Child Adolesc Psychiatry*. 2018;27(6):683-700. doi:10.1007/s00787-017-1046-3

12. Hartmann JA, Yuen HP, McGorry PD, et al. Declining transition rates to psychotic disorder in "ultra-high risk" clients: Investigation of a dilution effect. *Schizophr Res*. 2016;170(1):130-136. doi:10.1016/j.schres.2015.11.026

13. Fusar-Poli P, Palombini E, Davies C, et al. Why transition risk to psychosis is not declining at the OASIS ultra high risk service: The hidden role of stable pretest risk enrichment. *Schizophr Res*. 2018;192:385-390. doi:10.1016/j.schres.2017.06.015

14. Bakker G, Caan MWA, Vingerhoets WAM, et al. Cortical Morphology Differences in Subjects at Increased Vulnerability for Developing a Psychotic Disorder: A Comparison between Subjects with Ultra-High Risk and 22q11.2 Deletion Syndrome. *PLOS ONE*. 2016;11(11):e0159928. doi:10.1371/journal.pone.0159928

15. Zikidi K, Gajwani R, Gross J, et al. Grey-matter abnormalities in clinical high-risk participants for psychosis. *Schizophr Res*. Published online November 15, 2019. doi:10.1016/j.schres.2019.08.034

16. Velakoulis D, Wood SJ, Wong MTH, et al. Hippocampal and Amygdala Volumes According to Psychosis Stage and Diagnosis: A Magnetic Resonance Imaging Study of Chronic Schizophrenia, First-Episode Psychosis, and Ultra–High-Risk Individuals. *Arch*

Gen Psychiatry. 2006;63(2):139. doi:10.1001/archpsyc.63.2.139

17.   Sun D, Phillips L, Velakoulis D, et al. Progressive brain structural changes mapped as
psychosis develops in 'at risk' individuals. *Schizophrenia Research*. 2009;108(1-3):85-92.
doi:10.1016/j.schres.2008.11.026

18.   Fornito A, Yung AR, Wood SJ, et al. Anatomic Abnormalities of the Anterior Cingulate
Cortex Before Psychosis Onset: An MRI Study of Ultra-High-Risk Individuals. *Biological
Psychiatry*. 2008;64(9):758-765. doi:10.1016/j.biopsych.2008.05.032

19.   Takahashi T, Wood SJ, Yung AR, et al. Insular cortex gray matter changes in individuals
at ultra-high-risk of developing psychosis. *Schizophrenia Research*. 2009;111(1-3):94-
102. doi:10.1016/j.schres.2009.03.024

20.   Tomyshev AS, Lebedeva IS, Akhadov TA, Omelchenko MA, Rumyantsev AO, Kaleda
VG. Alterations in white matter microstructure and cortical thickness in individuals at ultra-
high risk of psychosis: A multimodal tractography and surface-based morphometry study.
Psychiatry Res Neuroimaging. 2019;289:26-36. doi:10.1016/j.pscychresns.2019.05.002

21.   Kwak YB, Kim M, Cho KIK, Lee J, Lee TY, Kwon JS. Reduced cortical thickness in
subjects at clinical high risk for psychosis and clinical attributes. *Aust N Z J Psychiatry*.
2019;53(3):219-227. doi:10.1177/0004867418807299

22.   Yassin W, Nakatani H, Zhu Y, et al. Machine-learning classification using neuroimaging
data in schizophrenia, autism, ultra-high risk and first-episode psychosis. *Transl
Psychiatry*. 2020;10(1):278. doi:10.1038/s41398-020-00965-5

23.   Nakamura M, Takahashi T, Takayanagi Y, et al. Surface morphology of the orbitofrontal
cortex in individuals at risk of psychosis: a multicenter study. *Eur Arch Psychiatry Clin
Neurosci*. 2019;269(4):397-406. doi:10.1007/s00406-018-0890-6

24.   Sasabayashi D, Takayanagi Y, Takahashi T, et al. Increased Occipital Gyrification and
Development of Psychotic Disorders in Individuals With an At-Risk Mental State:
A Multicenter Study. *Biol Psychiatry*. 2017;82(10):737-745.
doi:10.1016/j.biopsych.2017.05.018

25.   Iwashiro N, Suga M, Takano Y, et al. Localized gray matter volume reductions in the pars
triangularis of the inferior frontal gyrus in individuals at clinical high-risk for psychosis and
first episode for schizophrenia. *Schizophr Res*. 2012;137(1-3):124-131.
doi:10.1016/j.schres.2012.02.024

26.   Nakamura K, Takahashi T, Nemoto K, et al. Gray matter changes in subjects at high risk
for developing psychosis and first-episode schizophrenia: a voxel-based structural MRI
study. *Front Psychiatry*. 2013;4:16. doi:10.3389/fpsyt.2013.00016

27.   Takayanagi Y, Kulason S, Sasabayashi D, et al. Reduced Thickness of the Anterior
Cingulate Cortex in Individuals With an At-Risk Mental State Who Later Develop
Psychosis. *Schizophr Bull*. 2017;43(4):907-913. doi:10.1093/schbul/sbw167

28.   Takahashi T, Tsugawa S, Nakajima S, et al. Thalamic and striato-pallidal volumes in
schizophrenia patients and individuals at risk for psychosis: A multi-atlas segmentation
study. *Schizophr Res*. Published online May 21, 2020. doi:10.1016/j.schres.2020.04.016

29.   Buechler R, Wotruba D, Michels L, et al. Cortical Volume Differences in Subjects at Risk
for Psychosis are Driven by Surface Area. *Schizophr Bull*. Published online May 28,
2020. doi:10.1093/schbul/sbaa066

30.   Chung Y, Allswede D, Addington J, et al. Cortical abnormalities in youth at clinical high-
risk for psychosis: Findings from the NAPLS2 cohort. *Neuroimage Clin*. 2019;23:101862.
doi:10.1016/j.nicl.2019.101862

31.   Chung Y, Addington J, Bearden CE, et al. Use of Machine Learning to Determine
Deviance in Neuroanatomical Maturity Associated With Future Psychosis in Youths at
Clinically High Risk. *JAMA Psychiatry*. 2018;75(9):960-968.
doi:10.1001/jamapsychiatry.2018.1543

32.   Koutsouleris N, Meisenzahl EM, Davatzikos C, et al. Use of neuroanatomical pattern

classification to identify subjects in at-risk mental states of psychosis and predict disease
transition. *Arch Gen Psychiatry*. 2009;66(7):700-712.
doi:10.1001/archgenpsychiatry.2009.62

33. Koutsouleris N, Riecher-Rössler A, Meisenzahl EM, et al. Detecting the psychosis
prodrome across high-risk populations using neuroanatomical biomarkers. *Schizophr*
Bull. 2015;41(2):471-482. doi:10.1093/schbul/sbu078

34. Koutsouleris N, Kambeitz-Ilankovic L, Ruhrmann S, et al. Prediction Models of Functional
Outcomes for Individuals in the Clinical High-Risk State for Psychosis or With Recent-
Onset Depression: A Multimodal, Multisite Machine Learning Analysis. *JAMA Psychiatry*.
2018;75(11):1156-1172. doi:10.1001/jamapsychiatry.2018.2165

35. Klauser P, Zhou J, Lim JKW, et al. Lack of Evidence for Regional Brain Volume or
Cortical Thickness Abnormalities in Youths at Clinical High Risk for Psychosis: Findings
From the Longitudinal Youth at Risk Study. *Schizophr Bull*. 2015;41(6):1285-1293.
doi:10.1093/schbul/sbv012

36. Ziermans TB, Durston S, Sprong M, et al. No evidence for structural brain changes in
young adolescents at ultra high risk for psychosis. *Schizophr Res*. 2009;112(1-3):1-6.
doi:10.1016/j.schres.2009.04.013

37. Takahashi T, Yücel M, Yung AR, et al. Adhesio interthalamica in individuals at high-risk
for developing psychosis and patients with psychotic disorders. *Prog*
Neuropsychopharmacol Biol Psychiatry. 2008;32(7):1708-1714.
doi:10.1016/j.pnpbp.2008.07.007

38. Cannon TD, Chung Y, He G, et al. Progressive reduction in cortical thickness as
psychosis develops: a multisite longitudinal neuroimaging study of youth at elevated
clinical risk. *Biol Psychiatry*. 2015;77(2):147-157. doi:10.1016/j.biopsych.2014.05.023

39. Sakuma A, Obara C, Katsura M, et al. No regional gray matter volume reduction
observed in young Japanese people at ultra-high risk for psychosis: A voxel-based
morphometry study. *Asian J Psychiatr*. 2018;37:167-171. doi:10.1016/j.ajp.2018.09.009

40. Fusar-Poli P, Borgwardt S, Crescini A, et al. Neuroanatomy of vulnerability to psychosis:
a voxel-based meta-analysis. *Neurosci Biobehav Rev*. 2011;35(5):1175-1185.
doi:10.1016/j.neubiorev.2010.12.005

41. Mechelli A, Riecher-Rössler A, Meisenzahl EM, et al. Neuroanatomical abnormalities that
predate the onset of psychosis: a multicenter study. *Arch Gen Psychiatry*.
2011;68(5):489-495. doi:10.1001/archgenpsychiatry.2011.42

42. Farris MS, Devoe DJ, Addington J. Attrition rates in trials for adolescents and young
adults at clinical high-risk for psychosis: A systematic review and meta-analysis. *Early*
Interv Psychiatry. 2020;14(5):515-527. doi:10.1111/eip.12864

43. Schlosser DA, Jacobson S, Chen Q, et al. Recovery from an at-risk state: clinical and
functional outcomes of putatively prodromal youth who do not develop psychosis.
Schizophr Bull. 2012;38(6):1225-1233. doi:10.1093/schbul/sbr098

44. Rutigliano G, Valmaggia L, Landi P, et al. Persistence or recurrence of non-psychotic
comorbid mental disorders associated with 6-year poor functional outcomes in patients at
ultra high risk for psychosis. *J Affect Disord*. 2016;203:101-110.
doi:10.1016/j.jad.2016.05.053

45. Simon AE, Borgwardt S, Riecher-Rössler A, Velthorst E, de Haan L, Fusar-Poli P. Moving
beyond transition outcomes: meta-analysis of remission rates in individuals at high clinical
risk for psychosis. *Psychiatry Res*. 2013;209(3):266-272.
doi:10.1016/j.psychres.2013.03.004

46. Button KS, Ioannidis JPA, Mokrysz C, et al. Power failure: why small sample size
undermines the reliability of neuroscience. *Nat Rev Neurosci*. 2013;14(5):365-376.
doi:10.1038/nrn3475

47. Consideration of Sample Size in Neuroscience Studies. *J Neurosci*. 2020;40(21):4076-

4077. doi:10.1523/JNEUROSCI.0866-20.2020
48. Addington J, Cadenhead KS, Cornblatt BA, et al. North American Prodrome Longitudinal
Study (NAPLS 2): Overview and recruitment. *Schizophrenia Research*. 2012;142(1):77-
82. doi:10.1016/j.schres.2012.09.012
49. Addington J, Liu L, Brummitt K, et al. North American Prodrome Longitudinal Study
(NAPLS 3): Methods and baseline description. *Schizophrenia Research*. Published online
April 18, 2020. doi:10.1016/j.schres.2020.04.010
50. Tognin S, van Hell HH, Merritt K, et al. Towards Precision Medicine in Psychosis:
Benefits and Challenges of Multimodal Multicenter Studies-PSYSCAN: Translating
Neuroimaging Findings From Research into Clinical Practice. *Schizophr Bull*.
2020;46(2):432-441. doi:10.1093/schbul/sbz067
51. Jones P, Rodgers B, Murray R, Marmot M. Child development risk factors for adult
schizophrenia in the British 1946 birth cohort. *Lancet*. 1994;344(8934):1398-1402.
doi:10.1016/s0140-6736(94)90569-x
52. Lauronen E, Miettunen J, Veijola J, Karhu M, Jones PB, Isohanni M. Outcome and its
predictors in schizophrenia within the Northern Finland 1966 Birth Cohort. *Eur Psychiatry*.
2007;22(2):129-136. doi:10.1016/j.eurpsy.2006.07.001
53. Giedd JN, Blumenthal J, Jeffries NO, et al. Brain development during childhood and
adolescence: a longitudinal MRI study. *Nature Neuroscience*. 1999;2(10):861-863.
doi:10.1038/13158
54. Raznahan A, Shaw P, Lalonde F, et al. How Does Your Cortex Grow? *Journal of
Neuroscience*. 2011;31(19):7174–7177.
55. Sowell ER. Longitudinal Mapping of Cortical Thickness and Brain Growth in Normal
Children. *Journal of Neuroscience*. 2004;24(38):8223–8231.
56. Shaw P, Lerch J, Greenstein D, et al. Longitudinal mapping of cortical thickness and
clinical outcome in children and adolescents with attention-deficit/hyperactivity disorder.
Arch Gen Psychiatry. 2006;63(5):540-549. doi:10.1001/archpsyc.63.5.540
57. van Soelen ILC, Brouwer RM, van Baal GCM, et al. Genetic influences on thinning of the
cerebral cortex during development. *Neuroimage*. 2012;59(4):3871-3880.
doi:10.1016/j.neuroimage.2011.11.044
58. Vijayakumar N, Allen NB, Youssef G, et al. Brain development during adolescence: A
mixed-longitudinal investigation of cortical thickness, surface area, and volume. *Hum
Brain Mapp*. 2016;37(6):2027-2038. doi:10.1002/hbm.23154
59. Tamnes CK, Herting MM, Goddings A-L, et al. Development of the Cerebral Cortex
across Adolescence: A Multisample Study of Inter-Related Longitudinal Changes in
Cortical Volume, Surface Area, and Thickness. *J Neurosci*. 2017;37(12):3402-3412.
doi:10.1523/JNEUROSCI.3302-16.2017
60. Mills KL, Goddings A-L, Herting MM, et al. Structural brain development between
childhood and adulthood: Convergence across four longitudinal samples. *Neuroimage*.
2016;141:273–281.
61. Mills KL, Goddings A-L, Clasen LS, Giedd JN, Blakemore S-J. The developmental
mismatch in structural brain maturation during adolescence. *Dev Neurosci*. 2014;36(3-
4):147-160. doi:10.1159/000362328
62. Mills KL, Tamnes CK. Methods and considerations for longitudinal structural brain
imaging analysis across development. *Dev Cogn Neurosci*. 2014;9:172-190.
doi:10.1016/j.dcn.2014.04.004
63. Mills KL, Lalonde F, Clasen LS, Giedd JN, Blakemore S-J. Developmental changes in the
structure of the social brain in late childhood and adolescence. *Soc Cogn Affect Neurosci*.
2014;9(1):123-131. doi:10.1093/scan/nss113
64. Bridgwater M, Bachman P, Tervo-Clemmens B, et al. Developmental influences on
symptom expression in antipsychotic-naïve first-episode psychosis. *Psychol Med*.

734   Published online October 6, 2020:1-12. doi:10.1017/S0033291720003463

65. Schimmelmann BG, Michel C, Martz-Irngartinger A, Linder C, Schultze-Lutter F. Age matters in the prevalence and clinical significance of ultra-high-risk for psychosis symptoms and criteria in the general population: Findings from the BEAR and BEARS-kid studies. *World Psychiatry*. 2015;14(2):189–197.

66. Gerstenberg M, Theodoridou A, Traber-Walker N, et al. Adolescents and adults at clinical high-risk for psychosis: age-related differences in attenuated positive symptoms syndrome prevalence and entanglement with basic symptoms. *Psychol Med*. 2016;46(5):1069-1078. doi:10.1017/S0033291715002627

67. Schultze-Lutter F, Hubl D, Schimmelmann BG, Michel C. Age effect on prevalence of ultra-high risk for psychosis symptoms: replication in a clinical sample of an early detection of psychosis service. *Eur Child Adolesc Psychiatry*. 2017;26(11):1401-1405. doi:10.1007/s00787-017-0994-y

68. Schultze-Lutter F, Schimmelmann BG, Flückiger R, Michel C. Effects of age and sex on clinical high-risk for psychosis in the community. *World J Psychiatry*. 2020;10(5):101-124. doi:10.5498/wjp.v10.i5.101

69. van Erp TGM, Hibar DP, Rasmussen JM, et al. Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. *Molecular Psychiatry*. 2016;21(4):547-553. doi:10.1038/mp.2015.63

70. van Erp TGM, Walton E, Hibar DP, et al. Cortical Brain Abnormalities in 4474 Individuals With Schizophrenia and 5098 Control Subjects via the Enhancing Neuro Imaging Genetics Through Meta Analysis (ENIGMA) Consortium. *Biol Psychiatry*. 2018;84(9):644-654. doi:10.1016/j.biopsych.2018.04.023

71. Okada N, Fukunaga M, Yamashita F, et al. Abnormal asymmetries in subcortical brain volume in schizophrenia. *Mol Psychiatry*. 2016;21(10):1460-1466. doi:10.1038/mp.2015.209

72. Ching CRK, Gutman BA, Sun D, et al. Mapping Subcortical Brain Alterations in 22q11.2 Deletion Syndrome: Effects of Deletion Size and Convergence With Idiopathic Neuropsychiatric Illness. *AJP*. Published online February 12, 2020:appi.ajp.2019.1. doi:10.1176/appi.ajp.2019.19060583

73. Sun D, Ching CRK, Lin A, et al. Large-scale mapping of cortical alterations in 22q11.2 deletion syndrome: Convergence with idiopathic psychosis and effects of deletion size. *Mol Psychiatry*. Published online June 13, 2018. doi:10.1038/s41380-018-0078-5

74. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194. doi:10.1001/jama.2013.281053

75. Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface reconstruction. *Neuroimage*. 1999;9(2):179–194.

76. Fischl B, Sereno MI, Dale AM. Cortical surface-based analysis. II: Inflation, flattening, and a surface-based coordinate system. *Neuroimage*. 1999;9(2):195–207.

77. Fischl B, Dale AM. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc Natl Acad Sci U S A*. 2000;97(20):11050-11055. doi:10.1073/pnas.200033797

78. Fischl B, Salat DH, Busa E, et al. Whole brain segmentation: automated labeling of neuroanatomical structures in the human brain. *Neuron*. 2002;33(3):341–355.

79. Hibar DP, Westlye LT, Doan NT, et al. Cortical abnormalities in bipolar disorder: an MRI analysis of 6503 individuals from the ENIGMA Bipolar Disorder Working Group. *Mol Psychiatry*. 2018;23(4):932-942. doi:10.1038/mp.2017.73

80. van Rooij D, Anagnostou E, Arango C, et al. Cortical and Subcortical Brain Morphometry Differences Between Patients With Autism Spectrum Disorder and Healthy Individuals Across the Lifespan: Results From the ENIGMA ASD Working Group. *Am J Psychiatry*.

785    2018;175(4):359-369. doi:10.1176/appi.ajp.2017.17010100

786 81. Hoogman M, Bralten J, Hibar DP, et al. Subcortical brain volume differences in
787    participants with attention deficit hyperactivity disorder in children and adults: a cross-
788    sectional mega-analysis. *Lancet Psychiatry*. 2017;4(4):310–319.

789 82. Boedhoe PSW, Schmaal L, Abe Y, et al. Cortical Abnormalities Associated With Pediatric
790    and Adult Obsessive-Compulsive Disorder: Findings From the ENIGMA Obsessive-
791    Compulsive Disorder Working Group. *Am J Psychiatry*. 2018;175(5):453-462.
792    doi:10.1176/appi.ajp.2017.17050485

793 83. Radua J, Vieta E, Shinohara R, et al. Increased power by harmonizing structural MRI site
794    differences with the ComBat batch adjustment method in ENIGMA. *Neuroimage*.
795    2020;218:116956. doi:10.1016/j.neuroimage.2020.116956

796 84. Johnson WE, Li C, Rabinovic A. Adjusting batch effects in microarray expression data
797    using empirical Bayes methods. *Biostatistics*. 2007;8(1):118-127.
798    doi:10.1093/biostatistics/kxj037

799 85. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
800    Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
801    *(Methodological)*. 1995;57(1):289-300. doi:https://doi.org/10.1111/j.2517-
802    6161.1995.tb02031.x

803 86. Efron B, Stein C. The Jackknife Estimate of Variance. *Ann Statist*. 1981;9(3):586-596.
804    doi:10.1214/aos/1176345462

805 87. Efron B. *The Jackknife, the Bootstrap, and Other Resampling Plans*. Society for Industrial
806    and Applied Mathematics (SIAM, 3600 Market Street, Floor 6, Philadelphia, PA 19104);
807    1982.

808 88. Lakens D. Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-
809    Analyses. *Soc Psychol Personal Sci*. 2017;8(4):355-362.
810    doi:10.1177/1948550617697177

811 89. Hastie T, Tibshirani R. Generalized Additive Models. *Statist Sci*. 1986;1(3):297-310.
812    doi:10.1214/ss/1177013604

813 90. Hastie T, Tibshirani R. *Generalized Additive Models*. CRC Press; 1990.

814 91. Jalbrzikowski M, Larsen B, Hallquist MN, Foran W, Calabro F, Luna B. Development of
815    White Matter Microstructure and Intrinsic Functional Connectivity Between the Amygdala
816    and Ventromedial Prefrontal Cortex: Associations With Anxiety and Depression. *Biol*
817    *Psychiatry*. 2017;82(7):511-521. doi:10.1016/j.biopsych.2017.01.008

818 92. Marek S, Hwang K, Foran W, Hallquist MN, Luna B. The Contribution of Network
819    Organization and Integration to the Development of Cognitive Control. *PLoS Biol*.
820    2015;13(12):e1002328. doi:10.1371/journal.pbio.1002328

821 93. Schmaal L, Veltman DJ, van Erp TGM, et al. Subcortical brain alterations in major
822    depressive disorder: findings from the ENIGMA Major Depressive Disorder working
823    group. *Mol Psychiatry*. 2016;21(6):806-812. doi:10.1038/mp.2015.69

824 94. Rimol LM, Nesvåg R, Hagler DJ, et al. Cortical volume, surface area, and thickness in
825    schizophrenia and bipolar disorder. *Biol Psychiatry*. 2012;71(6):552-560.
826    doi:10.1016/j.biopsych.2011.11.026

827 95. Writing Committee for the Attention-Deficit/Hyperactivity Disorder, Autism Spectrum
828    Disorder, Bipolar Disorder, et al. Virtual Histology of Cortical Thickness and Shared
829    Neurobiology in 6 Psychiatric Disorders. *JAMA Psychiatry*. Published online August 26,
830    2020. doi:10.1001/jamapsychiatry.2020.2694

831 96. Del Re EC, Stone WS, Bouix S, et al. Baseline Cortical Thickness Reductions in Clinical
832    High Risk for Psychosis: Brain Regions Associated with Conversion to Psychosis Versus
833    Non-Conversion as Assessed at One-Year Follow-Up in the Shanghai-At-Risk-for-
834    Psychosis (SHARP) Study. *Schizophr Bull*. Published online September 14, 2020.
835    doi:10.1093/schbul/sbaa127

836   97.   Borgwardt SJ, McGuire PK, Aston J, et al. Structural brain abnormalities in individuals
837   with an at-risk mental state who later develop psychosis. *Br J Psychiatry Suppl*.
838   2007;51:s69-75. doi:10.1192/bjp.191.51.s69
839   98.   Pantelis C, Velakoulis D, Mcgorry PD, et al. Neuroanatomical abnormalities before and
840   after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *Lancet*.
841   2003;361(9354):281–288.
842   99.   Takahashi T, Wood SJ, Yung AR, et al. Progressive gray matter reduction of the superior
843   temporal gyrus during transition to psychosis. *Arch Gen Psychiatry*. 2009;66(4):366-376.
844   doi:10.1001/archgenpsychiatry.2009.12
845   100.   Lee CU, Shenton ME, Salisbury DF, et al. Fusiform gyrus volume reduction in first-
846   episode schizophrenia: a magnetic resonance imaging study. *Arch Gen Psychiatry*.
847   2002;59(9):775-781. doi:10.1001/archpsyc.59.9.775
848   101.   Onitsuka T, Shenton ME, Kasai K, et al. Fusiform gyrus volume reduction and facial
849   recognition in chronic schizophrenia. *Arch Gen Psychiatry*. 2003;60(4):349-355.
850   doi:10.1001/archpsyc.60.4.349
851   102.   Takahashi T, Suzuki M, Zhou S-Y, et al. Temporal lobe gray matter in schizophrenia
852   spectrum: a volumetric MRI study of the fusiform gyrus, parahippocampal gyrus, and
853   middle and inferior temporal gyri. *Schizophr Res*. 2006;87(1-3):116-126.
854   doi:10.1016/j.schres.2006.04.023
855   103.   van Lutterveld R, van den Heuvel MP, Diederen KMJ, et al. Cortical thickness in
856   individuals with non-clinical and clinical psychotic symptoms. *Brain*. 2014;137(Pt
857   10):2664-2669. doi:10.1093/brain/awu167
858   104.   Sasabayashi D, Takayanagi Y, Takahashi T, et al. Subcortical Brain Volume
859   Abnormalities in Individuals With an At-risk Mental State. *Schizophr Bull*. 2020;46(4):834-
860   845. doi:10.1093/schbul/sbaa011
861   105.   Pontious A, Kowalczyk T, Englund C, Hevner RF. Role of intermediate progenitor cells in
862   cerebral cortex development. *Dev Neurosci*. 2008;30(1-3):24-32. doi:10.1159/000109848
863   106.   Rakic P. Specification of cerebral cortical areas. *Science*. 1988;241(4862):170–176.
864   107.   Grasby KL, Jahanshad N, Painter JN, et al. The genetic architecture of the human
865   cerebral cortex. *Science*. 2020;367(6484). doi:10.1126/science.aay6690
866   108.   Huttenlocher PR. Synaptic density in human frontal cortex - developmental changes and
867   effects of aging. *Brain Res*. 1979;163(2):195-205. doi:10.1016/0006-8993(79)90349-4
868   109.   Rakic P, Bourgeois JP, Eckenhoff MF, Zecevic N, Goldman-Rakic PS. Concurrent
869   overproduction of synapses in diverse regions of the primate cerebral cortex. *Science*.
870   1986;232(4747):232-235. doi:10.1126/science.3952506
871   110.   Petanjek Z, Judas M, Kostović I, Uylings HBM. Lifespan alterations of basal dendritic
872   trees of pyramidal neurons in the human prefrontal cortex: a layer-specific pattern. *Cereb
873   Cortex*. 2008;18(4):915-929. doi:10.1093/cercor/bhm124
874   111.   Natu VS, Gomez J, Barnett M, et al. Apparent thinning of human visual cortex during
875   childhood is associated with myelination. *Proc Natl Acad Sci U S A*. 2019;116(41):20750-
876   20759. doi:10.1073/pnas.1904931116
877   112.   Pantelis C, Yücel M, Wood SJ, et al. Structural brain imaging evidence for multiple
878   pathological processes at different stages of brain development in schizophrenia.
879   *Schizophr Bull*. 2005;31(3):672-696. doi:10.1093/schbul/sbi034
880   113.   Sun D, Stuart GW, Jenkinson M, et al. Brain surface contraction mapped in first-episode
881   schizophrenia: a longitudinal magnetic resonance imaging study. *Mol Psychiatry*.
882   2009;14(10):976-986. doi:10.1038/mp.2008.34
883   114.   Sellgren CM, Gracias J, Watmuff B, et al. Increased synapse elimination by microglia in
884   schizophrenia patient-derived models of synaptic pruning. *Nat Neurosci*. 2019;22(3):374-
885   385. doi:10.1038/s41593-018-0334-7
115.   Whitaker KJ, Vértes PE, Romero-Garcia R, et al. Adolescence is associated with

genomically patterned consolidation of the hubs of the human brain connectome. *Proc*
Natl Acad Sci U S A. 2016;113(32):9105-9110. doi:10.1073/pnas.1601745113

116.   Parker N, Patel Y, Jackowski AP, et al. Assessment of Neurobiological Mechanisms of
Cortical Thinning During Childhood and Adolescence and Their Implications for
Psychiatric Disorders. *JAMA Psychiatry*. Published online June 17, 2020.
doi:10.1001/jamapsychiatry.2020.1495

117.   Perkins DO, Olde Loohuis L, Barbee J, et al. Polygenic Risk Score Contribution to
Psychosis Prediction in a Target Population of Persons at Clinical High Risk. *Am J*
Psychiatry. 2020;177(2):155-163. doi:10.1176/appi.ajp.2019.18060721

118.   Mei C, Nelson B, Hartmann J, Spooner R, McGorry PD. Chapter 4 - Transdiagnostic early
intervention, prevention, and prediction in psychiatry. In: Baune BT, ed. *Personalized*
Psychiatry. Academic Press; 2020:27-37. doi:10.1016/B978-0-12-813176-3.00004-3

119.   McGorry PD, Hartmann JA, Spooner R, Nelson B. Beyond the "at risk mental state"
concept: transitioning to transdiagnostic psychiatry. *World Psychiatry*. 2018;17(2):133-
142. doi:10.1002/wps.20514

120.   Dazzan P, Soulsby B, Mechelli A, et al. Volumetric abnormalities predating the onset of
schizophrenia and affective psychoses: an MRI study in subjects at ultrahigh risk of
psychosis. *Schizophr Bull*. 2012;38(5):1083-1091. doi:10.1093/schbul/sbr035

121.   Ziermans TB, Schothorst PF, Schnack HG, et al. Progressive structural brain changes
during development of psychosis. *Schizophr Bull*. 2012;38(3):519-530.
doi:10.1093/schbul/sbq113

122.   Walter A, Studerus E, Smieskova R, et al. Hippocampal volume in subjects at high risk of
psychosis: a longitudinal MRI study. *Schizophr Res*. 2012;142(1-3):217-222.
doi:10.1016/j.schres.2012.10.013

123.   Valli I, Marquand AF, Mechelli A, et al. Identifying Individuals at High Risk of Psychosis:
Predictive Utility of Support Vector Machine using Structural and Functional MRI Data.
Front Psychiatry. 2016;7:52. doi:10.3389/fpsyt.2016.00052

Table and Figure Legends

Table 1. Demographic descriptives for every site in HC and CHR, and CHR-PS+ and CHR-PS-. The Ns for CHR-PS+ and CHR-PS- do not always sum to the sample CHR N because most sites lose individuals to follow-up (CHR-UNK).

Figure 1. Effect sizes for mega-analysis of group and conversion status. **A.** Overview of effect sizes for HC vs. CHR. The top row reflects the results of the overall generalized linear model. A deeper purple color indicates a greater effect of group (HC vs. CHR) in this region. We observed the greatest effects of group in cortical thickness measures. The second row indicates the pairwise effect sizes for HC vs. CHR, in regions that were statistically significant ($q<0.05$) in the overall comparison (top row of **A.**). Regions that were not statistically significant in the overall comparison are in gray. In comparison to HC, CHR exhibited lower cortical thickness across the cortex. Red color indicates that HC has a larger value in comparison to CHR for this region. **B.** Overview of effect sizes for HC vs. CHR-PS+ vs. CHR-PS-. The top row reflects the results of the overall generalized linear model. A deeper purple color indicates a greater effect of conversion status (HC vs. CHR-PS+ vs. CHR-PS-) in this region. The second and third rows indicate the pairwise effect sizes for HC vs. CHR-PS+ and CHR-PS- vs. CHR-PS+, respectively. Pairwise comparisons are presented in regions that were statistically significant ($q<0.05$) in the overall comparison (top row of **B.**). Regions that were not statistically significant in the overall comparison are in gray. Regions that CHR-PS+ had lower cortical thickness in comparison to HC and CHR-PS-, CHR-PS+ are highlighted in yellow. **C.** Bar graphs for regions in which CHR-PS+ (pink) had lower CT in comparison to CHR-PS- (purple) and HC (green).

Figure 2. Age effects of regions that exhibited an effect of conversion status. HC are in green, CHR-PS+ are pink, and CHR-PS- are purple. Each graph has a line of best fit for the effect of age on the respective neuroimaging measures. Shading around the line indicates the standard error. The bars underneath the age plots reflect the derivative of the slope, i.e., the rate of change taking place at a particular age, scaled as a pseudo *t*-statistic, based on the posterior simulation. Age effects are plotted for A. fusiform cortical thickness, B. paracentral cortical thickness, and C. superior temporal cortical thickness.

Figure 3. Cortical thickness alterations in CHR-PS+, schizophrenia, and 22q.11DS **A.** Magnitude of cortical thickness alterations for the three groups. Whole-brain cortical thickness effect sizes were not reported for SZ and 22q.11DS in original publications; values shown are the hemispheres with the smallest effect size. Colored bars represent standard error. **B.** Spearman's rank correlations between cortical thickness alterations in CHR-PS+ and individuals with schizophrenia (top) and 22q.11 DS with psychosis (bottom).

A

Cortical Thickness   Surface Area   Subcortical Volume

L   R   L   R   L   R

HC vs. CHR   *F* 0 5 10 15 20 25

HC vs. CHR   *d* −0.1 0.0 0.1

B

Cortical Thickness   Surface Area   Subcortical Volume

L   R   L   R   L   R

HC vs. CHR-PS+ vs. CHR-PS−   *F* 0 5 10 15

HC vs. CHR-PS+   *d* −0.3 −0.2 −0.1 0.0 0.1 0.2 0.3

CHR-PS− vs. CHR-PS+

C

HC (N=1377)
CHR-PS− (N=1234)
CHR-PS+ (N=253)

Fusiform Cortical Thickness   Paracentral Cortical Thickness   Superior Temporal Cortical Thickness

+ *p* <.10   * *p* <.05   ** *p* ≤0.01   *** *p* ≤.001

A Age vs. Fusiform Cortical Thickness

B Age vs. Paracentral Thickness

C Age vs. Superior Temporal Thickness

Table 1. Age and sex information for healthy controls (HC) and clinical high risk for psychosis (CHR) participants at each site

Site	HC				CHR				CHR-PS+				CHR-PS-				Transition Rate	Mean Follow-up Length (SD)
	N	% F	Mean Age (SD)	Age Range	N	% F	Mean Age (SD)	Age Range	N	% F	Mean Age (SD)	Age Range	N	% F	Mean Age (SD)	Age Range		
Amsterdam	23	57	23.4 (2.8)	19-32	16	62	23.6 (2.5)	20-30	0	•	•	•	16	62	23.6 (2.5)	20-30	•	24 (0)
Barcelona-HSJD	44	41	15.6 (1.6)	13-18	55	58	15.2 (1.6)	11-18	7	43	15.7 (2.4)	12-18	35	66	15.2 (1.5)	11-18	16.7%	17.4 (4.8)
Bern	17	35	20.2 (6.1)	13-34	43	51	18.5 (5.1)	10-35	0	•	•	•	19	58	16.8 (3.8)	10-26	•	15.8 (5.7)
Columbia1	9	22	24.4 (4.1)	20-33	17	53	22.9 (5)	15-31	3	67	19.4 (7.3)	15-28	14	50	23.6 (4.3)	16-31	17.6%	20.6 (10.3)
Columbia2	15	60	26.2 (2.2)	22-30	19	37	23.3 (3.6)	18-31	•	•	•	•	•	•	•	•	•	•
Columbia3	37	32	22.9 (3.6)	15-30	58	26	21 (4)	14-29	14	29	21.1 (4.8)	15-29	44	25	20.9 (3.8)	14-27	24.1%	24 (0)
Copenhagen	59	49	24.8 (3.3)	20-35	163	53	24.2 (4.2)	18-38	13	38	23.1 (3.4)	18-29	95	57	24.5 (4.6)	18-38	12%	11.8 (1.2)
CSU	59	42	21.5 (3.1)	15-30	52	46	19.5 (5)	13-35	21	57	19.4 (5.1)	13-35	31	39	19.5 (5)	13-30	40.4%	14.7 (7.6)
Glasgow	46	65	22.8 (3.6)	18-32	80	79	22.2 (4.8)	17-34	6	83	18.5 (1.9)	17-22	74	78	22.5 (4.8)	17-34	7.5%	20.2 (9.9)
Heidelberg	33	48	15.7 (0.9)	14-17	22	59	15.1 (1.1)	14-17	0	•	•	•	18	67	15.1 (1)	14-17	•	9.7 (3.3)
IDIBAPS	54	65	15.9 (1.6)	11-18	74	66	15.3 (1.7)	10-18	17	65	15.1 (1.4)	13-17	42	67	15.5 (1.9)	10-18	28.8%	14.3 (5.5)
ISMMS	12	42	27.9 (3.7)	23-35	25	52	23.4 (5.4)	17-35	1	0	26.3 (0)	26-26	12	58	23.6 (5.5)	17-35	7.7%	6 (0)
London	29	34	24.5 (4.7)	20-36	81	31	22.6 (4.4)	18-38	6	0	24 (4.4)	18-29	62	32	22.3 (4.6)	18-38	8.8%	18.8 (8.7)
Maastricht	38	32	25.6 (5.7)	18-39	48	29	20.2 (4.1)	12-29	6	33	20.5 (5.2)	15-26	25	32	18.6 (3.4)	12-27	19.4%	•
Melbourne	92	46	21.9 (3.6)	15-30	249	50	19.7 (3.4)	14-29	71	44	19.5 (3.5)	14-29	165	52	19.8 (3.4)	14-28	30.1%	71.4 (53.7)
Mexico City	38	26	20.9 (3.4)	15-28	33	21	19.5 (4.1)	13-31	7	29	20.4 (6)	15-31	26	19	19.3 (3.6)	13-27	21.2%	24 (0)
MHRC	51	0	22.2 (2.8)	16-27	38	0	20.1 (2.5)	16-28	3	0	19 (2.4)	16-21	35	0	20.2 (2.5)	17-28	7.9%	30.2 (6.6)
MPRC	20	40	17.8 (4.3)	12-24	31	52	17 (3.2)	12-22	3	33	16 (4.4)	13-21	10	50	15.8 (2.7)	12-20	23.1%	7.1 (1.1)
Newcastle	17	76	20.3 (1.9)	17-24	45	53	19.6 (2.2)	15-24	1	0	17.4 (0)	17-17	41	54	19.6 (2.2)	15-24	2.4%	21.5 (12)
Oslo Region	63	38	19.8 (3.6)	15-29	21	38	19.9 (3.6)	16-29	2	50	20 (5.4)	16-24	18	39	19.9 (3.7)	16-29	10%	12.5 (1.4)
Pitt	65	40	23 (5.9)	14-40	26	54	20.8 (5.3)	12-36	2	50	17.1 (0.8)	16-18	11	64	19.5 (6.4)	12-36	15.4%	13.1 (3.6)
RUMC	29	66	23.9 (3.2)	18-30	67	46	22.7 (3.9)	16-30	4	50	24.5 (2.9)	21-27	60	47	22.6 (3.9)	16-30	6.2%	8.3 (4.5)
Singapore	53	47	22 (4.2)	14-30	100	32	21.9 (3.6)	14-30	11	27	20.1 (3.1)	15-26	88	32	22.1 (3.6)	14-30	11.1%	18.9 (5.9)
SNUH	74	32	21.2 (2.5)	17-27	74	26	20.7 (3.8)	15-34	9	33	22 (5)	16-33	46	24	20.4 (3.7)	15-34	16.4%	30.9 (17.2)
Stavanger	33	48	17 (3.1)	13-27	37	59	16.6 (2.4)	13-25	5	80	16.2 (2.7)	13-19	31	58	16.7 (2.3)	14-25	13.9%	24 (0)
Toho	16	50	23.2 (2.9)	18-28	40	70	23.7 (6.9)	13-39	4	75	19 (4.4)	14-24	36	69	24.2 (7)	13-39	10%	12 (0)
Tokyo	25	48	22.1 (2.8)	16-25	39	46	20.9 (3.5)	14-29	3	0	22.7 (4.7)	19-28	25	60	20.5 (3.1)	14-27	10.7%	26.5 (12.1)
Toronto	39	41	25.5 (5.2)	18-38	27	44	20.8 (1.8)	18-27	0	•	•	•	4	50	19.7 (1.1)	18-21	•	24 (0)
Toyama	141	48	25.1 (4.2)	18-38	79	44	18.5 (4)	13-31	10	30	20.8 (5.7)	15-31	69	46	18.1 (3.7)	13-30	12.7%	37.4 (34.2)
UCSF	103	42	23.7 (7.5)	13-40	71	48	19.3 (4.4)	12-32	13	38	21.4 (4.4)	16-29	47	49	19 (4.1)	12-29	21.7%	21 (5.4)
Zurich	43	51	22.2 (5.6)	13-36	62	40	19.1 (4.9)	13-35	11	18	18.5 (3.4)	14-24	35	40	19.1 (5.3)	13-35	23.9%	

Follow-up times reported in months. "•" Indicates that data was not available, either because no CHR subjects transitioned to a psychotic disorder or follow-up data was not available. Site name abbreviations as follows: Amsterdam = Vrije Universiteit Amsterdam, The Netherlands; Barcelona-HSJD

= Hospital Sant Joan de Déu Barcelona, Barcelona, Spain; Bern = University Hospital of Child and Adolescent Psychiatry and Psychotherapy, University of Bern, Switzerland; Columbia1 = New York State Psychiatric Institute, Columbia University, USA; Columbia2 = New York State Psychiatric Institute, Columbia University, USA; Columbia3 = New York State Psychiatric Institute, Columbia University, USA; Copenhagen = Mental Health Center Copenhagen and CINS, Mental Health Center Glostrup, University of Copenhagen, Denmark; CSU = Central South University, China; Glasgow = Institute of Neuroscience and Psychology, University of Glasgow, Scotland; Heidelberg = Heidelberg University Hospital, Germany; IDIBAPS = August Pi i Sunyer Biomedical Research Institute, Barcelona, Spain; ISMMS = Icahn School of Medicine at Mount Sinai, USA; London = Institute of Psychiatry, Psychology & Neuroscience, King's College London, UK; Maastricht = Maastricht University, The Netherlands; Melbourne = University of Melbourne, Australia; Mexico City = Instituto Nacional de Neurología y Neurocirugía, Mexico City, Mexico; MHRC = Mental Health Research Center Moscow, Russia; MPRC = Maryland Psychiatric Research Center, University of Maryland School of Medicine, USA; Newcastle = University of Newcastle, Australia; Oslo Region = NORMENT, University of Oslo and Oslo University Hospital, Norway; Pitt = University of Pittsburgh, USA; RUMC = Rush University Medical Center, Chicago, IL USA; Singapore = Institute of Mental Health, Singapore and National University of Singapore, Singapore; SNUH = Seoul National University, South Korea; Stavanger = Stavanger University Hospital, Norway; Toho = Department of Neuropsychiatry, Toho University School of Medicine, Japan; Tokyo = Department of Neuropsychiatry, Graduate School of Medicine, The University of Tokyo; Toronto = Centre for Addiction and Mental Health, University of Toronto, Canada; Toyama = University of Toyama Graduate School of Medicine and Pharmaceutical Sciences, Japan; UCSF = University of California, San Francisco, USA; Zurich = Psychiatric Hospital, University of Zurich, Switzerland