Convergence of Comorbidity and COVID-19 Infection to Fatality: An Investigation Based on Concurrent Health Status Evaluation among the Elderly Population in Kerala

Sindhu Joseph[1,*], Jijo Pulickiyil Ulahannan[2,3]

[1]Department of Travel and Tourism Management, Govinda Pai Memorial Government College, Kannur University, Kerala, India

[2]Department of Physics, Government College Kasaragod, Kannur University, Kerala, India

[3] Collective for Open Data Distribution-Keralam (CODD-K)

Abstract

The COVID-19 infection rapidly spread globally, mostly affecting the extremely vulnerable category of the elderly with comorbidities. There are inconsistencies in the findings on the type of comorbidity of the elderly and its association with fatalities. In this context, this research investigated the impact of comorbidity in the fatality of elderly COVID-19 patients in Kerala based on their healthcare status, functionality, and morbidity profiles. A concurrent mixed-method approach was adopted for the study to achieve the objectives, where the quantitative and qualitative data had been collected in the COVID-19 situation, from June to November 2020. This study's findings have been further triangulated with the COVID-19 elderly fatality data, which is available from the crowdsourced dashboard of the research team and two other volunteering dashboards. This paper establishes that comorbidities can predict potential fatality among elderly COVID-19 patients. While facing an epidemic like the present zoonotic disease, better knowledge of these high-risk factors will help clinicians to pinpoint the situation and

implement therapeutic and preventive methodologies and interventions. The comorbidity level of the elderly in Kerala matches with the profile of COVID-19 death cases where heart disease, diabetes, cancer, and hypertension are the significant predictors of COVID-19 elderly fatality in Kerala.

Word count: Abstract: 200, Main text: 8445

Keywords: COVID-19, Kerala, Elderly, Comorbidity, Mortality, Fatality, Convergence

*Corresponding Author: Sindhu Joseph Ph.D., GPM Government College Manjeshwar, Kannur University, Kerala, India 671323, sjsindhu@gmail.com, +91.9400034634

INTRODUCTION

It is already acknowledged that health conditions deteriorate when age increases and aging goes hand in hand with many psychological, physical, and behavioral issues and disabilities, contagious infections, lifestyle diseases, and chronic diseases. In later ages physical immobilities will increase due to degenerative diseases. Aging associated 'multi-morbidity' is of higher risk where multiple chronic conditions are experienced simultaneously.[1] Consequently, the rate of health-care utilization and health care expenses may be significantly greater among elderly[2] who might receive aggressive and expensive treatments to prolong their lives. Therefore, the impact of multi-morbidity on various aspects such as quality of life, functionality and risk of mortality becomes a matter of present discussions worldwide.

The latest research reveals that the elderly suffers from non-communicable diseases (NCDs).[3] The elderly face many health issues such as loss of appetite, gastric pain, cataracts, toothache and body pain, arthritis, uterine problems, fever, and general weakness.[4] Muscular-skeletal diseases have been reported as a common problem of the elderly, mainly from African countries.[5] Hypertension becomes the common morbidity, followed by respiratory diseases, Diabetes Mellitus and stroke[6] and the same findings have been reported from Korea and Malaysia.[7] Table 1 shows the prevalence of different levels of disability among the elderly (over 60 years) in both developed as well as developing countries.[8]

In India, geriatric care has been adversely affected by the population size leading to the scarcity of resources and the inadequacy of the public healthcare system in providing

equitable healthcare. In 2001, in India, around 33.1 percent of the elderly population reported with unmet healthcare needs after losing their spouses.[9] Prevailing threats due to nutritional, maternal, infectious, and perinatal conditions were added with the "triple burden of disease" involving infectious and chronic illnesses and gender-based violence and injury.[10] Elderly in India faces NCDs' threat, including cancers, cardiovascular diseases, respiratory diseases, eyesight problems, and diabetes. In India, particularly in urban areas, one out of every two elderly suffers from at least one chronic disease requiring life-long medication.[11] Tiwari et al.,[12] found that a great majority of the inmates in nursing homes were suffering from mental health issues, including depression, and had one or more related physical morbidity(ies). Arokiasamy,[13] stated that, in India, one-third of the elderly suffer from either short term ailments or endure hospitalization where short-duration ailments are common (4 out of 10 urban elderly) among elderly women, and they increase with age.

Kerala has the highest outpatient doctor consultation per 1,000 population compared to rest of the states in India, and this trend has a more robust economic status gradient for the older people than their conterparts.[14] Higher hospitalization rates indicate higher morbidity levels among the elderly in Kerala.[15] The prominent reasons for the highest rate of morbidity are the high literacy rate induced healthcare awareness, better health care access due to the existence of the private and public sector, and larger share of the elderly population due to demographic transition.[16] In a 2004 study, about 25 percent of the elderly population reported with poor health status, while only 5 percent were reported with excellent health status.[13] The common ailments of the elderly in

Kerala, in a 2010 study, were paralysis and urinary problems followed by heart disease and cancer.[17] However, in 2011, hypertension was the most widely prevalent illness, particularly among women, followed by diabetes mellitus, while cataract and heart disease were found mostly among men.[15] Public Health of India[18] reported that CVD (Cardiovascular Disease) was the most prevalent morbidity among more than 22% of the males and 20% of the females in 2016, followed by diabetes, musculoskeletal, and respiratory disease. A study on the prevalence of NCDs during 2016–2017 also found that diabetes, abdominal obesity, and hypertension were the most prevalent morbidities.[19] To support this, in 2020, a report found that about 70 lakh people aged more than 65 years have cancer, diabetes, hypertension, kidney disease, heart diseases, and respiratory tract infections.[20] Based on all the previous studies, it can be assumed that the significant prevalent diseases in Kerala are Hypertension, Diabetes, Heart Diseases, Cancer, and Respiratory Diseases.

ELDERLY AND THE COVID-19 FATALITY

The outspread of COVID-19 affected mostly the elderly. The Case Fatality Rate (CFR) among them is disproportionately higher.[21,22] Guan et al.,[23] analyzed the composite end-points of 1590 confirmed hospitalized COVID-19 cases in China and found that patients with any comorbidity generated worse clinical outcomes than their counterparts. Patients with chronic comorbidities, including malignancy, cardiovascular disease, kidney disease, diabetes, hypertension, and respiratory disease are prone to the severe or fatal outcome of COVID-19 infection.[24–26]

Using seven databases, Wang et al.[27] performed a systematic review of the literature with meta-analysis and found that the most common pre-existing comorbidities in COVID-19 patients were hypertension. Similar results have been obtained by others.[23] It is also found that those with pre-existing lungs and respiratory disease would be susceptible to severe SARS-CoV-2 infection as it can progress to Acute Respiratory Distress Syndrome (ARDS), and potentially even death.[28] However, in China, a study of 548 COVID-19 patients revealed that the prevalence of asthma was just 0.90% .[29] In New York and China, diabetic patients were seriously infected with the COVID-19 virus[25,30,31] as they are more vulnerable to develop pneumonia, and they account for 11.7% of severe COVID-19 cases.[28] Nevertheless Zhou et al.,[25] found that among the comorbidities leading to higher mortality rate, cardiovascular disease was most prevalent among COVID-19 patients. Various studies[25,32,33] found that for cardiovascular disease and hypertension , the use of renin-angiotensin system inhibitors may accelerate the susceptibility to SARS-CoV-2 infection. Emami et al.,[34] had done a meta-analysis of 76,993 patients and found that the susceptibility to SARS-CoV-2 of cancer patients is low as malignancy was prevalent in just 0.92% of reported comorbidities. Similarly, in China, a countrywide analysis reported that only 1.1% of COVID-19 patients had active cancer.[35]

Despite the outnumbered post-COVID-19 studies describing comorbidities and the consequent poor clinical outcomes leading to fatality, the results are inconsistent. Throughout the COVID-19 pandemic spread, wide variations in CFR and IFR estimates have been noted, which are misleading.[36] Likewise, the study findings related to the

health status, comorbidity profile, and the NCDs scenario of the elderly population in Kerala are inconsistent as different reports give varied results over a decade. In this context, this study was intended to fill the above gaps and planned to relate the healthcare status of the elderly, their disease profile, and functionality level in the post-COVID-19 scenario to the mortality rate of COVID-19, which will help to identify the severity of risks. Kerala is a state that has gained acclaim for its healthcare indicators and preventive and curative measures for COVID-19. Thus, it is essential to draw a clear picture of the health status and morbidity levels of the elderly amidst the pandemic and to assess whether the comorbidity level has converged with the COVID-19 infection.

This study is unique as it measured the healthcare status and the comorbidity level of the elderly in Kerala, while the pandemic is ongoing with simultaneous assessment of the elderly COVID-19 deaths due to the suspected comorbidity levels. To check and identify a consistent pattern in the association between age and COVID-19 morbidity or mortality, an extensive review of literature was carried. The study insights will help to shape the geriatric treatment policies and preventive measures for infectious diseases. This study has implications for the state healthcare system and policy efforts to expand the healthcare facilities and programs for the elderly, who contribute about 20% of the total population of Kerala[37] and are projected to grow at a growth rate of 2.45 percent.[38]

METHODOLOGY

This research is based on mixed methodology, using two different data collection methods based on positivist and interpretivist paradigms. The study used a concurrent timing strategy where both the quantitative and qualitative strands occurred during a single phase, from June 2020 to November 2020. Quantitative data was collected from 405 elderly from 36 panchayats of Kerala, and 102 in-depth interviews were carried out with the elderly from all over Kerala.

Quantitative data had been collected using a structured questionnaire adopting multi-stage sampling technique. The study area was divided into three, South Zone (Thiruvananthapuram, Kollam, Alappuzha, and Pathanamthitta districts), Central Zone (Idukki, Kottayam, Ernakulam, Thrissur, and Palakkad districts), and North Zone (Malappuram, Wayanad, Kozhikode, Kannur, and Kasaragod districts). From each of the zones mentioned above, three districts had been selected randomly, and hence Kasaragod, Wayanad and Malappuram (North Zone), Kottayam, Idukki, and Ernakulam (Central Zone) and Kollam, Thiruvananthapuram, and Alappuzha (Southern Zone) districts were selected at random. Four panchayats from each of these nine districts were selected by drawing lots, making 36 panchayats of Kerala under study. From each selected panchayath, 11 households with elderly persons were selected randomly. The respondents to the survey questionnaire included any one of the resident members above the age of 60, who could respond to the questionnaire. When there were two older adults in one household, only one person was included in the study. The data were analyzed with SPSS 2.0.

Using a panchayat wise list of the elderly population, received from the *Anganvadi* workers, *Asha* workers, and *Vayomithram* co-ordinators, elderly persons above 60 were selected randomly for the qualitative interview. The interviews were conducted at their households for about 30 - 45 minutes. An average of seven interviews per district was conducted making 102 in-depth interviews from all the fourteen districts of Kerala. The participants were met between June to November 2020. They were asked about the present ailments and health behavior (exercise, diet, smoking, use of pan and alcohol, morbidities). To further enrich the data, 12 interviews were conducted with *Vayomithram* (a government project for holistic care of elderly in Kerala) district coordinators of 12 districts who assume the charge to ensure good health and quality of life to the elderly.

The COVID-19 outspread was progressing in Kerala from January 2020 and has been continuing with a higher mortality rate in the later phases, particularly from June 2020. The Government of Kerala dashboard (https://dashboard.kerala.gov.in) and the Directorate Health Services (DHS) website (http://dhs.kerala.gov.in) give comprehensive updates and daily bulletins about the COVID-19 spread. However, access to primary data available on the website is not complete, and critical factors such as age, gender, comorbidity status of patients are not complete, and therefore, a multi-disciplinary team of experts compiled data through a citizen science initiative, led by the researchers of this study. Publicly available and crowd-verified data have been used to track the outbreak of COVID-19 along with the help of government bulletins

(tps://dashboard.kerala.gov.in/deaths.php) and media reports to produce reusable datasets (https://doi.org/10.5281/zenodo.3818096).[39,40]

To supplement the information available from the above sources, another volunteering dashboard (https://t.co/5e70Wu2dZz) was also analyzed for getting information that is not available on the official website of the government or elsewhere. This website gives accurate information on the actual fatality rate, especially when it has been widely argued that the official data on death cases attributed to COVID-19 are seemingly an underestimation because the official mortality data do not incorporate the cases which are not directly attributable to coronavirus 2 (SARS-CoV-2).

A systematic analysis of the above mentioned three dashboards providing reliable information on CFR and comorbidities was done. The data was extracted from the death cases reported in Kerala with COVID-19 infection between January 2020 and December 27, 2020 (Fig. 1), including a total of 2232 confirmed elderly COVID-19 patient deaths (Table 8). The quantitative and qualitative strands were mixed and integrated at the point of interpretation and drawn inferences by comparing and synthesizing the results of quantitative as well as qualitative data on elderly healthcare status, disease profile, and functionality level with the fatality of the elderly from the meta-analysis of COVID-19 dashboards.

This study adopted the definition of WHO[41] for COVID-19 death "as a death resulting from a clinically compatible illness in a probable or confirmed COVID-19 case unless

there is a clear alternative cause of death that cannot be related to COVID-19 disease (e.g. trauma)." Before the qualitative and quantitative data collection, the participants were briefed about the study's purpose, and informed consent was obtained. COVID-19 protocol was adhered to while collecting the data.

RESULTS

Quantitative Study

Socio-Demographic Variables

Out of the 405-elderly surveyed (see Table 2), 224 (55.3%) were males, and 181 (44.7%) were females. Around 50.1% (203) were in the 60-69 age category, followed by 31.6% (128) in the 70 -79 category. 77.5% (314) were married, 2.7% (11) were not married, and 18.5% (65) lost their life partners. The educational status of the elderly showed that 18.3% (74) does not have any formal education, and a large majority (267) has school education (65.9%). Around 25.9% (125) have no income. 56.3% (228) has a monthly income of less than INR 10,000. 52.1% (211) have some pension, particularly the old-age pension, INR 1,400 per month.

Health Condition and Healthy Behaviors

The self-evaluation of the health condition of the elderly is very significant as it is associated with physical as well as mental impacts. Health status and disability levels are essential factors to determine the overall well-being of the elderly.[15] 7.7% of the elderly population are in 'excellent' health condition, 20.7 % are in 'good' health, 44.9 %

have 'satisfactory' health, 20.7% have 'bad' health, and 5.9% have 'very bad' health condition. It shows that the majority of the elderly perceive their health as above average.

Health Behavior

Health behavior is a predictor of good health. Table 3 shows that only 15.8 % exercise regularly, 16.8% exercise frequently, and 12.8% exercise sometimes, 9.9% exercise rarely, and 44.4% do not exercise at all. In dietary habits, 31.9% have a regular healthy diet, 28.9% have frequently followed a healthy diet, and 13.6% do not bother about having a healthy diet in their lives. The risky habits such as smoking, consumption of alcohol, chewing tobacco have not impacted their lives much as 82% never smoke, 85.4% never consume alcohol, and 90.9% never chew tobacco. The previous study reported that 31.6% had alcohol consumption, 21.3% had smoked regularly, and 2.3% had the habit of chewing tobacco[19] which is almost consistent with the present study findings and the small variance occurred as the latter study had been carried out among the entire population of Kerala.

Functionality Level

Quality of life depends mainly on health status and functionality level. This construct was studied using two latent constructs such as Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL). ADLs are regular activities people do every day, usually without assistance when they are in good health. Basically, there are six basic ADLs: eating, bathing, toileting, mobility, getting dressed, and continence. Instrumental Activities of Daily Living (IADLs) is the ability of a person to care for herself

or himself, and that includes the set of activities for independent living and are important for assessing the early-stage disease of a person.

ASSISTANCE REQUIRED – ADL

Table 4 shows that most of the elderly can do their own daily personal activities. About 78-90% of the elderly can perform their basic needs, such as eating, dressing, and toileting, without any assistance.

ASSISTANCE REQUIRED - IADLs

As shown in Table 5, about 40-65% of the elderly do not require any assistance for their instrumental activities. Most of them can use telephones (52%). The elderly need assistance in activities that require physical strength. Some of the activities get higher responses (e.g., cooking), perhaps due to more men's inclusion in the study. Hence it can be concluded that the elderly in Kerala have an above-average functionality.

ELDERLY DISEASE PROFILE

Table 4 shows that an average of 64% of the elderly suffers from Hypertension, 51.4% from Diabetes, 42% from Rheumatic Disease, 37.5% from Cataract, 31.4% from Urinary Disease, 27.4 % from Abdominal Disease, 22.7% from Heart Disease, 19.5% from Asthma, 18.3% from Depression 6.7% from Cancer. Many of them have two or more diseases (Comorbidities).

Results-Qualitative Data

This study's key findings were also based on qualitative responses of the elderly to the interview questions considering their disease profile. Analysis of data collected used a grounded theory research design,[42] where the approach intends to develop concepts, insights, or models grounded in the participant's data.[43]

Most of the participants in the study eat food in minimal quantities. They were practicing diet control, as most of them had lifestyle diseases like blood pressure (Hypertension), Diabetes, Heart Disease, Cancer, and Cholesterol. They try to avoid sugar and fried items from their daily menu.

Vayomithram coordinators also shared the same view about the disease profile of the elderly. The following excerpts validate the finding;

……, most of them come to the clinic to check their BP (Hypertension) and sugar (Diabetes)……….

……….many are having a cardiac disease which we (do not give medicines), and they have Diabetes, BP, and cholesterol. We do not give them medicine for cancer….only palliative care.

…..

Covid-19 Deaths in Kerala-Age wise

Fig. 1 shows that age plays a vital role in the increased mortality rate. Till December 27, 2020, 2976 COVID-19 deaths happened in Kerala, of which 2232 (75.0%) deaths were elderly, where the majority belongs to the 60-69 age group.

Elderly COVID-19 Deaths and Comorbidity

As of September 30, 2020, out of the 525 elderly persons who died, 209 of them (¬40%) had comorbidities. Since the comorbidity data is not available for the total infected, the study used the available data till the above period. There are many elderly persons with three or more morbidities. Due to the absence of specific data, the study included the bedridden and unspecified cases also in this category.

As per the epidemiology data on COVID-19, for the entire population of Kerala, till 27, December 2020, CFR=0.44%, IFR=0.40%, CMR=0.008%. However, the granular data of the recovered elderly is not available. Hence, the CFR calculation is not possible as per the WHO's[36] guidelines, where to calculate CFR of an ongoing pandemic, CFR is the number of deaths divided by the number of deaths plus the number of recoveries, which is free from underestimations (Table 8).

DISCUSSION

The emergence of the Corona Virus Disease-2019 has been spread all over the world with surges and resurges. This paper intended to associate Kerala's elderly health status, in the post-COVID-19 scenario, with COVID-19 comorbidity and fatality rates, using a concurrent mixed-method approach. The study finding shows that (Fig. 1), age is a predictor of increased mortality rate in Kerala when the first covid-19 death case itself was reported from this category (69-year-old from Ernakulam District on 28 March 2020).[21] This is consistent with the findings of several international studies.[25,44–47] In Kerala, the COVID-19 fatality among the elderly population has been mostly reported from the 60-69 age category. This can be explained by using Kerala's

socio-demographic profile, where the majority of the elderly belong to the same category (Table 2).

The effect of this pandemic can be studied by using CFR, CMR, and IFR.[36,48] CFR helps to recognize the severity of this disease, identify the risk of the population, and assess the quality of the healthcare system,[36] while CMR estimates the probability of any individual in a population dying from the disease.[36,48] IFR estimations give the proportion of fatality among all infected people. As per the available published COVID-19 epidemiology data, in Kerala, the CFR=0.44%, IFR=0.40%, and CMR=0.008% till 27 December 2020. The CFR of an ongoing pandemic is calculated using the formula.[36]

$$CFR(\%) = \frac{Total\ number\ of\ elderly\ deaths}{Number\ of\ elderly\ deaths + Number\ of\ elderly\ recoveries} \times 100\%$$

As per this formula, the CFR of the elderly COVID-19 death cannot be estimated as the number of infected and recovered elderly data is not available. The elderly IFR=0.30%, and CMR=0.0532% (till 27 December 2020) which clearly shows that the elderly population has been adversely affected by the pandemic.

The inferences of Tables 3, 4, and 5 show the health behavior and functionality (ADL and IADL) level of the elderly. Many of them are bedridden due to attributable age-related diseases. This is consistent with the previous report of Public Health Foundation of India[18] that, in the last two decades, the morbidity burden in Kerala has grown faster than the rest of India wherein the Proportion of Ailing Population (PAP) in Kerala during 1995-96 was 109 against the national average of 55 which was increased to 251 against the national average in 2004, and to 308 against the national average in 2014. It shows that during the 2004 -2014 period, Kerala witnessed an increase of 57

points in the overall morbidity rate when the national average increased just seven points.

Previous studies on Middle East Respiratory Syndrome Coronavirus (MERS-CoV) also found that chronic patients with comorbidity were more likely to be affected by MERS-CoV.[48,49] In West Bengal, comorbidity accounts for 90 percent of the COVID-19 deaths.[50] Therefore, it can be assumed that a high proportion of elderly persons with high NCDs prevalence leading to multi-morbidities can be attributed to the increased fatality of COVID-19 in Kerala.

Existing evidence on the health profile suggests that Kerala has been going through an epidemiological transition with an unprecedented increase in the NCDs burden. Table 6 shows that the most prevalent comorbidities among the elderly in Kerala are hypertension (64%), diabetes (51.4%), urinary disease (31.4%), abdominal disease (27.4 %), heart disease (22.7%), lungs disease (19.5%), and cancer (6.7%). Qualitative data also validate this finding (Table 7). This is akin to the previous demographic study results of Kerala over a decade.[15,18–20] This disease profile also validates the comorbidity status of the elderly COVID-19 death cases in Kerala where the most prevalent comorbidities were heart disease, diabetes, COPD (Chronic Obstructive Pulmonary Disease), kidney disease, hypertension, cancer, and many of them had multi-morbidities. Many previous studies on COVID-19 have obtained the same results.[23,27,48,51–55] However, this finding discards the findings that there is no significant

relationship between COVID-19 infection and diabetes[28,55] and the findings of Wang et al.,[26] who found a negative relation between malignancy and COVID-19 infection.

To be more specific, the major comorbidity found in Kerala's COVID-19 death cases is heart disease, followed by diabetes, which is consistent with the previous studies on COVID-19 fatality.[28,30,56–58] At this juncture, it sets an alarm for the whole society that, in Kerala, the incidence of type 2 diabetes (T2DM) is 21.9%.[59] According to WHO (2015),[1] above 80% of cardiovascular and diabetes deaths and more than 90% of COPD deaths occurred in middle and low income nations. This can be attributable to risky health behaviors such as lack of exercise and an unhealthy diet.

Multi-morbidity gives rise to multiple interactions between one condition and the treatment recommendations for another condition. In a multi-morbidity situation, multiple drug use is inevitable for patients, and the impact of simultaneous use of multiple drugs leads to complications. The susceptibility of the elderly to COVID-19 is explained by *immunosenescence* where the innate immune cells' function is impaired when there is a decrease in the production of naïve T as well as B cells, consequently leading to a situation where the innate immunity will not get activated well to fight the infection.[60] One more characteristic of aging immunity is the CSSI (Chronic Subclinical Systemic Inflammation), which results in an elevation of inflammatory cytokines in serum due to the failure to resolve severe inflammation, which is a critical pathogenic mechanism in COVID-19, contribute to the more unsatisfactory clinical outcome in elderly COVID-19 patients.[61] This phenomenon called cytokine storm/ hypercytokinemia, associated

impairment begins with the damage of the lungs' epithelial barrier. Subsequently, this initiates a cascade of tissue damage in other vital human organs, including the heart, kidneys,brain and blood vessels, leading to a condition called Multiple Organ Dysfunction Syndrome (MODS), which may be even more fatal.[62]

Studies report that most of the patients who recovered from COVID-19 infection experienced different manifestations even 20 days after the PCR reported negative for SARS-CoV-2. The severity of such post-COVID-19 manifestations was positively correlated to the severity of the condition, which in turn associated with the prevalence of comorbidities and age.[63] Hence, post COVID-19 clinical care and close observation of the elderly are important to avoid fatalities.[64]

Earlier, in 2013, a team of researchers found that the convergence of infectious diseases and NCDs escalate the consequences of morbidity and mortality, which will affect large numbers of people in the decades to come.[65] NCDs eliminate 41 million people every year, equal to 71% of global deaths, where 15 million deaths happen in the 30-69 age bracket; above 85% of these deaths are "premature" deaths in low-middle-income nations.[66] A clear understanding of these risk factors among the elderly will help the healthcare system, particularly the clinicians, to identify better and implement the protocols to mitigate the fatal outcomes. In this context, increased investments in public primary and secondary health care facilities are necessary to ensure quality healthcare access,[14,67] and handle post-COVID-19 syndrome situations. Since epidemiology has a holistic approach focused on wellness and maintaining it among a specific demographic population, elderly healthcare must become a priority

issue before the governments. Therefore, priority must be given to the elderly in the COVID-19 vaccine distribution phase.

Organized preventive and curative care for infectious diseases and NCDs must be ensured to the elderly in the line of the 'Vayomithram' project of Kerala, which provides free health check-ups and medicines for the elderly. These interventions for the elderly are significant when, in India, the healthcare system's unaccountability and inefficiency paved the way for privatizing the healthcare industry, turning to unregulated and expensive treatments, making the common man vulnerable to high health expenditures and poverty.[68] According to the 2011 census data, In Kerala, one in eight persons were above 60 years and above, and out of the total elderly, 11 percent of the population are 'old-old' category, 80 years and above.[38] Therefore, geriatric care must be expanded on evidence-based, innovative interventions, in the light of possible syndemic interactions of chronic diseases and their treatments.[69]

Aging is inevitable, and the elderly are affected by the health emergencies to a more considerable extent, determined by their living conditions and health status, irrespective of the nation's development. As a result, amidst Covid-19, the elderly were the worst-affected even in more advanced countries with an efficient healthcare system and disaster response. An effective healthcare system has to advance its facilities, including geriatricians, to ensure healthcare access to the elderly. Often, the promotional and preventative aspects of geriatric care are neglected with a strong notion that it is 'unavoidable' and 'genetically determined' and the increased risk of disease is a part of 'normal aging,' neglecting the importance of healthier lifestyle which can lead to

decreased health care expenditure, even in later stages of life.[70] To ensure geriatric care, there must be an attitudinal change and subsequent efforts from the policymakers to ensure 'quality aging.'

Furthermore, the public health care system should ensure documentation of incidents and causes of death to estimate 'excess deaths' occurring during such outbreaks or pandemics, wherein it becomes difficult to estimate the CFR. Weekly/ monthly counts of death can be collated with trends over the years to ascertain whether the death count is significantly higher than the expected count in that specific period. This estimation can provide information about the potential burden of mortality and fatality, associated with COVID-19 infection, directly or indirectly.[71]

The study finding on elderly comorbidity and their subsequent mortality determine the etiology of COVID-19 and, therefore, the containment and prevention strategies can be implemented to avoid further spread and increased fatalities. Moreover, the study findings will add to the existing knowledge realm on the spectrum of comorbidities in the elderly population and its converged impact on the phase of epidemics spread. It will help to formulate policies and protocols for this most vulnerable group, and therefore, the welfare of the elderly and their equitable healthcare access may be the next priorities before the governments.

The study has its limitations. Firstly, this study was carried out among the elderly in Kerala, and therefore, the results may not be suitable in an international context. Secondly, getting authentic and correct primary data limited the study, particularly in

calculating CFR, which is an estimate and may not be accurate. Thirdly, for the COVID-19, as in the case of previous infectious diseases, the accurate level of transmission is often underestimated as a considerable proportion of the infected population goes undetected for infection due to asymptomatic or mild symptomatic cases failing to test for the infection.[72,73] Fourthly, there will be vulnerable segments that are either neglected or under-served and under-detected who are likely to keep themselves away from healthcare access for testing and treatment.[74]

Conclusion

It is found that patients with chronic diseases, particularly those belonging to the elderly population, are more susceptible to an epidemic, particularly COVID-19 infection due its convergence. Knowledge of these risk factors necessitates a more focused approach to introduce a spectrum of interventions to protect the lives of the elderly when there is a convergence of epidemics and NCDs. The study indicates the need for having timely and accurate data of a population on their demographic profile, which can easily guide the healthcare system and authorities to more efficient prevention and treatment methodologies in health care emergencies. Having evidenced the severity of COVID-19 to the elderly, health commodities such as preventing vaccines and diagnostics must be made available to them at the earliest possible.

Ethical approval

The entire study was conducted following the guidelines and ethical approval received from the Indian Council for Social Science Research (F.No.

IMPRESS/P1132/428/2018-19/ICSSR). Informed consent was obtained from the respondents before collecting their data.

Acknowledgement

The first author acknowledges financial support from the Indian Council for Social Science Research under the IMPRESS grant (F.No. IMPRESS/P1132/428/2018-19/ICSSR). The authors acknowledge Dr. Arun N. M. for granting permission to use the volunteering database on COVID-19 in Kerala. The authors acknowledge discussions with Dr. Anupama M Pai, Dr. Shabeesh Balan, and Dr. Maya Madhavan.

Reference:

1. World Health Organization. *World Report on Ageing and Health*.; 2015.

2. Marengoni A, Angleman S, Melis R, et al. Aging with multimorbidity: A systematic review of the literature. *Ageing Res Rev*. 2011;10(4):430-439.

3. UNFPA and HelpAge International. *Ageing in the Twenty-First Century: A Celebration and A Challenge*. UNFPA and HelpAge International; 2012. Accessed December 30, 2020. https://www.unfpa.org/publications/ageing-twenty-first-century

4. Biswas P, Nahar Kabir Z, Nilsson J, Zaman S. Dynamics of Health Care Seeking Behaviour of Elderly People in Rural Bangladesh. *International Journal of Ageing and Later Life*. 2006;1(1):69-89.

5. Clausen F, Sandberg E, Ingstad B, Hjortdahl P. Morbidity and health care utilisation among elderly people in Mmankgodi village, Botswana. *Journal of Epidemiology & Community Health*. 2000;54(1):58-63.

6. Soe Moe , Kyi Tha , Daw Khin Saw Naing , Maung Maung Than Htike. Health Seeking Behaviour of Elderly in Myanmar. *International Journal of Collaborative Research on Internal Medicine & Public Health*. 2012;4(8):1538-1544.

7. Woo E-K, Han C, Jo SA, et al. Morbidity and related factors among elderly people in South Korea: results from the Ansan Geriatric (AGE) cohort study. *BMC Public Health*. 2007;7(1):10.

8.   World Health Organization. *World Report on Disability*.; 2011.

9.   Rajan SI, Aiyar S. Population ageing in India. In: Rajan SI, Risseeuw C, Perera M, eds. *Institutional Provisions and Care for the Aged: Perspectives from Asia and Europe*. Anthem Press; 2008:39-54.

10. Bloom DE, Cafiero-Fonseca ET, Candeias V, et al. *Economics of Non-Communicable Diseases in India: The Costs and Returns on Investment of Interventions to Promote Healthy Living and Prevent, Treat, and Manage NCDs*. World Economic Forum and the Harvard School of Public Health; 2014. http://www3.weforum.org/docs/WEF_EconomicNonCommunicableDiseasesIndia_Report_2014.pdf

11. Rajan SI. *Population Aging, Health and Social Security in India Population Aging, Health and Social Security in India Population Aging, Health and Social Security in India*.; 2007. https://www.econ.osaka-cu.ac.jp/CREI/discussion/2006/CREI_DP003.pdf

12. Tiwari SC, Pandey NM, Singh N. Mental health problems among inhabitants of old age homes: A preliminary study. *Indian J Psychiatry*. 2012;54(2):144.

13. Arokiasamy P. G. Giridhar, K. M. Sathyanarayana, Sanjay Kumar, K. S. James and Moneer Alam (eds), Population Ageing in India. *Ageing and Society*. 2016;36(2):445-447.

14. Mukherjee S, Levesque J-F. Morbidity and Outpatient Care for the Elderly in Kerala, South India: Evidence from a National Population based Survey. *J Popul Ageing*. 2012;5(3):177-192.

15. Alam M, James KS, Giridhar G, et al. *Building a Knowledge Base on Population Aging in India: Report on the Status of Elderly in Select States of India*. United Nations Population Fund; 2013. https://ideas.repec.org/p/ess/wpaper/id5209.html

16. Dilip TR. The burden of ill health among elderly in Kerala. *Man in India*. 2003;83(1-2):195-205.

17. Sandhya RS. Sociological analysis of the Problems of the oldest old (A study conducted in Kerala). *International Research Journal of Social Sciences*. 2010;3(1):57-79.

18. Karan A, Srivastava S, Chakraborty A, Matela H. *Key Indicators of Health and Morbidity - Kerala 2014*. Public health foundation of India; 2016.

19. Madore A, Rosenberg J, Dreisbach T, Weintraub R. *Positive Outlier: Health Outcomes in Kerala, India over Time | The Global Health Delivery Project*. Harvard Business Publishing; 2018. https://www.globalhealthdelivery.org/files/ghd/files/ghd-042_kerala_as_a_positive_outlier_2018_09_05.pdf

20. BusinessLine. Covid-19: Underlying ailments behind most deaths in Kerala - The Hindu BusinessLine. *The Hindu Business Line*. https://www.thehindubusinessline.com/news/covid-19-underlying-ailments-behind-most-deaths-in-kerala/article31801111.ece. Published June 11, 2020.

21. Mohan N. In Kerala COVID mortality very low, but elderly die mostly | Kerala News | Manorama. *Malayala Manorama*. https://www.onmanorama.com/news/kerala/2020/07/27/in-kerala-covid-mortality-very-low-but-elderly-die-mostly.html. Published July 27, 2020.

22. Pal R, Bhadada SK. COVID-19 and non-communicable diseases. *Postgrad Med J*. 2020;96(1137):429-430.

23. Guan W-J, Liang W-H, Zhao Y, et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J*. 2020;55(5):2000547.

24. Callender LA, Curran M, Bates SM, Mairesse M, Weigandt J, Betts CJ. The Impact of Pre-existing Comorbidities and Therapeutic Interventions on COVID-19. *Frontiers in Immunology*. Published online 2020. doi:10.3389/fimmu.2020.01991

25. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-1062.

26. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061.

27. Wang B, Li R, Lu Z, Huang Y. Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging* . 2020;12(7):6049-6057.

28. Li J, Guo T, Dong D, et al. Defining heart disease risk for death in COVID-19 infection. *QJM: An International Journal of Medicine*. Published online August 13, 2020. doi:10.1093/qjmed/hcaa246

29. Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *N Engl J Med*. 2020;382(13):1199-1207.

30. Guo W, Li M, Dong Y, et al. Diabetes is a risk factor for the progression and prognosis of COVID-19. *Diabetes Metab Res Rev*. 2020;36(7). doi:10.1002/dmrr.3319

31. Petrilli CM, O'Donnell L, Horwitz LI, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: Prospective cohort study. *BMJ*. Published online 2020.

32. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *The Lancet Respiratory*

Medicine. 2020;8(4):e21.

33. Pal R, Bhansali A. COVID-19, diabetes mellitus and ACE2: The conundrum. *Diabetes Res Clin Pract*. 2020;162:108132.

34. Emami A, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. *Archives of academic emergency medicine*. Published online 2020. doi:10.22037/aaem.v8i1.600

35. Liang W, Guan W, Chen R, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *The Lancet Oncology*. 2020;21(3):335-337. doi:10.1016/S1470-2045(20)30096-6

36. World Health Organization. Estimating mortality from COVID-19: Scientific brief, 4 August 2020. WHO Scientific brief. Published 2020. https://www.who.int/publications/i/item/WHO-2019-nCoV-Sci-Brief-Mortality-2020.1

37. Census of India. *D1: Population Classified by Place of Birth and Sex - 2011*. Office of the Registrar General & Census Commissioner, India; 2011.

38. Rajan SI, Shajan A, Sunitha S. Ageing and Elderly Care in Kerala. *China Rep*. 2020;56(3):354-373.

39. Ulahannan J, Narayanan N, Suresh S, et al. Covid19Kerala.info-Data: A collective open dataset of COVID-19 outbreak in the south Indian state of Kerala. Published online September 5, 2020. doi:10.5281/zenodo.3818096

40. Ulahannan JP, Narayanan N, Thalhath N, et al. A citizen science initiative for open data and visualization of COVID-19 outbreak in Kerala, India. *J Am Med Inform Assoc*. 2020;00(0):1-8.

41. World Health Organization. *Coronavirus Disease 2019 (COVID-19) Situation Report – 82*. World Health Organization; 2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200411-sitrep-82-covid-19.pdf

42. Glaser BG, Strauss AL. *The Discovery of Grounded Theory: Strategies for Qualitative Research*. Aldine.; 1967.

43. Charmaz K. *Constructing Grounded Theory | SAGE Publications Ltd*. SAGE Publications; 2006:189.

44. Yanez ND, Weiss NS, Romand JA, Treggiari MM. COVID-19 mortality risk for older men and women. *BMC Public Health*. 2020;20(1):1742.

45. Wang L, He W, Yu X, et al. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect*.

2020;80(6):639-645.

46. Weekly CC. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. *China CDC Weekly*. February 1, 2020:113-122.

47. Kang SJ, Jung SI. Age-related morbidity and mortality among patients with COVID-19. *Infect Chemother*. 2020;52(2):154-164.

48. Roser M, Ritchie H, Ortiz-Ospina E, Hasell J. Coronavirus Pandemic (COVID-19). *Our World in Data*. Published online 2020. https://ourworldindata.org/coronavirus

49. Badawi A, Ryoo SG. Prevalence of comorbidities in the Middle East respiratory syndrome coronavirus (MERS-CoV): a systematic review and meta-analysis. *Int J Infect Dis*. Published online 2016. https://www.sciencedirect.com/science/article/pii/S1201971216311006

50. Kumar DV. Kerala COVID report: Death high among people without travel history, stress on comorbidities- The New Indian Express. *The New Indian Express*. https://www.newindianexpress.com/states/kerala/2020/aug/29/kerala-covid-report-death-high-among-people-without-travel-history-stress-on-comorbidities-2189845.html. Published August 29, 2020.

51. Paudel SS. A meta-analysis of 2019 novel corona virus patient clinical characteristics and comorbidities. *Research Square*. Published online April 8, 2020. doi:10.21203/rs.3.rs-21831/v1

52. Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA - Journal of the American Medical Association*. 2020;323(18):1775-1776. doi:10.1001/jama.2020.4683

53. Franki R. Comorbidities the rule in New York's COVID-19 deaths | The Hospitalist. The Hospitalist. Published 2020. https://www.the-hospitalist.org/hospitalist/article/220457/coronavirus-updates/comorbidities-rule-new-yorks-covid-19-deaths

54. Sanyaolu A, Okorie C, Marinkovic A, et al. Comorbidity and its Impact on Patients with COVID-19. *SN Comprehensive Clinical Medicine*. 2020;2(8):1069-1076.

55. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory Medicine*. 2020;8(5):475-481.

56. Thankappan K. Combating corona virus disease 2019 and comorbidities: The Kerala experience for the first 100 days. *International Journal of Noncommunicable Diseases*. Published online 2020. doi:10.4103/jncd.jncd_26_20

57. Clerkin KJ, Fried JA, Raikhelkar J, et al. COVID-19 and Cardiovascular Disease.

Circulation. 2020;141(20):1648-1655. doi:10.1161/CIRCULATIONAHA.120.046941

58. Nishiga M, Wang DW, Han Y, Lewis DB, Wu JC. COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives. *Nature Reviews Cardiology*. 2020;17(9):543-558. doi:10.1038/s41569-020-0413-9

59. Vijayakumar G, Manghat S, Vijayakumar R, et al. Incidence of type 2 diabetes mellitus and prediabetes in Kerala, India: results from a 10-year prospective cohort. *BMC Public Health*. 2019;19(1):140.

60. Castle SC. Clinical relevance of age-related immune dysfunction. *Clin Infect Dis*. 2000;31(2):578-585.

61. Bonafè M, Prattichizzo F, Giuliani A, Storci G, Sabbatinelli J, Olivieri F. Inflamm-aging: Why older men are the most susceptible to SARS-CoV-2 complicated outcomes. *Cytokine Growth Factor Rev*. 2020;53:33-37.

62. Bhaskar Sonu, Sinha Akansha, Banach Maciej, Mittoo Shikha, Weissert Robert, Kass Joseph S., Rajagopal Santhosh, Pai Anupama R., Kutty Shelby. Cytokine storm in COVID-19—immunopathological mechanisms, clinical considerations, and therapeutic approaches: the REPROGRAM consortium position paper. *Frontiers in Immunology*. 2020;11:1648.

63. Xiaochen Li, Shuyun Xu, Muqing Yu, Ke Wang, Yu Tao, Ying Zhou, Jing Shi, Min Zhou, Bo Wu, Zhenyu Yang, Cong Zhang, Junqing Yue, Zhiguo Zhang, Harald Renz, Xiansheng Liu, Jungang Xie, Min Xie, Jianping Zhao. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *Journal of Allergy and Clinical Immunology*. 2020;146(1):110-118.

64. Kamal M, Abo Omirah M, Hussein A, Saeed H. Assessment and characterisation of post-COVID-19 manifestations. *Int J Clin Pract*. Published online September 29, 2020:e13746.

65. Marais BJ, Lönnroth K, Lawn SD, et al. Tuberculosis comorbidity with communicable and non-communicable diseases: integrating health services and control efforts. *Lancet Infect Dis*. 2013;13(5):436-448.

66. World Health Organization. *WHO | Noncommunicable Diseases Country Profiles 2018*. World Health Organization; 2018. http://www.who.int/nmh/publications/ncd-profiles-2018/en/

67. Levesque JF. Outpatient care utilization in urban Kerala, India. *Health Policy Plan*. 2006;21(4):289-301.

68. Pal R. Analysing Catastrophic OOP Health Expenditure in India: Concepts, Determinants and Policy Implications. *Working Papers*. Published online 2010. https://ideas.repec.org/p/ess/wpaper/id2420.html

69. National Academies of Sciences EAM. *The Convergence of Infectious Diseases and Noncommunicable Diseases*. (Ogawa VA, Shah CM, Negussie Y, Nicholson A, eds.). National Academies Press; 2019.

70. Prabhu P, Singh FMA. Nutrition the Geriatric Population. In: *Geriatric Care*. Brookdale Foundation; 2002:56-65.

71. Centers for Disease Control and Prevention. Excess deaths associated with COVID-19: Provisional Death Counts for Coronavirus Disease (COVID-19). National Center for Health Statistics. Published December 23, 2020. Accessed December 29, 2020. https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm

72. Kim GU, Kim MJ, Ra SH, et al. Clinical characteristics of asymptomatic and symptomatic patients with mild COVID-19. *Clin Microbiol Infect*. Published online 2020. doi:10.1016/j.cmi.2020.04.040

73. Nishiura H, Kobayashi, Jung T, et al. Communicating the Risk of Death from Novel Coronavirus Disease (COVID-19). Journal of Clinical Medicine, 9(2), 580. *J Clin Med Res*. Published online 2020. https://www.ncbi.nlm.nih.gov/pubmed/32098019

74. Lau H, Khosrawipour T, Kocbach P, Ichii H, Bania J, Khosrawipour V. Evaluating the massive underreporting and undertesting of COVID-19 cases in multiple global epicenters. *Pulmonology*. Published online 2020. doi:10.1016/j.pulmoe.2020.05.015

75. Department of Health & Family Welfare. *COVID-19 Death Audit Report*. Government of Kerala; 2020. http://dhs.kerala.gov.in/wp-content/uploads/2020/10/Death-Audit-August-2020.docx .pdf

Table 1. Disability among the elderly population

Source:

Disease	Developed Countries (millions)	Developing Countries (millions)
Visual Impairment	15.0	94.2
Hearing Loss	18.5	43.9
Osteoarthritis	8.1	19.4
Ischaemic Heart Disease	2.2	11.9
Dementia	6.2	7.0
Chronic Obstructive Pulmonary Disease	4.8	8.0
Cerebra-vascular Disease	2.2	4.9
Depression	0.5	4.8
Rheumatoid Arthritis	1.7	3.7

Table 2 Socio-Demographic Variables

Variable	Category	Frequency	Percent
Gender	Female	181	44.7
	Male	224	55.3
Age	60-69	203	50.1
	70-79	128	31.6
	80-89	60	14.8
	Above 90	9	2.2
Marital Status	Married	314	77.5
	Not Married	11	2.7
	Partner Died	75	18.5
	Divorced	3	0.7
Education	No formal Education	74	18.3
	School	267	65.9

	College/University	53	13.1
	No response	4	1
Monthly income	No income	125	25.9
	Less than INR 10,000	228	56.3
	INR 10,000-20,000	37	9.1
	INR 21,000-30,000	20	4.9
	Above INR 30,000	5	1.2
Source of Income	Job	18	4.4
	Pension	211	52.1
	Income from assets	29	7.2
	Business	35	8.6
	Income from agriculture	49	12.1
	Income from other sources	15	3.7

Table 3. Health Behaviour of the elderly in Kerala

Variables	Regularly	Frequently	Sometimes	Rarely	Never
Exercise	64 (15.8%)	68 (16.8%)	52 (12.8%)	40 (9.9%)	180 (44.4%)
Diet	129 (31.9%)	117 (28.9%)	70 (17.3%)	34 (8.4%)	55 (13.6%)
Smoking	30 (7.4%)	19 (4.7%)	9 (2.2%)	15 (3.7%)	332 (82%)
Alcohol	1 (0.2%)	15 (3.7%)	26 (6.4%)	17 (4.2%)	346 (85.4%)
Chewing tobacco	14 (3.5%)	6 (1.5%)	10 (2.5%)	7 (1.7%)	368 (90.9%)

Table 4. Assistance required in IADL

Activities	Assistance Not Required N (%)	Assistance required Sometimes N (%)	Full Assistance Required N (%)
Eating	365 (90.1%)	28 (6.9%)	12 (3%)
Get-up/ sit down	303 (74.8%)	82 (20.2%)	20 (4.9%)
Walking	317 (78.3%)	66 (16.3%)	22 (5.4%)
Toileting	367 (90.6%)	22 (5.4%)	16 (4%)
Dressing	365 (90.1%)	20 (4.9%)	20 (4.9%)
Bathing	360 (88.9%)	26 (6.4%)	19 (4.7%)

Table.5 Assistance required for IADL

Activities	Assistance not Required N (%)	Assistance required Sometimes N (%)	Full Assistance Required N (%)
Telephone	214 (52.8%)	70 (17.3%)	121 (29.9%)
Shopping	198 (48.9%)	63 (15.6%)	144 (35.6%)
Cooking	199 (49.1%)	43 (10.6%)	163 (40.2%)
Household work	188 (46.4%)	57 (14.1%)	160 (39.5%)
Washing	200 (49.4%)	51 (12.65)	154 (38%)
Travelling	194 (47.9%)	70 (17.3%)	141 (34.8)
Medication	262 (64.7%)	62 (15.3%)	81 (20%)
Finance	266 (65.7%)	51 (12.6%)	88 (21.7%)

Table 6 Elderly Disease Profile in Kerala

Type of Disease	Regularly	Frequently	Sometimes	Rarely	Never
Diabetes	156 (38.5%)	37 (9.1%)	9 (2.2%)	6 (1.5%)	197 (48.6%)
Hypertension	145 (35.8%)	66 (16.3)	29 (7.2%)	19 (4.7%)	146 (36%)
Cataract	56 (13.8%)	30 (7.4%)	35 (8.6%)	30 (7.4%)	253 (62.5%)
Urinary Disease	24 (5.9%)	27 (6.7%)	33 (8.1%)	43 (10.6%)	278 (68.6%)
Lungs Disease	32 (7.9%)	8 (2%)	14 (3.5%)	25 (6.2%)	326 (80.5%)
Heart Disease	46 (11.4%)	18 (4.4%)	19 (4.7%)	9 (2.2%)	313 (77.3%)
Depression	12 (3%)	12 (3%)	13 (3.2%)	37 (9.1%)	331 (81.7%)
Cancer	17 (4.2%)	4 (1%)	3 (0.7%)	3 (0.7%)	378 (93.3%)
Rheumatic Disease	57 (14.1%)	35 (8.6%)	36 (8.9%)	42 (10.4%)	235 (58%)
Abdominal Disease	25 (6.2%)	13 (3.2%)	31 (7.7%)	42 (10.4%)	294 (72.6%)

Table 7. Disease and health behavior of elderly - Qualitative data

Attribute	Responses (District-participant ID, Gender, Age)
Disease	..I have Diabetes for the past 35 years, severe joint pain, back pain... (Palakkad, F, 70-75) …there are number of diseases in this age…headache, back pain, blood pressure, disk problem, cholesterol(Pathanamthitta, F, 60-65) I am taking medicines for blood pressure regularly…. (Pathanamthitta, M, 60-65) ….I had a block in my heart, for which cardiac surgery was done. Check-up is done every three months. Then I had prostate cancer… follow up visits once in six month. I am a diabetic patient too...(Thrissur, M, 65-70) …angioplasty was done ….taking medicines for blood pressure twice a day." (Kozhikode, F, 70-75) "I am a cardiac patient, using pacemaker….(Palakkad, F, 70-75) "Taking medicines for thyroid, sugar, BP daily… (Thiruvananthapuram, F, 65-70) "My current health problems are Asthma, Cholesterol, Sugar, blood pressure, thyroid, Cardiac diseases……(Kollam, F, 70-75) I have cancer, BP, Heart problem , doing chemotherapy(Malappuram, M, 70-75)
Healthy Behaviour (Exercise)	…Many years before, I learned a special Japan exercise. Since I worked in companies, I got training for warm up exercise. That I

	continue for half an hour… I walk daily 4 kilometres…. (Pathanamthitta, M, 60-65)
	……my exercise is my house work There is lots of work in the house. That itself is an exercise…..(Alappuzha, M, 60-65)
	'I have a morning walk regularly '(Thrissur, M, 60-65)
	…I go for rubber tapping daily by walking 2 kilometres in the morning daily. After that I have to feed my cattle. All these are my exercises……(Kozhikode, M, 60-65)
	…Doing exercise for the past two years regularly, suggested by doctor after one surgery……(Thirvananthapuram, F, 70-75)
	….Daily exercise is there for 1 hour. Every evening @ 5 pm I walk for one hour. (Palakkad, F, 65-70)
Diet	I do not eat sugar, no spicy food, no oil in preparing food, just eating vegetables half cooked and fruits (Kasaragod, F, 60-65)
	Food control is maintained (Thrissur, M, 63-65)
	Restrict myself from eating fried items (Kozhikode, M, 60-65)
	Eat food in usual quantities. Do not use sugar and meat…….(Wayanad, M, 70-75).

Table. 8 COVID-19 Indicators (CFR, CMR, IFR)

Note: Kerala population as per the Census data of 2011: 35,699,443. The number of the elderly population as per the Census data of 2011: 4, 193,393. The data on the comorbidity of the elderly death cases are not available after August 2020 even when there was a sudden curve in the fatality rate in Kerala

Type of data	No. of infected cases	No of elderly with comorbidity	No of deaths	CFR %	CMR %	IFR %
DHS Dashboard (*as on 27 December 2020*)	740516	No Data	2976	0.44	0.008	0.40
Crowdsourced dashboard (*as on 27 December 2020*)	740516	No Data	2976	0.41	0.008	0.37
Volunteering dashboard (*as on 26 December 2020*)	735611	Data available till September 30	4396	0.65	0.012	0.60

Fig. 1 Covid-19 Deaths in Kerala-Age wise till December 27, 2020

Source: DHS, Dashboard

Fig. 2 Types of Comorbidities Prevalent in Fatality Cases of Covid-19

Source: Death Audit Report, DHS, Government of Kerala[75]

Fig. 3 Types of Elderly Comorbidities Prevalent in Fatality Cases of Covid-19

Source: Source: Compiled from dashboards