Convergence of Comorbidity and COVID-19 Infection to Fatality: An Investigation Based on Concurrent Health Status Evaluation among Older Adults

Sindhu Joseph[1,*], Jijo Pulickiyil Ulahannan[2,3]

[1]Department of Travel and Tourism Management, Govinda Pai Memorial Government College, Kannur University, Kerala, India.

[2]Department of Physics, Government College Kasaragod, Kannur University, Kerala, India.

[3] Collective for Open Data Distribution-Keralam (CODD-K).

Abstract

Background

The COVID-19 infection rapidly spread globally, mostly affecting the extremely vulnerable category of the older people with comorbidities. There are inconsistencies in the findings on the type of comorbidity of the older people and its association with fatalities.

Objective

To investigate the impact of age and comorbidity in the fatality of older COVID-19 patients in the state of Kerala, India, based on their health status, health behaviour, functionality, and morbidity profile.

Design

A cross sectional study

Methods

Using a concurrent mixed method approach, 405 older people were surveyed, and 102 in-depth interviews were carried out at the households of the older population between June to November 2020. The results of the study have been further triangulated with the COVID-19 fatality data of the older people, available from the citizen-science dashboard of the research team and two other dashboards.

Results

It is found that age is a predictor of COVID-19 fatality and heart disease, diabetes, COPD, kidney disease, hypertension and cancer are the significant predictors of COVID-19 fatality of older people in Kerala and comorbidity of the COVID-19 older death cases matches with the current disease profile of the older population in Kerala.

Conclusions

Comorbidities can predict potential fatality among older COVID-19 patients. While facing an epidemic like the present zoonotic disease, better knowledge of these high-risk factors will help clinicians and healthcare systems to implement therapeutic and preventive methodologies and interventions for the older population.

Keywords: COVID-19 Mortality, Kerala, Elderly, Comorbidity, Fatality, Convergence

*Corresponding Author: Sindhu Joseph Ph.D., GPM Government College

Manjeshwar, Kannur University, Kerala, India 671323, Email: sjsindhu@gmail.com

INTRODUCTION

It is already acknowledged that health conditions deteriorate when age increases and ageing goes hand in hand with many psychological, physical, and behavioural issues and disabilities, contagious infections, lifestyle and chronic diseases. Consequently, the rate of health-care utilisation and healthcare expenses may be significantly greater[1]. Therefore, the impact of multi-morbidity on various aspects such as quality of life, functionality and risk of mortality becomes a matter of present discussions worldwide. The latest research reveals that the older people suffer from non-communicable diseases (NCDs)[2]. They face health issues such as loss of appetite, gastric and body pain, cataract, arthritis, uterine problems, hypertension, Diabetes Mellitus, general weakness, stroke, muscular-skeletal and respiratory diseases[3-6].

In India, geriatric care has been adversely affected by the population size leading to inadequacy of the public healthcare system in providing equitable healthcare. In 2001, in India, around 33.1% of the older people reported with unmet healthcare needs[7]. They face NCDs' threat, including cancers, CVD (Cardiovascular Disease), respiratory diseases, and diabetes and one out of every two older people suffers from at least one chronic disease requiring life-long medication, particularly in urban areas[8] and mental health issues, including depression[9].

Kerala has the highest outpatient doctor consultation per 1,000 population compared to the rest of the states in India[10]. Higher hospitalisation rates indicate higher morbidity levels[11]. In a 2004 study, about 25% of the older people reported with poor health status[12].

Their common ailments were paralysis, urinary problems, CVD and cancer in 2010[13], hypertension, diabetes mellitus, cataract and heart disease in 2011[11], CVD, diabetes, musculoskeletal, and respiratory disease in 2016[14] and diabetes, abdominal obesity, and hypertension in 2016-17[15]. In 2020, about 70 lakh older people have cancer, diabetes, hypertension, respiratory tract infections, kidney and heart diseases[16]. Based on these previous studies, it can be assumed that the significant prevalent diseases of older people are Hypertension, Diabetes, CVD, Cancer, and Respiratory Diseases.

THE COVID-19 FATALITY

The outspread of COVID-19 affected mostly the older people. The Case Fatality Rate (CFR) among them is disproportionately higher[17,18]. Patients with chronic comorbidities, including malignancy, CVD, diabetes, hypertension, kidney and respiratory disease are prone to the fatal outcome of COVID-19 infection[19-27]. Various studies[20,26,27] found that for CVD and hypertension, the use of renin-angiotensin system inhibitors may accelerate the susceptibility to SARS-CoV-2 infection. However, the susceptibility to SARS-CoV-2 of cancer patients is low[28,29] is also reported.

Despite the outnumbered post-COVID-19 studies describing comorbidities and the poor clinical outcomes leading to fatality, the results are inconsistent. Throughout the COVID-19 outbreak, wide variations in CFR and IFR (Infection Fatality Rate) estimates have been noted, which are misleading[30]. Likewise, the study findings related to the health status, comorbidity profile, and the NCDs scenario of the older people over a decade in Kerala are inconsistent[11,13-16]. In this context, this study is intended to fill the above gaps and

planned to relate the health status of the older people in the post-COVID-19 scenario to the mortality rate of COVID-19, which will help to identify the severity of risks. Kerala has gained acclaim for its healthcare indicators and preventive and curative measures for COVID-19. Thus, it is essential to draw a clear picture of the health status and morbidity levels of the older people amidst the pandemic and to assess whether the comorbidity has converged with the COVID-19 infection.

This study is unique as it measured the disease profile and the comorbidity level of the older people, while the pandemic is ongoing with simultaneous assessment of the older people COVID-19 deaths due to the suspected comorbidity levels. To check and identify a consistent pattern in the association between age and COVID-19 morbidity or mortality, an extensive review of the literature was carried out. The study insights will help to shape the geriatric treatment policies and preventive measures for infectious diseases.

METHODOLOGY

This research is based on a mixed methodology, based on a pragmatic paradigm. The study used a concurrent timing strategy where both the quantitative and qualitative strands occurred during a single phase, from June 2020 to November 2020. Quantitative data was collected from 405 older people from 36 panchayats of Kerala adopting multistage sampling techniques, using a structured questionnaire. The study area was divided into three zones, South, Central and North. From each zone, three districts were selected at random. Four panchayats from each district were selected by drawing lots, making 36 panchayats of Kerala under study. From each selected panchayath, 11

households with older people were selected randomly. The data were analysed with SPSS 2.0.

Using a panchayat-wise list, received from the *Vayomithram* coordinators (a government project for holistic care of older people), Anganvadi and *Asha* workers, older people above 60 years of age were selected randomly for the qualitative interview. The interviews were conducted at their households for about 30 - 45 minutes. An average of seven interviews per district were conducted and therefore 102 in-depth interviews were undertaken from all the fourteen districts of Kerala. They were asked about the present ailments and health behaviour (exercise, diet, smoking, use of pan and alcohol and morbidities). To further enrich the data, 12 interviews were conducted with *Vayomithram* district coordinators of 12 districts in Kerala.

The Government of Kerala dashboard (https://dashboard.kerala.gov.in) and the Directorate Health Services (DHS) website (http://dhs.kerala.gov.in) give comprehensive updates about the COVID-19 spread. However, access to primary the data is not complete, and therefore, a multi-disciplinary team of experts compiled data through a citizen science initiative, led by the researchers of this study (https://doi.org/10.5281/zenodo.3818096)[31,32]. To supplement the information available from the above three sources, another volunteering dashboard (https://t.co/5e70Wu2dZz) was also analysed for getting information that is not available on the official website of the government or elsewhere.

A systematic analysis of the above mentioned three dashboards on CFR and comorbidities was done. The data was extracted from the death cases reported in Kerala with COVID-19 infection between January 2020 and December 27, 2020 (Fig. 1). The quantitative and qualitative strands were mixed and integrated at the point of interpretation and drawn inferences.

This study adopted the definition of WHO[33 for] COVID-19 death. Before the qualitative and quantitative data collection, the participants were briefed about the study's purpose, and informed consent was obtained. COVID-19 protocol was adhered to while collecting the data.

RESULTS

Quantitative Study

The socio-demographic profile (Table I) shows that 55.3% were males, and 44.7% were females. Around 50.1% were in the 60-69 age category, followed by 31.6% in the 70-79 category.

Health Behaviour and Functionality Level

Health behaviour is a predictor of good health. Quality of life depends mainly on health status and functionality level. The health status of the older people shows that 28.4% are in good health condition (Table II), and only 15.8% exercise regularly. In dietary habits, only 31.9% have a regular healthy diet (Table III). The risky habits like smoking, alcohol consumption and chewing tobacco have not impacted their lives much.

Functionality level was studied using two latent constructs such as Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL). ADLs are regular activities like eating, bathing, toileting, mobility, getting dressed, and continence. IADLs are the activities for independent living and are important for assessing the early-stage disease of a person. Results show that about 78-90% of the older people can perform ADLs while about 40-65% can perform IADLs (Tables IV and V).

DISEASE PROFILE

Table VI shows that an average of 64% of the older people suffers from Hypertension, 51.4% from Diabetes, 42% from Rheumatic Disease, 37.5% from Cataract, 31.4% from Urinary Disease, 27.4 % from Abdominal Disease, 22.7% from Heart Disease, 19.5% from Asthma, 18.3% from Depression 6.7% from Cancer. Many of them have two or more diseases (comorbidities).

Results-Qualitative Data

This study's key findings were also based on qualitative responses of the older people to the interview questions (Table VII). Analysis of data used a grounded theory research design[34], where the approach intends to develop concepts, insights, or models grounded in the participant's data[35]. Interviews showed that most of the older people had lifestyle diseases like blood pressure (Hypertension), Diabetes, Heart Disease, Cancer, and Cholesterol (Table VII). *Vayomithram* coordinators also shared the same view.

Covid-19 Deaths in Kerala-Age wise

Fig. 1 shows that age is a predictor of increased mortality rate. Till December 27, 2020, 2976 COVID-19 deaths happened in Kerala, of which 2232 (75.0%) deaths were older people, where the majority belongs to the 60-69 age group (Table VIII).

Elderly COVID-19 Deaths and Comorbidity

As of September 30, 2020, out of the 525 older people who died, 209 of them (~40%) had comorbidities. Since the comorbidity data is not available for the total infected, the study used available data till the above period. There are many older people with comorbidities (Fig. 2). As per the epidemiology data on COVID-19, for the entire population of Kerala, till 27, December 2020, CFR=0.44%, IFR=0.40%, CMR (Crude Mortality Rate) =0.008%.

DISCUSSION

The emergence of the COVID-19 disease has been spread with surges and resurges. In this context, this paper is intended to associate older people's health profile of Kerala with COVID-19 comorbidity and fatality rates, using a concurrent mixed-method approach.

The effect of this pandemic can be studied by using CFR, CMR, and IFR[30,36]. CFR helps to recognise the disease severity, risks, and healthcare system quality[30], while CMR estimates the probability of any individual in a population dying from the disease[30,36]. IFR estimations give the proportion of fatality among all infected. As per the available published COVID-19 epidemiology data, in Kerala, the CFR=0.44%, IFR=0.40%, and CMR=0.008% till 27 December 2020 (Table VIII). The CFR of an ongoing pandemic is calculated using the formula[30].

$$CFR(\%) = \frac{Total\ number\ of\ elderly\ deaths}{Number\ of\ elderly\ deaths\ +\ Number\ of\ elderly\ recoveries} \times 100\%$$

As per this formula, the CFR of the older people COVID-19 death cannot be estimated as the number of infected and recovered older people data is not available. The older people IFR=0.30%, and CMR=0.0532% (till 27 December 2020) clearly show that the older population has been adversely affected by the pandemic, which is consistent with the international studies[20,37-40]. The fatality has been mostly reported from the 60-69 age category, which can be explained by using Kerala's socio-demographic profile (Table I).

Many of the older people are bedridden due to attributable age-related diseases (Tables II-V). This is consistent with the previous report[14] that, in the last two decades, the morbidity burden has grown faster than the rest of India wherein the Proportion of Ailing Population (PAP) in Kerala during 1995-96 was 109 against the national average of 55 which was increased to 251 in 2004, and to 308 in 2014, showing, an increase of 57 points in the overall morbidity rate against national average increase of just seven points during the 2004-2014 period. Existing evidence on the health profile suggests that Kerala has been going through a demographic transition with an unprecedented increase in the NCDs burden (Table VI). Qualitative data also validate this finding (Table VII). This is akin to the previous demographic study results of Kerala over a decade[11,14-16]. This disease profile validates the comorbidity status of the older COVID-19 death cases in Kerala where the most prevalent comorbidities were heart disease, diabetes, COPD (Chronic Obstructive Pulmonary Disease), kidney disease, hypertension, cancer, and many of them had multi-morbidities. Many previous studies on COVID-19 have obtained the same results[22,36,41-46]. To be more specific, the major comorbidity found in Kerala's COVID-19 death cases is heart disease, followed by diabetes which are major predictors of COVID-

19 fatality[23,24,47-49]. It is found that, in Kerala, the incidence of type 2 diabetes (T2DM) is 21.9%[50]. This can be attributable to risky health behaviours such as lack of exercise and an unhealthy diet (Table II) which necessitates urgent implementation of healthy behaviour policy initiative.

Previous studies on Middle East Respiratory Syndrome (MERS-CoV) also found comorbidity leads to MERS-CoV infection[36,51]. In West Bengal, comorbidity accounts for 90% of the COVID-19 deaths[52]. Therefore, it can be assumed that a high proportion of older people with NCDs prevalence leading to multi-morbidities can be attributed to the increased older people fatality of COVID-19 in Kerala. However, this study discards the findings that there is no significant relationship between COVID-19 infection and diabetes[23,46] and cancer[21].

Multi-morbidity gives rise to multiple interactions between one condition and the treatment recommendations for another which necessitates simultaneous multiple drug use leading to complications. The susceptibility of the older people to COVID-19 is explained by *immunosenescence* where the innate immune cells' function is impaired when there is a decrease in the naïve T as well as B cells production, consequently leading to a situation where the innate immunity cannot fight the infection[53]. Another characteristic of ageing immunity is the CSSI (Chronic Subclinical Systemic Inflammation), which results in an elevation of inflammatory cytokines in serum due to the failure to resolve severe inflammation, which is a critical pathogenic mechanism in COVID-19, contribute to poor clinical outcome in older people[54]. This phenomenon called cytokine storm/ hypercytokinemia, associated impairment begins with the damage of the lungs' epithelial

barrier. Subsequently, this initiates a cascade of tissue damage in other vital human organs, including the heart, kidneys, brain and blood vessels, leading Multiple Organ Dysfunction Syndrome (MODS), which may be even more fatal[55].

Studies report that the majority of recovered patients experienced different manifestations even 20 days after the PCR reported negative for SARS-CoV-2. The severity of such post-COVID-19 manifestations was positively correlated to the severity of the condition, which in turn was associated with the prevalence of comorbidities and age[56]. Hence, post COVID-19 clinical care and close observation of the older people are important to avoid fatalities[57].

Earlier, in 2013, researchers alarmed about the threat of convergence of infectious diseases and NCDs, which will affect many in the decades to come[58]. A clear understanding of these risk factors among the older people will help the healthcare system, particularly the clinicians, to identify and implement the protocols to mitigate the fatal outcomes. In this context, increased investments in public healthcare facilities are necessary to ensure quality healthcare access[10,59], and handle post-COVID-19 syndrome situations.

Organized preventive and curative care for infectious diseases and NCDs must be ensured to the older people in the line of the 'Vayomithram' project of Kerala, where free health check-ups and medicines are available. These interventions are significant when, in India, the healthcare system's unaccountability and inefficiency resulted in privatisation of healthcare, making the common man vulnerable to high healthcare expenditures and poverty[60]. According to the 2011 census data, older people contribute about 20% of the

total population of Kerala[61] and are projected to grow at a growth rate of 2.45%[62]. One in eight persons were above 60 years, and out of the total older people, 11% of the population are 'old-old' category, 80 years and above[62]. Therefore, geriatric care must be expanded with a pool of geriatricians in the light of possible syndemic interactions of chronic diseases and their treatments[63]. Since epidemiology has a holistic approach on wellness and maintenance, older people's healthcare must become a priority before the governments, including in the vaccine distribution phase.

Ageing is inevitable, and often, the promotional and preventative aspects of geriatric care are neglected with a notion that it is 'unavoidable' and 'genetically determined' and the increased disease risk is attributed to 'normal ageing,' neglecting impact of healthier lifestyle to decreased healthcare expenditure[64]. To ensure geriatric care, there must be an attitudinal change and subsequent efforts from the policymakers to ensure 'quality ageing.' Furthermore, the public healthcare system should ensure documentation of incidents and causes of death to estimate 'excess deaths' occurring during outbreaks, wherein it becomes difficult to estimate the CFR. Weekly/ monthly death counts can be collated with trends over the years to ascertain whether it is significantly higher than the expected count. This estimation can provide information about the potential burden of mortality and fatality, associated with the infection, directly or indirectly[65].

The study finding on older people's comorbidity and their subsequent mortality determine the aetiology of COVID-19 and, therefore, prevention strategies can be implemented to avoid further spread and increased fatalities. Moreover, the study findings will add to the

existing knowledge realm on the spectrum of comorbidities among the older people and its converged impact on the phase of epidemics spread.

The study has its limitations. First, this study was carried out among the older people in Kerala, and therefore, the results may not be suitable in an international context. Second, getting authentic and correct primary data limited the study, particularly in calculating CFR, which is an estimate. Third, for the COVID-19, as in the case of previous infectious diseases, the accurate level of transmission is often underestimated as a considerable proportion of the infected population goes undetected due to asymptomatic or mild symptomatic cases failing to test for the infection[66,67]. Fourth, there will be vulnerable segments that are either neglected or under-served and under-detected who are likely to keep themselves away from testing and treatment[68].

Conclusion

It is found that older patients with chronic diseases are more susceptible to COVID-19 infection. Knowledge of these risk factors necessitates a more focused approach to introduce a spectrum of interventions to protect the lives of the older people when there is a convergence of epidemics and NCDs. The study indicates the need for having timely and accurate data of a population on their demographic profile, which can easily guide the healthcare system and authorities to more efficient prevention and treatment methodologies in health care emergencies. Having evidenced the severity of COVID-19 to the older people, health commodities such as preventing vaccines and diagnostics must be made available to them at the earliest possible.

Ethical approval

The entire study was conducted following the guidelines of the Indian Council for Social Science Research (F. No. IMPRESS/P1132/428/2018-19/ICSSR). The study was approved by the Institutional Ethics committee of the parent institution (Order No. B1/RP-ICSSR-02/ dated 22/02/2018). Informed consent was obtained from every participant prior to the start of the study after briefing about the study.

Acknowledgement

The first author acknowledges financial support from the Indian Council for Social Science Research under the IMPRESS grant (F.No. IMPRESS/P1132/428/2018-19/ICSSR). The authors acknowledge Dr Arun N. M. for granting permission to use the volunteering database on COVID-19 in Kerala. The authors acknowledge discussions with Dr Anupama M Pai, Dr Shabeesh Balan, and Dr Maya Madhavan.

Reference:

1. Marengoni A, Angleman S, Melis R *et al.* Aging with multimorbidity: A systematic review of the literature. *Ageing Res Rev* 2011;**10**:430–9.

2. UNFPA and HelpAge International. *Ageing in the Twenty-First Century: A Celebration and A Challenge*. UNFPA and HelpAge International, 2012.

3. Biswas P, Nahar Kabir Z, Nilsson J *et al.* Dynamics of Health Care Seeking Behaviour of Elderly People in Rural Bangladesh. *International Journal of Ageing and Later Life* 2006;**1**:69–89.

4. Clausen F, Sandberg E, Ingstad B *et al.* Morbidity and health care utilisation among elderly people in Mmankgodi village, Botswana. *Journal of Epidemiology & Community Health* 2000;**54**:58–63.

5. Soe Moe , Kyi Tha , Daw Khin Saw Naing , Maung Maung Than Htike. Health Seeking Behaviour of Elderly in Myanmar. *International Journal of Collaborative Research on Internal Medicine & Public Health* 2012;**4**:1538–44.

6. Woo E-K, Han C, Jo SA *et al.* Morbidity and related factors among elderly people in South Korea: results from the Ansan Geriatric (AGE) cohort study. *BMC Public Health* 2007;**7**:10.

7. Rajan SI, Aiyar S. Population ageing in India. In: Rajan SI, Risseeuw C, Perera M (eds.). *Institutional Provisions and Care for the Aged: Perspectives from Asia and Europe*. New Delhi: Anthem Press, 2008, 39–54.

8. Rajan SI. *Population Aging, Health and Social Security in India Population Aging, Health and Social Security in India Population Aging, Health and Social Security in India.*, 2007.

9. Tiwari SC, Pandey NM, Singh N. Mental health problems among inhabitants of old age homes: A preliminary study. *Indian J Psychiatry* 2012;**54**:144.

10. Mukherjee S, Levesque J-F. Morbidity and Outpatient Care for the Elderly in Kerala, South India: Evidence from a National Population based Survey. *J Popul Ageing* 2012;**5**:177–92.

11. Alam M, James KS, Giridhar G *et al. Building a Knowledge Base on Population Aging in India: Report on the Status of Elderly in Select States of India.* United Nations Population Fund, 2013.

12. Arokiasamy P. G. Giridhar, K. M. Sathyanarayana, Sanjay Kumar, K. S. James and Moneer Alam (eds), Population Ageing in India. *Ageing and Society* 2016;**36**:445–7.

13. Sandhya RS. Sociological analysis of the Problems of the oldest old (A study

conducted in Kerala). *International Research Journal of Social Sciences* 2010;**3**:57–79.

14. Karan A, Srivastava S, Chakraborty A *et al. Key Indicators of Health and Morbidity - Kerala 2014*. Public health foundation of India, 2016.

15. Madore A, Rosenberg J, Dreisbach T *et al. Positive Outlier: Health Outcomes in Kerala, India over Time | The Global Health Delivery Project*. Boston, MA: Harvard Business Publishing, 2018.

16. BusinessLine. Covid-19: Underlying ailments behind most deaths in Kerala - The Hindu BusinessLine. *The Hindu Business Line*. https://www.thehindubusinessline.com/news/covid-19-underlying-ailments-behind-most-deaths-in-kerala/article31801111.ece. Published June 11, 2020.

17. Mohan N. In Kerala COVID mortality very low, but elderly die mostly | Kerala News | Manorama. *Malayala Manorama*. https://www.onmanorama.com/news/kerala/2020/07/27/in-kerala-covid-mortality-very-low-but-elderly-die-mostly.html. Published July 27, 2020.

18. Pal R, Bhadada SK. COVID-19 and non-communicable diseases. *Postgrad Med J* 2020;**96**:429–30.

19. Callender LA, Curran M, Bates SM *et al.* The Impact of Pre-existing Comorbidities and Therapeutic Interventions on COVID-19. *Frontiers in Immunology* 2020, DOI: 10.3389/fimmu.2020.01991.

20. Zhou F, Yu T, Du R *et al.* Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;**395**:1054–62.

21. Wang D, Hu B, Hu C *et al.* Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA* 2020;**323**:1061.

22. Guan W-J, Liang W-H, Zhao Y *et al.* Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J* 2020;**55**:2000547.

23. Li J, Guo T, Dong D *et al.* Defining heart disease risk for death in COVID-19 infection. *QJM: An International Journal of Medicine* 2020, DOI: 10.1093/qjmed/hcaa246.

24. Guo W, Li M, Dong Y *et al.* Diabetes is a risk factor for the progression and prognosis of COVID-19. *Diabetes Metab Res Rev* 2020;**36**, DOI: 10.1002/dmrr.3319.

25. Petrilli CM, O'Donnell L, Horwitz LI *et al.* Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: Prospective cohort study. *BMJ* 2020.

26. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes

mellitus at increased risk for COVID-19 infection? *The Lancet Respiratory Medicine* 2020;**8**:e21.

27. Pal R, Bhansali A. COVID-19, diabetes mellitus and ACE2: The conundrum. *Diabetes Res Clin Pract* 2020;**162**:108132.

28. Emami A, Javanmardi F, Pirbonyeh N *et al.* Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: a Systematic Review and Meta-Analysis. *Archives of academic emergency medicine* 2020, DOI: 10.22037/aaem.v8i1.600.

29. Liang W, Guan W, Chen R *et al.* Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *The Lancet Oncology* 2020;**21**:335–7.

30. World Health Organization. Estimating mortality from COVID-19: Scientific brief, 4 August 2020. *WHO Scientific brief* 2020.

31. Ulahannan J, Narayanan N, Suresh S *et al.* Covid19Kerala.info-Data: A collective open dataset of COVID-19 outbreak in the south Indian state of Kerala. 2020, DOI: 10.5281/zenodo.3818096.

32. Ulahannan JP, Narayanan N, Thalhath N *et al.* A citizen science initiative for open data and visualization of COVID-19 outbreak in Kerala, India. *J Am Med Inform Assoc* 2020;**00**:1–8.

33. World Health Organization. *Coronavirus Disease 2019 (COVID-19) Situation Report – 82*. World Health Organization, 2020.

34. Glaser BG, Strauss AL. *The Discovery of Grounded Theory: Strategies for Qualitative Research*. Chicago: Aldine., 1967.

35. Charmaz K. *Constructing Grounded Theory | SAGE Publications Ltd*. London: SAGE Publications, 2006:189.

36. Roser M, Ritchie H, Ortiz-Ospina E *et al.* Coronavirus Pandemic (COVID-19). *Our World in Data* 2020.

37. Yanez ND, Weiss NS, Romand JA *et al.* COVID-19 mortality risk for older men and women. *BMC Public Health* 2020;**20**:1742.

38. Wang L, He W, Yu X *et al.* Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect* 2020;**80**:639–45.

39. Weekly CC. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. *China CDC Weekly*. February 1, 2020:113–22.

40. Kang SJ, Jung SI. Age-related morbidity and mortality among patients with COVID-

19. *Infect Chemother* 2020;**52**:154–64.

41. Paudel SS. A meta-analysis of 2019 novel corona virus patient clinical characteristics and comorbidities. *Research Square* 2020, DOI: 10.21203/rs.3.rs-21831/v1.

42. Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA - Journal of the American Medical Association* 2020;**323**:1775–6.

43. Franki R. Comorbidities the rule in New York's COVID-19 deaths | The Hospitalist. *The Hospitalist* 2020.

44. Wang B, Li R, Lu Z *et al.* Does comorbidity increase the risk of patients with COVID-19: evidence from meta-analysis. *Aging* 2020;**12**:6049–57.

45. Sanyaolu A, Okorie C, Marinkovic A *et al.* Comorbidity and its Impact on Patients with COVID-19. *SN Comprehensive Clinical Medicine* 2020;**2**:1069–76.

46. Yang X, Yu Y, Xu J *et al.* Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory Medicine* 2020;**8**:475–81.

47. Thankappan K. Combating corona virus disease 2019 and comorbidities: The Kerala experience for the first 100 days. *International Journal of Noncommunicable Diseases* 2020, DOI: 10.4103/jncd.jncd_26_20.

48. Clerkin KJ, Fried JA, Raikhelkar J *et al.* COVID-19 and Cardiovascular Disease. *Circulation* 2020;**141**:1648–55.

49. Nishiga M, Wang DW, Han Y *et al.* COVID-19 and cardiovascular disease: from basic mechanisms to clinical perspectives. *Nature Reviews Cardiology* 2020;**17**:543–58.

50. Vijayakumar G, Manghat S, Vijayakumar R *et al.* Incidence of type 2 diabetes mellitus and prediabetes in Kerala, India: results from a 10-year prospective cohort. *BMC Public Health* 2019;**19**:140.

51. Badawi A, Ryoo SG. Prevalence of comorbidities in the Middle East respiratory syndrome coronavirus (MERS-CoV): a systematic review and meta-analysis. *Int J Infect Dis* 2016.

52. Kumar DV. Kerala COVID report: Death high among people without travel history, stress on comorbidities- The New Indian Express. *The New Indian Express*. https://www.newindianexpress.com/states/kerala/2020/aug/29/kerala-covid-report-death-high-among-people-without-travel-history-stress-on-comorbidities-2189845.html. Published August 29, 2020.

53. Castle SC. Clinical relevance of age-related immune dysfunction. *Clin Infect Dis*

2000;**31**:578–85.

54. Bonafè M, Prattichizzo F, Giuliani A *et al.* Inflamm-aging: Why older men are the most susceptible to SARS-CoV-2 complicated outcomes. *Cytokine Growth Factor Rev* 2020;**53**:33–7.

55. Bhaskar Sonu, Sinha Akansha, Banach Maciej, Mittoo Shikha, Weissert Robert, Kass Joseph S., Rajagopal Santhosh, Pai Anupama R., Kutty Shelby. Cytokine storm in COVID-19—immunopathological mechanisms, clinical considerations, and therapeutic approaches: the REPROGRAM consortium position paper. *Frontiers in Immunology* 2020;**11**:1648.

56. Xiaochen Li, Shuyun Xu, Muqing Yu, Ke Wang, Yu Tao, Ying Zhou, Jing Shi, Min Zhou, Bo Wu, Zhenyu Yang, Cong Zhang, Junqing Yue, Zhiguo Zhang, Harald Renz, Xiansheng Liu, Jungang Xie, Min Xie, Jianping Zhao. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *Journal of Allergy and Clinical Immunology* 2020;**146**:110–8.

57. Kamal M, Abo Omirah M, Hussein A *et al.* Assessment and characterisation of post-COVID-19 manifestations. *Int J Clin Pract* 2020:e13746.

58. Marais BJ, Lönnroth K, Lawn SD *et al.* Tuberculosis comorbidity with communicable and non-communicable diseases: integrating health services and control efforts. *Lancet Infect Dis* 2013;**13**:436–48.

59. Levesque JF. Outpatient care utilization in urban Kerala, India. *Health Policy Plan* 2006;**21**:289–301.

60. Pal R. Analysing Catastrophic OOP Health Expenditure in India: Concepts, Determinants and Policy Implications. *Working Papers* 2010.

61. Census of India. *D1: Population Classified by Place of Birth and Sex - 2011*. New Delhi: Office of the Registrar General & Census Commissioner, India, 2011.

62. Rajan SI, Shajan A, Sunitha S. Ageing and Elderly Care in Kerala. *China Rep* 2020;**56**:354–73.

63. National Academies of Sciences, Engineering. *The Convergence of Infectious Diseases and Noncommunicable Diseases*. Ogawa VA, Shah CM, Negussie Y, et al. (eds.). Washington, D.C.: National Academies Press, 2019.

64. Prabhu P, Singh FMA. Nutrition the Geriatric Population. *Geriatric Care*. Cochin: Brookdale Foundation, 2002, 56–65.

65. Centers for Disease Control and Prevention. Excess deaths associated with COVID-19: Provisional Death Counts for Coronavirus Disease (COVID-19). *National Center for Health Statistics* 2020.

66. Kim GU, Kim MJ, Ra SH *et al.* Clinical characteristics of asymptomatic and symptomatic patients with mild COVID-19. *Clin Microbiol Infect* 2020, DOI: 10.1016/j.cmi.2020.04.040.

67. Nishiura H, Kobayashi, Jung T *et al.* Communicating the Risk of Death from Novel Coronavirus Disease (COVID-19). Journal of Clinical Medicine, 9(2), 580. *J Clin Med Res* 2020.

68. Lau H, Khosrawipour T, Kocbach P *et al.* Evaluating the massive underreporting and undertesting of COVID-19 cases in multiple global epicenters. *Pulmonology* 2020, DOI: 10.1016/j.pulmoe.2020.05.015.

69. Department of Health & Family Welfare. *COVID-19 Death Audit Report*. Government of Kerala, 2020.

Table I. Socio-Demographic Variables

Variable	Category	Frequency	Percent
Gender	Female	181	44.7
	Male	224	55.3
Age	60-69	203	50.1
	70-79	128	31.6
	80-89	60	14.8
	Above 90	9	2.2
Marital Status	Married	314	77.5
	Not Married	11	2.7
	Partner Died	75	18.5
	Divorced	3	0.7
Education	No formal Education	74	18.3
	School	267	65.9

	College/University	53	13.1
	No response	4	1
Monthly income	No income	125	25.9
	Less than INR 10,000	228	56.3
	INR 10,000-20,000	37	9.1
	INR 21,000-30,000	20	4.9
	Above INR 30,000	5	1.2
Source of Income	Job	18	4.4
	Pension	211	52.1
	Income from assets	29	7.2
	Business	35	8.6
	Income from agriculture	49	12.1
	Income from other sources	15	3.7

Table II. Health Status

Category	Frequency	Percentage
Very Good	31	7.7
Good	84	20.7
Satisfactory	182	44.9
Bad	84	20.7
Very Bad	24	5.9

Table III. Health Behaviour of the elderly in Kerala

Variables	Regularly	Frequently	Sometimes	Rarely	Never
Exercise	64 (15.8%)	68 (16.8%)	52 (12.8%)	40 (9.9%)	180 (44.4%)
Diet	129 (31.9%)	117 (28.9%)	70 (17.3%)	34 (8.4%)	55 (13.6%)
Smoking	30 (7.4%)	19 (4.7%)	9 (2.2%)	15 (3.7%)	332 (82%)
Alcohol	1 (0.2%)	15 (3.7%)	26 (6.4%)	17 (4.2%)	346 (85.4%)
Chewing tobacco	14 (3.5%)	6 (1.5%)	10 (2.5%)	7 (1.7%)	368 (90.9%)

Table IV. Assistance required in ADL

Activities	Assistance Not Required N (%)	Assistance required Sometimes N (%)	Full Assistance Required N (%)
Eating	365 (90.1%)	28 (6.9%)	12 (3%)

Get-up/ sit down	303 (74.8%)	82 (20.2%)	20 (4.9%)
Walking	317 (78.3%)	66 (16.3%)	22 (5.4%)
Toileting	367 (90.6%)	22 (5.4%)	16 (4%)
Dressing	365 (90.1%)	20 (4.9%)	20 (4.9%)
Bathing	360 (88.9%)	26 (6.4%)	19 (4.7%)

Table V. Assistance required for IADL

Activities	Assistance not Required N (%)	Assistance required Sometimes N (%)	Full Assistance Required N (%)
Telephone	214 (52.8%)	70 (17.3%)	121 (29.9%)
Shopping	198 (48.9%)	63 (15.6%)	144 (35.6%)
Cooking	199 (49.1%)	43 (10.6%)	163 (40.2%)
Household work	188 (46.4%)	57 (14.1%)	160 (39.5%)
Washing	200 (49.4%)	51 (12.65)	154 (38%)
Travelling	194 (47.9%)	70 (17.3%)	141 (34.8)
Medication	262 (64.7%)	62 (15.3%)	81 (20%)
Finance	266 (65.7%)	51 (12.6%)	88 (21.7%)

Table VI. Elderly Disease Profile in Kerala

Type of Disease	Regularly	Frequently	Sometimes	Rarely	Never
Diabetes	156 (38.5%)	37 (9.1%)	9 (2.2%)	6 (1.5%)	197 (48.6%)
Hypertension	145 (35.8%)	66 (16.3)	29 (7.2%)	19 (4.7%)	146 (36%)
Cataract	56 (13.8%)	30 (7.4%)	35 (8.6%)	30 (7.4%)	253 (62.5%)
Urinary Disease	24 (5.9%)	27 (6.7%)	33 (8.1%)	43 (10.6%)	278 (68.6%)
Lungs Disease	32 (7.9%)	8 (2%)	14 (3.5%)	25 (6.2%)	326 (80.5%)
Heart Disease	46 (11.4%)	18 (4.4%)	19 (4.7%)	9 (2.2%)	313 (77.3%)
Depression	12 (3%)	12 (3%)	13 (3.2%)	37 (9.1%)	331 (81.7%)
Cancer	17 (4.2%)	4 (1%)	3 (0.7%)	3 (0.7%)	378 (93.3%)
Rheumatic Disease	57 (14.1%)	35 (8.6%)	36 (8.9%)	42 (10.4%)	235 (58%)

Abdominal Disease	25 (6.2%)	13 (3.2%)	31 (7.7%)	42 (10.4%)	294 (72.6%)

Table VII. Disease and health behaviour of elderly - Qualitative data

Attribute	Responses (District-participant ID, Gender, Age)
Disease	..I have Diabetes for the past 35 years, severe joint pain, back pain... (Palakkad, F, 70-75) …there are number of diseases in this age…headache, back pain, blood pressure, disk problem, cholesterol (Pathanamthitta, F, 60-65) I am taking medicines for blood pressure regularly…. (Pathanamthitta, M, 60-65) ….I had a block in my heart, for which cardiac surgery was done. Check-up is done every three months. Then I had prostate cancer… follow up visits once in six months. I am a diabetic patient too... (Thrissur, M, 65-70) …angioplasty was done ….taking medicines for blood pressure twice a day." (Kozhikode, F, 70-75) "I am a cardiac patient, using pacemaker…. (Palakkad, F, 70-75) "Taking medicines for thyroid, sugar, BP daily… (Thiruvananthapuram, F, 65-70) "My current health problems are Asthma, Cholesterol, Sugar, blood pressure, thyroid, Cardiac diseases…… (Kollam, F, 70-75) I have cancer, BP, Heart problem, doing chemotherapy (Malappuram, M, 70-75)

Health Behaviour (Exercise)	…Many years before, I learned a special Japan exercise. Since I worked in companies, I got training for warm up exercise. That I continue for half an hour… I walk daily 4 kilometres…. (Pathanamthitta, M, 60-65) ……my exercise is my housework There is lots of work in the house. That itself is an exercise….. (Alappuzha, M, 60-65) 'I have a morning walk regularly '(Thrissur, M, 60-65) …I go for rubber tapping daily by walking 2 kilometres in the morning daily. After that I have to feed my cattle. All these are my exercises…… (Kozhikode, M, 60-65) …Doing exercise for the past two years regularly, suggested by doctor after one surgery…… (Thiruvananthapuram, F, 70-75) …. Daily exercise is there for 1 hour. Every evening @ 5 pm I walk for one hour. (Palakkad, F, 65-70)
Diet	I do not eat sugar, no spicy food, no oil in preparing food, just eating vegetables half cooked and fruits (Kasaragod, F, 60-65) Food control is maintained (Thrissur, M, 63-65) Restrict myself from eating fried items (Kozhikode, M, 60-65) Eat food in usual quantities. Do not use sugar and meat…….(Wayanad, M, 70-75).

Table VIII. COVID-19 Indicators (CFR, CMR, IFR)

Note: Kerala population as per the Census data of 2011: 35,699,443. The number of the elderly population as per the Census data of 2011: 4, 193,393. The data on the comorbidity of the elderly death cases are not available after August 2020 even when there was a sudden curve in the fatality rate in Kerala

Type of data	No. of infected cases	No of elderly with comorbidity	No of deaths	CFR %	CMR %	IFR %
DHS Dashboard (*as on 27 December 2020*)	740516	No Data	2976	0.44	0.008	0.40
Crowdsourced dashboard (*as on 27 December 2020*)	740516	No Data	2976	0.41	0.008	0.37
Volunteering dashboard (*as on 26 December 2020*)	735611	Data available till September 30	4396	0.65	0.012	0.60

Figure 1. Covid-19 Deaths in Kerala-Age wise till December 27, 2020

Source: DHS, Dashboard

Figure 2. Types of Comorbidities Prevalent in Fatality Cases of COVID-19

Source: Death Audit Report, DHS, Government of Kerala (Department of Health &

Family Welfare 2020)

Figure 3. Types of Elderly Comorbidities Prevalent in Fatality Cases of COVID-19

Source: Source: Compiled from dashboards