Polygenic risk for immuno-metabolic markers and specific depressive symptoms: A multi-sample network analysis study

Nils Kappelmann[1,2], Darina Czamara[1], Nicolas Rost[1,2], Sylvain Moser[1,2], Vanessa Schmoll[1],

Lucia Trastulla[1], Jan Stochl[3,4], Susanne Lucae[5], CHARGE inflammation working group,

Elisabeth B. Binder[1], Golam M. Khandaker[3,6,7,8,*], & Janine Arloth[1,9,*]

[1]Department of Translational Research in Psychiatry, Max-Planck-Institute of Psychiatry, Munich, Germany
[2]International Max Planck Research School for Translational Psychiatry (IMPRS-TP), Munich, Germany
[3]Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom
[4]Department of Kinanthropology, Charles University, Prague, Czech Republic
[5]Max-Planck-Institute of Psychiatry, Munich, Germany
[6]Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, United Kingdom
[7]MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom
[8]Centre for Academic Mental Health, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom
[9]Institute of Computational Biology, Helmholtz Zentrum Munich, Neuherberg, Germany

*joint senior authors

Corresponding author:

Nils Kappelmann
Max-Planck-Institute of Psychiatry
Kraepelinstraße 2-10
80804 Munich, Germany

Email: nils_kappelmann@psych.mpg.de; n.kappelmann@gmail.com

Abstract word count: 399

Manuscript word count: 6,306

Tables and Figures: 2 Tables and 3 Figures

Online Supplementary Tables and Figures: 7 Tables and 9 Figures

ABSTRACT

Background

About every fourth patient with major depressive disorder (MDD) shows evidence of systemic

inflammation. Previous studies have shown inflammation-depression associations of multiple

serum inflammatory markers and multiple specific depressive symptoms. It remains unclear,

however, if these associations extend to genetic/lifetime predisposition to higher inflammatory

marker levels and what role metabolic factors such as Body Mass Index (BMI) play. It is also

unclear whether inflammation-symptom associations reflect direct or indirect associations, which

can be disentangled using network analysis.

Methods

This study examined associations of polygenic risk scores (PRSs) for immuno-metabolic markers

(C-reactive protein [CRP], interleukin [IL]-6, IL-10, tumour necrosis factor [TNF]-α, BMI) with

seven depressive symptoms in one general population sample, the UK Biobank study

(n=110,010), and two patient samples, the Munich Antidepressant Response Signature (MARS,

n=1,058) and Sequenced Treatment Alternatives to Relieve Depression (STAR*D, n=1,143)

studies. Network analysis was applied jointly for these samples using fused graphical least

absolute shrinkage and selection operator (FGL) estimation as primary analysis and, individually,

using unregularized model search estimation. Stability of results was assessed using

bootstrapping and three consistency criteria were defined to appraise robustness and replicability

of results across estimation methods, network bootstrapping, and samples.

Results

Network analysis results displayed to-be-expected PRS-PRS and symptom-symptom associations

(termed edges), respectively, that were mostly positive. Using FGL estimation, results further

suggested 28, 29, and six PRS-symptom edges in MARS, STAR*D, and UK Biobank samples,

respectively. Unregularized model search estimation suggested three PRS-symptom edges in the

UK Biobank sample. Applying our consistency criteria to these associations indicated that only

the association of higher CRP PRS with greater changes in appetite fulfilled all three criteria.

Four additional associations fulfilled at least two consistency criteria; specifically, higher CRP

PRS was associated with greater fatigue and reduced anhedonia, higher TNF-α PRS was

associated with greater fatigue, and higher BMI PRS with greater changes in appetite and

anhedonia. Associations of the BMI PRS with anhedonia, however, showed an inconsistent

valence across estimation methods.

Conclusions

Genetic predisposition to higher systemic inflammatory markers are primarily associated with

somatic/neurovegetative symptoms of depression such as changes in appetite and fatigue,

consistent with previous studies based on circulating levels of inflammatory markers. We extend

these findings by providing evidence that associations are direct (using network analysis) and

extend to genetic predisposition to immuno-metabolic markers (using PRSs). Our findings can

inform selection of patients with inflammation-related symptoms into clinical trials of immune-

modulating drugs for MDD.

Keywords*: Depression; Depressive Symptoms; Inflammation; C-reactive protein; Body Mass

Index; Interleukin 6; Interleukin 10; Tumour Necrosis Factor-α; Network Analysis

INTRODUCTION

Recent findings suggest that every fourth patient with Major Depressive Disorder (MDD) shows evidence of systemic, low-grade inflammation as indicated by elevated (>3mg/L) C-reactive protein (CRP) concentrations (Osimo et al., 2019). This association has been supported by cross-sectional case-control studies synthesised in multiple meta-analyses (Dowlati et al., 2010; Goldsmith et al., 2016; Haapakoski et al., 2015; Howren et al., 2009; Köhler et al., 2017) as well as longitudinal studies (Khandaker et al., 2014; Lamers et al., 2020; Mac Giollabhui et al., 2020). Clinically, patients with evidence of inflammation do not respond as well to standard monoaminergic and psychotherapeutic treatments (Liu et al., 2020; Lopresti, 2017). These patients may, however, benefit from alternative treatment with immune-modulating drugs (Kappelmann et al., 2018; Köhler-Forsberg et al., 2019; Wittenberg et al., 2020). To prioritise drug and patient selection for clinical trials, it is crucial to further understand immunological and clinical complexity of inflammation-symptom associations, which may allow shortlisting of promising immunotherapeutic drug targets and could highlight patients with a profile of inflammation-related depression.

Regarding immunological complexity, studies have reported various associations of serum inflammatory proteins with depression, including among others CRP, interleukin (IL)-6, IL-10, and tumour necrosis factor (TNF)-α (Goldsmith et al., 2016; Haapakoski et al., 2015; Köhler et al., 2017). Evidence from in-depth immunophenotyping further suggests that there may be distinct subgroups of inflammation-related depression as shown by immune cell count clustering and transcriptome analyses (Cattaneo et al., 2020; Lynall et al., 2020). These studies suggest that elevated serum levels of inflammatory markers are associated with depression, but associations of depression with genetic/lifetime predisposition to higher inflammatory markers has been studied

less frequently and primarily for CRP (Badini et al., 2020; Kappelmann et al., 2021; Milaneschi

et al., 2017b, 2016). Elevated serum levels of inflammatory markers also conflate tonic and

phasic levels of inflammatory markers while genetic/lifetime predisposition to inflammatory

markers specifically maps their tonic levels. This differentiation could be relevant as highlighted

by research into tonic versus phasic dopamine levels (see Bilder et al., 2004), whereby tonic

levels regulate the amplitude of the phasic response, which has unique consequences for

downstream signalling. Lastly, inflammatory markers such as CRP are influenced by metabolic

factors (Timpson et al., 2011), which may causally underlie some inflammation-symptom

associations (Kappelmann et al., 2021), so a combined investigation of immuno-metabolic factors

is needed to disentangle their etiological roles.

Regarding clinical complexity, most prior research has restricted its investigation of the

inflammation-depression association to complexity on one side, that is focusing on multiple

immune markers (e.g., cell counts/ serum cytokine levels) while studying a composite depression

phenotype (Goldsmith et al., 2016; Haapakoski et al., 2015; Köhler et al., 2017) or focusing on

multiple depressive symptoms or symptom groups in the context of a single inflammatory marker

(mostly CRP) (Badini et al., 2020; Jokela et al., 2016; Köhler-Forsberg et al., 2017; Lamers et al.,

2020, 2019; White et al., 2017). Among studies focusing on individual symptoms, results have

highlighted associations of inflammatory markers with specific depressive symptoms of fatigue,

changes in appetite, anhedonia, and suicidality (Badini et al., 2020; Chu et al., 2019; Jokela et al.,

2016; Kappelmann et al., 2021; Köhler-Forsberg et al., 2017; Lamers et al., 2020, 2018;

Milaneschi et al., 2017a; Simmons et al., 2018; White et al., 2017). However, most of these

studies have considered associations of inflammatory markers with each depressive symptom in

isolation (Chu et al., 2019; Jokela et al., 2016; Kappelmann et al., 2021; Köhler-Forsberg et al.,

2017; Lamers et al., 2018; White et al., 2017). Although these prior approaches have led to

important findings, they cannot address potential causal interactions between symptoms, thus

conflate evidence for indirect and direct associations. For example, analyses of isolated

symptoms could hypothetically provide evidence for associations of CRP with both fatigue and

sleep problems even if CRP was only indirectly associated with fatigue *via* its effect on sleep

problems. A network-based approach provides one means of disentangling such direct from

indirect inflammation-symptom associations.

Network theory and related analysis techniques have recently been put forward to accommodate

the symptomatic complexity of mental disorders (Borsboom, 2017). Network theory proposes

putative causal interactions between symptoms (e.g., fatigue *causing* concentration problems

causing low mood), which could result in self-reinforcing vicious symptom cycles triggering and

maintaining mental disorders. Such associations have been investigated in an increasing amount

of studies on psychological symptom networks (Contreras et al., 2019; Robinaugh et al., 2020).

To accommodate etiological factors beyond symptoms, however, recent work has proposed an

expansion of symptom networks to so-called 'multi-plane' networks, for instance also including

genetic, metabolic, immunological, or environmental variables (Guloksuz et al., 2017). To our

knowledge, so far, two studies have evaluated such multi-plane networks in the context of

inflammation and depression by jointly analysing serum CRP (plus IL-6 & TNF-α in the study of

Fried et al., 2019), BMI, and potential covariates with individual depressive symptoms (Fried et

al., 2019; Moriarity et al., 2020a). The most consistently replicated findings between these two

studies suggested unique associations of CRP with fatigue and changes in appetite. A third study

has recently also provided evidence that the symptom structure itself was a function of CRP

levels; that is, interconnections between symptoms were moderated by CRP (Moriarity et al.,

2020b). All of these previous studies were based on serum markers for inflammatory proteins,

however, reflective of acutely elevated inflammatory activity. Therefore, it remains unclear if

inflammation-symptom associations generalise to genetic/lifetime predisposition to higher

immuno-metabolic marker levels.

In the present study, we explored associations of polygenic risk scores (PRSs) for four major pro-

and anti-inflammatory markers (i.e., CRP, IL-6, IL-10, & TNF-α) and Body Mass Index (BMI),

as a metabolic marker, with individual depressive symptoms using a multi-sample, multi-plane

network analysis approach. We evaluated associations in three large samples including the

inpatient Munich Antidepressant Response Signature (MARS) study (n=1,058), the outpatient

Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study (n=1,143), and the

general population UK Biobank cohort (n=110,010) (Hennings et al., 2009; Rush et al., 2004;

Sudlow et al., 2015). This investigation aimed to contribute to the study of inflammation and

depression by simultaneously addressing (i) combined immunological and symptom complexity

(using network analysis), (ii) unclarity regarding the influence of genetic/lifetime predisposition

to higher immuno-metabolic marker levels on depression (defining immuno-metabolic markers

using PRSs), and (iii) issues of replicability and generalisability (testing associations in one large

general population and two clinical samples).

METHODS

An overview of the study design and analytic procedure is presented in Figure 1.

-Figure 1-

Study samples

The Munich Antidepressant Response Signature (MARS) study was a naturalistic, observational study of inpatients with major depressive disorder (MDD) or bipolar disorder conducted between 2000 and 2015 in three Southern German hospitals (Hennings et al., 2009). Based on an original sample of 1,411 patients, the present study included 1,058 patients of European decent with an ICD diagnosis of MDD (F32 and F33 codes) and genetic and depressive symptom data.

The STAR*D trial (identifier: NCT00021528) was a multisite, multistep, randomised controlled trial (RCT), conducted from 2000 to 2004, evaluating different treatment options and sequences for outpatients suffering from DSM-IV MDD without psychotic features (Rush et al., 2004). Based on an original sample of 1,953 patients who took part in the STAR*D genetics study, the present study included 1,143 individuals of European decent with genetic and depressive symptom data.

The UK Biobank is a general population cohort including more than 500,000 individuals, recruited from 2006 to 2010, with genotyping and in-depth phenotyping information (Bycroft et al., 2018). More than 150,000 individuals from the initial sample took part in a follow-up mental health survey (Davis et al., 2020) and we included a subset of 110,010 individuals that were of European decent and had available genetic and depressive symptom data.

Ethics approval and informed consent

MARS received local ethics approval from Ludwig Maximilians University Munich (Hennings et

al., 2009). STAR*D received ethics approval from 14 participating institutional review boards, a

National Coordinating Center, a Data Coordinating Center, and the Data Safety and Monitoring

Board at the National Institute of Mental Health (Rush et al., 2006, 2004). The UK Biobank study

received ethics approval from North West Centre for Research Ethics Committee and Human

Tissue Authority research tissue bank (Bycroft et al., 2018); this project was approved under

project no. 26999. All three studies collected informed consent from participants prior to study

participation.

Depressive symptom assessment

Depressive symptoms were assessed differently across the three samples. MARS and STAR*D

studies used the observer-rated Hamilton Rating Scale for Depression (HAM-D) (Hamilton,

1986) while the UK Biobank study used the self-report Patient Heath Questionnaire (PHQ)-9

(Löwe et al., 2004). From these questionnaires, we selected seven depressive symptoms for joint

analyses across samples. These symptoms included completely overlapping symptoms of

depressed mood, anhedonia, fatigue, and suicidality, but also partially overlapping symptoms of

sleep problems, changes in appetite, and psychomotor changes. Supplementary Table 1 provides

an item-level overview of depressive symptoms and Supplementary Table 2 displays symptom

coding, where this differed from original Likert scale ratings.

Regarding partially overlapping symptoms of sleep problems, changes in appetite, and

psychomotor changes, the PHQ-9 only assesses information on conflated symptoms (e.g.,

insomnia and hypersomnia are conflated to sleep problems) while the HAM-D incorporates

disaggregated symptoms. To harmonise these symptom data for retention in network analyses,

we conflated HAM-D symptoms of psychomotor retardation and agitation to "psychomotor

changes". For sleep problems and changes in appetite (available in the PHQ-9), only insomnia

and loss of appetite are available in the HAM-D, so we included both conflated and

unidirectional symptoms in network analyses as previous studies have specifically highlighted

associations of inflammation with these symptoms (Jokela et al., 2016; Milaneschi et al., 2017b).

We reasoned that comparative appraisal of associations, for example with changes in appetite and

loss of appetite, could give further indications on potential specificity of associations to symptom

directions, as observed in previous reports (Kappelmann et al., 2021; Milaneschi et al., 2021b,

2021a).

We also note that we have not included items of "guilt or self-blame" from the respective studies

in our analyses as we considered the item content of HAM-D and PHQ-9 too distinct.

Specifically, the HAM-D conflates feelings of guilt with delusions of guilt and death, thus

moving towards psychotic symptomatology. Contrary to this, the PHQ-9 also includes "feelings

of inadequacy" about oneself, which are not covered by the HAM-D item.

Genotyping, quality control and imputation

We provide detailed information on genotyping, quality control and imputation procedures in the

Supplementary Methods. Briefly, genotyping in the MARS study was conducted using three

genotyping arrays across the recruitment period (see Supplementary Figure 1), the Illumina 610k

(n=548), Illumina OmniExpress (n=284) and Illumina GSA (n=226) arrays. In STAR*D,

genotyping was conducted using the Affymetrix Human Mapping 500K Array Set (n=979) and

the Affymetrix Genome-Wide Human SNP Array 5.0 (n=969) that displayed a concordance of

>99%; described in detail by Garriock *and colleagues* (2010). In the UK Biobank study, samples

were genotyped on the UK BiLEVE Axiom Array or the Affymetrix UK Biobank Axiom Array

(Bycroft et al., 2018). Following imputation in all samples, single nucleotide polymorphisms

(SNPs) with info-metric>0.6, minor allele frequency (MAF)>1%, genotyping missingness<2%,

and no deviation from Hardy-Weinberg Equilibrium (MARS & STAR*D: P>1e^{-5}; UK Biobank:

P>1e^{-7}) were retained.

Polygenic risk scores

Immuno-metabolic marker selection and GWAS data sources

PRSs for CRP, IL-6, IL-10, TNF-α, and BMI were computed based on available summary

statistics from genome-wide association studies (GWAS; Ahola-Olli et al., 2017; Ligthart et al.,

2018; Locke et al., 2015). These inflammatory markers were selected, because (i) they showed

robust differences in case-control studies; (ii) CRP, IL-6, and TNF-α have been the most

frequently investigated inflammatory markers overall in the context of depression; and (iii) IL-10

was the most frequently studied anti-inflammatory cytokine, so could be informative on direction

of associations between depressive symptoms and innate immune activity (Köhler et al., 2017;

Osimo et al., 2019). BMI was selected as the most frequently investigated metabolic marker.

GWAS data for CRP were obtained from a large GWAS of 88 studies including 204,402

individuals of European decent (Ligthart et al., 2018). GWAS data for IL-6, IL-10, and TNF-α

were obtained from a GWAS of 8,293 Finns (Ahola-Olli et al., 2017); of note, Finns have

Siberian ancestry (Lamnidis et al., 2018), which leads to a divergence from European ancestry of

our analytic samples. GWAS data for BMI were obtained from the Genetic Investigation of

Anthropometric Traits (GIANT) consortium that included up to 322,154 individuals of European

decent (Locke et al., 2015).

PRS computation

PRSs can be computed by summing the GWAS association estimates of risk alleles for each

individual. Classically, this summation is done using an approach termed "clumping and

thresholding" (C+T), which first reduces summary statistics to independent SNPs and then

applies one or multiple thresholds (usually based on P-values) to restrict summation to SNPs with

high evidence for associations with phenotypes (Choi et al., 2020). As the optimal threshold for

the C+T approach is unknown and should ideally be estimated in a separate dataset with available

phenotype data, we computed PRSs using the Bayesian regression and continuous shrinkage

priors (PRS-CS) approach, which has been shown to perform similar to or outperform other PRS

computation approaches such as C+T (Ge et al., 2019; Ni et al., 2020).

PRS-CS takes a linkage disequilibrium (LD) reference panel into account (we used European

ancestry data from 1000 Genomes Project phase 3 samples) to update SNP effect sizes in a

blocked fashion, thus providing accurate LD adjustment. We pre-specified the global shrinkage

parameter $\square$ using suggested defaults for less polygenic ($\square=1e^{-4}$) and more polygenic ($\square=1e^{-2}$)

phenotypes as $\square=1e^{-4}$ for CRP, IL-6, IL-10, and TNF-α, and as $\square=1e^{-2}$ for BMI; see details in

Supplementary Methods. Following PRS computation in individual samples, polygenic scores

were corrected for age, sex, and the first two genotyping principal or multidimensional scaling

(MDS) components using linear regression; two genotyping principal or MDS components were

selected as visual inspection of component inter-correlations did not suggest evidence for

population stratification. Genotyping MDS components were computed based on raw Hamming-

distances in MARS and STAR*D, and using principal component analysis on high-quality,

unrelated individuals in the UK Biobank sample (Bycroft et al., 2018). PRSs in MARS were

additionally corrected for the genotyping array. Following computation, higher PRSs reflect

higher genetic predisposition to respective immuno-metabolic phenotype levels.

PRS evaluation

In Supplementary Table 3, we provide the number of SNPs included in PRS computation in each

sample, which was approximately around one million SNPs for each phenotype-sample

combination. The proportion of SNP overlap between samples (for the same phenotype) was

>0.89 suggesting that mostly overlapping SNPs contributed to PRSs (Supplementary Table 4).

Taking these overlapping SNP sets, correlations between the posterior SNP effect sizes between

samples were large for CRP (Pearson's *r* range: 0.69-0.76) and BMI (Pearson's *r* range: 0.79-

0.80) and relatively smaller for IL-6, IL-10, and TNF-α (Pearson's *r* range: 0.41-0.46; see

Supplementary Table 5). This suggests polygenic risk was quantified more similarly across

samples for CRP and BMI as compared to IL-6, IL-10, and TNF-α.

We quantified the impact that pre-specification of the hyperparameter □ had on resulting PRSs,

which was likely small (Supplementary Table 6). Specifically, PRSs with pre-specified □

exhibited large correlations with PRSs based on automatic learning of □ from GWAS summary

data (termed PRS-CS-auto in the literature; Pearson's *r* range: 0.82-0.98). Furthermore,

moderate-to-large correlations remained to PRSs based on extreme grid search boundary values

of □ (Pearson's *r* range: 0.47-0.93).

Since MARS utilised three different genotyping arrays, we verified that our approach of

combining data from these arrays into one sample was justified before proceeding with the main

analysis (see Supplementary Methods and Supplementary Figures 2 and 3).

Network analysis

Estimation

Network analysis was conducted using *R* software (version 4.0.3; R Core Team, 2017). In

network analysis, unique associations between variables reflect partial correlations and are

termed 'edges'. Variables in the network are referred to as 'nodes'.

Network models can be broadly categorised into regularized and unregularized models, that have

distinct advantages and disadvantages. Regularised models apply penalties that shrink edges

towards zero. This has the advantage that it results in sparser and more parsimonious network

models as small edges can be set exactly to zero. Contrary to this, non-regularized models do not

apply such a penalty while still controlling the false positive rate and recent studies have

suggested that unregularized models perform better in estimating psychological symptom

networks and multi-plane immunopsychiatric networks than regularized network models

(Moriarity et al., 2020a; Williams et al., 2019). A disadvantage of unregularized models,

however, is that they are currently only suitable for network estimation of individual

samples/datasets. Contrary to this, regularised models have recently been adapted for application

in multi-sample contexts using so-called fused graphical LASSO (FGL) estimation. FGL

estimation allows synthesising data across multiple samples, which increases statistical power.

Based on these respective advantages and disadvantages, we have decided to use a regularized

network model as primary analysis, which maximises statistical power due to the multi-sample

design of our study. As unregularized models are preferable for estimation of individual samples

and may be better suited to retrieve multi-plane edges, however, we also apply unregularized

network estimation as secondary analysis.

In primary analyses, networks were estimated using FGL estimation as implemented in the

EstimateGroupNetwork package (version 0.2.2; Costantini et al., 2020, 2019; Danaher et al.,

2014). FGL estimation relies on the two tuning parameters λ_1, which penalizes network density,

and λ_2, which penalizes edge differences across samples. Values for these tuning parameters were

selected using 10-fold cross-validation to optimise the Bayesian Information Criterion (BIC). As

recommended, we set weights for the importance of each sample as 'equal' to ascertain that a

single sample would not dominate estimation (Danaher et al., 2014).

As secondary analysis, we estimated unregularized networks for each sample individually using

the gaussian graphical stepwise model selection ("ggModSelect") algorithm implemented in the

qgraph package (version 1.6.5; Epskamp et al., 2012) based on Spearman correlations and

starting from an empty model. Throughout results, we refer to this estimation strategy as

"unregularized model search" or "model search" for simplification.

Node predictability

We also estimated node predictability, which describes the amount of variance in a node that is

explained by all other nodes in the network, so can be interpreted akin to R^2 (Haslbeck and Fried,

2017). Node predictability cannot be inferred from FGL or model search networks as it requires a

node-wise estimation approach. Therefore, we used a mixed graphical model as a third estimation

strategy as implemented in the *mgm* package (version 1.2-10), selecting tuning parameter λ based

on BIC optimisation in 10-fold cross-validation (Haslbeck and Waldorp, 2020). Of note, this

model was only used to infer node predictability, which provides additional information on

network density and sample comparability. However, we do not report any individual edge

estimates based on this model as FGL estimation and unregularized model search are better

suited for our study aims.

Visualisation

Networks were visualised with the *qgraph* package using an average layout estimated with the

Fruchterman-Reingold algorithm for the FGL networks. This algorithm places nodes close to

each other that are connected by large edges (Epskamp et al., 2012). While this simplifies

network appraisal, it is important to note that nodes and edges should not be interpreted based on

their relative position within the network, which can be unstable.

Stability

To evaluate stability of estimated networks, we assessed accuracy of edge estimates using

bootstrapping strategies. Specifically, for FGL networks 500 bootstrapped samples with

replacement were drawn, and FGL networks re-estimated, using the implementation in the

EstimateGroupNetwork package (Costantini et al., 2020). For unregularized model search

estimation, the same procedure was applied using non-parametric bootstrapping procedures

implemented in the *bootnet* package (version 1.4.3; Epskamp et al., 2018).

Interpretation

We interpreted estimated networks based on the presence, stability, and replicability of edges as

defined using three consistency criteria. First, we tested if edges were nonzero in FGL networks

as well as nonzero and directionally consistent in >50% of bootstrapped analyses (consistency

criterion 1) akin to a previous PRS-symptom network study in psychosis by Isvoranu *and*

colleagues (2020). Second, we tested if edges between PRSs and symptoms replicated (according

to criterion 1) across FGL networks of the three samples (consistency criterion 2). Third, we

tested if edges were present in secondary analyses using unregularized model search estimation,

again confirmed in >50% of bootstrapped estimations exhibiting directionally consistent

estimates (consistency criterion 3).

Availability of data and materials

Data from original studies is not openly available, but can be requested; see details in

Supplementary Table 7. GWAS summary data for IL-6, IL-10, and TNF-α is openly available

from the original publication by Ahola-Olli *and colleagues* (2017), for BMI from the GIANT

consortium, and can be requested for CRP from the CHARGE inflammation working group. We

provide analysis scripts and estimated network matrices (including bootstrapped network

matrices) on the Open Science Platform (OSF) under https://osf.io/q4vw9/.

RESULTS

Baseline characteristics of study populations are displayed in Table 1.

-Table 1-

Network analysis

We conducted network analyses of five immuno-metabolic PRSs (CRP, IL-6, IL-10, TNF-α, &

BMI) and seven depressive symptoms using two estimation techniques (FGL & unregularized

model search estimation) in three samples (MARS, STAR*D & UK Biobank). Bootstrap

analyses were conducted to assess stability of networks and node predictability estimated using a

mixed graphical model. We defined three consistency criteria to assess robustness and

replicability of our results across estimation techniques, bootstrapping, and samples. Focus of this

network investigation were unique associations (termed edges in network analysis) between PRSs

and symptoms, which are summarised in Table 2.

Fused Graphical LASSO (FGL) estimation suggests four consistent PRS-symptom edges
according to criteria 1 & 2

Using FGL estimation, we obtained networks that are visualised in Figure 2. PRS-symptom edge

bootstrapping results are displayed in Figure 3 with PRS-PRS and symptom-symptom edge

bootstrapping results shown in Supplementary Figures 4 and 5.

As expected, nodes within the same plane displayed relatively stronger within-plane (i.e.,

symptom-symptom & PRS-PRS) than between-plane (i.e., PRS-symptom) associations. Among

PRSs, CRP displayed associations with BMI (edge weight range across samples: 0.16-0.19) while

IL-6, IL-10, and TNF-α (based on the same GWAS) were associated with each other (edge

weight range across samples: 0.08-0.52). Associations of BMI and CRP with IL-6, IL-10, and

TNF-α were largely absent or very small (edge weight range across samples: -0.02-0.01). Among

symptoms, the largest associations were present between the core symptoms depressed mood and

anhedonia (edge weight range across samples: 0.14-0.55), which is to-be-expected in clinical

samples where these symptoms form the basis of the MDD diagnosis. Supplementary Figure 4

also illustrates interesting edge differences between samples that are likely arising from the

diverging symptom definitions in individual samples. For instance, edges of fatigue with changes

in appetite (edge weight=0.21) and sleep problems (edge weight=0.33) were relatively larger in

the UK Biobank, assessing composite symptoms of changes in appetite and sleep problems, but

substantially smaller in MARS (fatigue-changes in appetite: edge weight=0.09; fatigue-sleep

problems: edge weight=0.10) and STAR*D (fatigue-changes in appetite: edge weight=-0.01;

fatigue-sleep problems: edge weight=-0.01), assessing loss of appetite and insomnia.

Regarding PRS-symptom edges, FGL estimation surprisingly resulted in a much larger number of

PRS-symptom edges in MARS and STAR*D samples compared to the UK Biobank sample with

28 (MARS), 29 (STAR*D), and 6 (UK Biobank) nonzero PRS-symptom edges. 26 (MARS), 28

(STAR*D), and 5 (UK Biobank) of these edges fulfilled criterion 1 (nonzero edges are nonzero

and directionally consistent in >50% of bootstraps). Although the difference between samples

could have resulted from network differences of clinical versus general population-based

samples, it may also reflect some degree of inconsistency or even noise as edge estimates often

exhibited unstable directions of association in clinical samples (see Table 2).

Applying consistency criterion 2 (consistency of results across samples), we observed replicable

edges of the CRP PRS with anhedonia (negative edge weight), changes in appetite, and fatigue

and of the TNF-α PRS with fatigue; these edges were manually unfaded in Figure 2. It is

important to note that the edge between the CRP PRS and changes in appetite has a diverging

valence in individual samples; in MARS and STAR*D (assessing loss of appetite) the edge

weight was negative and in the UK Biobank study (assessing changes in appetite) the edge

weight was positive.

383   -Figure 2-

384   -Figure 3-

Unregularized model search estimation suggests three consistent PRS-symptom edges
according to criterion 3.

Using unregularized model search estimation, we again observed networks with relatively larger

within-plane (i.e., PRS-PRS & symptom-symptom) than between-plane (i.e., PRS-symptom)

edges. Networks were comparable to FGL estimation, but generally sparser than those using FGL

estimation; see network graphs in Supplementary Figure 6 and bootstrapping results in

Supplementary Figures 7-9.

Regarding PRS-symptom edges, only three edges were estimated, which were all observed in the

UK Biobank sample and fulfilled consistency criterion 3 (nonzero edges are also nonzero and

directionally consistent in >50% of bootstraps); these edges have been manually unfaded in

Supplementary Figure 6. The specific PRS-symptom edges were between the BMI PRS and

changes in appetite and anhedonia and between the CRP PRS and changes in appetite.

Comparing these edges to FGL estimation, the edge of the CRP PRS with changes in appetite

replicated one of the edges fulfilling consistency criteria 1 and 2 while the two edges observed

for the BMI PRS were only fulfilling consistency criterion 1 (presence in FGL estimation and

>50% of bootstraps). Moreover, the BMI PRS association with anhedonia was negative using

unregularized model search estimation, but positive using FGL estimation.

402   -Table 2-

Node predictability

Average node predictability was similar across samples for PRS nodes with 16% (UK Biobank),

17% (MARS), and 19% (STAR*D) of variance explained by all other nodes in the network.

Contrary to this, average node predictability for symptom nodes differed with 38% of variance

explained by all other nodes in the UK Biobank sample and only 9% in both clinical samples.

These findings highlight differences in network density of symptoms in the UK Biobank (using

the PHQ-9) and clinical samples (using the HAM-D).

DISCUSSION

The present study investigated associations of PRSs for immuno-metabolic markers with depressive symptoms using a multi-plane, multi-sample network analysis approach. Based on three consistency criteria emphasising robustness and replicability of network analysis results across statistical bootstraps, samples, and estimation methods, we observed a unique association between the CRP PRS and changes in appetite that met all three consistency criteria. In addition to this association, we observed five additional PRS-symptom associations that met two consistency criteria. These included edges of the CRP PRS with anhedonia (negative association) and fatigue, the TNF-α PRS with fatigue, and the BMI PRS with anhedonia and changes in appetite. However, the BMI PRS-anhedonia association switched association direction depending on the estimation method, so may not be fully consistent despite fulfilling our consistency criteria. Due to the novelty of our analysis approach, we highlight several methodological considerations below, which we hope provides a helpful framework to the discussion of our findings afterwards.

Methodological challenges and opportunities

Combining PRSs with psychological symptom networks is a relatively recent extension of network analysis and, to our knowledge, has only been applied in one previous investigation incorporating a schizophrenia PRS into a psychotic symptom network (Isvoranu et al., 2020). Therefore, it is important to emphasise the unique challenges and opportunities of this approach.

First, as noted by Isvoranu *and colleagues* (2020), statistical power is potentially the greatest challenge of PRS-symptom network analysis. Network analysis itself requires relatively large sample sizes for psychological symptom networks (Epskamp et al., 2018; Fried and Cramer, 2017), which should be in the hundreds or thousands depending on the number of nodes in the

network. Inclusion of PRSs into psychological symptom networks, and especially of potential

pathomechanistic (e.g., inflammatory) rather than main illness (e.g., depression/schizophrenia)

scores into these networks, aggravates the sample size requirements for network analysis as PRSs

only explain a fraction of variance in the heritable component of their target phenotypes (Choi et

al., 2020; Wray et al., 2020).

Second, and because PRSs only measure a fraction of variance in their target phenotype, unique

associations observed in network analyses are inevitably smaller than actual target phenotype-

symptom associations. Taking this study as an example, absolute sizes of CRP PRS-symptom

associations were 5- to 10-fold smaller than those from a prior network investigation using serum

CRP concentrations by Moriarity *and colleagues* (2020a). Therefore, PRS-symptom associations

are unlikely to give meaningful insights into size of association with the target phenotype, but

should, in our opinion, be interpreted based on robust presence/absence of specific associations.

Third, the large statistical power requirements and difficulty quantifying such power for a given

study may lead to biased result interpretations. Absence of PRS-symptom associations could be

interpreted as false negatives while presence of association may be interpreted as true positives.

Such divergence in interpretation necessarily biases the literature towards hypothesis

confirmation. Consequently, any associations observed in PRS-symptom network analyses

should be followed up by- and interpreted in line with- evidence from other studies, thus adhering

to the recommended triangulation of evidence approach (Lawlor et al., 2017; Ohlsson and

Kendler, 2019).

Despite these challenges, PRS-symptom networks also provide multiple opportunities. First,

PRSs reflect estimates of genetic liability to phenotype expression, so can give an indication on

the influence of lifelong predisposition to higher phenotype levels on the symptom level. In this

way, PRS-symptom associations also provide an indication regarding temporality of association,

which Bradford-Hill defined as one of the viewpoints for causality (Bradford Hill, 1965). It is

important to note, however, that evidence for a unidirectional temporal association does not

preclude bi-directionality. Moreover, PRSs combine information from a multitude of genetic

variants (in our case from ~1 million SNPs) that are not restricted to functional SNPs, can include

false positive associations (i.e., noise), and can also tag information of pleiotropic environmental

confounding factors. Therefore, causal inferences should rely on separate evidence from clinical

trials and/or more focused genetic approaches such as Mendelian Randomisation studies (Lawlor

et al., 2008).

Second, the PRS-symptom network analysis approach allows the concurrent investigation of

multiple immuno-metabolic markers with multiple symptoms. Thereby, immunological and

clinical complexity is addressed concurrently, which is an advantage to previous investigations.

Furthermore, network analyses usually estimate partial/unique associations, so any emerging

associations could suggest direct causal paths from PRS phenotypes to individual symptoms, so

may pinpoint so-called 'bridge symptoms' that act as etiological docking sites of risk effects on

the symptom plane.

Third, large-scale population-based or patient cohort studies, commonly used in network

analysis, often do not have detailed immunophenotyping data available. If at all, studies mostly

have data available for serum CRP, but rarely for more specific cytokines. Conversely, the advent

of large GWAS investigations has produced a substantial amount of large cohort databases with

in-depth genotyping and phenotyping information. Combining such databases with GWAS

summary statistics from more focused investigations, such as on individual cytokines (Ahola-Olli

et al., 2017), enables the investigation of a diverse range of immunopsychiatric research

questions.

Fourth, PRS-symptom networks could be extended, for instance, by adding serum inflammatory

markers to these networks, which could provide additional insights into associations between

genetic/lifelong predisposition to, and acute levels of, immuno-metabolic markers with individual

symptoms.

Associations of immuno-metabolic markers with depressive symptoms

Network analysis results showed consistent associations of the CRP PRS with changes in

appetite, which was the only association that fulfilled all of our quality criteria. The BMI PRS

showed similar associations with changes in appetite, but only fulfilled two quality criteria.

Importantly, both of these associations were positive in the UK Biobank sample, which assessed

changes in appetite, and negative in MARS and STAR*D samples, which assessed loss of

appetite. Previous studies reporting results from cross-sectional, longitudinal, genetic correlation,

PRS, and Mendelian randomisation analyses have also consistently reported associations of

CRP/BMI with changes in appetite (Fried et al., 2019; Jokela et al., 2016; Kappelmann et al.,

2021; Moriarity et al., 2020a). Importantly, whenever studies disaggregated appetite symptoms

into decreased versus increased appetite, associations of CRP/BMI were specific to increased

appetite (Lamers et al., 2018; Milaneschi et al., 2021b, 2021a, 2017b; Pistis et al., 2021;

Simmons et al., 2018). In light of these findings, our results provide indirect support for an

immune-metabolic contribution to increased appetite specifically.

In addition to these PRS associations with changes in appetite, we also observed associations of

higher CRP PRS with lower anhedonia and greater fatigue and of higher TNF-α PRS with greater

fatigue. Fatigue in particular has long been considered to have a neuroimmune basis (Dantzer et

al., 2014), is common across other medical illnesses characterised by chronic inflammation, and

has been reliably associated with inflammatory markers in previous studies including two

network investigations (Fried et al., 2019; Jokela et al., 2016; Lamers et al., 2020; Moriarity et

al., 2020a; van Eeden et al., 2020; White et al., 2017). While there have also been some studies

suggesting associations of inflammatory markers with anhedonia (Köhler-Forsberg et al., 2017;

van Eeden et al., 2020), it is important to note that associations of the CRP PRS with anhedonia

observed in the present report were negative, so do not offer straightforward replication of these

findings. Nonetheless, we have recently shown in Mendelian randomisation analyses that BMI

could be a potential causal factor for both fatigue and anhedonia (Kappelmann et al., 2021), so

continued investigation of these symptoms is warranted.

Together, our findings add to the notion of an immuno-metabolic subtype of depression

characterised by neurovegetative symptoms of changes in appetite and fatigue (Dantzer et al.,

2008; Milaneschi et al., 2020). We also expand upon previous work by showing that

genetic/lifetime predisposition to higher inflammation and metabolic dysregulation increases risk

for depression and, based on network analysis results, these etiological factors may specifically

confer their risk on the broader depression syndrome through symptoms such as changes in

appetite and fatigue. These results can inform the design of clinical trials of anti-inflammatory

approaches and metabolic interventions by specifically selecting patients with an atypical,

neurovegetative symptom presentation. As clinical trials for immune-modulating drugs are

currently still characterised by relatively small sample sizes (Husain et al., 2020; Khandaker et

al., 2018; McIntyre et al., 2019; Nettis et al., 2021; Raison et al., 2013), it may be worthwhile to

pilot new interventions with neurovegetative symptoms/phenotypes as outcome variables. This

might increase statistical power and sensitivity to detect effects for these proof-of-concept trials

and could then be followed up by larger trials testing broader clinical efficacy measures.

Strengths and limitations

Strength of this study include availability of large general population-based and patient samples

(maximising replicability and generalisability), polygenic definition of immuno-metabolic risk

variables (indexing lifetime predisposition to higher immuno-metabolic marker levels), and

application of network analysis (addressing immunological and clinical complexity

concurrently). We have addressed some of the more general limitations of combined PRS-

symptom network analysis above, but there are three more specific limitations that warrant

mentioning.

First, data used in the current study included inpatients, outpatients, and individuals from the

general population and was based on different scales to measure depressive symptoms.

Depressive symptom structure varies between acutely ill patients versus those in remission (van

Borkulo et al., 2015), which may have influenced PRS-symptom associations. Moreover, two of

the seven symptoms used in the present report only overlap partially; the UK Biobank study

includes conflated items on sleep problems and changes in appetite while MARS and STAR*D

include items on insomnia and loss of appetite, respectively. This difference may explain some of

the inconsistencies observed in the current report such as the diverging valence of edge estimates

between CRP and changes in appetite. However, this may have also reduced statistical power to

detect associations. Study questionnaires also differed regarding the method of assessment as the

HAM-D is observer-rated and the PHQ-9 self-reported. By definition, inflammation-symptom

research is affected from modality-specific measurement variability (Moriarity and Alloy, 2021)

and in our study this is aggravated through the added variability unique to the method of

symptom assessment (Möller, 2000). Future studies would benefit from inclusion of studies with

the same questionnaire and disaggregated symptom measures.

Second, the combination of clinical and general population samples poses unique challenges. The

application of a clinical depression measure in the UK Biobank study could have resulted in

potential floor effects for some symptoms while specific selection of MDD patients into MARS

and STAR*D studies could have resulted in ceiling effects for core symptoms of depressed mood

and anhedonia as these are required for a diagnosis. Selection of clinical populations in network

studies can also result in Berkson/collider bias (de Ron et al., 2019), which can induce negative

correlations. This again warrants replication of our results in independent samples.

Third, PRSs are based on GWAS with highly diverging samples sizes as a large number of

individuals were included in the GWAS for BMI and CRP (>200 thousand individuals) and

smaller numbers of individuals (~8 thousand individuals) for IL-6, IL-10, and TNF-α.

Consistency of effect sizes following the PRS-CS approach was also larger for CRP and BMI as

compared to IL-6, IL-10, and TNF-α. This is likely to have shifted the balance of statistical

power towards detection of PRS-symptom associations to BMI and CRP rather than IL-6, IL-10,

and TNF-α. Therefore, our findings require replication once larger individual cytokine GWAS

become available.

Conclusion

The present investigation studied associations between four major pro- and anti-inflammatory

markers, BMI, and depressive symptoms by applying network analysis across one large general

population and two patient samples. Defining immuno-metabolic markers using polygenic risk

scores expanded previous reports by suggesting direct associations of genetic/lifetime

predisposition to immune-metabolic markers with depressive symptoms and provided evidence

for temporality of association. Despite methodological restrictions of the presented approach, we

observed associations of polygenic risk for CRP with changes in appetite and fatigue, for TNF-α

with fatigue, and similar associations for BMI. These findings align with recent

conceptualisations of an immuno-metabolic subgroup of depressed patients characterised by

atypical, neurovegetative symptom profiles. Results can inform future clinical trials of anti-

inflammatory approaches by prioritising these patients for selection into clinical trials.

DECLARATION OF INTERESTS

The authors do not have any competing interests.

FUNDING

This study is funded by the Max Planck Institute of Psychiatry. NK, NR, and SM are supported by the International Max Planck Research School of Translational Psychiatry (IMPRS-TP). NR received funding from the Bavarian Ministry of Economic Affairs, Regional Development and Energy (BayMED, PBN_MED-1711-0003). GMK acknowledges funding support from the Wellcome Trust (grant code: 201486/Z/16/Z), the MQ: Transforming Mental Health (grant code: MQDS17/40), the Medical Research Council, UK (grant code: MC_PC_17213 and grant code: MR/S037675/1), and the BMA Foundation (J Moulton grant 2019). JA received support by a NARSAD Young Investigator Grant from Brain and Behavior Research Foundation.

ACKNOWLEDGMENTS

We are grateful to all original authors, technical assistants and patients who contributed to the MARS study. We are grateful for the National Institute of Mental Health (NIMH) and the NIMH Repository and Genomics Resource (NRGR) for the possibility of analysing the STAR*D data. We are also grateful to the original STAR*D authors, and particularly for the contributions of all patients and families who participated in the study. Data were obtained from the limited access datasets distributed from the NIH-supported 'Sequenced Treatment Alternatives to Relieve Depression' (STAR*D). The study was supported by NIMH Contract No. N01MH90003 to the University of Texas Southwestern Medical Center. The ClinicalTrials.gov identifier is NCT00021528. This research has been conducted using the UK Biobank Resource. We are grateful for all scientists and participants who made this large-scale effort and resource possible.

REFERENCES

Ahola-Olli, A. V, Würtz, P., Havulinna, A.S., Aalto, K., Pitkänen, N., Lehtimäki, T., Kähönen, M., Lyytikäinen, L.-P., Raitoharju, E., Seppälä, I., Sarin, A.-P., Ripatti, S., Jalkanen, S., Maksimow, M., Salomaa, V., Salmi, M., Kettunen, J., Raitakari, O.T., 2017. Genome-wide Association Study Identifies 27 Loci Influencing Concentrations of Circulating Cytokines and Growth Factors. Am. J. Hum. Genet. 100, 40–50. https://doi.org/10.1016/j.ajhg.2016.11.007

Badini, I., Coleman, J.R., Hagenaars, S.P., Hotopf, M., Breen, G., Lewis, C.M., Fabbri, C., 2020. Depression with atypical neurovegetative symptoms shares genetic predisposition with immuno-metabolic traits and alcohol consumption. Psychol. Med. 1–11. https://doi.org/10.1017/S0033291720002342

Bilder, R.M., Volavka, J., Lachman, H.M., Grace, A.A., 2004. The Catechol-O-Methyltransferase Polymorphism: Relations to the Tonic–Phasic Dopamine Hypothesis and Neuropsychiatric Phenotypes. Neuropsychopharmacology 29, 1943–1961. https://doi.org/10.1038/sj.npp.1300542

Borsboom, D., 2017. A network theory of mental disorders. World Psychiatry 16, 5–13. https://doi.org/10.1002/wps.20375

Bradford Hill, A., 1965. The environment and disease: association or causation? Proc. R. Soc. Med. 58, 295.

Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A., Vukcevic, D., Delaneau, O., O'Connell, J., Cortes, A., Welsh, S., Young, A., Effingham, M., McVean, G., Leslie, S., Allen, N., Donnelly, P., Marchini, J., 2018. The UK Biobank resource with

deep phenotyping and genomic data. Nature 562, 203–209. https://doi.org/10.1038/s41586-018-0579-z

Cattaneo, A., Ferrari, C., Turner, L., Mariani, N., Enache, D., Hastings, C., Kose, M., Lombardo, G., McLaughlin, A.P., Nettis, M.A., Nikkheslat, N., Sforzini, L., Worrell, C., Zajkowska, Z., Cattane, N., Lopizzo, N., Mazzelli, M., Pointon, L., Cowen, P.J., Cavanagh, J., Harrison, N.A., de Boer, P., Jones, D., Drevets, W.C., Mondelli, V., Bullmore, E.T., Pariante, C.M., 2020. Whole-blood expression of inflammasome- and glucocorticoid-related mRNAs correctly separates treatment-resistant depressed patients from drug-free and responsive patients in the BIODEP study. Transl. Psychiatry 10, 232. https://doi.org/10.1038/s41398-020-00874-7

Choi, S.W., Mak, T.S.-H., O'Reilly, P.F., 2020. Tutorial: a guide to performing polygenic risk score analyses. Nat. Protoc. 15, 2759–2772. https://doi.org/10.1038/s41596-020-0353-1

Chu, A.L., Stochl, J., Lewis, G., Zammit, S., Jones, P.B., Khandaker, G.M., 2019. Longitudinal association between inflammatory markers and specific symptoms of depression in a prospective birth cohort. Brain. Behav. Immun. 76, 74–81. https://doi.org/10.1016/j.bbi.2018.11.007

Contreras, A., Nieto, I., Valiente, C., Espinosa, R., Vazquez, C., 2019. The Study of Psychopathology from the Network Analysis Perspective: A Systematic Review. Psychother. Psychosom. 88, 71–83. https://doi.org/10.1159/000497425

Costantini, G., Kappelmann, N., Epskamp, S., 2020. EstimateGroupNetwork: Perform the Joint Graphical Lasso and Selects Tuning Parameters.

Costantini, G., Richetin, J., Preti, E., Casini, E., Epskamp, S., Perugini, M., 2019. Stability and

variability of personality networks. A tutorial on recent developments in network

psychometrics. Pers. Individ. Dif. 136, 68–78. https://doi.org/10.1016/j.paid.2017.06.011

Danaher, P., Wang, P., Witten, D.M., 2014. The joint graphical lasso for inverse covariance

estimation across multiple classes. J. R. Stat. Soc. Ser. B (Statistical Methodol. 76, 373–397.

https://doi.org/10.1111/rssb.12033

Dantzer, R., Heijnen, C.J., Kavelaars, A., Laye, S., Capuron, L., 2014. The neuroimmune basis of

fatigue. Trends Neurosci. 37, 39–46. https://doi.org/10.1016/j.tins.2013.10.003

Dantzer, R., O'Connor, J.C., Freund, G.G., Johnson, R.W., Kelley, K.W., 2008. From

inflammation to sickness and depression: when the immune system subjugates the brain.

Nat. Rev. Neurosci. 9, 46–56. https://doi.org/10.1038/nrn2297

Davis, K.A.S., Coleman, J.R.I., Adams, M., Allen, N., Breen, G., Cullen, B., Dickens, C., Fox,

E., Graham, N., Holliday, J., Howard, L.M., John, A., Lee, W., McCabe, R., McIntosh, A.,

Pearsall, R., Smith, D.J., Sudlow, C., Ward, J., Zammit, S., Hotopf, M., 2020. Mental health

in UK Biobank – development, implementation and results from an online questionnaire

completed by 157 366 participants: a reanalysis. BJPsych Open 6, e18.

https://doi.org/10.1192/bjo.2019.100

de Ron, J., Fried, E.I., Epskamp, S., 2019. Psychological networks in clinical populations:

investigating the consequences of Berkson's bias. Psychol. Med. 1–9.

https://doi.org/10.1017/S0033291719003209

Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E.K., Lanctôt, K.L., 2010.

A Meta-Analysis of Cytokines in Major Depression. Biol. Psychiatry 67, 446–457.

https://doi.org/10.1016/j.biopsych.2009.09.033

Epskamp, S., Borsboom, D., Fried, E.I., 2018. Estimating psychological networks and their accuracy: A tutorial paper. Behav. Res. Methods 50, 195–212. https://doi.org/10.3758/s13428-017-0862-1

Epskamp, S., Cramer, A.O.J., Waldorp, L.J., Schmittmann, V.D., Borsboom, D., 2012. qgraph: Network Visualizations of Relationships in Psychometric Data. J. Stat. Softw. 48, 1–18. https://doi.org/10.18637/jss.v048.i04

Fried, E.I., Cramer, A.O.J., 2017. Moving forward: Challenges and direction for psychopathological network theory and methodology. Perspect. Psychol. Sci. 12, 999–1020. https://doi.org/10.1177/1745691617705892

Fried, E.I., von Stockert, S., Haslbeck, J.M.B., Lamers, F., Schoevers, R.A., Penninx, B.W.J.H., 2019. Using network analysis to examine links between individual depressive symptoms, inflammatory markers, and covariates. Psychol. Med. 1–9. https://doi.org/10.1017/S0033291719002770

Garriock, H.A., Kraft, J.B., Shyn, S.I., Peters, E.J., Yokoyama, J.S., Jenkins, G.D., Reinalda, M.S., Slager, S.L., McGrath, P.J., Hamilton, S.P., 2010. A Genomewide Association Study of Citalopram Response in Major Depressive Disorder. Biol. Psychiatry 67, 133–138. https://doi.org/10.1016/j.biopsych.2009.08.029

Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C.A., Smoller, J.W., 2019. Polygenic prediction via Bayesian regression and continuous shrinkage priors. Nat. Commun. 10, 1776. https://doi.org/10.1038/s41467-019-09718-5

Goldsmith, D.R., Rapaport, M.H., Miller, B.J., 2016. A meta-analysis of blood cytokine network alterations in psychiatric patients: comparisons between schizophrenia, bipolar disorder and

depression. Mol. Psychiatry 21, 1696–1709. https://doi.org/10.1038/mp.2016.3

Guloksuz, S., Pries, L.-K., van Os, J., 2017. Application of network methods for understanding
mental disorders: pitfalls and promise. Psychol. Med. 47, 2743–2752.
https://doi.org/10.1017/S0033291717001350

Haapakoski, R., Mathieu, J., Ebmeier, K.P., Alenius, H., Kivimäki, M., 2015. Cumulative meta-
analysis of interleukins 6 and 1β, tumour necrosis factor α and C-reactive protein in patients
with major depressive disorder. Brain. Behav. Immun. 49, 206–215.
https://doi.org/10.1016/j.bbi.2015.06.001

Hamilton, M., 1986. The Hamilton Rating Scale for Depression, in: Sartorius, N., Ban, T.A.
(Eds.), Assessment of Depression. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 143–
152. https://doi.org/10.1007/978-3-642-70486-4_14

Haslbeck, J.M.B., Fried, E.I., 2017. How predictable are symptoms in psychopathological
networks? A reanalysis of 18 published datasets. Psychol. Med. 47, 2767–2776.
https://doi.org/10.1017/S0033291717001258

Haslbeck, J.M.B., Waldorp, L.J., 2020. mgm☐: Estimating Time-Varying Mixed Graphical
Models in High-Dimensional Data. J. Stat. Softw. 93. https://doi.org/10.18637/jss.v093.i08

Hennings, J.M., Owashi, T., Binder, E.B., Horstmann, S., Menke, A., Kloiber, S., Dose, T.,
Wollweber, B., Spieler, D., Messer, T., Lutz, R., Kü Nzel, H., Bierner, T., Pollmächer, T.,
Pfister, H., Nickel, T., Sonntag, A., Uhr, M., Ising, M., Holsboer, F., Lucae, S., 2009.
Clinical characteristics and treatment outcome in a representative sample of depressed
inpatients – Findings from the Munich Antidepressant Response Signature (MARS) project.
J. Psychiatr. Res. 43, 215–229. https://doi.org/10.1016/j.jpsychires.2008.05.002

Howren, M.B., Lamkin, D.M., Suls, J., 2009. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. Psychosom. Med. 71, 171–186. https://doi.org/10.1097/PSY.0b013e3181907c1b

Husain, M.I., Chaudhry, I.B., Khoso, A.B., Husain, M.O., Hodsoll, J., Ansari, M.A., Naqvi, H.A., Minhas, F.A., Carvalho, A.F., Meyer, J.H., Deakin, B., Mulsant, B.H., Husain, N., Young, A.H., 2020. Minocycline and celecoxib as adjunctive treatments for bipolar depression: a multicentre, factorial design randomised controlled trial. The Lancet Psychiatry 7, 515–527. https://doi.org/10.1016/S2215-0366(20)30138-3

Isvoranu, A.-M., Guloksuz, S., Epskamp, S., van Os, J., Borsboom, D., 2020. Toward incorporating genetic risk scores into symptom networks of psychosis. Psychol. Med. 50, 636–643. https://doi.org/10.1017/S003329171900045X

Jokela, M., Virtanen, M., Batty, G., Kivimäki, M., 2016. Inflammation and specific symptoms of depression. JAMA Psychiatry 73, 87–88. https://doi.org/10.1001/jamapsychiatry.2015.1977

Kappelmann, N., Arloth, J., Georgakis, M.K., Czamara, D., Rost, N., Ligthart, S., Khandaker, G.M., Binder, E.B., 2021. Dissecting the Association Between Inflammation, Metabolic Dysregulation, and Specific Depressive Symptoms. JAMA Psychiatry 78, 161–170. https://doi.org/10.1001/jamapsychiatry.2020.3436

Kappelmann, N., Lewis, G., Dantzer, R., Jones, P.B., Khandaker, G.M., 2018. Antidepressant activity of anti-cytokine treatment: a systematic review and meta-analysis of clinical trials of chronic inflammatory conditions. Mol. Psychiatry 23, 335–343. https://doi.org/10.1038/mp.2016.167

Khandaker, G.M., Oltean, B.P., Kaser, M., Dibben, C.R.M., Ramana, R., Jadon, D.R., Dantzer,

R., Coles, A.J., Lewis, G., Jones, P.B., 2018. Protocol for the insight study: a randomised controlled trial of single-dose tocilizumab in patients with depression and low-grade inflammation. BMJ Open 8, e025333. https://doi.org/10.1136/bmjopen-2018-025333

Khandaker, G.M., Pearson, R.M., Zammit, S., Lewis, G., Jones, P.B., 2014. Association of serum interleukin 6 and C-reactive protein in childhood with depression and psychosis in young adult life: a population-based longitudinal study. JAMA Psychiatry 71, 1121–1128. https://doi.org/10.1001/jamapsychiatry.2014.1332

Köhler-Forsberg, O., Buttenschøn, H.N., Tansey, K.E., Maier, W., Hauser, J., Dernovsek, M.Z., Henigsberg, N., Souery, D., Farmer, A., Rietschel, M., McGuffin, P., Aitchison, K.J., Uher, R., Mors, O., 2017. Association between C-reactive protein (CRP) with depression symptom severity and specific depressive symptoms in major depression. Brain. Behav. Immun. 62, 344–350. https://doi.org/10.1016/j.bbi.2017.02.020

Köhler-Forsberg, O., Nicolaisen Lydholm, C., Hjorthøj, C., Nordentoft, M., Mors, O., Benros, M.E., 2019. Efficacy of anti-inflammatory treatment on major depressive disorder or depressive symptoms: Meta-analysis of clinical trials. Acta Psychiatr. Scand. 0–2. https://doi.org/10.1111/acps.13016

Köhler, C.A., Freitas, T.H., Maes, M., de Andrade, N.Q., Liu, C.S., Fernandes, B.S., Stubbs, B., Solmi, M., Veronese, N., Herrmann, N., Raison, C.L., Miller, B.J., Lanctôt, K.L., Carvalho, A.F., 2017. Peripheral cytokine and chemokine alterations in depression: a meta-analysis of 82 studies. Acta Psychiatr. Scand. 135, 373–387. https://doi.org/10.1111/acps.12698

Lamers, F., Milaneschi, Y., de Jonge, P., Giltay, E.J., Penninx, B.W.J.H., 2018. Metabolic and inflammatory markers: associations with individual depressive symptoms. Psychol. Med. 48,

1102–1110. https://doi.org/10.1017/S0033291717002483

Lamers, F., Milaneschi, Y., Smit, J.H., Schoevers, R.A., Wittenberg, G., Penninx, B.W.J.H., 2019. Longitudinal Association Between Depression and Inflammatory Markers: Results From the Netherlands Study of Depression and Anxiety. Biol. Psychiatry 85, 829–837. https://doi.org/10.1016/j.biopsych.2018.12.020

Lamers, F., Milaneschi, Y., Vinkers, C.H., Schoevers, R.A., Giltay, E.J., Penninx, B.W.J.H., 2020. Depression profilers and immuno-metabolic dysregulation: Longitudinal results from the NESDA study. Brain. Behav. Immun. 88, 174–183. https://doi.org/10.1016/j.bbi.2020.04.002

Lamnidis, T.C., Majander, K., Jeong, C., Salmela, E., Wessman, A., Moiseyev, V., Khartanovich, V., Balanovsky, O., Ongyerth, M., Weihmann, A., Sajantila, A., Kelso, J., Pääbo, S., Onkamo, P., Haak, W., Krause, J., Schiffels, S., 2018. Ancient Fennoscandian genomes reveal origin and spread of Siberian ancestry in Europe. Nat. Commun. 9, 5018. https://doi.org/10.1038/s41467-018-07483-5

Lawlor, D.A., Harbord, R.M., Sterne, J.A.C., Timpson, N., Davey Smith, G., 2008. Mendelian randomization: Using genes as instruments for making causal inferences in epidemiology. Stat. Med. 27, 1133–1163. https://doi.org/10.1002/sim.3034

Lawlor, D.A., Tilling, K., Davey Smith, G., 2017. Triangulation in aetiological epidemiology. Int. J. Epidemiol. 45, dyw314. https://doi.org/10.1093/ije/dyw314

Ligthart, S., Vaez, A., Võsa, U., Stathopoulou, M.G., de Vries, P.S., Prins, B.P., Van der Most, P.J., Tanaka, T., Naderi, E., Rose, L.M., Wu, Y., Karlsson, R., Barbalic, M., Lin, H., Pool, R., Zhu, G., Macé, A., Sidore, C., Trompet, S., Mangino, M., Sabater-Lleal, M., Kemp, J.P.,

Abbasi, A., Kacprowski, T., Verweij, N., Smith, A. V, Huang, T., Marzi, C., Feitosa, M.F., Lohman, K.K., Kleber, M.E., Milaneschi, Y., Mueller, C., Huq, M., Vlachopoulou, E., Lyytikäinen, L.-P., Oldmeadow, C., Deelen, J., Perola, M., Zhao, J.H., Feenstra, B., Amini, M., Lahti, J., Schraut, K.E., Fornage, M., Suktitipat, B., Chen, W.-M., Li, X., Nutile, T., Malerba, G., Luan, J., Bak, T., Schork, N., Del Greco M., F., Thiering, E., Mahajan, A., Marioni, R.E., Mihailov, E., Eriksson, Joel, Ozel, A.B., Zhang, W., Nethander, M., Cheng, Y.-C., Aslibekyan, S., Ang, W., Gandin, I., Yengo, L., Portas, L., Kooperberg, C., Hofer, E., Rajan, K.B., Schurmann, C., den Hollander, W., Ahluwalia, Tarunveer S., Zhao, J., Draisma, H.H.M., Ford, I., Timpson, N., Teumer, A., Huang, H., Wahl, S., Liu, Y., Huang, J., Uh, H.-W., Geller, F., Joshi, P.K., Yanek, L.R., Trabetti, E., Lehne, B., Vozzi, D., Verbanck, M., Biino, G., Saba, Y., Meulenbelt, I., O'Connell, J.R., Laakso, M., Giulianini, F., Magnusson, P.K.E., Ballantyne, C.M., Hottenga, J.J., Montgomery, G.W., Rivadineira, F., Rueedi, R., Steri, M., Herzig, K.-H., Stott, D.J., Menni, C., Frånberg, M., St. Pourcain, B., Felix, S.B., Pers, T.H., Bakker, S.J.L., Kraft, P., Peters, A., Vaidya, D., Delgado, G., Smit, J.H., Großmann, V., Sinisalo, J., Seppälä, I., Williams, S.R., Holliday, E.G., Moed, M., Langenberg, C., Räikkönen, K., Ding, J., Campbell, H., Sale, M.M., Chen, Y.-D.I., James, A.L., Ruggiero, D., Soranzo, N., Hartman, C.A., Smith, E.N., Berenson, G.S., Fuchsberger, C., Hernandez, D., Tiesler, C.M.T., Giedraitis, V., Liewald, D., Fischer, K., Mellström, D., Larsson, A., Wang, Y., Scott, W.R., Lorentzon, M., Beilby, J., Ryan, K.A., Pennell, C.E., Vuckovic, D., Balkau, B., Concas, M.P., Schmidt, R., Mendes de Leon, C.F., Bottinger, E.P., Kloppenburg, M., Paternoster, L., Boehnke, M., Musk, A.W., Willemsen, G., Evans, D.M., Madden, P.A.F., Kähönen, M., Kutalik, Z., Zoledziewska, M., Karhunen, V., Kritchevsky, S.B., Sattar, N., Lachance, G., Clarke, R., Harris, T.B., Raitakari, O.T., Attia, J.R., van Heemst, D., Kajantie, E., Sorice, R., Gambaro, G., Scott, R.A., Hicks, A.A.,

Ferrucci, L., Standl, M., Lindgren, C.M., Starr, J.M., Karlsson, M., Lind, L., Li, J.Z., Chambers, J.C., Mori, T.A., de Geus, E.J.C.N., Heath, A.C., Martin, N.G., Auvinen, J., Buckley, B.M., de Craen, A.J.M., Waldenberger, M., Strauch, K., Meitinger, T., Scott, R.J., McEvoy, M., Beekman, M., Bombieri, C., Ridker, P.M., Mohlke, K.L., Pedersen, N.L., Morrison, A.C., Boomsma, D.I., Whitfield, J.B., Strachan, D.P., Hofman, A., Vollenweider, P., Cucca, F., Jarvelin, M.-R., Jukema, J.W., Spector, T.D., Hamsten, A., Zeller, T., Uitterlinden, André G., Nauck, M., Gudnason, V., Qi, L., Grallert, H., Borecki, I.B., Rotter, J.I., März, W., Wild, P.S., Lokki, M.-L., Boyle, M., Salomaa, V., Melbye, M., Eriksson, J.G., Wilson, J.F., Penninx, B.W.J.H., Becker, D.M., Worrall, B.B., Gibson, G., Krauss, R.M., Ciullo, M., Zaza, G., Wareham, N.J., Oldehinkel, A.J., Palmer, L.J., Murray, S.S., Pramstaller, P.P., Bandinelli, S., Heinrich, J., Ingelsson, E., Deary, I.J., Mägi, R., Vandenput, L., van der Harst, P., Desch, K.C., Kooner, J.S., Ohlsson, C., Hayward, C., Lehtimäki, T., Shuldiner, A.R., Arnett, D.K., Beilin, L.J., Robino, A., Froguel, P., Pirastu, M., Jess, T., Koenig, W., Loos, R.J.F., Evans, D.A., Schmidt, H., Davey Smith, G., Slagboom, P.E., Eiriksdottir, G., Morris, A.P., Psaty, B.M., Tracy, R.P., Nolte, I.M., Boerwinkle, E., Visvikis-Siest, S., Reiner, A.P., Gross, M., Bis, J.C., Franke, L., Franco, O.H., Benjamin, E.J., Chasman, D.I., Dupuis, Josée, Snieder, H., Dehghan, A., Alizadeh, B.Z., Alizadeh, B.Z., Boezen, H.M., Franke, L., van der Harst, P., Navis, G., Rots, M., Snieder, H., Swertz, M., Wolffenbuttel, B.H.R., Wijmenga, C., Benjamin, E., Chasman, D.I., Dehghan, A., Ahluwalia, Tarunveer Singh, Meigs, J., Tracy, R., Alizadeh, B.Z., Ligthart, S., Bis, J., Eiriksdottir, G., Pankratz, N., Gross, M., Rainer, A., Snieder, H., Wilson, J.G., Psaty, B.M., Dupuis, Josee, Prins, B., Vaso, U., Stathopoulou, M., Franke, L., Lehtimaki, T., Koenig, W., Jamshidi, Y., Siest, S., Abbasi, A., Uitterlinden, Andre G., Abdollahi, M., Schnabel, R., Schick, U.M., Nolte, I.M., Kraja, A., Hsu, Y.-H., Tylee, D.S.,

Zwicker, A., Uher, R., Davey Smith, G., Morrison, A.C., Hicks, A., van Duijn, C.M., Ward-Caviness, C., Boerwinkle, E., Rotter, J., Rice, K., Lange, L., Perola, M., de Geus, E., Morris, A.P., Makela, K.M., Stacey, D., Eriksson, Johan, Frayling, T.M., Slagboom, E.P., 2018. Genome Analyses of >200,000 Individuals Identify 58 Loci for Chronic Inflammation and Highlight Pathways that Link Inflammation and Complex Disorders. Am. J. Hum. Genet. 103, 691–706. https://doi.org/10.1016/j.ajhg.2018.09.009

Liu, J.J., Wei, Y. Bin, Strawbridge, R., Bao, Y., Chang, S., Shi, L., Que, J., Gadad, B.S., Trivedi, M.H., Kelsoe, J.R., Lu, L., 2020. Peripheral cytokine levels and response to antidepressant treatment in depression: a systematic review and meta-analysis. Mol. Psychiatry 25, 339–350. https://doi.org/10.1038/s41380-019-0474-5

Locke, A.E., Kahali, B., Berndt, S.I., Justice, A.E., Pers, T.H., Day, F.R., Powell, C., Vedantam, S., Buchkovich, M.L., Yang, J., Croteau-Chonka, D.C., Esko, T., Fall, T., Ferreira, T., Gustafsson, S., Kutalik, Z., Luan, J., Mägi, R., Randall, J.C., Winkler, T.W., Wood, A.R., Workalemahu, T., Faul, J.D., Smith, J.A., Hua Zhao, J., Zhao, W., Chen, J., Fehrmann, R., Hedman, Å.K., Karjalainen, J., Schmidt, E.M., Absher, D., Amin, N., Anderson, D., Beekman, M., Bolton, J.L., Bragg-Gresham, J.L., Buyske, S., Demirkan, A., Deng, G., Ehret, G.B., Feenstra, B., Feitosa, M.F., Fischer, K., Goel, A., Gong, J., Jackson, A.U., Kanoni, S., Kleber, M.E., Kristiansson, K., Lim, U., Lotay, V., Mangino, M., Mateo Leach, I., Medina-Gomez, C., Medland, S.E., Nalls, M.A., Palmer, C.D., Pasko, D., Pechlivanis, S., Peters, M.J., Prokopenko, I., Shungin, D., Stančáková, A., Strawbridge, R.J., Ju Sung, Y., Tanaka, Toshiko, Teumer, A., Trompet, S., van der Laan, S.W., van Setten, J., Van Vliet-Ostaptchouk, J. V., Wang, Z., Yengo, L., Zhang, W., Isaacs, A., Albrecht, E., Ärnlöv, J., Arscott, G.M., Attwood, A.P., Bandinelli, S., Barrett, A., Bas, I.N., Bellis, C., Bennett, A.J.,

Berne, C., Blagieva, R., Blüher, M., Böhringer, S., Bonnycastle, L.L., Böttcher, Y., Boyd, H.A., Bruinenberg, M., Caspersen, I.H., Ida Chen, Y.-D., Clarke, R., Warwick Daw, E., de Craen, A.J.M., Delgado, G., Dimitriou, M., Doney, A.S.F., Eklund, N., Estrada, K., Eury, E., Folkersen, L., Fraser, R.M., Garcia, M.E., Geller, F., Giedraitis, V., Gigante, B., Go, A.S., Golay, A., Goodall, A.H., Gordon, S.D., Gorski, M., Grabe, H.-J., Grallert, H., Grammer, T.B., Gräßler, J., Grönberg, H., Groves, C.J., Gusto, G., Haessler, J., Hall, P., Haller, T., Hallmans, G., Hartman, C.A., Hassinen, M., Hayward, C., Heard-Costa, N.L., Helmer, Q., Hengstenberg, C., Holmen, O., Hottenga, J.-J., James, A.L., Jeff, J.M., Johansson, Å., Jolley, J., Juliusdottir, T., Kinnunen, L., Koenig, W., Koskenvuo, M., Kratzer, W., Laitinen, J., Lamina, C., Leander, K., Lee, N.R., Lichtner, P., Lind, L., Lindström, J., Sin Lo, K., Lobbens, S., Lorbeer, R., Lu, Y., Mach, F., Magnusson, P.K.E., Mahajan, A., McArdle, W.L., McLachlan, S., Menni, C., Merger, S., Mihailov, E., Milani, L., Moayyeri, A., Monda, K.L., Morken, M.A., Mulas, A., Müller, G., Müller-Nurasyid, M., Musk, A.W., Nagaraja, R., Nöthen, M.M., Nolte, I.M., Pilz, S., Rayner, N.W., Renstrom, F., Rettig, R., Ried, J.S., Ripke, S., Robertson, N.R., Rose, L.M., Sanna, S., Scharnagl, H., Scholtens, S., Schumacher, F.R., Scott, W.R., Seufferlein, T., Shi, J., Vernon Smith, A., Smolonska, J., Stanton, A. V., Steinthorsdottir, V., Stirrups, K., Stringham, H.M., Sundström, J., Swertz, M.A., Swift, A.J., Syvänen, A.-C., Tan, S.-T., Tayo, B.O., Thorand, B., Thorleifsson, G., Tyrer, J.P., Uh, H.-W., Vandenput, L., Verhulst, F.C., Vermeulen, S.H., Verweij, N., Vonk, J.M., Waite, L.L., Warren, H.R., Waterworth, D., Weedon, M.N., Wilkens, L.R., Willenborg, C., Wilsgaard, T., Wojczynski, M.K., Wong, A., Wright, A.F., Zhang, Q., Brennan, E.P., Choi, M., Dastani, Z., Drong, A.W., Eriksson, P., Franco-Cereceda, A., Gådin, J.R., Gharavi, A.G., Goddard, M.E., Handsaker, R.E., Huang, J., Karpe, F., Kathiresan, S., Keildson, S., Kiryluk, K., Kubo, M., Lee, J.-Y., Liang, L., Lifton,

R.P., Ma, B., McCarroll, S.A., McKnight, A.J., Min, J.L., Moffatt, M.F., Montgomery, G.W., Murabito, J.M., Nicholson, G., Nyholt, D.R., Okada, Y., Perry, J.R.B., Dorajoo, R., Reinmaa, E., Salem, R.M., Sandholm, N., Scott, R.A., Stolk, L., Takahashi, A., Tanaka, Toshihiro, van't Hooft, F.M., Vinkhuyzen, A.A.E., Westra, H.-J., Zheng, W., Zondervan, K.T., Heath, A.C., Arveiler, D., Bakker, S.J.L., Beilby, J., Bergman, R.N., Blangero, J., Bovet, P., Campbell, H., Caulfield, M.J., Cesana, G., Chakravarti, A., Chasman, D.I., Chines, P.S., Collins, F.S., Crawford, D.C., Adrienne Cupples, L., Cusi, D., Danesh, J., de Faire, U., den Ruijter, H.M., Dominiczak, A.F., Erbel, R., Erdmann, J., Eriksson, J.G., Farrall, M., Felix, S.B., Ferrannini, E., Ferrières, J., Ford, I., Forouhi, N.G., Forrester, T., Franco, O.H., Gansevoort, R.T., Gejman, P. V., Gieger, C., Gottesman, O., Gudnason, V., Gyllensten, U., Hall, A.S., Harris, T.B., Hattersley, A.T., Hicks, A.A., Hindorff, L.A., Hingorani, A.D., Hofman, A., Homuth, G., Kees Hovingh, G., Humphries, S.E., Hunt, S.C., Hyppönen, E., Illig, T., Jacobs, K.B., Jarvelin, M.-R., Jöckel, K.-H., Johansen, B., Jousilahti, P., Wouter Jukema, J., Jula, A.M., Kaprio, J., Kastelein, J.J.P., Keinanen-Kiukaanniemi, S.M., Kiemeney, L.A., Knekt, P., Kooner, J.S., Kooperberg, C., Kovacs, P., Kraja, A.T., Kumari, M., Kuusisto, J., Lakka, T.A., Langenberg, C., Le Marchand, L., Lehtimäki, T., Lyssenko, V., Männistö, S., Marette, A., Matise, T.C., McKenzie, C.A., McKnight, B., Moll, F.L., Morris, A.D., Morris, A.P., Murray, J.C., Nelis, M., Ohlsson, C., Oldehinkel, A.J., Ong, K.K., Madden, P.A.F., Pasterkamp, G., Peden, J.F., Peters, A., Postma, D.S., Pramstaller, P.P., Price, J.F., Qi, L., Raitakari, O.T., Rankinen, T., Rao, D.C., Rice, T.K., Ridker, P.M., Rioux, J.D., Ritchie, M.D., Rudan, I., Salomaa, V., Samani, N.J., Saramies, J., Sarzynski, M.A., Schunkert, H., Schwarz, P.E.H., Sever, P., Shuldiner, A.R., Sinisalo, J., Stolk, R.P., Strauch, K., Tönjes, A., Trégouët, D.-A., Tremblay, A., Tremoli, E., Virtamo, J., Vohl, M.-C., Völker, U., Waeber, G., Willemsen, G., Witteman, J.C., Carola Zillikens, M.,

Adair, L.S., Amouyel, P., Asselbergs, F.W., Assimes, T.L., Bochud, M., Boehm, B.O., Boerwinkle, E., Bornstein, S.R., Bottinger, E.P., Bouchard, C., Cauchi, S., Chambers, J.C., Chanock, S.J., Cooper, R.S., de Bakker, P.I.W., Dedoussis, G., Ferrucci, L., Franks, P.W., Froguel, P., Groop, L.C., Haiman, C.A., Hamsten, A., Hui, J., Hunter, D.J., Hveem, K., Kaplan, R.C., Kivimaki, M., Kuh, D., Laakso, M., Liu, Y., Martin, N.G., März, W., Melbye, M., Metspalu, A., Moebus, S., Munroe, P.B., Njølstad, I, Oostra, B.A., Palmer, C.N.A., Pedersen, N.L., Perola, M., Pérusse, L., Peters, U., Power, C., Quertermous, T., Rauramaa, R., Rivadeneira, F., Saaristo, T.E., Saleheen, D., Sattar, N., Schadt, E.E., Schlessinger, D., Eline Slagboom, P., Snieder, H., Spector, T.D., Thorsteinsdottir, U., Stumvoll, M., Tuomilehto, J., Uitterlinden, A.G., Uusitupa, M., van der Harst, P., Walker, M., Wallaschofski, H., Wareham, N.J., Watkins, H., Weir, D.R., Wichmann, H.-E., Wilson, J.F., Zanen, P., Borecki, I.B., Deloukas, P., Fox, C.S., Heid, I.M., O'Connell, J.R., Strachan, D.P., Stefansson, K., van Duijn, C.M., Abecasis, G.R., Franke, L., Frayling, T.M., McCarthy, M.I., Visscher, P.M., Scherag, A., Willer, C.J., Boehnke, M., Mohlke, K.L., Lindgren, C.M., Beckmann, J.S., Barroso, I., North, K.E., Ingelsson, E., Hirschhorn, J.N., Loos, R.J.F., Speliotes, E.K., 2015. Genetic studies of body mass index yield new insights for obesity biology. Nature 518, 197–206. https://doi.org/10.1038/nature14177

Lopresti, A.L., 2017. Cognitive behaviour therapy and inflammation: A systematic review of its relationship and the potential implications for the treatment of depression. Aust. New Zeal. J. Psychiatry 51, 565–582. https://doi.org/10.1177/0004867417701996

Löwe, B., Unützer, J., Callahan, C.M., Perkins, A.J., Kroenke, K., 2004. Monitoring Depression Treatment Outcomes With the Patient Health Questionnaire-9. Med. Care 42, 1194–1201. https://doi.org/10.1097/00005650-200412000-00006

Lynall, M.-E., Turner, L., Bhatti, J., Cavanagh, J., de Boer, P., Mondelli, V., Jones, D., Drevets, W.C., Cowen, P., Harrison, N.A., Pariante, C.M., Pointon, L., Clatworthy, M.R., Bullmore, E., 2020. Peripheral Blood Cell–Stratified Subgroups of Inflamed Depression. Biol. Psychiatry 88, 185–196. https://doi.org/10.1016/j.biopsych.2019.11.017

Mac Giollabhui, N., Ng, T.H., Ellman, L.M., Alloy, L.B., 2020. The longitudinal associations of inflammatory biomarkers and depression revisited: systematic review, meta-analysis, and meta-regression. Mol. Psychiatry. https://doi.org/10.1038/s41380-020-00867-4

McIntyre, R.S., Subramaniapillai, M., Lee, Y., Pan, Z., Carmona, N.E., Shekotikhina, M., Rosenblat, J.D., Brietzke, E., Soczynska, J.K., Cosgrove, V.E., Miller, S., Fischer, E.G., Kramer, N.E., Dunlap, K., Suppes, T., Mansur, R.B., 2019. Efficacy of Adjunctive Infliximab vs Placebo in the Treatment of Adults With Bipolar I/II Depression. JAMA Psychiatry 1–8. https://doi.org/10.1001/jamapsychiatry.2019.0779

Milaneschi, Y., Kappelmann, N., Ye, Z., Lamers, F., Moser, S., Jones, P.B., Burgess, S., Penninx, B.W.J.H., Khandaker, G.M., 2021a. Association of Inflammation with Depression and Anxiety: Evidence for Symptom-Specificity and Potential Causality from UK Biobank and NESDA Cohorts. medRxiv. https://doi.org/10.1101/2021.01.08.20248710

Milaneschi, Y., Lamers, F., Berk, M., Penninx, B.W.J.H., 2020. Depression Heterogeneity and Its Biological Underpinnings: Toward Immunometabolic Depression. Biol. Psychiatry 88, 369–380. https://doi.org/10.1016/j.biopsych.2020.01.014

Milaneschi, Y., Lamers, F., Bot, M., Drent, M.L., Penninx, B.W.J.H., 2017a. Leptin Dysregulation Is Specifically Associated With Major Depression With Atypical Features: Evidence for a Mechanism Connecting Obesity and Depression. Biol. Psychiatry 81, 807–

814. https://doi.org/10.1016/j.biopsych.2015.10.023

Milaneschi, Y., Lamers, F., Penninx, B.W.J.H., 2021b. Dissecting Depression Biological and
Clinical Heterogeneity—The Importance of Symptom Assessment Resolution. JAMA
Psychiatry 362, 2021. https://doi.org/10.1001/jamapsychiatry.2020.4373

Milaneschi, Y., Lamers, F., Peyrot, W., Abdellaoui, A., Willemsen, G., Hottenga, J.-J., Jansen,
R., Mbarek, H., Dehghan, A., Lu, C., 2016. Polygenic dissection of major depression
clinical heterogeneity. Mol. Psychiatry 21, 516–522. https://doi.org/10.1038/mp.2015.86

Milaneschi, Y., Lamers, F., Peyrot, W.J., Baune, B.T., Breen, G., Dehghan, A., Forstner, A.J.,
Grabe, H.J., Homuth, G., Kan, C., Lewis, C., Mullins, N., Nauck, M., Pistis, G., Preisig, M.,
Rivera, M., Rietschel, M., Streit, F., Strohmaier, J., Teumer, A., Van der Auwera, S., Wray,
N.R., Boomsma, D.I., Penninx, B.W.J.H., 2017b. Genetic Association of Major Depression
With Atypical Features and Obesity-Related Immunometabolic Dysregulations. JAMA
Psychiatry 74, 1214. https://doi.org/10.1001/jamapsychiatry.2017.3016

Möller, H.., 2000. Rating depressed patients: observer- vs self-assessment. Eur. Psychiatry 15,
160–172. https://doi.org/10.1016/S0924-9338(00)00229-7

Moriarity, D.P., Alloy, L.B., 2021. Back to Basics: The Importance of Measurement Properties in
Biological Psychiatry. Neurosci. Biobehav. Rev. 123, 72–82.
https://doi.org/10.1016/j.neubiorev.2021.01.008

Moriarity, D.P., Horn, S.R., Kautz, M.M., Haslbeck, J.M.B., Alloy, L.B., 2020a. How handling
extreme C-reactive protein (CRP) values and regularization influences CRP and depression
criteria associations in network analyses. Brain. Behav. Immun.
https://doi.org/10.1016/j.bbi.2020.10.020

Moriarity, D.P., van Borkulo, C., Alloy, L.B., 2020b. Inflammatory phenotype of depression symptom structure: A network perspective. Brain. Behav. Immun. 87, S48. https://doi.org/10.1016/j.bbi.2020.12.005

Nettis, M.A., Lombardo, G., Hastings, C., Zajkowska, Z., Mariani, N., Nikkheslat, N., Worrell, C., Enache, D., McLaughlin, A., Kose, M., Sforzini, L., Bogdanova, A., Cleare, A., Young, A.H., Pariante, C.M., Mondelli, V., 2021. Augmentation therapy with minocycline in treatment-resistant depression patients with low-grade peripheral inflammation: results from a double-blind randomised clinical trial. Neuropsychopharmacology 1–10. https://doi.org/10.1038/s41386-020-00948-6

Ni, G., Zeng, J., Revez, J.A., Wang, Y., Ge, T., Restaudi, R., Kiewa, J., Nyholt, D.R., Coleman, J.R.I., Smoller, J.W., Consortium, S.W.G. of the P.G., Yang, J., Visscher, P.M., Wray, N.R., 2020. A comprehensive evaluation of polygenic score methods across cohorts in psychiatric disorders. medRxiv 1–31.

Ohlsson, H., Kendler, K.S., 2019. Applying Causal Inference Methods in Psychiatric Epidemiology. JAMA Psychiatry 1–8. https://doi.org/10.1001/jamapsychiatry.2019.3758

Osimo, E.F., Baxter, L.J., Lewis, G., Jones, P.B., Khandaker, G.M., 2019. Prevalence of low-grade inflammation in depression: a systematic review and meta-analysis of CRP levels. Psychol. Med. 49, 1958–1970. https://doi.org/10.1017/S0033291719001454

Pistis, G., Milaneschi, Y., Vandeleur, C.L., Lasserre, A.M., Penninx, B.W.J.H., Lamers, F., Boomsma, D.I., Hottenga, J., Marques-Vidal, P., Vollenweider, P., Waeber, G., Aubry, J., Preisig, M., Kutalik, Z., 2021. Obesity and atypical depression symptoms: findings from Mendelian randomization in two European cohorts. Transl. Psychiatry 11, 96.

https://doi.org/10.1038/s41398-021-01236-7

R Core Team, 2017. R: A language and environment for statistical computing.

Raison, C.L., Rutherford, R.E., Woolwine, B.J., Shuo, C., Schettler, P., Drake, D.F., Haroon, E.,
Miller, A.H., 2013. A randomized controlled trial of the tumor necrosis factor antagonist
infliximab for treatment-resistant depression: the role of baseline inflammatory biomarkers.
JAMA Psychiatry 70, 31–41. https://doi.org/10.1001/2013.jamapsychiatry.4

Robinaugh, D.J., Hoekstra, R.H.A., Toner, E.R., Borsboom, D., 2020. The network approach to
psychopathology: a review of the literature 2008–2018 and an agenda for future research.
Psychol. Med. 50, 353–366. https://doi.org/10.1017/S0033291719003404

Rush, A.J., Fava, M., Wisniewski, S.R., Lavori, P.W., Trivedi, M.H., Sackeim, H.A., Thase,
M.E., Nierenberg, A.A., Quitkin, F.M., Kashner, T.M., Kupfer, D.J., Rosenbaum, J.F.,
Alpert, J., Stewart, J.W., McGrath, P.J., Biggs, M.M., Shores-Wilson, K., Lebowitz, B.D.,
Ritz, L., Niederehe, G., for the STAR*D Investigators Group, 2004. Sequenced treatment
alternatives to relieve depression (STAR*D): rationale and design. Control. Clin. Trials 25,
119–142. https://doi.org/10.1016/S0197-2456(03)00112-0

Rush, A.J., Trivedi, M.H., Wisniewski, S.R., Nierenberg, A.A., Stewart, J.W., Warden, D.,
Niederehe, G., Thase, M.E., Lavori, P.W., Lebowitz, B.D., McGrath, P.J., Rosenbaum, J.F.,
Sackeim, H.A., Kupfer, D.J., Luther, J., Fava, M., 2006. Acute and Longer-Term Outcomes
in Depressed Outpatients Requiring One or Several Treatment Steps: A STAR*D Report.
Am. J. Psychiatry 163, 1905–1917. https://doi.org/10.1176/appi.ajp.163.11.1905

Simmons, W.K., Burrows, K., Avery, J.A., Kerr, K.L., Taylor, A., Bodurka, J., Potter, W.,
Teague, T.K., Drevets, W.C., 2018. Appetite changes reveal depression subgroups with

distinct endocrine, metabolic, and immune states. Mol. Psychiatry 1–12.

https://doi.org/10.1038/s41380-018-0093-6

Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P.,

Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A.,

Sprosen, T., Peakman, T., Collins, R., 2015. UK Biobank: An Open Access Resource for

Identifying the Causes of a Wide Range of Complex Diseases of Middle and Old Age.

PLOS Med. 12, e1001779. https://doi.org/10.1371/journal.pmed.1001779

Timpson, N.J., Nordestgaard, B.G., Harbord, R.M., Zacho, J., Frayling, T.M., Tybjærg-Hansen,

A., Davey Smith, G., 2011. C-reactive protein levels and body mass index: elucidating

direction of causation through reciprocal Mendelian randomization. Int. J. Obes. 35, 300–

308. https://doi.org/10.1038/ijo.2010.137

van Borkulo, C., Boschloo, L., Borsboom, D., Penninx, B.W.J.H., Waldorp, L.J., Schoevers,

R.A., 2015. Association of Symptom Network Structure With the Course of Depression.

JAMA Psychiatry 72, 1219–1226. https://doi.org/10.1001/jamapsychiatry.2015.2079

van Eeden, W.A., van Hemert, A.M., Carlier, I.V.E., Penninx, B.W.J.H., Lamers, F., Fried, E.I.,

Schoevers, R., Giltay, E.J., 2020. Basal and LPS-stimulated inflammatory markers and the

course of individual symptoms of depression. Transl. Psychiatry 10, 235.

https://doi.org/10.1038/s41398-020-00920-4

White, J., Kivimäki, M., Jokela, M., Batty, G.D., 2017. Association of inflammation with specific

symptoms of depression in a general population of older people: The English Longitudinal

Study of Ageing. Brain. Behav. Immun. 61, 27–30.

https://doi.org/10.1016/j.bbi.2016.08.012

Williams, D.R., Rhemtulla, M., Wysocki, A.C., Rast, P., 2019. On Nonregularized Estimation of
Psychological Networks. Multivariate Behav. Res. 0, 1–23.
https://doi.org/10.1080/00273171.2019.1575716

Wittenberg, G.M., Stylianou, A., Zhang, Y., Sun, Y., Gupta, A., Jagannatha, P.S., Wang, D., Hsu,
B., Curran, M.E., Khan, S., Chen, G., Bullmore, E.T., Drevets, W.C., 2020. Effects of
immunomodulatory drugs on depressive symptoms: A mega-analysis of randomized,
placebo-controlled clinical trials in inflammatory disorders. Mol. Psychiatry 25, 1275–1285.
https://doi.org/10.1038/s41380-019-0471-8

Wray, N.R., Lin, T., Austin, J., McGrath, J.J., Hickie, I.B., Murray, G.K., Visscher, P.M., 2020.
From Basic Science to Clinical Application of Polygenic Risk Scores: A Primer. JAMA
Psychiatry. https://doi.org/10.1001/jamapsychiatry.2020.3049

TABLES & FIGURES

Tables

Table 1. Baseline characteristics of MARS, STAR*D, and UK Biobank samples

	MARS	**STAR*D**	**UK Biobank**
N	1,058	1,143	110,010
Sex			
Women, N (%)	563 (53.2%)	676 (59.1%)	61,212 (55.6%)
Men, N (%)	495 (46.8%)	467 (40.9%)	48,798 (44.4%)
Age in years			
Mean (SD)	47.8 (14.4)	43.2 (13.6)	56.2 (7.7)
Range	18-87	18-75	39-72
Study location	Germany	United States	United Kingdom
Study population	MDD inpatients	MDD outpatients	General population
CRP PRS			
Mean (SD)	-0.01 (1.00)	0.00 (1.00)	-0.01 (1.00)
Range	-3.14-3.15	-3.21-2.94	-4.58-4.21
IL-6 PRS			
Mean (SD)	0.00 (1.00)	0.00 (1.00)	0.00 (1.00)
Range	-3.33-3.34	-3.41-3.94	-4.07-3.92
IL-10 PRS			
Mean (SD)	0.00 (1.00)	0.00 (1.00)	0.00 (1.00)
Range	-3.11-3.24	-3.28-3.13	-4.35-4.60
TNF-a PRS			
Mean (SD)	0.00 (1.00)	0.00 (1.00)	0.00 (1.00)
Range	-3.61-3.67	-3.35-3.52	-4.17-4.51
BMI PRS			
Mean (SD)	-0.02 (1.01)	0.00 (1.00)	-0.04 (1.00)
Range	-3.02-4.06	-3.04-3.20	-4.53-3.96
PHQ-9 sum-score			
Mean (SD)	-	-	2.7 (3.6)
Range	-	-	0-27
Missing, N (%)	-	-	351 (0.3)
HAM-D sum-score			
Mean (SD)	23.8 (5.9)	22.4 (4.9)	-
Range	5-42	13-38	-
Missing, N (%)	6 (0.6%)	0 (0%)	-
Depressed mood			
Mean (SD)	3.05 (0.88)	2.59 (0.77)	0.23 (0.56)
Range	0-4	0-4	0-3
Anhedonia			
Mean (SD)	3.63 (0.68)	2.54 (0.76)	0.26 (0.56)
Range	0-4	0-4	0-3
Sleep problems			
Mean (SD)	3.46 (2.00)	3.42 (1.77)	0.71 (0.99)
Range	0-6	0-6	0-3

	MARS	STAR*D	UK Biobank
Fatigue			
Mean (SD)	1.49 (0.68)	1.68 (0.55)	0.66 (0.81)
Range	0-2	0-2	0-3
Changes in appetite			
Mean (SD)	0.77 (0.66)	0.68 (0.80)	0.25 (0.62)
Range	0-2	0-2	0-3
Psychomotor changes			
Mean (SD)	1.44 (0.96)	1.42 (0.72)	0.07 (0.34)
Range	0-4	0-4	0-3
Suicidality			
Mean (SD)	1.35 (1.15)	0.94 (0.85)	0.05 (0.28)
Range	0-4	0-4	0-3

Note: MDD=Major Depressive Disorder, SD=Standard deviation, PHQ-9=Patient Health Questionnaire-9, HAM-D=Hamilton Rating Scale for Depression.

Table 2. PRS-symptom edge consistency criteria (C) across network analyses

PRS-symptom edges	MARS		STAR*D		UK Biobank		FGL consistency (C2)
	FGL (C1)	Model search (C3)	FGL (C1)	Model search (C3)	FGL (C1)	Model search (C3)	
CRP							
Anhedonia	-0.016 (67%)		-0.043 (95%)		-0.002 (60%)		Yes
Depressed mood	-0.009 (57%)		0.045 (94%)				
Sleep problems*	-0.02 (75%)		0.031 (84%)				
Fatigue	0.053 (98%)		0.025 (79%)		0.011 (100%)		Yes
Changes in appetite*	-0.034 (89%)		-0.043 (94%)		0.003 (91%)	0.013 (73%)	Yes
Psychomotor changes	0.039 (91%)		0.001 (53%)				
Suicidality	-0.02 (74%)		-0.038 (88%)				
IL-6							
Anhedonia	-0.047 (97%)						
Depressed mood	-0.001 (52%)						
Sleep problems*			-0.02 (82%)				
Fatigue	0.012 (72%)						
Changes in appetite*	-0.032 (87%)		0.032 (85%)				
Psychomotor changes							
Suicidality	0.053 (95%)		-0.029 (83%)				
IL-10							
Anhedonia	0.008 (67%)		-0.005 (66%)				
Depressed mood			-0.002 (52%)				
Sleep problems*			0.014 (72%)				
Fatigue			0.007 (66%)				
Changes in appetite*	0.021 (80%)						
Psychomotor changes	0.015 (66%)		0.033 (90%)				
Suicidality			0.054 (99%)				
TNF-α							
Anhedonia	0.054 (96%)		0.013 (66%)				
Depressed mood	-0.017 (66%)		-0.017 (75%)				
Sleep problems*	-0.005 (57%)		0.005 (68%)				
Fatigue	0.016 (65%)		0.032 (91%)		0.002 (58%)		Yes
Changes in appetite*	-0.015 (70%)		0.023 (75%)				

PRS-symptom edges	MARS FGL (C1)	MARS Model search (C3)	STAR*D FGL (C1)	STAR*D Model search (C3)	UK Biobank FGL (C1)	UK Biobank Model search (C3)	FGL consistency (C2)
Psychomotor changes	0.008 (63%)		0.008 (59%)				
Suicidality			0.047 (94%)				
BMI							
Anhedonia	0.036 (91%)		0.033 (86%)			-0.010 (63%)	
Depressed mood							
Sleep problems*	0.06 (99%)		0.055 (97%)				
Fatigue	-0.016 (74%)						
Changes in appetite*			-0.031 (87%)		0.054 (100%)	0.066 (100%)	
Psychomotor changes	0.003 (55%)		-0.023 (83%)				
Suicidality	0.021 (75%)		0.04 (91%)				

Note: Cell values reflect edge weights (i.e., partial correlation coefficients) and the percentage of 500 bootstrap estimations that edges were present. Estimates are restricted to those edges, for which >50% of bootstrapped samples were non-zero and directionally consistent (i.e., criteria 1 & 3). *Changes in appetite and sleep problems are measured as composite symptoms in UK Biobank, but as loss of appetite and insomnia in MARS and STAR*D samples.

Figure Titles and Legends

Figure 1. Study design and analysis pipeline

Legend: BIC=Bayesian information criterion; CV=cross-validation; PC=principal component (or multi-dimensional scaling component used for MARS & STAR*D); □=PRS-CS tuning parameter.

Figure 2. Estimated FGL networks across samples

Legend: Networks are visualised with the *qgraph* package. Blue lines indicate positive and red lines negative associations, respectively, with larger associations displayed with thicker lines. Circles around nodes display node predictability, which can be interpreted similar to explained variance. Maximum size of edge associations is 0.55. As the primary focus of this investigation was to identify consistent PRS-symptom associations, we manually unfaded edges between PRSs and symptoms if these edges met quality criteria 1 and 2 (see Table 2). Changes in appetite and sleep problems are measured as composite symptoms in UK Biobank, but as loss of appetite and insomnia in MARS and STAR*D samples.

Figure 3. Bootstrapped 95% quantile intervals of PRS-symptom edges using FGL estimation

Legend: Bootstrapped 95% quantile intervals (i.e., 95% of the distribution of raw bootstrapped edge estimates) are highlighted as shaded area for each edge. Black points indicate the raw FGL sample estimate while red points indicate the raw bootstrapped mean estimate. Edges are indicated on the y-axis and sorted by mean edge weight across samples in descending order.

Sample	**MARS** n=1,058	**STAR*D** n=1,143	**UK Biobank** n=110,010

Data Preparation

Polygenic Risk Score Computation (PRS-CS)

1. CRP
2. IL-6
3. IL-10
4. TNF-α
5. BMI

$\phi = 1e^{-4}$

$\phi = 1e^{-2}$

Residual-correction:
PRS ~ age + sex + PC1 + PC2 + array*
*correction for array only in MARS

Symptom Data Harmonisation

1. Depressed mood
2. Anhedonia
3. Sleep problems*
4. Fatigue
5. Changes in appetite*
6. Psychomotor changes*
7. Suicidality

* only partially overlapping

Network Analysis

Primary analysis

Fused Graphical LASSO (FGL)

- *EstimateGroupNetwork* package
- 10-fold CV (BIC optimisation)
- equal sample weighting
- 500 bootstraps

Node Predictability

- *mgm* package

Secondary analysis

Unregularised Model Search

- *qgraph* & *bootnet* packages
- *ggModSelect* algorithm
- Spearman correlations
- 500 bootstraps

- 10-fold CV (BIC optimisation)
- 'OR'-rule

Consistency Evaluation

Criterion 1

- PRS-symptom edge present in individual sample
- Edge present and directionally consistent in >50% bootstraps

Criterion 2

- Criterion 1 replicates across samples

Criterion 3

- PRS-symptom edge present in individual sample
- Edge present and directionally consistent in >50% bootstraps

MARS

STAR*D

UK Biobank

A: Depressive Symptoms
- 1: Depressed mood
- 2: Anhedonia
- 3: Sleep problems
- 4: Fatigue
- 5: Changes in appetite
- 6: Psychomotor changes
- 7: Suicidality

B: Polygenic Risk Scores
- 8: CRP
- 9: IL-6
- 10: IL-10
- 11: TNF-a
- 12: BMI