1   Title: CENTER-BASED EXPERIENCES IMPLEMENTING STRATEGIES TO REDUCE RISK OF

2   HORIZONTAL TRANSMISSION OF SARS-COV-2: POTENTIAL FOR COMPROMISE OF NEONATAL

3   MICROBIOME ASSEMBLAGE

4
Running Title: COVID-19 infection policies and the neonatal microbiome

Joann Romano-Keeler, MD, MS[1,*], Dana Fiszbein, DO[1], Jilei Zhang, PhD[2], Joseph Horowitz,

MD[3], Karen Hayani, MD[4], Irina Buhimschi, MD[5], Christina Lopez, MS, RNC[6], Zaynab Kadhem[1],

James Berman, MD[7], Phornphat Rasamimari, MD[1], Aarti Raghavan, MD, FAAP, MS[1], De-Ann

M. Pillers, MD, PhD[1], Jun Sun, PhD[2,8*]

[1] Section of Neonatology, Department of Pediatrics, University of Illinois at Chicago, Chicago,

IL; [2] Division of Gastroenterology and Hepatology, Department of Medicine, University of Illinois

at Chicago, Chicago, IL; [3] Department of Pediatrics, University of Illinois at Chicago, Chicago, IL;

[4] Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Illinois at

Chicago; [5] Department of Obstetrics and Gynecology, University of Illinois at Chicago, Chicago,

IL; [6] Department of Neonatal Nursing, University of Illinois Hospital and Health Services;

Chicago, IL [7] Division of Pediatric Gastroenterology, Department of Pediatrics, University of

Illinois at Chicago, Chicago, IL [8] University of Illinois at Chicago Cancer Center, University of

Illinois at Chicago, Chicago, IL

* **Corresponding Authors**

Joann Romano-Keeler, MD, MS

University of Illinois at Chicago

840 S. Wood Street (MC 856)

Chicago, IL 60612

jromanok@uic.edu

Jun Sun, PhD, AGAF, FAPS, Professor

Division of Gastroenterology and Hepatology, Department of Medicine,

University of Illinois at Chicago,

840 S Wood Street,

Room 704 CSB, MC716,

Chicago, IL, 60612, USA

Tel: 312-996-5020,

E-mail: junsun7@uic.edu

Abstract

Perinatal transmission of COVID-19 is poorly understood and many neonatal intensive care

units' (NICU) policies minimize mother-infant contact to prevent transmission. We present our

unit's approach and ways it may impact neonatal microbiome acquisition. We attended COVID-

19 positive mothers' deliveries from March-August 2020. Delayed cord clamping and skin-to-

skin were avoided and infants were admitted to the NICU. No parents' visits were allowed and

discharge was arranged with COVID-19 negative family members. Maternal breast milk was

restricted in the NICU. All twenty-one infants tested negative at 24 and 48 hours and had

average hospital stays of nine days. 40% of mothers expressed breastmilk and 30% of infants

were discharged with COVID-19 negative caregivers. Extended hospital stays, no skin-to-skin

contact, limited maternal milk use, and discharge to caregivers outside primary residences,

potentially affect the neonatal microbiome. Future studies are warranted to explore how ours

and other centers' similar policies influence this outcome.

.

Introduction

Since its presentation in Wuhan, China, coronavirus disease of 2019 (COVID-19) caused by

severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has evolved into a global

pandemic with significant mortality and morbidity. The number of confirmed cases has climbed

to over thirty million with one million deaths in more than 200 countries and territories.(1, 2)

While initially associated with fever, cough and shortness of breath, presentation of this virus

has progressed to include gastrointestinal (GI) symptoms either in isolation or in conjunction

with respiratory symptoms in 10-15% of COVID-19 positive patients.(3, 4) GI symptoms may

herald COVID-19 by up to 5 days before admission and these patients may have more

complicated courses with longer hospital stays.(5, 6)

Neonatal intensive care units (NICU) and newborn nurseries have developed strategies to

minimize risk of infection to the relatively immunocompromised newborn population born to

COVID-19 positive mothers. American Academy of Pediatrics (AAP) recommendations

addressed this issue based on best available evidence at that time, experiences from other

institutions, and analysis of the risk/benefit to the mother and child.(7) Our guidelines were

based on limited evidence available at the early stages of the pandemic about neonatal COVID-

19 infections and our concerns about transmission in an open bay unit with limited space to

socially distance.(8, 9) Our feeding polices reflected our not having a dedicated milk preparation

room or a barcoded milk storage system. No existing recommendations address potential

impact that perinatal management may have on early life microbial colonization and subsequent

newborn and childhood health outcomes.(10, 11)

Intrinsic and extrinsic factors drive early life assembly of the intestinal microbiome, including

maternal genetics and nutrition during pregnancy.(12, 13) Postnatal variables including infants'

diet, antibiotic use, and environmental exposures can shape bacterial colonization until "adult-

like" gut microbiome is established.(14, 15) Virome-microbiome crosstalk during this period is

poorly understood.(16) No studies have investigated interactions between COVID-19 and the

nascent microbiome or how policies to prevent perinatal COVID transmission may affect

newborn microbial colonization **(Figure 1)**.

Twenty-one patients born to COVID-19 positive mothers were admitted to the NICU per

University of Illinois at Chicago (UIC) guidelines. In this paper, we review NICU courses,

including the incidence viral transmission. We explore ways that our guidelines, designed to

reduce risk of viral transmission to infants, may impact composition of the newborn microbiome.

Finally, we discuss future directions for NICU management that minimize perinatal viral

transmission, promote normal early life microbiome assemblage, and optimize short and long-

term health outcomes.

Materials/Subjects and Methods

This study was approved by the UIC Institutional Review Board (protocol no. 2020-0443).

Neonatal teams attended deliveries of all COVID-19 positive mothers from March- August 2020.

Deidentified maternal data collected included age, reported race, dates and types of COVID-19

testing, COVID-related symptoms at delivery, and details on peripartum courses including

intrapartum antibiotics and timing of rupture of membranes (**Table 1**).

Delayed cord clamping and skin-to-skin contact were avoided, including physical contact with

family members and infants were admitted to the NICU with contact isolation. Metadata

obtained on infants included delivery indication and mode, Apgar scores, gestational age, birth

weight, and gender. Respiratory and/or gastrointestinal symptoms were recorded throughout

hospitalization and POCT was completed at 24 and 48 hours of life. Visitors were not allowed

during NICU stays. Discharges were arranged whenever possible with COVID-19 negative

caregivers. No expressed breastmilk was used in the NICU but mothers were encouraged to

pump and store milk and instructed on how to safely breastfeed at home. Information on feeding

regimen after discharge was recorded when available from outpatient clinic notes.

Results

Subjects

Twenty-one COVID-19 positive mothers positive delivered during this study with a mean (range)

age of 26 (17-42) years. Approximately 62% of mothers were African American (n=13) and 5%

were Latinx (n=5). COVID-19 antigen detection via rapid point of care testing (POCT) within 72

hours of delivery or as an inpatient in labor and delivery was positive in 90% of mothers (n=19).

The remaining two mothers had positive viral PCR tests collected prior to delivery.

Clinical Characteristics of COVID-19 mothers

Sixteen COVID-19 positive mothers (76%) were asymptomatic and five (24%) were

symptomatic, including one with gastrointestinal symptoms. Group B Streptococcus (GBS)

screening was positive in 48% of mothers (n=10) and prolonged rupture of membranes (>18

hours prior to delivery) complicated one delivery. Intrapartum antibiotics were administered in

57% (n=11) of mothers, including five who were GBS positive and seven who received cefazolin

perioperatively. In 38% of mothers (n=8) Caesarean sections (C-section) were performed for

fetal heart rate decelerations or arrest of dilation (n=5), breech presentation (n=1), repeat C-

sections (n=1), or COVID-19 related maternal morbidities (n=1).

Infant demographics and Hospital Course

Mean [range] gestational age and birth weight was 36 [31-40] weeks and 2800 [1300-3500]

grams, respectively. Prematurity complicated two deliveries, including one case where an infant

was delivered early due to the severity of mother's COVID-19 infection. In 40% of cases (n=8),

delayed cord clamping occurred and in 10% (n=2) of deliveries, mothers were allowed skin-to-

skin contact. Approximately 70% of infants were male with a mean one minute Apgar score of 7

and a 5 minute score of 9. POCT was negative at 24 and 48 hours for all infants. In addition,

two infants had a third POCT due to persistent oxygen requirements, which were also negative.

More than half of patients (n=11) required respiratory support including mechanical ventilation

(n=1), continuous positive airway pressure (n=8), and nasal cannula (n=2). At the time of

discharge, one patient remained on oxygen.

Breast milk usage

Lactation counselors met with mothers to instruct on safe techniques for nursing and pumping,

including pumping during NICU stays and either discarding or storing milk. All infants received

formula in the NICU, except for one premature infant who qualified for donor breastmilk. In 30%

of newborns, mothers provided expressed milk at discharge or at the time of the first outpatient

appointment. No infants received breast milk exclusively.

Length of Stay and Discharge Coordination

Average length of stay was nine (5-52) days, however, when excluding two premature infants,

length of stay decreased to seven days. Discharge to COVID-19 negative family members

outside the primary residence occurred in 60% of cases. In 40% of cases, discharge was

arranged with fathers who were negative, mothers who were >10 days from their positive test

results, or COVID-19 positive parents who could not identify uninfected family members.

Discussion

We describe infection control measurees that resulted in no vertical or horizontal transmission of

SARS-CoV-2. Other centers reporting no perinatal transmission adopted similar infection control

measures.(17) While measures may decrease transmission, they may also interrupt exchange

of maternal flora with newborns. We also report secondary outcomes associated with our

policies, including increased length of hospital stay, decreased breastmilk use, and discharge to

households outside principal residences.

While C-section rates were higher in our cohort when compared with institutional (29% in 2019)

and national rates (31.9% in 2018), our sample size limits interpretation of this result.(18)

Maternal complications from COVID-19 may contribute to C-sections. However, in our study,

this was the case in only one patient. Other studies of COVID-19 and pregnancy outcomes

report an increases in C-sections due to fetal distress even in asymptomatic mothers, which

may impact the neonatal microbiome.(17, 19)

Vaginal deliveries represent one of the earliest exposures of newborns to microbes.(20, 21)

These pioneer bacteria initiate intestinal eubiosis and impart optimal health outcomes, including

decreased incidence of otitis media, atopic disease, and obesity.(22, 23) Without exposure to

the vaginal microbiome, the microbiota of infants born via C-sections shifts towards colonization

with flora from maternal skin, hospital staff, and hospital environment.(24) Further research is

needed to understand factors contributing to an increased incidence of C-sections, particularly

in asymptomatic COVID-19 positive women.(25)

Newborns in our cohort were isolated post-partum and then exposed to households outside

their parents. New environmental exposures could alter intestinal bacterial colonization early in

life. We were not able to evaluate the gut microbiome in our cohort but could be incorporated

into future research. Over the first year, newborn microbiota is influenced by mothers'

microbiome and evolves to have a similar composition.(15) Interruptions in this microbial

exchange may have unintended consequences later in life, including inflammatory bowel

disease, atopic disorders, and obesity.

It is unclear whether separation of infants from mothers had unintended consequences on

maternal milk use. Consistent with AAP recommendations, we provided lactation support to

encourage expression and storage of milk.(26) However, given ambiguity about COVID-19

transmission through breastmilk and our unit's milk cataloging system, we restricted maternal

breastmilk use in the NICU, including breastfeeding. Data on breastmilk transmission is

emerging and a recent study reports no detection of SARS-COV-2 in breast milk from nine

infected mothers.(27) However, in one pre-print report, SARS-COV-2 was detected in

breastmilk from a symptomatic mother whose infant also subsequently tested positive.(28)

Benefits of breastmilk are well understood and implications of not using breastmilk are

significant, particularly as it relates to the neonatal microbiome.

In conclusion, while adopting approaches to minimize viral transmission, we and other

institutions may have affected various aspects of newborn health, including assembly of the

neonatal microbiome. With new information on COVID-19 perinatal transmission, the AAP has

revised recommendations, including the safety of delayed cord clamping, mothers and infants

rooming together, and breastfeeding.(29) As of November, we revised our policies to allow

COVID-19 positive mothers and their offspring to room together without NICU admission. We

continue to test infants after delivery and track postnatal outcomes. Meanwhile, a better

understanding of the gut-lung axis dysbiosis in COVID-19 is necessary to elucidate potential

microbiome-virome cross-talk in utero and postnatally, impact of COVID-19 infection control

policies on the neonatal gut microbiome, and ways to safeguard pediatric health outcomes

during the pandemic.(30, 31)

Author contributions

JRK, JS, and DMP conceptualized and drafted the article. JRK, DF, JH, and ZK acquired and

analyzed data. JRK and JZ developed figures. All authors revised the article and gave final

approval for its content.

199
200 **Acknowledgements**

201
202 We would like to acknowledge NIDDK/National Institutes of Health grants NIDDKR01

203 DK105118 and R01DK114126 (JS).

205 **Compliance with ethical standards**

206 **Conflict of interest**

207 No authors have actual or potential conflicts of interest in relationship to this publication.

References

1. Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, et al. A new genomic blueprint of the human gut microbiota. Nature. 2019;568(7753):499-504.

2. Sender R, Fuchs S, Milo R. Revised Estimates for the Number of Human and Bacteria Cells in the Body. PLoS Biol. 2016;14(8):e1002533.

3. Mao R, Qiu Y, He JS, Tan JY, Li XH, Liang J, et al. Manifestations and prognosis of gastrointestinal and liver involvement in patients with COVID-19: a systematic review and meta-analysis. Lancet Gastroenterol Hepatol. 2020.

4. Zhang JG, S.; Sun, J. Gastrointestinal symptoms, pathophysiology, and treatment in COVID-19. Genes & Diseases. 2020(https://doi.org/10.1016/j.gendis.2020.08.013).

5. Jin X, Lian JS, Hu JH, Gao J, Zheng L, Zhang YM, et al. Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. Gut. 2020;69(6):1002-9.

6. Buscarini E, Manfredi G, Brambilla G, Menozzi F, Londoni C, Alicante S, et al. GI symptoms as early signs of COVID-19 in hospitalised Italian patients. Gut. 2020.

7. Pediatrics AAo. FAQs: Management of Infants Born to Mothers with Suspected or Confirmed COVID-19. AAP Publications. April 2020.

8. Procianoy RS, Silveira RC, Manzoni P, Sant'Anna G. Neonatal COVID-19: little evidence and the need for more information. J Pediatr (Rio J). 2020;96(3):269-72.

9. Piersigilli F, Carkeek K, Hocq C, van Grambezen B, Hubinont C, Chatzis O, et al. COVID-19 in a 26-week preterm neonate. Lancet Child Adolesc Health. 2020;4(6):476-8.

10. Tamburini S, Shen N, Wu HC, Clemente JC. The microbiome in early life: implications for health outcomes. Nat Med. 2016;22(7):713-22.

11. Arrieta MC, Stiemsma LT, Dimitriu PA, Thorson L, Russell S, Yurist-Doutsch S, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. Sci Transl Med. 2015;7(307):307ra152.

12. Romano-Keeler J, Moore DJ, Wang C, Brucker RM, Fonnesbeck C, Slaughter JC, et al. Early life establishment of site-specific microbial communities in the gut. Gut Microbes. 2014;5(2):192-201.

13. Romano-Keeler J, Weitkamp JH. Maternal influences on fetal microbial colonization and immune development. Pediatr Res. 2015;77(1-2):189-95.

14. Romano-Keeler J, Zhang, J, Sun, J. The Life-long Role of Nutrition on the Gut Microbiome and Gastrointestinal Disease. Gastroenterology Clinics of North America. 2020.

15. Backhed F, Roswall J, Peng Y, Feng Q, Jia H, Kovatcheva-Datchary P, et al. Dynamics and Stabilization of the Human Gut Microbiome during the First Year of Life. Cell Host Microbe. 2015;17(6):852.

16. Duranti S, Lugli GA, Mancabelli L, Armanini F, Turroni F, James K, et al. Maternal inheritance of bifidobacterial communities and bifidophages in infants through vertical transmission. Microbiome. 2017;5(1):66.

17. Zaigham M, Andersson O. Maternal and perinatal outcomes with COVID-19: A systematic review of 108 pregnancies. Acta Obstet Gynecol Scand. 2020;99(7):823-9.

18. Martin JA, Hamilton BE, Osterman MJK, Driscoll AK. Births: Final Data for 2018. Natl Vital Stat Rep. 2019;68(13):1-47.

19.   Di Mascio D, Khalil A, Saccone G, Rizzo G, Buca D, Liberati M, et al. Outcome of
coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic
review and meta-analysis. Am J Obstet Gynecol MFM. 2020;2(2):100107.
20.   Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, Fierer N, et al.
Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body
habitats in newborns. Proc Natl Acad Sci U S A. 2010;107(26):11971-5.
21.   Chu DM, Ma J, Prince AL, Antony KM, Seferovic MD, Aagaard KM. Maturation of the
infant microbiome community structure and function across multiple body sites and in relation to
mode of delivery. Nat Med. 2017;23(3):314-26.
22.   Ajslev TA, Andersen CS, Gamborg M, Sorensen TI, Jess T. Childhood overweight after
establishment of the gut microbiota: the role of delivery mode, pre-pregnancy weight and early
administration of antibiotics. Int J Obes (Lond). 2011;35(4):522-9.
23.   Metsala J, Lundqvist A, Virta LJ, Kaila M, Gissler M, Virtanen SM. Prenatal and post-
natal exposure to antibiotics and risk of asthma in childhood. Clin Exp Allergy. 2015;45(1):137-
45.
24.   Bokulich NA, Chung J, Battaglia T, Henderson N, Jay M, Li H, et al. Antibiotics, birth
mode, and diet shape microbiome maturation during early life. Sci Transl Med.
2016;8(343):343ra82.
25.   Gensollen T, Iyer SS, Kasper DL, Blumberg RS. How colonization by microbiota in
early life shapes the immune system. Science. 2016;352(6285):539-44.
26.   Pediatrics AAo. Breastfeeding Guidance Post Hospital Discharge for Mothers or Infants
with Suspected or Confirmed SARS-Co V-2 Infection. AAP Publications. 2020.
27.   Lackey KA, Pace RM, Williams JE, Bode L, Donovan SM, Jarvinen KM, et al. SARS-
CoV-2 and human milk: what is the evidence? medRxiv. 2020.
28.   Kimberlin DW SS. Can SARS-CoV-2 infection be acquired in utero?: more definitive
evidence is needed. JAMA. 2020;Doi: 10.1001/jama.2020.4868.
29.   Pediatrics AAo. FAQs: Management of Infants Born to Mothers with Suspected or
Confirmed COVID-19. July 22, 2020.
30.   Aktas B, Aslim B. Gut-lung axis and dysbiosis in COVID-19. Turk J Biol.
2020;44(3):265-72.
31.   Conte L, Toraldo DM. Targeting the gut-lung microbiota axis by means of a high-fibre
diet and probiotics may have anti-inflammatory effects in COVID-19 infection. Ther Adv Respir
Dis. 2020;14:1753466620937170.
288

	Normal	**COVID-19 Positive**
Before hospitalization	Maternal GI eubiosis Transfer of materal microbes to fetus	Maternal GI dysbiosis Placental transfer of coronavirus, maternal antibodies Decidual arteriopathy and maternal vascular malperfusion of placenta
Delivery	**Delayed cord clamping** Transmission of blood, antibodies and microbes **Vaginal delivery** Colonization of newborn with maternal vaginal flora	No delayed cord clamping No skin-to-skin contact during delivery Increased risk of Caesarean section delivery, no maternal transfer of vaginal microbes to newborn
Post parturition	Kangaroo care/skin flora transfer Breastmilk microbes/metabolites Microbes from parents and their home environment	Mother-newborn separation, restricted visitors Limited or absent breastmilk exposure Discharge to homes different than parents

Characteristic	N=21 (%)	
Age (years)	26 (17-42)	
Ethnicity	Black-African American	13 (62)
	Latinx	5 (24)
	Caucasian	1 (5)
	Other	2 (9)
Maternal symptoms	Asymptomatic	16 (76)
	Respiratory symptoms	4 (19)
	Gastrointestinal symptoms	1 (5)
GBS status	Positive	10 (48)
	Negative/Unknown	11 (52)
Intrapartum antibiotics	10 (47)	
Mode of delivery	Cesarean section	8 (38)
	Vaginal delivery	13 (62)
Gestational age (weeks)	36 (31- 40)	
Birth weight (grams)	2800 (1300- 3500)	
Length of stay (days)	9 (5-52)	
Type of feeds	Breastmilk + formula	6 (2)
	Formula	15 (71)
Discharge plan	Principal residence	13 (62)
	Secondary residence	8 (3)

Figure 1: Potential interruptions in early life microbial colonization due to COVID-19 and infection control policies. Placental exchange of the maternal microbiome with the fetus during pregnancy contributes to early bacterial colonization. In contrast, placentas of women with COVID-19 have increased rates of decidual arteriopathy and maternal vascular malperfusion which may affect normal bacterial exchange.(33) At delivery, mothers with critical illness or fetal distress secondary to COVID-19 may be more likely to deliver via Cesarean section depriving the newborn of normal bacterial colonization. Institutional policies to avoid delayed cord clamping in deliveries of COVID-19 mothers may also impact microbial transfer to the newborn. Another infection control measure restricts skin-to-skin care between COVID-19 positive mothers and their infants which interferes with newborn's inheritance of maternal oral and skin flora. In addition, we recommended discharge of the newborn to COVID negative family members if possible until COVID-19 parents complete 10 days of separation, which often resulted in discharge home with family members outside the traditional household.

Tables

Table 1. Clinical Characteristics of COVID positive mother-infant dyads