The importance of non-pharmaceutical interventions during the COVID-19 vaccine rollout

Nicolò Gozzi[1]   Paolo Bajardi[2,*]   Nicola Perra[1]

[1] Networks and Urban Systems Centre, University of Greenwich, London, UK
[2] ISI Foundation, Turin, Italy
* paolo.bajardi@isi.it

Abstract

The promise of efficacious vaccines against SARS-CoV-2 is fulfilled and vaccination campaigns have started worldwide. However, the fight against the pandemic is far from over. Here, we propose an age-structured compartmental model to study the interplay of disease transmission, vaccines rollout, and behavioural dynamics. We investigate, via in-silico simulations, individual and societal behavioural changes, possibly induced by the start of the vaccination campaigns, and manifested as a relaxation in the adoption of non-pharmaceutical interventions. We explore different vaccine efficacy, vaccination rollout speeds, prioritization strategies, as well as multiple behavioural responses. We apply our model to six countries worldwide (Egypt, Peru, Serbia, Ukraine, Canada, and Italy) selected to sample diverse socio-demographic and socio-economic contexts. To isolate the effects of age-structures and contacts patterns from the particular pandemic history of each location, we first study the model considering the same hypothetical initial epidemic scenario in all countries. We then calibrate the model using real epidemiological and mobility data for the different countries. Our findings suggest that early relaxation of safe behaviours can jeopardize the benefits brought by the vaccine in the short term: a fast vaccine distribution and policies aimed at keeping high compliance of individual safe behaviours are key to mitigate disease resurgence.

Introduction

The COVID-19 pandemic has been largely fought with non-pharmaceutical interventions (NPIs). Bans of events and social gatherings, limitations in national and international travels, school closures, shifts towards remote working, curfews, closure of pubs and restaurants, cordon sanitaires, national and regional lockdowns are examples of governmental interventions implemented around the world to curb the spreading of SARS-CoV-2 [1–5]. While extremely effective, such top-down NPIs induce profound behavioural changes, bring many social activities to a halt, and thus have huge socio-economic costs. Hence, alongside these measures, governments nudged and/or mandated populations to adopt another set of NPIs. Social distancing, face masks, and increased hygiene are examples [6]. Although far from being cost-free, they are more feasible, sustainable, and allow for higher levels of socio-economic activity. As such, they have been the leitmotif of the post first COVID-19 wave in many countries. Unfortunately, awareness, adoption, and compliance with these NPIs have been spotty [6]. Furthermore, they have not been complemented with sufficiently aggressive test and trace programs. As result, many countries experienced marked disease resurgences after the summer 2020 and some had to resort to new lockdowns [7]. As we write, we have turned a crucial corner in hampering the resurgence, diffusion and severe outcomes of the disease. Several vaccines have shown great results of their phase 3 trials, and have been authorized for emergency use by several regulating agencies and tens of others are in the pipeline [8–10]. As result, we have witnessed the start of vaccination campaigns around the world. However, the logistical issues linked to the production, delivery, and administration of billions of doses on a global scale impose caution when evaluating the impact vaccines will have on the pandemic in the short and medium term. They will be a scarce resource and it will take time to vaccinate the fraction of the population necessary for herd immunity [11, 12]. Furthermore, vaccines are not perfect and there are many other unknowns [13]. The results show high levels, around 95% [14, 15], of direct protection against the disease. Despite the first signs of the vaccine impact in the real world are extremely encouraging [16, 17], these figures might end up to be lower, it is still unclear to what extent vaccines limit further transmission, and how long the immunity lasts [13]. Finally, vaccines' acceptance is a complex challenge. A recent survey among $13,426$

participants in 19 countries shows that, while 71.5% of the sample is very or somewhat likely to take the vaccine, there are large heterogeneities [18]. Acceptance rates vary from 90% in China to less than 55% in Russia. Furthermore, they are linked to socio-economic, socio-demographic features, and education attainment [18]. Arguably, vaccines alone will not be able to contain the spreading of the virus, at least in the short term [19]. Social distancing, face masks, hygiene measures, and other NPIs will be still key during the delivery of such vaccination programs, especially considering the emergence and fast spread of variants of concern [20].

In this context, an important question emerges. What will happen to adoption and compliance to NPIs as vaccination campaigns progress? Their arrival and delivery might induce individual and collective behavioural changes. Some might see this milestone as the official end of the emergency and as result relax their COVID-safe behaviours. Somehow paradoxically, vaccines might have, at least initially, a net negative effect. According to the health-belief model, one of the most commonly used psychological theories to characterize health-related behaviours, beliefs, perceptions, barriers to take action, and other modifying variables such as socio-demographic and socio-economic factors are key ingredients driving behavioural changes [21–23]. A recent study may offer an empirical evidence of this tendency [24]. By means of surveys delivered before the vaccination rollout in December 2020, authors found that vaccine information reduce adherence to social distancing, hygiene measures, and the willingness to stay at home. Several surveys conducted during the COVID-19 pandemic, well before any concrete hope for a vaccine, confirm this picture, provide hints of how the arrival of vaccines might corrode even more adoption, highlight how compliance is a complex multi-faced problem [5] and that risk-perception as well as NPIs adoption are indeed associated to several socio-economic determinants such as age, gender, wealth, urban-rural divide [25–40].

The literature aimed at estimating the epidemiological and societal impact of COVID-19 vaccines has been focused mainly on two very important points. The first line of research has been devoted to quantifying the effects of a vaccine on the evolution of the pandemic, considering different efficacy and coverage levels [41]. The second instead tackled the issue of vaccine allocation investigating strategies that target first different groups (i.e., age brackets, high-risk individuals) or particular occupations (i.e., doctors, nurses) [10, 42–44]. Very recently, the intuition that social distancing remains key during vaccination rollout stimulated few studies on the effects of a vaccine on the adoption of NPIs in specific settings [45–48], but the impact on real-world scenarios is still largely unexplored.

To tackle such limitation, we introduce an age-structured compartmental epidemic model capturing the possible relaxation of NPIs adoption in response to the vaccine rollout. We model different compliance levels as distinct compartments and consider different behavioural dynamics driving the relaxation of NPIs. We test extensively the effects of behaviour change on disease spreading for different prioritization strategies, vaccine efficacy, and vaccination rollout speeds, using real demographic data and contacts matrices for six countries: Egypt, Peru, Serbia, Ukraine, Canada, and Italy. We choose these countries sampling levels of economic development. Indeed, in the World Economic Situation and Prospects 2020 issued by the United Nations, Egypt and Peru are classified as *Developing Economies* [49], Serbia and Ukraine as *Economies in Transition*, Canada and Italy as *Developed Economies*. Furthermore, considering dissimilar countries allows us to explore the interplay between vaccination and behaviours also as a function of population pyramids and intra/inter-generational mixing observed around the world. High income countries are typically characterized by higher average age, but lower inter-generational interactions respects to mid/low income countries [50]. These observations, together with the dependence of COVID-19 fatality rates on age, point to the possibility of non-trivial dependencies which we aim to explore here. As a way to realistically account for the different epidemic trajectories, we also explore the model after calibrating it on COVID-19 weekly deaths in the period $2020/09/01 - 2020/12/31$ in these countries. To this end, we incorporate the timeline and effects of governmental restrictions on social contacts.

Our results provide quantitative insights on the interaction between sustained NPIs and an effective vaccination program. We show that an early relaxation of COVID-safe behaviours may lower, and even nullify, the advantages brought by the vaccine in the short term. Overall, the picture that emerges from the analysis of the different countries is consistent: a high level of compliance towards NPIs such as mask-wearing, social distancing, and avoidance of large gatherings, will be needed in the next months in order to avoid spoiling the great effort of the vaccination campaigns.

Results

Vaccine-Behaviour Model

We consider an age-structured epidemic model based on a *Susceptible-Latent-Infectious-Recovered* compartmentalization with the addition of pre-symptomatic and asymptomatic infection stages. Susceptible individuals (S), interacting with the infectious compartments, may enter the Latent stage (L). Disease progression let latent individuals evolve in the pre-symptomatic condition (P) and afterwards either in the asymptomatic (A) or symptomatic (I) stage. Both A and I individuals finally recover and enter the Recovered compartments (R_A, R_I). We also compute the number of daily deaths according to the age stratified Infection Fatality Rate (IFR) on the symptomatic infections. The individuals that are responsible for the new infections are I, A, and P, with the latest two characterized by a lower infectiousness. On top of the disease dynamics, we model both the vaccination rollout and the behavioural responses linked to it. After the start of the vaccination rollout, on each day a fraction of the susceptible population receives a vaccine that reduces the probability of infection proportional to vaccine efficacy VE and enters in the compartment V. We introduce the rollout speed r_V as the number of daily administered vaccine doses expressed as a percentage of the total population. We study and compare three different strategies of vaccine prioritization. The first considers distributing vaccines in decreasing order of age. Previous research on COVID-19 vaccines allocation has shown that this strategy is the most effective in reducing the number of deaths and overall severity [10, 42, 43]. It is also the main strategy currently deployed in the different vaccination campaigns across the globe. In the second and third strategy vaccines are either distributed homogeneously or first to the individuals in age brackets 20-49 [51] and then to the rest of the population. The last strategy seeks to reduce symptomatic transmission targeting the most social active part of the population [10]. For simplicity, we will refer to the three rollout approaches as *vaccine strategy 1*, *vaccine strategy 2*, and *vaccine strategy 3*. In parallel to the vaccination, individuals (both vaccinated and not) may start giving up safe behaviours and expose themselves to higher infection risks. We introduce the compartments S^{NC} and V^{NC} for susceptible and vaccinated individuals respectively, to describe different behavioural classes (for convenience "NC" refers to "non-compliant" to COVID-safe behaviours). The increased infection risk for these individuals is captured by the parameter $r > 1$ (for example, $r = 1.3$ indicates an increased risk of 30%). The model accounts also for possible transitions towards and from the non-compliance compartments. In particular, we model the transition from S and V towards riskier behavioural classes as a function of the fraction of the vaccinated population and the parameter α. In turn, we imagine that a worsening of the epidemiological conditions may push non-compliant individuals back to safer behaviours. Here, we consider the number of fatalities per $100,000$ individuals in the previous time step (i.e., day) and the parameter γ to control the second behavioural transition. This modeling choice aims to depict the adaptive nature of human behaviour where individual choices are influenced by vaccination and pandemic progression [52]. We refer the reader to the Materials and Methods section and the Supplementary Information for more details on the model. In the Supplementary Information, we also test another (simpler) mechanism where individuals relax their behaviours (i.e., transit to S^{NC} and V^{NC}) at a constant rate α. Similarly, non-compliant embrace COVID-safe behaviours again (i.e., transit back to S and V) at a constant rate γ. The overall picture emerging adopting this mechanism does not change significantly.

Interplay between NPIs adoption and vaccination campaign

We use data for six different countries: Egypt, Peru, Serbia, Ukraine, Canada, and Italy. As mentioned above, these countries have been selected to sample a range of demographics and socio-economic contexts. In Figure 1A, we represent some key characteristics of the demographic and the mixing patterns between age groups of these countries. First, we consider the fraction of people aged over 65. This is a measure of the epidemic fragility of a country: indeed 65+ individuals are at particularly high risk of mortality from COVID-19 disease [53]. Italy is the country showing the highest fraction of 65+ people (24%), while the developing economies show the lowest fraction (Egypt 5.3% and Peru 8.7%). Indeed, low/middle-income countries tend to have a younger population with respect to the high-income ones. Second, we present also the contacts intensity of different age groups. This is defined as the total number of contacts that an individual in a certain age group has, on average, with all the others in a day. We observe a typical decreasing trend, with younger people that tend to have more contacts. In particular, developing economies show a much higher number of daily contacts for individuals aged under 30. This can play an important role in the spreading even in developed economies. According to a recent study the resurgence of the COVID-19 Pandemic in the US after Summer 2020 was mainly sustained by younger

people [51]. Finally, we represent a measure of inter-generational mixing. We define it simply as the number of daily contacts that an individual in the age groups at high mortality risk from COVID-19 (65+) receives from individuals in the age brackets $0-49$. We observe that Egypt is the country showing the highest inter-generational mixing, followed closely by Peru.

It is important to notice how heterogeneities in health infrastructures, access to health care, and co-morbities are expected across the six countries under examination [50]. These features might induce variations in the IFR. For simplicity however, in the following, we use the same fatality rates for all countries. We leave the exploration of this important aspect to future work.

Prioritization strategies and vaccine efficacy. The proposed model allows to investigate the impact of the behavioural response under different conditions. We consider six populations matching the characteristics of the countries under consideration and explore the model with the same epidemic initial conditions. The experimental setup considers that each population has already experienced a previous wave of infections and that restrictive measures are in place to mitigate the spreading. Therefore, we set the basic reproductive number $R_0 = 1.15$, 1% of initially infected individuals, and 10% of immune individuals. In Figure 1B we show the relative deaths difference for the three vaccine prioritization strategies and three vaccine efficacy $VE = 50\%, 70\%, 90\%$. The relative deaths difference is defined as the fraction of deaths averted thanks to the vaccine with respect to a baseline simulation without vaccine and, therefore, no behavioural response. Moreover, we explore a range of behavioural responses running the simulations for a range of values of the parameter α. Starting from $\alpha = 0$ (which implies no behavioural response), we perform simulations with increasing α values (implying stronger reactions), while keeping constant the other behavioural parameter ($\gamma = 0.5$).

As a first observation, we note that, across the different countries and vaccine efficacy considered, the strategy aiming to reduce the severity of the pandemic (i.e., vaccine strategy 1) is indeed the most effective in reducing the number of deaths, followed by the homogeneous (i.e., vaccine strategy 2) and then the strategy targeting a reduction in transmission (i.e., vaccine strategy 3). This is consistent with the findings of previous studies focused on the allocation of vaccines among groups [42, 43]. As illustrative and concrete example let us consider the case of Canada. When $\alpha = 0$ (i.e., no behaviour response) and $VE = 90\%$ with this strategy the fraction of averted deaths is 0.68 with respect to a baseline without vaccine. This fraction reduces to 0.62 with the homogeneous, and to 0.59 with the strategy prioritizing younger individuals. Furthermore, the difference among strategies diminishes when lower vaccine efficacy are considered. In the previous example, when $VE = 70\%$ the fractions of averted deaths for the three strategies are 0.62, 0.58, 0.57, and when $VE = 50\%$ are 0.54, 0.53, 0.52.

When the behavioural response to the vaccine rollout (i.e., $\alpha > 0$) is considered, we note a consistent decreasing trend of the relative deaths difference for increasing values of α. This shows that the behavioural response impacts the unfolding of the epidemic, and that a relaxation of NPIs leads to a smaller fraction of averted deaths. Interestingly, a concerning effect also emerges: in some conditions, as non-compliance becomes larger, the benefit brought by the vaccine is nullified and the number of observed deaths increases with respect to the no-vaccine and no-behaviour change scenario (i.e., the relative deaths difference become negative). This is solely attributable to the behavioural reaction to the vaccination campaign which in turn is not efficient enough to balance behaviour relaxation. Indeed, this phenomenon is observed in particular when strategies that do not target a reduction in severity are employed, and when the vaccine efficacy is low. In the case of Serbia, with $VE = 90\%$ and strategy aimed at reducing severity, the fraction of averted deaths goes from 0.77 when $\alpha = 0$ to 0.56 when $\alpha = 10$, with a potential loss of 0.21 attributable to the NPIs relaxation. This effect is much more pronounced in the case of the vaccination strategy 2 (or 3): the fraction of averted deaths goes in this case from 0.67 (0.63) when $\alpha = 0$ to 0.34 (0.20) when $\alpha = 10$, with a potential loss of 0.33 (0.43) attributable to the NPIs relaxation. Similarly, lower vaccine efficacy are impacted more significantly by the behaviour. In the previous example, vaccines with an efficacy of 90%, 70%, and 50% would avert, in the case of the vaccination strategy 1, respectively 0.77, 0.69, 0.61 of deaths when $\alpha = 0$. When instead $\alpha = 10$, these figures drop to 0.56, 0.24, -0.19 with a drop of 0.21 in the first, of 0.45 in the second, and of 0.80 in the third case.

In the Supplementary Information we repeat the analysis considering the fraction of averted symptomatic infections instead of deaths. We find that the ranking of allocation strategies is inverted. Indeed, when considering averted infections the most efficient strategy is the one targeting the younger population first, followed by the homogeneous, and the strategy aimed at curbing severity. This is in line with previous findings [42]. Similarly to the case of deaths, also in this additional analysis we find that early behavioral relaxation reduces the fraction of averted cases, influences heterogeneously the prioritization

Figure 1: **Demographic, mixing patterns, and relative deaths difference for different vaccine efficacies and prioritization strategies.** A) Demographic characteristics and mixing patterns of the six countries considered are shown: percentage of the population aged over 65, contacts intensity for different age groups, and a measure of inter-generational mixing. B) Relative deaths difference is computed as the fraction of deaths that are avoided with a vaccine with respect to a baseline simulation without vaccine. We display results of the simulations for three vaccine efficacy and prioritization strategies. Other parameters used are $\gamma = 0.5$, $R_0 = 1.15$, $r = 1.5$, $r_V = 1\%$, 1% of initially infected, 10% of initially immune individuals, and simulations length is set to 1 year.

strategies, and impacts more significantly lower vaccine efficacy.

In the results presented in Figure 1B, we observe that the most disadvantaged countries are the developing economies, Egypt and Peru. At first, this may seem counter-intuitive. Indeed, these countries have a lower fraction of individuals aged over 65. A possible explanation is given by the high activity of young individuals combined with the high inter-generational mixing. Moreover, it is worth stressing that these results are to be intended in relative terms: a relative worst performance in averting deaths does not necessarily imply a worst absolute performance. We observe also that the difference between countries is more profound for the vaccination strategy aimed at reducing transmission. Furthermore, behavioural relaxation widens the distance between the countries. With $VE = 90\%$ and the rollout strategy targeting a reduction in transmission, the gap between the relative deaths difference between Italy and Egypt is 0.22 when $\alpha = 0$. This figure increases to 0.53 when $\alpha = 10$.

Figure 2: **Relative deaths difference for different vaccine rollout speed.** We display the relative deaths difference for three different rollout speed and prioritization strategies. Other parameters used are $\gamma = 0.5$, $R_0 = 1.15$, $r = 1.5$, $VE = 90\%$, 1% of initially infected, 10% of initially immune individuals, and simulations length is set to 1 year.

Vaccine rollout speed. In Figure 2, we investigate the interplay between NPIs relaxation and vaccine rollout speed, keeping the vaccine efficacy constant ($VE = 90\%$) and varying r_V between 0.1% and 1%. We choose these values to cover the spectrum of real vaccination rollout speeds of the vaccination campaigns across the globe. Peru, for example, administered on average 0.05 daily doses per 100 people in the week commencing on the 8^{th} of March, Italy administered 0.30, and Serbia 0.70 [54]. We observe that, besides allocation strategy and vaccine efficacy, rollout speed is a fundamental factor of the vaccination campaign. In the case of Egypt and a vaccination strategy that targets a reduction of severity, when $\alpha = 0$, the fraction of averted deaths spans between 0.54 with the fastest rollout considered ($r_V = 1\%$) and 0.23 with the slowest one ($r_V = 0.1\%$). This implies a drop of 0.31. When, instead, we consider also

behavioural response, this gap widens significantly. Indeed, for $\alpha = 10$, we go from a fraction of averted deaths of 0.31 with $r_V = 1\%$ to -1.12 with the ten times slower rollout $r_V = 0.1\%$, marking an absolute drop of 1.43. This pattern is observed consistently also for other countries, and the effect is more intense for higher values of α. Similarly to previous results, we find that Egypt and Peru are generally more disadvantaged while the difference between countries reduce when the homogeneous or the strategy that reduces severity are employed.

To further explore how to mitigate the impact of behavioural relaxation once that the vaccination campaign started, we systematically explored the interplay between vaccine efficacy and vaccination roll-out speed in Figure 3. The red dashed lines highlight the combinations of vaccine efficacy vs. rollout speed to achieve a 30% drop in observed deaths (taken as reference value), in the absence of a behavioural response (i.e., $\alpha = 0$). In most countries, it can be achieved with a vaccination rollout speed ranging from 0.15% to 0.25% when the available vaccine has an efficacy ranging widely from 90% to 60%. On the contrary, when even a mild behavioural response is considered ($\alpha = 1$), the rollout speed has to increase greatly when vaccine efficacy diminishes: for the case of Italy, a 30% drop in deaths would be achieved with a rollout speed of 0.47% when the vaccine efficacy is 90%, but in the case of $VE = 60\%$ the rollout speed r_V has to grow up to 1.1% to achieve the same result.

Figure 3: **Interplay between vaccine efficacy and rollout speed.** Colors refer to relative deaths difference in the scenario with a mild behavioural response ($\alpha = 1$) and a 30% reduction of deaths is highlighted with a dashed black line. The red dashed line highlight the 30% death drop achieved with a vaccination campaign without the behavioural component. Vaccination Strategy 1 is considered, parameters used are $\gamma = 0.5$, $R_0 = 1.15$, $r = 1.5$, 1% of initially infected, 10% of initially immune individuals, and simulations length is set to 1 year.

Epidemic trajectories and impact of behavioural relaxation. As a way to ground the model on more realistic epidemic scenarios, we calibrate it to the real epidemic trajectories of the six countries considered. The calibration is performed via an Approximate Bayesian Computation technique (ABC) [55] on weekly deaths for the period $2020/09/01 - 2020/12/31$. In doing so, we estimate some key epidemiological parameters matching the context of each country, thus defining realistic initial conditions for the model. To capture the effects of non-pharmaceutical interventions on the contacts between people we consider data from the Google Mobility Report [56] and the Oxford Coronavirus Government Response Tracker [57] until week 11 of 2021. After the calibration step, the model evolves from 2021/01/01 up to 2021/06/01 with the vaccination rollout for each of the countries. Note that in some of these countries, the vaccination campaigns officially started in the second half of December, though mostly symbolically.

Figure 4: **Giving up NPIs during rollout may nullify the benefits brought by the vaccine.** A) We display for the different countries the boxplot of the calibrated infection parameter β, the projected number of symptomatic infectious cases per $100'000$ and the fraction of recovered at the $2021/01/01$, start of the vaccination campaign in our simulation. We also report the ratio between the leading eigenvalue of the contacts matrix considering restrictions and of the baseline contacts matrix with no restrictions. B) We display the median relative deaths difference for the calibrated model in the different countries. We consider the three vaccination strategies and two possible rollout speed: $r_V = 1\%$ (faster rollout), and $r_V = 0.25\%$ (slower rollout). We run the model over the period $2021/01/01$-$2021/06/01$. Other parameters are $\gamma = 0.5$, $r = 1.5$, $VE = 90\%$.

In Figure 4A we represent key indicators resulting from the calibration. In particular, for the six countries we report the boxplot for the calibrated infection parameter β, and also the projected number of symptomatic infections per $100,000$ and the fraction of recovered individuals on the $2021/01/01$, which is the start of the vaccination campaign in our simulation. We acknowledge a significantly high acquired immunity in the case of Peru, where the estimated attack rate as of 1^{st} January, 2021, is around 38%. Nonetheless, this figure is line with other available estimates [54]. We observe how the calibration allows for an heterogeneous representation of the epidemiological conditions of the different countries. In Figure 4A we also represents the effects of restrictive measures on contacts. As a proxy, we consider the ratio between the leading eigenvalue of the contacts matrix considering the restrictions and the leading eigenvalue of the baseline contacts matrix without restrictions. The leading eigenvalue of the contacts matrix influences the reproductive number of the disease (see Materials and Methods), and more broadly it measures the intensity of contacts among people. By normalizing with respect to the baseline contacts matrix (with no restrictions) we can grasp the strictness of the measures in place. For example, a ratio of 0.3 would imply, in our simulations, a 70% reduction of the reproductive number with respect to the baseline without restrictions. The effect of restrictive measures on contacts varies over time up to week 11, 2021. In the case of Italy, for example, we observe the partial ease of the measures during January and February (i.e., $\rho(C_t)/\rho(C_b)$ increases) followed by the tightening of measures in March aimed at curbing the third wave of infections. Afterwards, we keep the mixing levels as those observed for week 11 (more details in the Materials and Methods section). This is a conservative assumption as we can imagine that the seasonality and impact of the vaccination campaign might induce a relaxation of NPIs as the number of cases and deaths, hopefully, will go down. In this scenario, the countries with the strictest measures in place after week 11, are Peru and Italy, while Egypt and Ukraine show the most permissive measures. In the Supplementary Information, we repeat the analysis considering scenarios in which we partially relax some of the measures. The results are in line with the observations presented below.

In Figure 4B we represent the relative deaths difference for the calibrated model in the six countries. We study two vaccination rates. A fast deployment that manages to vaccinate $r_V = 1\%$ of the population daily (aligned with the preparedness plan for Influenza pandemic [58] and vaccination rates achieved by countries like Israel or Chile), and a slower yet more realistic rollout $r_V = 0.25\%$ (aligned with current rates in European countries). Worryingly, in the different countries, even in a scenario with several restrictions and a successful vaccination campaign in place, it is possible to witness a significant increase (factors reach a level of 2) in deaths exclusively due to a relaxation of COVID-safe behaviours. As a result of the calibration step, the countries under investigation face different epidemiological conditions (e.g., effective reproductive number, immunity from previous waves, initial number of infected, estimated effects of restrictions). These translate into phenomenological differences with respect to the results of the previous analyses. In particular, Ukraine and Egypt are significantly impacted by behavioural relaxation in the case of the slower rollout $r_V = 0.25\%$. From Figure 4A we see that in both countries the estimated effect of restrictions on contacts (especially in the last weeks) is smaller with respect to the others. This underlines how, besides the characteristics of the vaccination campaign, also the epidemiological conditions and the measures in place are important factors influencing the behaviour-vaccine interplay. In the case of the faster rollout $r_V = 1\%$, the ranking of the countries is similar to the one obtained previously (Peru and Egypt are generally more disadvantaged), when we imposed the same conditions for all countries. A possible explanation is that when a fast rollout is employed the influence of epidemiological conditions and restrictions become less important (because of the efficiency of the rollout), and the effects observed in the exploratory analysis emerge more easily.

Figure 4B shows that while the vaccination strategy targeting a reduction of severity performs slightly better when no and mild behavioural responses are in place, it appears to be much more robust when strong behavioural responses are considered. Also, while in the case of a faster rollout ($r_V = 1\%$) behavioural changes impact mildly the number of avoided deaths, with a 4 times slower rollout ($r_V = 0.25\%$) we find that the number of deaths consistently increases due to a relaxation of COVID-safe behaviours. In fact, a relaxation of protective behaviors not backed by a fast enough rollout, might induce an increase of deaths with respect to scenarios without vaccines. Finally, in case of homogeneous vaccination or prioritization aimed at reducing transmission overconfident behaviours further increase such death toll. These findings are consistent across the different countries considered.

In order to provide a measure about vaccination strategy robustness against behavioural relaxation, we sum the relative deaths difference (or relative symptomatic cases difference) across the spectrum of considered α values. In Figure 5 we show for every country how these two key metrics would be affected by different vaccination strategies when a wide range of behavioural responses are considered.

Figure 5: **Robustness of vaccination strategies - Calibrated model.** For the different countries we represent the cumulative relative difference of deaths and symptomatic cases for α ranging from 0 to 10^3. Stars highlight the most effective strategy w.r.t. the measure of interest. We set $r_V = 1\%$, $VE = 90\%$, $\gamma = 0.5$, $r = 1.5$.

The plots show that vaccination strategy 1 is the most effective in reducing the number of deaths, while vaccination strategy 3 is the most effective in reducing the number of symptomatic cases, thus confirming that a potential behavioural response would not alter the already known vaccination prioritization strategies. Nonetheless, because of the unfolding of the epidemic in Egypt and Peru and the high inter-generational mixing, the vaccination strategy 2 (homogeneous across age classes) appears to be slightly better to minimize symptomatic cases when COVID-safe behaviours are lifted too soon. Information about vaccination strategy robustness against behavioural changes might be used to tune resilient roll-out campaigns. Policy-makers might also consider to optimize vaccination strategies with respect to a combination of multiple metrics. In this direction, our analysis provide additional insights. Interestingly, in Italy, Canada, Serbia and Ukraine, strategy 1 slightly outperforms strategy 3 in terms of robustness against behavioural changes to decrease the number of deaths, but strategy 3 greatly outperform strategy 1 in terms of robustness towards avoided infections, thus suggesting once again that behavioural changes play an important role that might modify our understanding of prioritization strategies.

Discussion

For almost a year, in the midst of a global pandemic, policymakers struggled to implement sustainable restrictions to slow SARS-CoV-2 spreading. Every non-pharmaceutical intervention was aimed at flattening the incidence curve and buying time for the development, test, production and distribution of vaccines that might ultimately protect the population. With an impressive scientific endeavor, several vaccines have been developed and an early distribution campaign was rolled out by the last days of December 2020. Besides the potential novel threats emerging from new virus strains [59], the current vaccination campaign represents the beginning of a new normal and a gigantic step towards complete virus suppression. However, we demonstrated that if the growth in vaccination uptake would lead to overconfident conducts inducing relaxation of COVID-safe behaviours, additional avoidable deaths will occur. We extended the literature proposing a mechanistic compartmental model able to simulate the unfolding of COVID-19, the vaccination dynamics and the compliance/non-compliance transition modulated by different behavioural mechanisms. Performing in-silico simulations allowed us to explore, from a theoretical standpoint, the interplay among different vaccination strategies, vaccine efficacy, rollout

speeds, and behavioural responses. We found that behaviour impacts non-linearly vaccine effectiveness. Indeed, early NPIs relaxation affects more significantly lower vaccine efficacy, slower rollouts, and allocation strategies that target reduction in transmission rather in severity. We included in our analysis six different countries (Egypt, Peru, Serbia, Ukraine, Canada, and Italy), representatives of various points of the spectrum of world economies. This allowed us to observe the effects of behaviour-vaccine relationships for various population pyramids and mixing patterns. We observed that the developing economies, characterized by a younger population, but higher contacts activity and inter-generational mixing, are generally more affected by behaviour change with respect to developed economies. Then, as a way to ground the model on more realistic epidemiological conditions, we calibrated it using real epidemic and mobility data for the six countries considered and simulated the unfolding of the first months of the vaccination campaign. In such realistic scenario, we observed that with restrictive measures in place and a successful vaccination campaign, it is possible to witness to non-negligible increases in deaths attributable to an early relaxation of COVID-safe behaviours. The calibration step allowed us to highlight that also the epidemiological conditions related to the country-specific unfolding of the disease are an important factor influencing the interplay behaviour-vaccine.

We acknowledge some limitations in the present study. First, the vaccine is modeled as protective towards infection, without modeling the protection against severe COVID-19 complications and potential reduction of viral shedding in infected vaccinated individuals. At the time of writing, such information are still matters of study though initial reports support high level of protection from severe outcomes and a significant reduction in transmission [16, 17, 60]. For the sake of simplicity, we considered the vaccines fully working immediately after the first dose and we neglected that vaccination campaigns are using a portfolio of vaccines rather than a single one. We have also studied three simple vaccination strategies that neglect the complexities of an unprecedented mass vaccination. As result, both the vaccination priorities, vaccine effects and vaccination rates are an approximation of reality. In the Supplementary Information, we studied a data-driven vaccination rollout for Italy, where vaccines are distributed to the various age brackets following the real daily administration data. The results we obtained are qualitatively similar to those presented in the main text. While the model calibration suggests that our approach can nicely capture national trends, the model is not meant to provide accurate forecasts of the local unfolding of the disease, but rather to test what-if scenarios in a comparative fashion. We have considered a simple age-structure compartmental model that does not capture spatio-temporal heterogeneity both in terms of spreading and of NPIs implementation which have instead been observed in the countries under investigation. Our model does not include new, more transmissible, variants of concern and assumes the same IFR across the six countries. Finally, we propose and model two potential mechanisms leading to behavioural changes, but data are not available to perform a quantitative validation of the behavioural components of the model.

Implementation of individual protective behaviours and adherence to NPIs have been vital in order to reduce the transmission of SARS-CoV-2 leading to substantial population-level effects [5, 61–72]. Behavioural science can provide valuable insights for managing policies, incentives, communication strategies and can help coordinate efforts to control threats and evaluate such interventions [73]. As during the first waves of COVID-19, when NPIs were the only available mitigation measures [74], the results of our paper call for adequate strategies to keep high the attention and compliance towards individual COVID-safe behaviours, such as mask-wearing, social distancing, and avoidance of large gatherings now that vaccines are finally available. Communication strategies and policies should keep targeting such non-pharmaceutical intervention to avoid frustrating the immense effort of the vaccination campaigns.

Materials and Methods

Epidemic Model

Model Definition. We propose a compartmental model that incorporates both the vaccination process and the behaviour dynamics possibly linked to it. See Figure 6 for a schematic representation. The virus transmission is modeled using the following approach. Susceptible individuals (S compartment) in contact with infectious compartments become latent (L compartment). After the latent period (ϵ^{-1}), L individuals enter the pre-symptomatic stage of the infection (P). Then, they can transition either in the asymptomatic (A) or the symptomatic stage (I) at rate ω (the length of time spent in the L and P compartments is the incubation period). The probability of being asymptomatic is f. After the infectious period (μ^{-1}), both I and A individuals enter the Recovered compartment (R). Alternatively,

I individuals can also die and transit to the D compartment. Note how we include a delay of Δ days from the time individuals enter the compartment D and die (D^o compartment, the superscript stands for "observed"). The infectious compartments are P, A, I. The transmission rate is β and the force of infection is dependent on the age-stratified contact matrix $\mathbf{C} \in \mathbb{R}^{K \times K}$, whose element C_{ij} represents the average number of contacts that an individual in age group i make with individuals in j per day. The matrix $\mathbf{C}$ has four location-specific contributions: contacts at home, workplace, school, and other locations. We adopt country-specific contacts matrices provided in Ref. [75]. We assume that P and I have lower infectiousness with respect to symptomatic I ($\beta\chi$, with $\chi < 1$). Similar approaches have been used in several modeling studies in the context of the current pandemic [67, 76].

On top of the disease dynamics, we model both the vaccination process and the behavioural change that is possibly coupled to the vaccination. More in detail, after the start of the vaccination campaign at t_V, at each time step, a fraction r_V of the susceptible population receive a vaccine and transit to compartment V. As mentioned above, we consider three vaccination strategies: one in which the vaccine is given in decreasing order of age (vaccination strategy 1 aimed at reducing the severity), one in which it is given homogeneously to the population (vaccination strategy 2), and one in which it is first given to individuals in age brackets 20-49 and then to the rest of the population (vaccination strategy 3 aimed at reducing the transmission of the virus) [10]. For convenience, we derive the model's equations in the homogeneous case, we refer the reader to the following sections for more details. We consider a "leaky" vaccine that reduces susceptibility with a certain efficacy VE. In other words, the infection rate for V individuals is $\beta(1 - VE)$. In parallel, we imagine a behavioural dynamics triggered by the presence of the vaccine. Indeed, susceptible individuals (both vaccinated and not) may start adopting less safe behaviours because reassured by the presence of an effective vaccine. This is encoded in the model with a transition from the compartment S (V) to a new compartment S^{NC} (V^{NC}) of individuals that protect less themselves and as a result get infected at a higher rate ($r\beta$, with $r > 1$). Note that NC stands for non-compliant. We propose two mechanisms to model this transition. In the first mechanism, the transition towards non-compliance happens at a rate α and it is catalyzed by the cumulative fraction of vaccinated individuals v_t. The opposite transition from S^{NC} (V^{NC}) to S (V) happens at rate γ and is catalyzed by the number of observed deaths per $100,000$ individuals in the previous time step d_{t-1}^o. Indeed, an increase in deaths is frequently used - especially by media - as an indicator of the severity of the current epidemiological situation. Existing literature suggests that risk perception (in the form of number of infected individuals or deaths) and communication of such risk significantly affect adherence to personal mitigation strategies such as social distancing and wearing face masks [5, 77]. Expressing the number of deaths in proportion to the population allows us to compare countries of different size. In the second mechanism, S (V) individuals transit to the non-compliant compartment S^{NC} (V^{NC}) at a constant rate α. We also account for the possibility of non-compliant going back to safer behaviours again at a constant rate γ. To simplify the narrative, we present the results considering this second mechanism only in the Supplementary Information. The overall picture discussed above does not change significantly. Note that in order to avoid issues with transition probabilities large than one we model the rates as $\lambda_{X \to X^{NC}} = 1 - \exp^{-g(\alpha)}$, $\lambda_{X^{NC} \to X} = 1 - \exp^{-h(\gamma)}$ with $X = [S, V]$ and where the specific expression of the exponent depend on the two mechanisms described above. Note how for small values of $g(\alpha)$, $h(\gamma)$ the rate converges to the usual mass-action law. The model can be written down as the following system of differential equations for individuals in age group k:

$$
\begin{aligned}
\frac{dS_k}{dt} =& -\beta \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} S_k - (1 - e^{-g(\alpha)})S_k + (1 - e^{-h(\gamma)})S_k^{NC} + \\
& - min(S_k, r_V N_k \frac{S_k}{S_k + S_k^{NC}})\delta[t - t_V] \\
\frac{dV_k}{dt} =& -\beta(1 - VE) \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} V_k - (1 - e^{-g(\alpha)})V_k + (1 - e^{-h(\gamma)})V_k^{NC} + \\
& + min(S_k, r_V N_k \frac{S_k}{S_k + S_k^{NC}})\delta[t - t_V] \\
\frac{dS_k^{NC}}{dt} =& -r\beta \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} S_k^{NC} + (1 - e^{-g(\alpha)})S_k - (1 - e^{-h(\gamma)})S_k^{NC} + \\
& - min(S_k^{NC}, r_V N_k \frac{S_k^{NC}}{S_k + S_k^{NC}})\delta[t - t_V] \\
\frac{dV_k^{NC}}{dt} =& -r\beta(1 - VE) \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} V_k^{NC} + (1 - e^{-g(\alpha)})V_k - (1 - e^{-h(\gamma)})V_k^{NC} + \\
& + min(S_k^{NC}, r_V N_k \frac{S_k^{NC}}{S_k + S_k^{NC}})\delta[t - t_V] \\
\frac{dL_k}{dt} =& +\beta \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} S_k + r\beta \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} S_k^{NC} + \\
& + \beta(1 - VE) \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} V_k + r\beta(1 - VE) \sum_{k'=1}^{K} C_{kk'} \frac{I_{k'} + \chi(P_{k'} + A_{k'})}{N_{k'}} V_k^{NC} - \epsilon L_k \\
\frac{dP_k}{dt} =& \epsilon L_k - \omega f P_k - \omega(1 - f)P_k \\
\frac{dI_k}{dt} =& \omega(1 - f)P_k - \mu(1 - IFR_k)I_k - \mu IFR_k I_k \\
\frac{dA_k}{dt} =& \omega f P_k - \mu A_k \\
\frac{dR_k}{dt} =& \mu(1 - IFR_k)I_k + \mu A_k \\
\frac{dD_k}{dt} =& \mu IFR_k I_k - \Delta^{-1} D_k \\
\frac{dD_k^o}{dt} =& \Delta^{-1} D_k
\end{aligned}
$$

$$(1)$$

Where $\delta[t - t_V]$ is a step function which is equal to 1 if $t > t_V$ and to 0 otherwise. The basic reproduction number is $R_0 = \rho(\tilde{C})[\frac{\beta\chi}{\omega} + \frac{\beta(1-f)}{\mu} + \frac{\beta\chi f}{\mu}]$, where $\tilde{C}$ is the contacts matrix weighted by the relative population in different age groups. In the Supplementary Information we provide details on its derivation. We adopt the model integrating numerically the equations, thus it is deterministic. However, it is worth stressing that when we calibrate the model to the real epidemic trajectory in the six countries, we use a probabilistic framework through an Approximate Bayesian Computation technique. Said differently, the calibrated parameters are characterized by posterior probability distributions rather than exact values. For this reason, the results presented in Figure 4 (i.e., the median of model's projections) are to be intended as an ensemble of multiple trajectories generated sampling from the posterior distribution of the fitted parameters. The model is implemented in the programming language python with the use of the libraries *scipy* [78], *numpy* [79], and *numba* [80]. The visualization of the results is realized with the library *matplotlib* [81].

Phase space exploration. In Figure 7 we explore the phase space of the behavioural parameters (α, γ) of the model. We set for the six countries $R_0 = 1.15$, 1% of initially infected individuals, and 10% of immune individuals, $r_V = 1\%$, and $VE = 90\%$. After setting the initial conditions, we let the model

Figure 6: **Compartmental model.** We consider an extension of the classic SLIR model adding compartments for presymptomatic (P) and asymptomatic (A). We also design compartments for vaccinated (V), dead (D describes individuals that will die with a delay of Δ^{-1} entering the compartment D^o), susceptible (S^{NC}) and vaccinated (V^{NC}) individuals that do not comply with COVID-safe behaviours. The vaccine offers a protection VE. The transmission rate for susceptible is β and for susceptible non-compliant $r\beta$ ($r > 1$). The parameter α regulates the transition from compliant to non-compliant behaviours, while γ regulates the opposite flow. Arrows describe the transitions between compartments. The dashed lines represent transitions mediated by interactions. For example, individuals move from S^{NC} to L by interacting with infectious individuals (P, A, I) with a transmission rate of $r\beta$. The compartmentalization is then extended to account for empirical age-structure and contact matrices.

evolve, individually for each country, for one year exploring a grid of the parameters α and γ. This allows us to observe the phase space of parameters regulating the behavioural transitions. In particular, for each (α, γ) pair we compute the relative deaths difference due to vaccines and behaviour change. As stated previously, the relative deaths difference is the fraction of deaths averted with respect to a baseline without vaccine and behavioural response. For the countries of focus, we consider two different values of the parameter r ($r = 1.3$, 1.5) which defines the increase in infection risk for individuals relaxing preventive behaviours. The obtained relative deaths difference varies from a maximum of 0.77 to a minimum of 0.26. This indicates that, in our simulations in the best scenario about 80% of deaths are averted thanks to the vaccine rollout. However, this potential gain reduces to around 25% in the worst case, with a potential waste of 55% of the benefit brought by the vaccine in terms of reduced mortality. Since α and γ are the only varying parameter in these simulations, such a reduction is only attributable to the relaxation of NPIs from the individuals. More in detail, across the different settings considered, a common pattern emerges. For α fixed, as γ grows we observe progressively an increase in the relative deaths difference. Indeed, if the population reacts promptly, non-compliant individuals turn back to COVID-safe behaviours, and the fraction of averted deaths benefits from this behaviour. On contrary, for a fixed γ an increase of α induces a stronger behavioural response causing more deaths otherwise avoided thanks to the vaccination. Furthermore, we observe that, for a given pair of the behavioural parameters, the fraction of averted deaths is lower when $r = 1.5$ with respect to $r = 1.3$. Indeed, in this case non-compliant individuals expose themselves to a higher risk of infection.

In the Results section, we kept for simplicity $\gamma = 0.5$ and we let vary α. This choice is informed by the maximum number of deaths observed in the countries of focus. In Italy, for example, the maximum number of deaths reported on a single day is around $1,000$. Therefore, this value of γ is such that, in a similar situation, non-compliant individuals would likely return to COVID-safe behaviours. As described in the previous paragraph, in the dynamic-rate behavioural mechanism we model the transition rate as $\lambda_{X^{NC} \to X} = 1 - \exp^{-\gamma d^o_{t-1}}$ ($X = [S, V]$), where d^o_{t-1} is the number of deaths per $100,000$ individuals observed at time $t - 1$. Hence at the peak of deaths the transition rate towards compliance is $\lambda_{X^{NC} \to X} \sim 0.6$. The sensitivity to this choice is discussed in the Supplementary Information.

Figure 7: **Impact of model parameters governing the behavioural response.** For the six countries, we explore different values of α and γ in terms of fraction of averted deaths with respect to a baseline without vaccine and no behavioural response (i.e. $r_V = 0\%$, $\alpha, \gamma = 0$). We consider two different values of the parameter r (1.3 and 1.5) and we set $r_V = 1\%$ and $VE = 90\%$ and we employ vaccine strategy 1.

Vaccination Strategies

As mentioned above, we consider three vaccination strategies. In the first one, the vaccine is given in decreasing order of age. Since the IFR of COVID-19 strongly correlates with age, many countries worldwide are adopting similar strategies prioritizing the vaccination among the elderly. Previous modeling works in the context of COVID-19 showed that this is the preferable strategy when considering the number of averted deaths [10, 42, 43]. In practice, this means that in our simulations we start giving the vaccine to the 75+ age bracket and we proceed in decreasing order only when everyone in this group is vaccinated. In the second strategy the vaccine is given homogeneously to the population respecting the age distribution. This means that, of the X vaccines available at step t, the fraction N_k/N is given to age group k (where N_k is the number of individuals in age bracket k, and N is the total number of individuals). In the third strategy, we prioritize age groups $20-49$. Indeed, individuals at high mortality risk from COVID-19 disease may be protected indirectly by vaccinating age brackets that sustain the transmission [82, 83]. In practice, in our simulations we start giving the vaccine homogeneously to the $20-49$ age brackets, and when they are all vaccinated we give it homogeneously to all other groups. In the previous two strategies, if, on a given time step, the number of people remained in the age group which is currently being vaccinated is smaller than the total number of vaccines available, the exceeding part is given to the next age group in line. This implies that the number of vaccines given in different time steps is always constant. In all cases, the vaccines are given only to the susceptible population.

Model Calibration

In this section we describe the methods used to calibrate the model to the real epidemic trajectories in the countries considered. We use an Approximate Bayesian Computation technique to calibrate the model on weekly deaths during the period $2020/09/01 - 2020/12/31$. We account for government-mandated restrictions and their effect on the contacts between individuals modifying the contacts matrices using data from the Google Mobility Report [56] and the Oxford Coronavirus Government Response Tracker [57].

Approximate Bayesian Computation. We calibrate the model for each country using the Approximate Bayesian Computation (ABC) rejection method [55, 84]. At each step of the rejection algorithm, a set of parameters θ is sampled from a prior distribution and an instance of the model is generated using θ. Then, an output quantity E' of the model is compared to the corresponding real quantity E using a distance measure $s(E', E)$. If this distance is greater (smaller) than a predefined tolerance ψ, then the sampled set of parameters is discarded (retained). After accepting N sets, the iteration stops and the

distribution of accepted parameters is an approximation of the real posterior distribution $P(\theta, E)$. The free parameters of our model and the related prior distributions are:

- The transmission rate β. We explore uniformly values of β such that the related R_0 is between 0.8 and 2.2. The basic reproduction number of SARS-CoV-2 is higher [85], but we consider lower values since our calibration starts on 2020/09/01 when restrictions were in place to mitigate the spreading.

- The delay in deaths $\Delta \sim [14, 25]$. Indeed, for COVID-19 the average time between symptoms onset and death is about 2 weeks [86] and we also account for possible additional delays in death reporting.

- The initial number of infected individuals. We explore uniformly values between 0.5 and 15 times the number of reported cases in the week before the start of the simulation. We then assign these individuals to the infected compartments (L, P, A, I) proportionally to the time spent there by individuals (ϵ^{-1} for L, ω^{-1} for P, and μ^{-1} for I, A), and we split between I and A individuals considering the fraction of asymptomatic f.

The model is calibrated on the period $2020/09/01 - 2020/12/31$ using the weighted mean absolute percentage error on weekly deaths as an error metric with a tolerance $\psi = 0.3$ ($\psi = 0.4$ for Egypt for convergence issues) and $5,000$ accepted parameters set.

Model Initialization. The number of individuals in different age groups is initialized considering the 2019 United Nation World Population Prospects [87]. We consider 16 age brackets of five-years, except for the last one that includes 75+ individuals. As specified above, the initial number of infected individuals is calibrated considering the total number of confirmed cases in the week before the start of the simulation according to the data issued from the European Centre for Disease Prevention and Control [88]. To initialize the number of non-susceptible individuals (placed in the R compartment) we compute the average of several publicly available projections of total COVID-19 infections up to 2020/09/01 (i.e., start of the simulation) from different modeling approaches [54]. Both the initial number of infected and of non-susceptibles is assigned homogeneously across age groups. Other parameters used are $\epsilon^{-1} = 3.7$ days, $\mu^{-1} = 2.5$ days, $\omega^{-1} = 1.5$ days $\chi = 0.55$, $f = 0.35$ in line with current estimates of COVID-19 infection dynamics parameters [89–93]. We use the age-stratified Infection Fatality Rate (IFR) from Ref. [53].

Modeling the Effects of NPIs on Contact Matrices. We incorporate in our model the implementation of top-down NPIs by changing the contacts patterns defined by the contacts matrix $\mathbf{C}$. As mentioned, we use the country-specific contacts matrices provided in Ref. [75]. These are made by four contributions: contacts that happen at school, workplace, home, and other locations. In a baseline scenario, the overall contacts matrix $\mathbf{C}$ is simply the sum of these four contributions. Following Ref. [94], we implement the reductions in contacts, due to the restrictions, multiplying the single contribution by a reduction factor $\eta_i(t)$. Thus, in general the overall contacts matrix at time t become:

$$C(t) = home + \eta_w(t) \cdot work + \eta_s(t) \cdot school + \eta_{ol}(t) \cdot other\ locations \tag{2}$$

For simplicity, we assume no changes to contacts at home, though lockdowns tend to increase them [61]. For the contacts locations *work* and *other locations* we use data from the Google Mobility Report [56]. In detail, we consider the fields `workplaces percent change from baseline` to model reduction in the contacts location *work* and the average of the fields `retail and recreation percent change from baseline` and `transit stations percent change from baseline` for *other locations*. A general entry of the report $p_l(t)$ represents the percentage change on day t of total visitors to a specific location l with respect to a pre-pandemic baseline. From $p_l(t)$ we derive a contacts reduction coefficient as follows: $\eta_l(t) = (1 + p_l(t)/100)^2$. Indeed, the number of potential contacts in a specific location is proportional to the square of the the number of visitors. We model contacts reduction in school considering instead the Oxford Coronavirus Government Response Tracker [57]. More in detail, we use the ordinal index `C1 School closing`. The index ranges from a minimum of 0 (no measures) to a maximum of 3 (require closing all levels). We turn this quantity into the contacts reduction coefficient in school as follows: $\eta_s(t) = (3 - \text{C1 School closing})/3$. We use these datasets to inform contacts reductions up to week 11, 2021. After, we assume that contacts remain at the same level of week 11. Note how we let vary both the transmission rate β and the contact matrix. The former describes the risk of infection given contacts

with infectious individuals. This is function of the disease (which is assumed to be the same, we don't consider multiple or emergent new strains possibly more transmissible) and of the protective behaviours such as social distancing and use of face masks. The latter describes variations to the number and types of contacts induced by top-down NPIs as for example remote working, schools closure and lockdowns. By splitting the contributions to the force of infection of transmission rate and contact matrix we are able to take into consideration different behavioural attitudes which, given the same number of contacts, might lead to higher or lower risks of infection. This allows us to consider explicitly both top-down and bottom-up NPIs.

Calibration Results. In Figure 8, we report the results of the calibration. It is important to stress how our goal is not to develop a predictive model aimed at forecasting the pandemic trajectory. The fit is used to ground the model and to define the epidemic conditions at the start of the vaccination campaign in the six countries. In fact, our aim is to understand the possible interplay between behaviours and vaccine rollout which is also function of the epidemic progression. In the Figure we report the official and simulated weekly number of deaths (median, 50% and 95% confidence interval). Despite its simplicity and approximations, the model is able to reproduce the evolution of the pandemic in the six countries capturing well its progression after the summer. In the Supplementary Information we also report the posterior distributions for the parameters.

Figure 8: **Calibration Results.** For each country we represent the observed and simulated weekly deaths (median, 50% and 95% confidence intervals).

Acknowledgements: All authors thank the High Performance Computing facilities at Greenwich University. NG acknowledges support from the Doctoral Training Alliance. PB acknowledges support from Intesa Sanpaolo Innovation Center. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions NG, PB, NP conceived the research and designed the analyses. NG conducted

the numerical simulations. All authors wrote the manuscript.

Competing Interests The authors declare that they have no competing financial interests.
Data Availability: All the data and code used in this paper are publicly available at `https://github.com/ngozzi/vaccine-behaviour`.

References

[1] Amelie Desvars-Larrive, Elma Dervic, Nils Haug, Thomas Niederkrotenthaler, Jiaying Chen, Anna Di Natale, Jana Lasser, Diana S Gliga, Alexandra Roux, Abhijit Chakraborty, et al. A structured open dataset of government interventions in response to covid-19. *medRxiv*, 2020.

[2] Benjamin J Cowling, Sheikh Taslim Ali, Tiffany WY Ng, Tim K Tsang, Julian CM Li, Min Whui Fong, Qiuyan Liao, Mike YW Kwan, So Lun Lee, Susan S Chiu, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in hong kong: an observational study. *The Lancet Public Health*, 2020.

[3] Seth Flaxman, Swapnil Mishra, Axel Gandy, H Juliette T Unwin, Thomas A Mellan, Helen Coupland, Charles Whittaker, Harrison Zhu, Tresnia Berah, Jeffrey W Eaton, et al. Estimating the effects of non-pharmaceutical interventions on covid-19 in europe. *Nature*, 584(7820):257–261, 2020.

[4] Nils Haug, Lukas Geyrhofer, Alessandro Londei, Elma Dervic, Amelie Desvars-Larrive, Vittorio Loreto, Beate Pinior, Stefan Thurner, and Peter Klimek. Ranking the effectiveness of worldwide covid-19 government interventions. *Nature Human Behavior*, 2020.

[5] Nicola Perra. Non-pharmaceutical interventions during the covid-19 pandemic: A review. *Physics Reports*, 2021.

[6] Holly Seale, Clare EF Dyer, Ikram Abdi, Kazi M Rahman, Yanni Sun, Mohammed O Qureshi, Alexander Dowell-Day, Jonathon Sward, and M Saiful Islam. Improving the impact of non-pharmaceutical interventions during covid-19: examining the factors that influence engagement and the impact on individuals. *BMC Infectious Diseases*, 20(1):1–13, 2020.

[7] Giulia Pullano, Laura Di Domenico, Chiara E Sabbatini, Eugenio Valdano, Clément Turbelin, Marion Debin, Caroline Guerrisi, Charly Kengne-Kuetche, Cécile Souty, Thomas Hanslik, et al. Underdetection of covid-19 cases in france threatens epidemic control. *Nature*, 2020.

[8] Ewen Callaway. What Pfizer's landmark COVID vaccine results mean for the pandemic. *Nature*, 2020.

[9] Ewen Callaway. COVID vaccine excitement builds as Moderna reports third positive result. *Nature*, 2020.

[10] Wei Wang, Qianhui Wu, Juan Yang, Kaige Dong, Xinghui Chen, Xufang Bai, Xinhua Chen, Zhiyuan Chen, Cécile Viboud, Marco Ajelli, et al. Global, regional, and national estimates of target population sizes for covid-19 vaccination: descriptive study. *bmj*, 371, 2020.

[11] Tom Britton, Frank Ball, and Pieter Trapman. A mathematical model reveals the influence of population heterogeneity on herd immunity to sars-cov-2. *Science*, 369(6505):846–849, 2020.

[12] Mark Channels Read. Eid: High contagiousness and rapid spread of severe acute respiratory syndrome coronavirus 2. *Emerg. Infect. Dis*, 26, 2020.

[13] Marc Lipsitch and Natalie E Dean. Understanding covid-19 vaccine efficacy. *Science*, 370(6518):763–765, 2020.

[14] Fernando P Polack, Stephen J Thomas, Nicholas Kitchin, Judith Absalon, Alejandra Gurtman, Stephen Lockhart, John L Perez, Gonzalo Pérez Marc, Edson D Moreira, Cristiano Zerbini, et al. Safety and efficacy of the bnt162b2 mrna covid-19 vaccine. *New England Journal of Medicine*, 2020.

[15] Lindsey R. Baden, Hana M. El Sahly, Brandon Essink, Karen Kotloff, Sharon Frey, Rick Novak, David Diemert, Stephen A. Spector, Nadine Rouphael, C. Buddy Creech, John McGettigan, Shishir Khetan, Nathan Segall, Joel Solis, Adam Brosz, Carlos Fierro, Howard Schwartz, Kathleen Neuzil, Larry Corey, Peter Gilbert, Holly Janes, Dean Follmann, Mary Marovich, John Mascola, Laura Polakowski, Julie Ledgerwood, Barney S. Graham, Hamilton Bennett, Rolando Pajon, Conor Knightly, Brett Leav, Weiping Deng, Honghong Zhou, Shu Han, Melanie Ivarsson, Jacqueline Miller, and Tal Zaks. Efficacy and safety of the mrna-1273 sars-cov-2 vaccine. *New England Journal of Medicine*, 0(0):null, 0.

[16] Noa Dagan, Noam Barda, Eldad Kepten, Oren Miron, Shay Perchik, Mark A Katz, Miguel A Hernán, Marc Lipsitch, Ben Reis, and Ran D Balicer. Bnt162b2 mrna covid-19 vaccine in a nationwide mass vaccination setting. *New England Journal of Medicine*, 2021.

[17] PHE monitoring of the early impact and effectiveness of COVID-19 vaccination in England. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/963532/COVID-19_vaccine_effectiveness_surveillance_report_February_2021_FINAL.pdf, 2021. Accessed: 2020-03-06.

[18] Jeffrey V Lazarus, Scott C Ratzan, Adam Palayew, Lawrence O Gostin, Heidi J Larson, Kenneth Rabin, Spencer Kimball, and Ayman El-Mohandes. A global survey of potential acceptance of a covid-19 vaccine. *Nature medicine*, pages 1–4, 2020.

[19] Malik Peiris and Gabriel M Leung. What can we expect from first-generation covid-19 vaccines? *Lancet (London, England)*, 2020.

[20] Rochelle P Walensky, Henry T Walke, and Anthony S Fauci. Sars-cov-2 variants of concern in the united states—challenges and opportunities. *JAMA*, 2021.

[21] Irwin M. Rosenstock. The health belief model and preventive health behavior. *Health Education Monographs*, 2(4):354–386, 1974.

[22] G.M. Hochbaum. *Public Participation in Medical Screening Programs: A Socio-psychological Study.* Public Health Service publication. U.S. Department of Health, Education, and Welfare, Public Health Service, Bureau of State Services, Division of Special Health Services, Tuberculosis Program, 1958.

[23] J. Hayden. *Introduction to Health Behavior Theory.* Jones & Bartlett Learning, 2013.

[24] Ola Andersson, Pol Campos-Mercade, Armando N. Meier, and Erik Wengström. Anticipation of covid-19 vaccines reduces social distancing. WorkingPaper 2020:29, Lund University, Department of Economics, 2020.

[25] Monica L Kasting, Katharine J Head, Jane A Hartsock, Lynne Sturm, and Gregory D Zimet. Public perceptions of the effectiveness of recommended non-pharmaceutical intervention behaviors to mitigate the spread of sars-cov-2. *PloS one*, 15(11):e0241662, 2020.

[26] Toby Wise, Tomislav D Zbozinek, Giorgia Michelini, Cindy C Hagan, and Dean Mobbs. Changes in risk perception and self-reported protective behaviour during the first week of the covid-19 pandemic in the united states. *Royal Society open science*, 7(9):200742, 2020.

[27] Vincenzo Galasso, Vincent Pons, Paola Profeta, Michael Becher, Sylvain Brouard, and Martial Foucault. Gender differences in covid-19 attitudes and behavior: Panel evidence from eight countries. *Proceedings of the National Academy of Sciences*, 117(44):27285–27291, 2020.

[28] Vladimir M. Cvetković, Neda Nikolić, Una Radovanović Nenadić, Adem Öcal, Eric K. Noji, and Miodrag Zečević. Preparedness and preventive behaviors for a pandemic disaster caused by covid-19 in serbia. *IJERPH*, 17(11):4124, jun 2020.

[29] Sarah J Barber and Hyunji Kim. Covid-19 worries and behavior changes in older and younger men and women. *The Journals of Gerontology: Series B*, may 2020.

[30] Daniel Lüdecke and Olaf von dem Knesebeck. Protective behavior in course of the COVID-19 outbreak. survey results from germany. *Front. Public Health*, 8, sep 2020.

[31] Kaori Muto, Isamu Yamamoto, Miwako Nagasu, Mikihito Tanaka, and Koji Wada. Japanese citizens behavioral changes and preparedness against covid-19: An online survey during the early phase of the pandemic. *PLoS ONE*, 15(6):e0234292, jun 2020.

[32] Shamsheer ul Haq, Pomi Shahbaz, and Ismet Boz. Knowledge, behavior and precautionary measures related to covid-19 pandemic among the general public of punjab province, pakistan. *The Journal of Infection in Developing Countries*, 14(08):823–835, 2020.

[33] Austin L. Wright, Konstantin Sonin, Jesse Driscoll, and Jarnickae Wilson. Poverty and economic dislocation reduce compliance with covid-19 shelter-in-place protocols. *Journal of Economic Behavior and Organization*, 180:544–554, dec 2020.

[34] Minha Lee, Jun Zhao, Qianqian Sun, Yixuan Pan, Weiyi Zhou, Chenfeng Xiong, and Lei Zhang. Human mobility trends during the early stage of the covid-19 pandemic in the united states. *PLoS One*, 15(11):e0241468, 2020.

[35] Joakim A. Weill, Matthieu Stigler, Olivier Deschenes, and Michael R. Springborn. Social distancing responses to covid-19 emergency declarations strongly differentiated by income. *Proc Natl Acad Sci USA*, 117(33):19658–19660, jul 2020.

[36] Giovanni Bonaccorsi, Francesco Pierri, Matteo Cinelli, Andrea Flori, Alessandro Galeazzi, Francesco Porcelli, Ana Lucia Schmidt, Carlo Michele Valensise, Antonio Scala, Walter Quattrociocchi, and Fabio Pammolli. Economic and social consequences of human mobility restrictions under covid-19. *Proc Natl Acad Sci USA*, 117(27):15530–15535, jun 2020.

[37] Nicolò Gozzi, Michele Tizzoni, Matteo Chinazzi, Leo Ferres, Alessandro Vespignani, and Nicola Perra. Estimating the effect of social inequalities in the mitigation of covid-19 across communities in santiago de chile. *medRxiv*, oct 2020.

[38] Serina Chang, Emma Pierson, Pang Wei Koh, Jaline Gerardin, Beth Redbird, David Grusky, and Jure Leskovec. Mobility network models of COVID-19 explain inequities and inform reopening.

[39] Laetitia Gauvin, Paolo Bajardi, Emanuele Pepe, Brennan Lake, Filippo Privitera, and Michele Tizzoni. Socioeconomic determinants of mobility responses during the first wave of covid-19 in italy: from provinces to neighbourhoods. *medRxiv*, 2020.

[40] Romain Garnier, Jan R Benetka, John Kraemer, and Shweta Bansal. Socio-economic disparities in social distancing during the covid-19 pandemic in the united states: an observational study. *Journal of medical Internet research*, 2020.

[41] Zindoga Mukandavire, Farai Nyabadza, Noble J Malunguza, Diego F Cuadros, Tinevimbo Shiri, and Godfrey Musuka. Quantifying early covid-19 outbreak transmission in south africa and exploring vaccine efficacy scenarios. *PloS one*, 15(7):e0236003, 2020.

[42] Kate M Bubar, Stephen M Kissler, Marc Lipsitch, Sarah Cobey, Yonatan Grad, and Daniel B Larremore. Model-informed covid-19 vaccine prioritization strategies by age and serostatus. *medRxiv*, 2020.

[43] Laura Matrajt, Julia Eaton, Tiffany Leung, and Elizabeth R Brown. Vaccine optimization for covid-19: who to vaccinate first? *medRxiv*, 2020.

[44] Engineering National Academies of Sciences, Medicine, et al. *Framework for equitable allocation of COVID-19 vaccine*. National Academies Press, 2020.

[45] Jay Love, Lindsay T Keegan, Frederick J Angulo, John McLaughlin, Kimberly M Shea, David L Swerdlow, Matthew H Samore, and Damon JA Toth. Continued need for non-pharmaceutical interventions after covid-19 vaccination in long-term-care facilities. *medRxiv*, 2021.

[46] Abba B Gumel, Enahoro A Iboi, Calistus N Ngonghala, and Gideon A Ngwa. Mathematical assessment of the roles of vaccination and non-pharmaceutical interventions on covid-19 dynamics: a multigroup modeling approach. *medRxiv*, pages 2020–12, 2021.

[47] Marta Galanti, Sen Pei, Teresa K Yamana, Frederick J Angulo, Apostolos Charos, David L Swerdlow, and Jeffrey Shaman. Social distancing remains key during vaccinations. *Science*, 371(6528):473–474, 2021.

[48] Sam Moore, Edward M Hill, Michael J Tildesley, Louise Dyson, and Matt J Keeling. Vaccination and non-pharmaceutical interventions for covid-19: a mathematical modelling study. *The Lancet Infectious Diseases*, 2021.

[49] World economic situation and prospects 2020. `https://www.un.org/development/desa/dpad/wp-content/uploads/sites/45/WESP2020_Annex.pdf`, 2020. Accessed: 2021-03-25.

[50] Patrick GT Walker, Charles Whittaker, Oliver J Watson, Marc Baguelin, Peter Winskill, Arran Hamlet, Bimandra A Djafaara, Zulma Cucunubá, Daniela Olivera Mesa, Will Green, et al. The impact of covid-19 and strategies for mitigation and suppression in low-and middle-income countries. *Science*, 369(6502):413–422, 2020.

[51] Mélodie Monod, Alexandra Blenkinsop, Xiaoyue Xi, Daniel Hebert, Sivan Bershan, Simon Tietze, Marc Baguelin, Valerie C. Bradley, Yu Chen, Helen Coupland, Sarah Filippi, Jonathan Ish-Horowicz, Martin McManus, Thomas Mellan, Axel Gandy, Michael Hutchinson, H. Juliette T. Unwin, Sabine L. van Elsland, Michaela A. C. Vollmer, Sebastian Weber, Harrison Zhu, Anne Bezancon, Neil M. Ferguson, Swapnil Mishra, Seth Flaxman, Samir Bhatt, Oliver Ratmann, and . Age groups that sustain resurging covid-19 epidemics in the united states. *Science*, 371(6536), 2021.

[52] Eli P. Fenichel, Carlos Castillo-Chavez, M. G. Ceddia, Gerardo Chowell, Paula A. Gonzalez Parra, Graham J. Hickling, Garth Holloway, Richard Horan, Benjamin Morin, Charles Perrings, Michael Springborn, Leticia Velazquez, and Cristina Villalobos. Adaptive human behavior in epidemiological models. *Proceedings of the National Academy of Sciences*, 108(15):6306–6311, 2011.

[53] Robert Verity, Lucy Okell, Ilaria Dorigatti, Peter Winskill, Charles Whittaker, Natsuko Imai, Gina Cuomo-Dannenburg, Hayley Thompson, Patrick Walker, Han Fu, Amy Dighe, Jamie Griffin, Marc Baguelin, Sangeeta Bhatia, Adhiratha Boonyasiri, Anne Cori, Zulma M. Cucunubá, Rich FitzJohn, Katy Gaythorpe, and Neil Ferguson. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet Infectious Diseases*, 20, 03 2020.

[54] Coronavirus pandemic (covid-19). `https://ourworldindata.org/coronavirus`, 2020. Accessed: 2020-11-30.

[55] Mikael Sunnåker, Alberto Giovanni Busetto, Elina Numminen, Jukka Corander, Matthieu Foll, and Christophe Dessimoz. Approximate bayesian computation. *PLOS Computational Biology*, 9(1):1–10, 01 2013.

[56] Google LLC "Google COVID-19 Community Mobility Reports". `https://www.google.com/covid19/mobility/`, 2020. Accessed: 2021-08-01.

[57] Oxford COVID-19 Government Response Tracker. `https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker#data`, 2020. Accessed: 2020-11-30.

[58] Neil M Ferguson, Derek AT Cummings, Christophe Fraser, James C Cajka, Philip C Cooley, and Donald S Burke. Strategies for mitigating an influenza pandemic. *Nature*, 442(7101):448–452, 2006.

[59] Nicholas G. Davies, Rosanna C. Barnard, Christopher I. Jarvis, Adam J. Kucharski, James Munday, Carl A. B. Pearson, Timothy W. Russell, Damien C. Tully, Sam Abbott, Amy Gimma, William Waites, Kerry LM Wong, Kevin van Zandvoort, Rosalind M. Eggo, Sebastian Funk, Mark Jit, Katherine E. Atkins, and W. John Edmunds. Estimated transmissibility and severity of novel sars-cov-2 variant of concern 202012/01 in england. *medRxiv*, 2020.

[60] Smriti Mallapaty. Can covid vaccines stop transmission? scientists race to find answers. *Nature*, 2021.

[61] Kaiyuan Sun, Wei Wang, Lidong Gao, Yan Wang, Kaiwei Luo, Lingshuang Ren, Zhifei Zhan, Xinghui Chen, Shanlu Zhao, Yiwei Huang, Qianlai Sun, Ziyan Liu, Maria Litvinova, Alessandro Vespignani, Marco Ajelli, Cécile Viboud, and Hongjie Yu. Transmission heterogeneities, kinetics, and controllability of sars-cov-2. *Science*, page eabe2424, nov 2020.

[62] Huaiyu Tian, Yonghong Liu, Yidan Li, Chieh-Hsi Wu, Bin Chen, Moritz U. G. Kraemer, Bingying Li, Jun Cai, Bo Xu, Qiqi Yang, Ben Wang, Peng Yang, Yujun Cui, Yimeng Song, Pai Zheng, Quanyi Wang, Ottar N. Bjornstad, Ruifu Yang, Bryan T. Grenfell, Oliver G. Pybus, and Christopher Dye. An investigation of transmission control measures during the first 50 days of the covid-19 epidemic in china. *Science*, 368(6491):638–642, mar 2020.

[63] Robert C Reiner, Ryan M Barber, James K Collins, Peng Zheng, Christopher Adolph, James Albright, Catherine M Antony, Aleksandr Y Aravkin, Steven D Bachmeier, Bree Bang-Jensen, et al. Modeling covid-19 scenarios for the united states. *Nat Med*, oct 2020.

[64] Frank Schlosser, Benjamin F. Maier, Olivia Jack, David Hinrichs, Adrian Zachariae, and Dirk Brockmann. Covid-19 lockdown induces disease-mitigating structural changes in mobility networks. *Proc Natl Acad Sci USA*, page 202012326, dec 2020.

[65] Moritz U. G. Kraemer, Chia-Hung Yang, Bernardo Gutierrez, Chieh-Hsi Wu, Brennan Klein, David M. Pigott, Louis du Plessis, Nuno R. Faria, Ruoran Li, William P. Hanage, John S. Brownstein, Maylis Layan, Alessandro Vespignani, Huaiyu Tian, Christopher Dye, Oliver G. Pybus, and Samuel V. Scarpino and. The effect of human mobility and control measures on the covid-19 epidemic in china. *Science*, 368(6490):493–497, mar 2020.

[66] Benjamin F. Maier and Dirk Brockmann. Effective containment explains subexponential growth in recent confirmed covid-19 cases in china. *Science*, 368(6492):742–746, apr 2020.

[67] Laura Di Domenico, Giulia Pullano, Chiara E Sabbatini, Pierre-Yves Boëlle, and Vittoria Colizza. Impact of lockdown on COVID-19 epidemic in Île-de-France and possible exit strategies. *BMC Medicine*, 18(1):240, 2020.

[68] Patrick G. T. Walker, Charles Whittaker, Oliver J Watson, Marc Baguelin, Peter Winskill, Arran Hamlet, Bimandra A. Djafaara, Zulma Cucunubá, Daniela Olivera Mesa, Will Green, Hayley Thompson, Shevanthi Nayagam, Kylie E. C. Ainslie, Sangeeta Bhatia, Samir Bhatt, Adhiratha Boonyasiri, Olivia Boyd, Nicholas F. Brazeau, Lorenzo Cattarino, Gina Cuomo-Dannenburg, Amy Dighe, Christl A. Donnelly, Ilaria Dorigatti, Sabine L. van Elsland, Rich FitzJohn, Han Fu, Katy A.M. Gaythorpe, Lily Geidelberg, Nicholas Grassly, David Haw, Sarah Hayes, Wes Hinsley, Natsuko Imai, David Jorgensen, Edward Knock, Daniel Laydon, Swapnil Mishra, Gemma Nedjati-Gilani, Lucy C. Okell, H. Juliette Unwin, Robert Verity, Michaela Vollmer, Caroline E. Walters, Haowei Wang, Yuanrong Wang, Xiaoyue Xi, David G Lalloo, Neil M. Ferguson, and Azra C. Ghani. The impact of covid-19 and strategies for mitigation and suppression in low- and middle-income countries. *Science*, page eabc0035, jun 2020.

[69] Kiesha Prem, Yang Liu, Timothy W Russell, Adam J Kucharski, Rosalind M Eggo, Nicholas Davies, Mark Jit, Petra Klepac, Stefan Flasche, Samuel Clifford, Carl A B Pearson, James D Munday, Sam Abbott, Hamish Gibbs, Alicia Rosello, Billy J Quilty, Thibaut Jombart, Fiona Sun, Charlie Diamond, Amy Gimma, Kevin van Zandvoort, Sebastian Funk, Christopher I Jarvis, W John Edmunds, Nikos I Bosse, and Joel Hellewell. The effect of control strategies to reduce social mixing on outcomes of the covid-19 epidemic in wuhan, china: a modelling study. *The Lancet Public Health*, 5(5):e261–e270, may 2020.

[70] Christopher I. Jarvis, , Kevin Van Zandvoort, Amy Gimma, Kiesha Prem, Petra Klepac, G. James Rubin, and W. John Edmunds. Quantifying the impact of physical distance measures on the transmission of covid-19 in the uk. *BMC Med*, 18(1), may 2020.

[71] Nicholas G Davies, Adam J Kucharski, Rosalind M Eggo, Amy Gimma, W John Edmunds, Thibaut Jombart, Kathleen O Reilly, Akira Endo, Joel Hellewell, Emily S Nightingale, Billy J Quilty, Christopher I Jarvis, Timothy W Russell, Petra Klepac, Nikos I Bosse, Sebastian Funk, Sam Abbott, Graham F Medley, Hamish Gibbs, Carl A B Pearson, Stefan Flasche, Mark Jit, Samuel Clifford, Kiesha Prem, Charlie Diamond, Jon Emery, Arminder K Deol, Simon R Procter, Kevin van Zandvoort, Yueqian Fiona Sun, James D Munday, Alicia Rosello, Megan Auzenbergs, Gwen Knight, Rein M G J Houben, and Yang Liu. Effects of non-pharmaceutical interventions on covid-19 cases, deaths, and demand for hospital services in the uk: a modelling study. *The Lancet Public Health*, 5(7):e375–e385, jul 2020.

[72] Falko F Sniehotta, Vera Araújo-Soares, Jamie Brown, Michael P Kelly, Susan Michie, and Robert West. Complex systems and individual-level approaches to population health: a false dichotomy? *The Lancet Public Health*, 2(9):e396–e397, 2017.

[73] Jay J Van Bavel, Katherine Baicker, Paulo S Boggio, Valerio Capraro, Aleksandra Cichocka, Mina Cikara, Molly J Crockett, Alia J Crum, Karen M Douglas, James N Druckman, et al. Using social and behavioural science to support covid-19 pandemic response. *Nature Human Behaviour*, pages 1–12, 2020.

[74] Robert West, Susan Michie, G James Rubin, and Richard Amlôt. Applying principles of behaviour change to reduce sars-cov-2 transmission. *Nature Human Behaviour*, pages 1–9, 2020.

[75] Kiesha Prem, Alex R. Cook, and Mark Jit. Projecting social contact matrices in 152 countries using contact surveys and demographic data. *PLOS Computational Biology*, 13(9):1–21, 09 2017.

[76] Xingjie Hao, Shanshan Cheng, Degang Wu, Tangchun Wu, Xihong Lin, and Chaolong Wang. Reconstruction of the full transmission dynamics of covid-19 in wuhan. *Nature*, 584(7821), August 2020.

[77] Merryn Voysey, Sue Ann Costa Clemens, Shabir A Madhi, Lily Y Weckx, Pedro M Folegatti, Parvinder K Aley, Brian Angus, Vicky L Baillie, Shaun L Barnabas, Qasim E Bhorat, et al. Safety and efficacy of the chadox1 ncov-19 vaccine (azd1222) against sars-cov-2: an interim analysis of four randomised controlled trials in brazil, south africa, and the uk. *The Lancet*, 2020.

[78] Pauli Virtanen, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, Pearu Peterson, Warren Weckesser, Jonathan Bright, Stéfan J. van der Walt, Matthew Brett, Joshua Wilson, K. Jarrod Millman, Nikolay Mayorov, Andrew R. J. Nelson, Eric Jones, Robert Kern, Eric Larson, C J Carey, İlhan Polat, Yu Feng, Eric W. Moore, Jake VanderPlas, Denis Laxalde, Josef Perktold, Robert Cimrman, Ian Henriksen, E. A. Quintero, Charles R. Harris, Anne M. Archibald, Antônio H. Ribeiro, Fabian Pedregosa, Paul van Mulbregt, and SciPy 1.0 Contributors. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods*, 17:261–272, 2020.

[79] Charles R. Harris, K. Jarrod Millman, St'efan J. van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J. Smith, Robert Kern, Matti Picus, Stephan Hoyer, Marten H. van Kerkwijk, Matthew Brett, Allan Haldane, Jaime Fern'andez del R'ıo, Mark Wiebe, Pearu Peterson, Pierre G'erard-Marchant, Kevin Sheppard, Tyler Reddy, Warren Weckesser, Hameer Abbasi, Christoph Gohlke, and Travis E. Oliphant. Array programming with NumPy. *Nature*, 585(7825):357–362, September 2020.

[80] Siu Kwan Lam, Antoine Pitrou, and Stanley Seibert. Numba: A llvm-based python jit compiler. In *Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC*, pages 1–6, 2015.

[81] J. D. Hunter. Matplotlib: A 2d graphics environment. *Computing in Science & Engineering*, 9(3):90–95, 2007.

[82] Derek Weycker, John Edelsberg, M. Elizabeth Halloran, Ira M. Longini, Azhar Nizam, Vincent Ciuryla, and Gerry Oster. Population-wide benefits of routine vaccination of children against influenza. *Vaccine*, 23(10):1284–1293, 2005.

[83] Jan Medlock and Alison P. Galvani. Optimizing influenza vaccine distribution. *Science*, 325(5948):1705–1708, 2009.

[84] Amanda Minter and Renata Retkute. Approximate bayesian computation for infectious disease modelling. *Epidemics*, 29:100368, 2019.

[85] Ying Liu, Albert A Gayle, Annelies Wilder-Smith, and Joacim Rocklöv. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *Journal of Travel Medicine*, 27(2), 02 2020. taaa021.

[86] Centers for Disease Control and Prevention, COVID-19 Pandemic Planning Scenarios. `https://www.cdc.gov/coronavirus/2019-ncov/hcp/planning-scenarios.html#table-1`, 2021. Accessed 2021/02/02.

[87] United nations, department of economic and social affairs, population division. world population prospects: The 2019 revision. `https://population.un.org/wpp/Download/Metadata/Documentation/`, 2020. Accessed: 2020-11-30.

[88] Daily number of new reported cases of covid-19 by country worldwide. `https://www.ecdc.europa.eu/en/publications-data/download-todays-data-geographic-distribution-covid-19-cases-worldwide`, 2020. Accessed: 2020-11-30.

[89] Stephen A. Lauer, Kyra H. Grantz, Qifang Bi, Forrest K. Jones, Qulu Zheng, Hannah Meredith, Andrew S. Azman, Nicholas G Reich, and Justin Lessler. The incubation period of coronavirus disease 2019 (covid-19) from publicly reported confirmed cases: Estimation and application. *Annals of Internal Medicine*, 172(9):577–582, 2020. PMID: 32150748.

[90] Luca Ferretti, Chris Wymant, Michelle Kendall, Lele Zhao, Anel Nurtay, Lucie Abeler-Dörner, Michael Parker, David Bonsall, and Christophe Fraser. Quantifying sars-cov-2 transmission suggests epidemic control with digital contact tracing. *Science*, 368(6491), 2020.

[91] Kenji Mizumoto, Katsushi Kagaya, Alexander Zarebski, and Gerardo Chowell. Estimating the asymptomatic proportion of coronavirus disease 2019 (covid-19) cases on board the diamond princess cruise ship, yokohama, japan, 2020. *Eurosurveillance*, 25(10), 2020.

[92] Jantien A Backer, Don Klinkenberg, and Jacco Wallinga. Incubation period of 2019 novel coronavirus (2019-ncov) infections among travellers from wuhan, china, 20–28 january 2020. *Eurosurveillance*, 25(5), 2020.

[93] Stephen M. Kissler, Christine Tedijanto, Edward Goldstein, Yonatan H. Grad, and Marc Lipsitch. Projecting the transmission dynamics of sars-cov-2 through the postpandemic period. *Science*, 368(6493):860–868, 2020.

[94] Nicolò Gozzi, Matteo Chinazzi, Jessica T Davis, Kunpeng Mu, Ana Pastore y Piontti, Marco Ajelli, Nicola Perra, and Alessandro Vespignani. Estimating the spreading and dominance of sars-cov-2 voc 202012/01 (lineage b. 1.1. 7) across europe. *medRxiv*, 2021.