Proteomic and Metabolomic Investigation of COVID-19 Patients with Elevated Serum Lactate Dehydrogenase

Haixi Yan[1,#], Xiao Liang[2,3,4#], Juping Du[1,#], Zebao He [1,#], Yu Wang[5], Mengge Lyu[2,3,4], Liang Yue[2,3,4], Fangfei Zhang [2,3,4], Zhangzhi Xue[2,3,4], Luang Xu[2,3,4], Guan Ruan[6], Jun Li[1], Hongguo Zhu [1], Jiaqin Xu [1], Shiyong Chen[1], Chao Zhang[1], Dongqing Lv[1], Zongmei Lin[1], Bo Shen[1], Yi Zhu[2,3,4†], Biyun Qian[5,†], Haixiao Chen[1,†], Tiannan Guo[2,3,4†]

[1] Taizhou Hospital of Zhejiang Province affiliated to Wenzhou Medical University, 150 Ximen Street, Linhai 317000, Zhejiang Province, China

[2] Key Laboratory of Structural Biology of Zhejiang Province, School of Life Sciences, Westlake University, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China

[3] Center for Infectious Disease Research, Westlake Laboratory of Life Sciences and Biomedicine, Hangzhou 310024, China

[4] Institute of Basic Medical Sciences, Westlake Institute for Advanced Study, 18 Shilongshan Road, Hangzhou 310024, Zhejiang Province, China

[5] Hongqiao International Institute of Medicine, Shanghai Tongren Hospital and Faculty of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

[6] Westlake Omics (Hangzhou) Biotechnology Co., Ltd. No.1, Yunmeng Road, Cloud Town, Xihu District, Hangzhou, Zhejiang Province, China

[#] These authors contributed equally to this study.

[†] Correspondence and requests for materials should be addressed to Z.Y. (zhuyi@westlake.edu.cn), B.Q. (qianbiyun@sjtu.edu.cn), H.C. (chenhx@enzemed.com), and T.G. (guotiannan@westlake.edu.cn).

Keywords: COVID-19; lactate dehydrogenase; proteomics; metabolomics; biomarker; prognosis

Abstract

Serum lactate dehydrogenase (LDH) has been established as a prognostic indicator given its differential expression in COVID-19 patients. However, the molecular mechanisms underneath remain poorly understood. In this study, 144 COVID-19 patients were enrolled to monitor the clinical and laboratory parameters over three weeks. Serum lactate dehydrogenase (LDH) was shown elevated in the COVID-19 patients on admission and declined throughout disease course, and its ability to classify patient severity outperformed other biochemical indicators. A threshold of 247 U/L serum LDH on admission was determined for severity prognosis. Next, we classified a subset of 14 patients into high- and low-risk groups based on serum LDH expression and compared their quantitative serum proteomic and metabolomic differences. The results found COVID-19 patients with high serum LDH exhibited differentially expressed blood coagulation and immune responses including acute inflammatory responses, platelet degranulation, complement cascade, as well as multiple different metabolic responses including lipid metabolism, protein ubiquitination and pyruvate fermentation. Specifically, activation of hypoxia responses was highlighted in patients with high LDH expressions. Taken together, our data showed that serum LDH levels are associated COVID-19 severity, and that elevated serum LDH might be consequences of hypoxia and tissue injuries induced by inflammation.

INTRODUCTION

COVID-19 is an ongoing global pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). A high viral transmission rate and the lack of effective therapy contributed to more than 83 million infected cases as of time January 3rd, 2021 [1].

To better diagnose COVID-19 and monitor the disease progress, multiple molecules have been proposed as prognostic indicators [2]. Lactate dehydrogenase (LDH) is an intracellular enzyme, catalyzing pyruvate fermentation and facilitating glycolysis. LDH is released into the blood after cell death and has been reported to increase in a variety of diseases including Severe acute respiratory syndrome (SARS) [3], diabetes [4], and cancers [5]. Serum LDH levels in COVID-19 patients are over-expressed [2], especially in severe and critical patients [6-9]. They decrease throughout disease course [7,10], in correlation with viral mRNA clearance [7]. Related studies have shown that serum LDH is well correlated with respiratory failure [11], lung injury, disease severity [12] and mortality [13] in COVID-19 patients.

However, the molecular mechanisms underlying the LDH's association with the COVID-19 disease progression remains poorly understood. Most studies attribute serum LDH elevation to its release from somatic tissue and organ damage caused by either viral attack [12] or inflammation [10]. These clinical assumptions lack molecular evidence, potentially leading to biased assessments. Moreover, these explanations failed to consider the metabolic role of LDH to balance excess lactate during hypoxia. Here we have systematically explored the proteome and metabolome of sera from COVID-19 patients with low and high serum LDH, and identified the specific host responses, which shed light on the pathogenesis and convalesce of COVID-19.

MATERIALS AND METHODS

Patient Information

We collected and curated the electronic medical records of patient information in Taizhou Hospital of Zhejiang Province, between January 17 and February 20, 2020. 212 patients met the criteria of suspected COVID-19, of which 145 patients were confirmed as COVID-19, based on the Government's Diagnosis and Treatment Guideline (5th version) [14].144 patients COVID-19 patients were included in this study after excluding one patient who had incomplete laboratory data. According to the admission period, we grouped them into two cohorts. Patients admitted to Taizhou Public Health Medical Center, Taizhou Hospital between January 17 and February 4, 2020 were included in the cohort 1 (n = 115). The cohort 2 (n = 29) contained COVID-19 patients admitted from February 4 to February 20, 2020. The end of the follow-up date was March 1, 2020. Disease severity was accessed according to the abovementioned guideline. We classified COVID-19 patients into two groups (severe and non-severe): the severe group included severe and critical patients, and the non-severe group included mild and typical patients. Briefly, those who had shortness of breath with respiratory rate $\geq$ 30 breaths/min, a ratio of arterial blood oxygen partial pressure to oxygen concentration $\leq$ 300 mmHg or saturation of oxygen $\leq$ 93% when resting were defined as severe patients. The other COVID-19 patients were grouped as non-severe patients. 125 healthy individuals were enrolled as controls. This study was conducted in accordance with the Helsinki Declaration and was approved by the institutional medical ethics review boards of Taizhou Hospital of Zhejiang Province and Westlake University (Approval ID: 20210119GTN001).

Laboratory tests

Samples were taken throughout disease course from patient admission to discharge. More details are described in Supplementary Table 1. For laboratory tests, 404 serial blood samples from 144 patients were collected and centrifuged at 1500 g for 10 min at room temperature.

For serum of COVID-19 patients, seven biochemical indicators were tested, namely lactate dehydrogenase (LDH), aspartate aminotransferase (AST), alanine aminotransferase (ALT), total bilirubin, total protein, creatinine, and creatine kinase (CK), with a Chemistry Analyzer (Beckman Coulter, AU5821). The serum LDH was measured for healthy controls.

For LDH isoform analyses, 32 serum samples from 32 COVID-19 patients (11 non-severe and 21 severe) were sent for electrophoresis with agarose gel electrophoretic analyzer (SEBIA, HYDRASYS 2). ISO-LDH substrate and blocking buffer were afterwards added to incubate for 20 min, respectively. The gel was scanned with the same analyzer.

Proteomic and metabolomic data set

The proteomic and metabolomic data were extracted from our previous publication [14]. Briefly, serum samples from COVID-19 patients were kept at 56°C for 30 min to inactivate potential SARS-CoV-2. For proteomic experiments, inactivated serum samples were processed into peptides, labeled with TMTpro 16plex chemical tags, fractionated to 40 aliquots, and analyzed by LC-MS/MS. The proteomics data were analyzed with Proteome Discoverer (Version 2.4.1.15, Thermo Fisher Scientific), with the default parameter and a protein database composed of the Homo sapiens fasta database (07 Jan 2020, UniProtKB), containing 20,412 reviewed protein sequences, and the SARS-CoV-2 virus fasta (version NC_045512.2, NCBI). Targeted false discovery rate (FDR) for peptide-spectrum match was set to 1% (strict) and 5% (relaxed). Normalization was performed against the total peptide intensity. 894 proteins were quantified altogether. For metabolomic experiments, inactivated serum samples were processed to collect metabolites, and divided into 4 fractions for 4 different modes of LC-MS/MS data acquisition, leading to characterization of 941 metabolites. The median coefficient of variance (CV) for proteomic and metabolomic data were 10% and 5%, respectively, determined by pooled control samples in each batch, as described previously [14]. The proteomics and metabolomics data could be referenced from the previously publication in ProteomeXchange Consortium (https://www.iprox.org/). Project ID: IPX0002106000 and IPX0002171000.

Statistical analysis

Statistical clinical data analyses were performed using SPSS software (version 22.0). Continuous variables were presented as median and interquartile range (IQR) values, while categorical variables were shown as frequency and percentage. An independent t-test was used for continuous variables when the data were normally distributed; otherwise, the Mann-Whitney test was used. Chi-square test or Fisher's exact test was used for categorical variables. Receiver operating characteristic (ROC) analysis was used for the selection of the best intercept point. Prediction of disease progression was obtained using the Cox proportional hazards model. Statistical proteomic data analysis was performed using R (version 3.6.3). Missing values in the proteomic data matrix were assigned as 0.01 unless otherwise mentioned. P values $\leq$ 0.05 were considered statistically significant unless otherwise mentioned. Differential protein expression was based on the cutoff: P values $\leq$ 0.05, $|\log_2 FC| > 0.25$. Plotting

150   was performed with R (version 3.6.3).
151

RESULTS AND DISCUSSION

Demographic, clinical, and laboratory characteristics
A total of 144 COVID-19 patients were enrolled in the study. Detailed
demographic, clinical, and laboratory characteristics of these patients on admission
were provided in Table 1. The median age was 47 years old, and 53.5% of them were
male. The severe patients account for 25% (36/144) of the group, were 10 years older
than the non-severe patients (55 vs. 45, $p < 0.001$), and were more likely to have fever
symptoms on admission ($p = 0.001$). Severe patients received higher percentages of
treatment in oxygen inhalation ($p < 0.001$), antibiotics ($p = 0.024$), glucocorticoid ($p <$
0.001), and intravenous gamma immunoglobulin ($p < 0.001$) than the non-severe
patients. These medications may have an impact on the patient blood metabolism and
may alter the COVID-19 microenvironment, which unfortunately could not be
rigorously examined in the current study due to the small sample size, awaiting future
investigations. The patients did not exhibit significant difference between severe and
non-severe groups in terms of gender unmentioned symptoms on admission, nor other
medical treatments ($p > 0.05$) as listed in Table 1.
Based on laboratory test results, a higher percentage of severe patients have
elevated levels of serum LDH that are above the upper limit of normal (ULN) value
than non-severe patients (58.3% vs. 7.4%, $p < 0.001$). Likewise, more severe patients
also showed higher level of alanine aminotransferase (ALT, $p = 0.023$), aspartate
aminotransferase (AST, $p < 0.001$), urea ($p = 0.004$), creatinine ($p = 0.029$), and
creatine kinase (CK, $p < 0.001$). We then compared their temporal changes at 7-day
intervals (Supplementary Table 1). LDH, CK, and creatinine showed continuous
decrease in the sera of severe patients, while only serum LDH showed a continuous
decrease in non-severe patients.
A Cox regression model was applied to evaluate the prognostic value of these
indicators (Supplementary Table 2). In the Cohort 1, LDH is the only indicator with
discrimination ability ($p < 0.001$) in the multivariate analysis, although LDH, AST,
and CK were determined in the univariate analysis with statistical significance ($p <$
0.001). We further validated the value of serum LDH in Cohort 2 and the data showed
significant discrimination ($p = 0.032$), suggesting that serum LDH could be a useful
prognostic biomarker.
184

Temporal characteristics of Serum LDH in COVID-19 patients
We then assessed the effect of age and sex on the temporal serum LDH expression.
COVID-19 patients older than 60 years showed a higher serum LDH expression
throughout the disease course than that in younger patients (Figure 1A), probably due
to ageing and underlying diseases. Males expressed higher serum LDH levels than
that from females only during the initial hospitalization stage (within the first week)
(Figure 1B). Serum LDH levels became comparable between COVID-19 patients and
healthy controls (median values: 177 U/L above 60 years old and 152 U/L below 60
years old; 154 U/L for males and 158.5 U/L for females) after three weeks.
We compared the median values of serum LDH levels from the initial admission
stage (1-3 days after admission) and discharge stage (1-3 days before discharge), in a
sub-cohort of recovered patients that were discharged by the end of follow-up. (total:
n=49; severe: n=13; non-severe: n=36) (Figure 1C). Serum LDH levels were
significantly higher in the severe patient group, particularly on admission ($p < 0.01$).

We analyzed serum LDH isoform expression from 32 patients upon admission (severe: n=22; non-severe: n=11) (Figure 1D) and found that LDH-2, LDH-4 and LDH-5 were significantly higher in the sera of severe COVID-19 patients. LDH-1 in contrast was not dysregulated. LDH-4 and LDH-5 mainly contribute to pyruvate fermentation under hypoxic conditions, while LDH-1 favors the opposite direction of the reversible reaction [15]. These data suggest anaerobic glycolysis metabolism in severe patients.

To better understand the temporal dynamics of serum LDH expression over the disease course, we next monitored the serum LDH level from day 1 on admission till day 21 at a 3-day interval (Figure 1E). As to the non-severe patient group, the serum LDH level was slightly higher on admission while declining slowly over the hospitalization period. The serum LDH levels in the severe group comparatively were significantly higher upon admission, with a prominent variance range. They dropped significantly from 3rd to 9th day as the patients were taking medical care and by the 21st day fell below the initial serum LDH levels in the non-severe group.

Classification of low- and high-risk patients based on serum LDH levels

To establish a cohort-specific serum LDH expression threshold as a risk indicator, we took patient severity (severe vs. non-severe) as the dichotomous variable and conducted the receiver operating characteristic (ROC) analysis on the serum LDH expression levels on admission. The area under curve (AUC) was 0.864 (Figure 1F), confirming the discriminative power of serum LDH. The serum LDH level corresponding to the maximum Youden index was determined as 247 U/L, within the threshold to determine serum LDH abnormality from past reports (240-253.2 U/L) [13].

Fourteen characteristic patients within the cohort were selected and divided into two groups for closer inspection (Figure 1H). The low-risk (LR) group consists of seven non-severe patients with serum LDH expressions below 247 U/L since admission except for one exceptional detection (LR4, 5th day, 306 U/L). The high-risk (HR) group are composed of six severe patients with on admission serum LDH levels above 247 U/L, and one severe patient with serum LDH levels below the threshold throughout hospitalization (HR1). We attribute this to the relatively late first sampling timepoint (8th day), given that serum LDH levels from all the patients started at a relatively high level and declined over time. Patient HR3 and patient HR7 in particular had exceptionally high serum LDH levels (> 450 U/L) upon admission but dropped dramatically within 10 days. We inspected the detailed medical records of the 14 patients (Table 2). COVID-19 patients with comorbidities including hypertension, chronic HBV infection, and diabetes tend to be severe patients in the HR group, in consistent with the literature [16,17].

Quantitative proteomics and metabolomics uncover dysregulated molecules associated with elevated serum LDH

The serum proteomic and metabolomic datasets of the HR and LR group patients were extracted from a collateral project [14]. 78.6% (11/14) of the patient sera were sampled during the first week on admission (Table 2), herein representing the stage when serum LDH levels exhibited sharp difference between the LR and HR groups. For the proteomic dataset, the Student's *t*-test highlighted 34 proteins as differentially expressed ($p < 0.05$) between HR and LR groups (Figure 2A, upper panel), 26 of which were up-regulated. Pathway enrichment analysis using Metascape [18] showed these proteins conduct three major immune-related activities including acute inflammatory responses (GO:0002526, $p < 0.001$), platelet degranulation

(GO:0002576, $p < 0.001$) and regulation of complement cascade (R-HSA-977606, $p <$
0.001) (Figure 2A and Supplementary Fig. 2A). Additional analyses using Ingenuity
Pathway Analysis (IPA) nominated acute phase response signaling as the most
activated immune-related pathway (Supplementary Fig. 2C) in HR patients. These
findings were in consistence with the prominent immune behaviors (Supplementary
Fig. 1B) as we have previously reported in COVID-19 patients with different severity
[14]. Moreover, blood coagulation (GO: 007596, $p < 0.001$) was significantly enriched
(Supplementary Fig. 2A and 2C). This pathway has been reported to be altered in
COVID-19, and associated with interleukin-6 (IL-6))[19]. Our data showed
upregulation of acute phase proteins (SAA1, ORM1, AGT, and SERPINA3),
complement subunits (C9, C6, and CFI), and LDH subtypes (LDHA and LDHB) in
the HR group (Figure 2B). Thirteen of the differentiated proteins were mapped into a
network wherein key regulators were focused (Figure 2C). Within them, pro-
inflammatory cytokine IL-6 has been widely recognized as a risk factor for COVID-
19 [19-23] and clinically observed to be positively correlated with serum LDH levels
[24]. IL-6 can activate TP53, which facilitates cell apoptosis and could enhance LDHA
expression in blood. Multiple COVID-19 studies involving IL-6 agree with our
profiling [19,25]. This network also includes CCAAT/enhancer-binding protein beta
(CEBPB) which mediates immune and inflammatory responses [26], and sterol
regulatory element binding transcription factor 1 (SREBF1) which regulates lipid
metabolisms that has been reported dysregulated in severe COVID-19 patients [14].
Taken together, the proteomic difference between LR and HR patients reflected
different host responses between the non-severe and severe COVID-19 patients.
Of the 34 differentially expressed metabolites listed in Supplementary Fig. 3A,
88.2% (30/34) were upregulated, and 52.9% (18/34) were lipids. Within them, 7-
hydroxycholesterol (beta) mediates oxidative stress and induces cell apoptosis. It was
elevated during hepatitis B virus (HBV) and hepatitis C virus (HCV) infections [27]
(Fig 2A and Fig 2B, lower panel). Further calibration and absolute quantification of
these lipids would enable in-depth characterization of lipids and their variants. The
protein-metabolite joint network generated using IPA (Supplementary Fig. 3B)
proposed two upstream molecules to regulate LDHA expression, including hypoxia-
inducible factor 1 (HIF1) that mediates hypoxic and inflammatory microenvironments
[28] and fibronectin 1 (FN1) that involves COVID-19 lung fibrosis [29]. The
dysregulated metabolites further consolidate disturbed host responses in association
with serum LDH increase uncovered by the proteomic data.
284
Protein and metabolite change in patients with exceptionally high serum LDH
Next, we narrowed our focus to the two patients with exceptionally high serum
LDH levels on admission (HR3 and HR7, HR outliers), and compared their proteomic
patterns with the other HR patients (Figure 3A). 38 proteins including LDHA and
LDHB were differentially expressed ($p < 0.05$, Figure 3A). Metascape pathway
enrichment nominated 23 proteins associated with immune system process
(GO:0002376, $p < 0.001$) and 21 proteins associated with metabolic process
(GO:0008152, $p < 0.001$) (Supplementary Fig 4B). The top enriched immune-related
pathways are the activation of immune response (GO:0002253, $p < 0.001$) and
humoral immunity response (GO:0006959, $p < 0.001$), while the top enriched
metabolic pathway relates to cofactors (R-HSA-89 78934, $p < 0.001$) (Supplementary
Fig. 4A). IPA analysis detailed the top metabolic functions as protein ubiquitination
pathway and pyruvate fermentation to lactate (Supplementary Fig 4C). Especially, the
HIF1α signaling pathway showed a drastic activation in the HR (outliers) group. As

for characteristic proteins in HR (outliers), our data uncovered a dysregulated protein group which includes five up-regulated proteasome subunits, namely PSMA3, PSMA4, PSMA5, PSMB1, and PSMB3 (Figure 3B), possibly due to cell apoptosis [30] from organ/tissue damage. They are the mediators of protein ubiquitination and associate with NF-κB signaling (GO:0038061, $p < 0.001$). Six of the seven differentially expressed immunoglobulin residues (IGHV3-43, IGHV3-30-5, IGKV1-5, IGLV1-36, IGKV1-17, and IGKV2-24) (Supplementary Fig. 5A) were up-regulated in HR (outliers), suggesting that humoral immunity at the point of detection was suppressed or not activated. For the other up-regulated proteins in HR (outliers) (Figure 3B), CES1 is a hepatic protein and its release in blood suggests liver injuries. Protein disulfide-isomerase (P4HB) was reported to up-rise in response to hypoxia [31]. GAPDH could enhance HIF activity [32] via NF-κB induction activated in hypoxia [33], which contributes to heat shock protein 90-alpha (HSP90AA1) upregulation [34] to form protein complexes with Hif1α. Dipeptidyl peptidase 4 (DPP4), also known as CD26, has been nominated as a potential critical marker in infection susceptibility [35], and its inhibition has been proposed to reduce COVID-19 patient severity [36]. DPP4 is also a downstream factor to mark HIF pathway induction [37]. Especially, the down-regulated protein in HR (outliers) includes CPB2 as a basic carboxypeptidase that suppresses complement system-mediated inflammation [38]. Its deficiency could lead to accelerated acute lung injuries [39]. 84.6% (11/13) of the dysregulated metabolites were lipids (Supplementary Fig. 5B), suggesting disturbed lipid metabolism accompanied with serum LDH changes. The network analysis (Supplementary Fig. 5C) further proposed LDH elevation to be associated with HSP90AA1 and proteasomes.

Serum LDH elevation might be driven by tissue injuries and hypoxia

Taking together all the perturbed molecules as highlighted above (Figure 2B and 3B), we propose a putative working model for the serum LDH elevation in COVID-19 patients (Figure 3C). On the one hand, the inflammation processes triggered by the host immune system induce apoptosis of the infected cells, leading to the release of intracellular LDH into the blood. In high-risk cases, these immune activities result in over-reactive inflammation processes (like "cytokine storm") [40], thereby releasing higher levels of serum LDH from multiple organs/tissues [10]. On the other hand, oxygen homeostasis was disturbed in severe COVID-19 patients [12]. Hypoxia reactions occur to accumulate lactate via glycolysis. LDH can balance lactate secretion via pyruvate fermentation and a series of metabolic regulation (Supplementary Fig. 4A and 4C) to maintain cellular homeostasis [41]. The activated NF-κB and HIF pathways in hypoxia conditions could also induce inflammatory responses [42].

CONCLUDING REMARKS

Here we systematically investigated serum LDH elevation in COVID-19 patients. We thoroughly inspected both clinical and molecular profiles of 144 COVID-19 patients. We confirmed serum LDH as the best independent risk indicator, and further optimized a threshold of 247 U/L for stratifying COVID-19 patients based on the serum LDH level. Our data showed that the serum LDH declined thereafter, patients with serum LDH levels higher than the threshold on admission are prone to severe conditions, hence determined as high-risk (HR) patients, and those lower than the threshold as low-risk (LR) patients.

Proteomic differences between LR and HR groups exposed a list of dysregulated host responses. Among them, acute inflammatory responses, platelet degranulation and complement cascade have been reported in previous studies comparing severe and non-severe COVID-19 patients [14,43]. Blood coagulation has been highlighted in another report that compares COVID-19 patients with high and low IL-6 levels [44]. Immune behaviors including activation of immune response and humoral immune response were further enriched during the intra-comparison within HR patients, suggesting that the immune behaviors are closely related to serum LDH expression. Proteomic profiling also highlighted a list of hypoxia related proteins and functions, including P4HB, DPP4, GAPDH, HSP90AA1, NF-κB and HIF signaling, suggesting that hypoxia might have contributed to elevated LDH. The metabolomic profiling complements findings on the proteomic level and further emphasizes dysregulated lipid metabolism. Taken together, we propose that elevation of serum LDH might attribute to inflammation-related tissue injuries and hypoxia-related metabolism.

This study is limited by several factors. Firstly, this is a single-center study with a relatively small patient cohort due to difficulties of sample collection, therefore subject to experimental bias. Also, the standard inactivation procedures for COVID-19 serum to minimize the risk of infection may have some impact on the characterized samples. Moreover, proteome data from only 14 individuals were acquired, from which only 2 patients were determined as high-risk group outliers for comparative analysis. We could not obtain the samples from different patients at the identical time points. And due to the small sample size, multiple testing wasn't performed for molecular analyses, therefore the statistical power from the proteomic data has to be interpreted with caution. It is worth noting that serum LDH elevation is not specific to COVID-19 disease [45]. Further studies should conduct clinical validation on larger cohorts, and compare the molecular differences including control patients with other diseases with similar symptoms. Targeted approaches would also be required to validate our findings for diagnostic purposes.

CONFLICT OF INTERESTS

The research group of T.G. is partly supported by Tencent, Thermo Fisher Scientific, SCIEX, and Pressure Biosciences Inc. T.G. is a shareholder of Westlake Omics Inc. G.R. is an employee of Westlake Omics Inc.

ACKNOWLEDGMENTS

This work was supported by grants from National Key R&D Program of China (No. 2020YFE0202200), National Natural Science Foundation of China (81972492, 21904107, 81672086, 82072333), Zhejiang Provincial Natural Science Foundation for Distinguished Young Scholars (LR19C050001), Hangzhou Agriculture and Society Advancement Program (20190101A04), Zhejiang Provincial Natural Science Foundation of China (LQ19H100001), Zhejiang Medical and Health Science and Technology Plan (2021KY394), Westlake Education Foundation, and Tencent Foundation. We thank Westlake University Supercomputer Center for assistance in data storage and computation.

References

[1] Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., . . . Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet (London, England), 395*(10223), 497-506. doi: 10.1016/S0140-6736(20)30183-5

[2] Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., . . . Zhang, L. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet (London, England), 395*(10223), 507-513. doi: 10.1016/S0140-6736(20)30211-7

[3] Hui, D. S.-C., Wong, P.-C., & Wang, C. (2003). SARS: clinical features and diagnosis. *Respirology (Carlton, Vic.), 8 Suppl*(Suppl 1), S20-S24. doi: 10.1046/j.1440-1843.2003.00520.x

[4] Huang, E.-J., Kuo, W.-W., Chen, Y.-J., Chen, T.-H., Chang, M.-H., Lu, M.-C., . . . Lee, S.-D. (2006). Homocysteine and other biochemical parameters in Type 2 diabetes mellitus with different diabetic duration or diabetic retinopathy. *Clinica Chimica Acta, 366*(1), 293-298. doi: https://doi.org/10.1016/j.cca.2005.10.025

[5] Ding, J., Karp, J. E., & Emadi, A. (2017). Elevated lactate dehydrogenase (LDH) can be a marker of immune suppression in cancer: Interplay between hematologic and solid neoplastic clones and their microenvironments. *Cancer Biomark, 19*(4), 353-363. doi: 10.3233/cbm-160336

[6] Wan, S., Xiang, Y., Fang, W., Zheng, Y., Li, B., Hu, Y., . . . Yang, R. (2020). Clinical features and treatment of COVID-19 patients in northeast Chongqing. *Journal of medical virology, 92*(7), 797-806. doi: 10.1002/jmv.25783

[7] Yuan, J., Zou, R., Zeng, L., Kou, S., Lan, J., Li, X., . . . Wang, Z. (2020). The correlation between viral clearance and biochemical outcomes of 94 COVID-19 infected discharged patients. *Inflammation research : official journal of the European Histamine Research Society ... [et al.], 69*(6), 599-606. doi: 10.1007/s00011-020-01342-0

[8] Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., . . . Huang, C.-L. (2020). Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *bioRxiv*.

[9] Chen, G., Wu, D., Guo, W., Cao, Y., Huang, D., Wang, H., . . . Ning, Q. (2020). Clinical and immunological features of severe and moderate coronavirus disease 2019. *The Journal of clinical investigation, 130*(5), 2620-2629. doi: 10.1172/JCI137244

[10] Shi, J., Li, Y., Zhou, X., Zhang, Q., Ye, X., Wu, Z., . . . Zhang, W. (2020). Lactate dehydrogenase and susceptibility to deterioration of mild COVID-19 patients: a multicenter nested case-control study. *BMC medicine, 18*(1), 168-168. doi: 10.1186/s12916-020-01633-7

[11] Poggiali, E., Zaino, D., Immovilli, P., Rovero, L., Losi, G., Dacrema, A., . . . Terracciano, C. (2020). Lactate dehydrogenase and C-reactive protein as predictors of respiratory failure in CoVID-19 patients. *Clinica chimica acta; international journal of clinical chemistry, 509*, 135-138. doi: 10.1016/j.cca.2020.06.012

[12] Han, Y., Zhang, H., Mu, S., Wei, W., Jin, C., Tong, C., . . . Gu, G. (2020). Lactate dehydrogenase, an independent risk factor of severe COVID-19 patients: a retrospective and observational study. *Aging, 12*(12), 11245-11258. doi: 10.18632/aging.103372

[13] Henry, B. M., Aggarwal, G., Wong, J., Benoit, S., Vikse, J., Plebani, M., & Lippi, G. (2020). Lactate dehydrogenase levels predict coronavirus disease 2019 (COVID-19) severity and mortality: A pooled analysis. *The American Journal of Emergency Medicine, 38*(9), 1722-1726. doi: 10.1016/j.ajem.2020.05.073

[14] Shen, B., Yi, X., Sun, Y., Bi, X., Du, J., Zhang, C., . . . Guo, T. (2020). Proteomic and Metabolomic Characterization of COVID-19 Patient Sera. *Cell, 182*(1), 59-72.e15. doi: 10.1016/j.cell.2020.05.032

[15] Augoff, K., Hryniewicz-Jankowska, A., & Tabola, R. (2015). Lactate dehydrogenase 5: An old friend and a new hope in the war on cancer. *Cancer Letters, 358*(1), 1-7. doi: https://doi.org/10.1016/j.canlet.2014.12.035

[16] Zheng, Z., Peng, F., Xu, B., Zhao, J., Liu, H., Peng, J., . . . Tang, W. (2020). Risk factors of critical
& mortal COVID-19 cases: A systematic literature review and meta-analysis. *J Infect, 81*(2),
e16-e25. doi: 10.1016/j.jinf.2020.04.021
[17] Zou, X., Fang, M., Li, S., Wu, L., Gao, B., Gao, H., . . . Huang, J. (2020). Characteristics of Liver
Function in Patients With SARS-CoV-2 and Chronic HBV Coinfection. *Clin Gastroenterol*
Hepatol. doi: 10.1016/j.cgh.2020.06.017
[18] Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., . . . Chanda, S.
K. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-
level datasets. *Nat Commun, 10*(1), 1523. doi: 10.1038/s41467-019-09234-6
[19] D'Alessandro, A., Thomas, T., Dzieciatkowska, M., Hill, R. C., Francis, R. O., Hudson, K. E., . . .
Hansen, K. C. (2020). Serum Proteomics in COVID-19 Patients: Altered Coagulation and
Complement Status as a Function of IL-6 Level. *Journal of Proteome Research, 19*(11),
4417-4427. doi: 10.1021/acs.jproteome.0c00365
[20] Tian, J., Yuan, X., Xiao, J., Zhong, Q., Yang, C., Liu, B., . . . Wang, Z. (2020). Clinical characteristics
and risk factors associated with COVID-19 disease severity in patients with cancer in
Wuhan, China: a multicentre, retrospective, cohort study. *The Lancet. Oncology, 21*(7),
893-903. doi: 10.1016/S1470-2045(20)30309-0
[21] Liu, F., Li, L., Xu, M., Wu, J., Luo, D., Zhu, Y., . . . Zhou, X. (2020). Prognostic value of interleukin-
6, C-reactive protein, and procalcitonin in patients with COVID-19. *Journal of clinical*
virology : the official publication of the Pan American Society for Clinical Virology, 127,
104370-104370. doi: 10.1016/j.jcv.2020.104370
[22] Santa Cruz, A., Mendes-Frias, A., Oliveira, A. I., Dias, L., Matos, A. R., Carvalho, A., . . . Silvestre,
R. (2021). Interleukin-6 Is a Biomarker for the Development of Fatal Severe Acute
Respiratory Syndrome Coronavirus 2 Pneumonia. *Frontiers in Immunology, 12*, 613422.
doi: 10.3389/fimmu.2021.613422
[23] Aziz, M., Fatima, R., & Assaly, R. (2020). Elevated interleukin-6 and severe COVID-19: A meta-
analysis. *Journal of medical virology, 92*(11), 2283-2285. doi: 10.1002/jmv.25948
[24] Liu, T., Zhang, J., Yang, Y., Ma, H., Li, Z., Zhang, J., . . . Yi, J. (2020). The role of interleukin-6 in
monitoring severe case of coronavirus disease 2019. *EMBO molecular medicine, 12*(7),
e12421-e12421. doi: 10.15252/emmm.202012421
[25] Hadjadj, J., Yatim, N., Barnabei, L., Corneau, A., Boussier, J., Smith, N., . . . Terrier, B. (2020).
Impaired type I interferon activity and inflammatory responses in severe COVID-19
patients. *Science, 369*(6504), 718-724. doi: 10.1126/science.abc6027
[26] Kinoshita, S., Akira, S., & Kishimoto, T. (1992). A member of the C/EBP family, NF-IL6 beta,
forms a heterodimer and transcriptionally synergizes with NF-IL6. *Proc Natl Acad Sci U S*
A, 89(4), 1473-1476. doi: 10.1073/pnas.89.4.1473
[27] Yoshida, Y., Kodai, S., Takemura, S., Minamiyama, Y., & Niki, E. (2008). Simultaneous
measurement of F2-isoprostane, hydroxyoctadecadienoic acid, hydroxyeicosatetraenoic
acid, and hydroxycholesterols from physiological samples. *Anal Biochem, 379*(1), 105-115.
doi: 10.1016/j.ab.2008.04.028
[28] Jahani, M., Dokaneheifard, S., & Mansouri, K. (2020). Hypoxia: A key feature of COVID-19
launching activation of HIF-1 and cytokine storm. *Journal of inflammation (London,*
England), 17, 33-33. doi: 10.1186/s12950-020-00263-3
[29] Xu, J., Xu, X., Jiang, L., Dua, K., Hansbro, P. M., & Liu, G. (2020). SARS-CoV-2 induces
transcriptional signatures in human lung epithelial cells that promote lung fibrosis. *Respir*
Res, 21(1), 182. doi: 10.1186/s12931-020-01445-6
[30] Weathington, N. M., & Mallampalli, R. K. (2013). New insights on the function of SCF ubiquitin
E3 ligases in the lung. *Cell Signal, 25*(9), 1792-1798. doi: 10.1016/j.cellsig.2013.05.003
[31] Ko, H. S., Uehara, T., & Nomura, Y. (2002). Role of ubiquilin associated with protein-disulfide
isomerase in the endoplasmic reticulum in stress-induced apoptotic cell death. *J Biol*
Chem, 277(38), 35386-35392. doi: 10.1074/jbc.M203412200

[32] Chiche, J., Pommier, S., Beneteau, M., Mondragon, L., Meynet, O., Zunino, B., . . . Ricci, J. E. (2015). GAPDH enhances the aggressiveness and the vascularization of non-Hodgkin's B lymphomas via NF-kappaB-dependent induction of HIF-1alpha. *Leukemia, 29*(5), 1163-1176. doi: 10.1038/leu.2014.324

[33] Li, Y., Yang, L., Dong, L., Yang, Z. W., Zhang, J., Zhang, S. L., . . . Li, S. Q. (2019). Crosstalk between the Akt/mTORC1 and NF-kappaB signaling pathways promotes hypoxia-induced pulmonary hypertension by increasing DPP4 expression in PASMCs. *Acta Pharmacol Sin, 40*(10), 1322-1333. doi: 10.1038/s41401-019-0272-2

[34] Zuehlke, A. D., Beebe, K., Neckers, L., & Prince, T. (2015). Regulation and function of the human HSP90AA1 gene. *Gene, 570*(1), 8-16. doi: 10.1016/j.gene.2015.06.018

[35] Senapati, S., Kumar, S., Singh, A. K., Banerjee, P., & Bhagavatula, S. (2020). Assessment of risk conferred by coding and regulatory variations of TMPRSS2 and CD26 in susceptibility to SARS-CoV-2 infection in human. *J Genet, 99*(1). doi: 10.1007/s12041-020-01217-7

[36] Solerte, S. B., Di Sabatino, A., Galli, M., & Fiorina, P. (2020). Dipeptidyl peptidase-4 (DPP4) inhibition in COVID-19. *Acta Diabetol, 57*(7), 779-783. doi: 10.1007/s00592-020-01539-z

[37] Dang, D. T., Chun, S. Y., Burkitt, K., Abe, M., Chen, S., Havre, P., . . . Dang, L. H. (2008). Hypoxia-inducible factor-1 target genes as indicators of tumor vessel response to vascular endothelial growth factor inhibition. *Cancer Res, 68*(6), 1872-1880. doi: 10.1158/0008-5472.can-07-1589

[38] Leung, L. L. K., & Morser, J. (2018). Carboxypeptidase B2 and carboxypeptidase N in the crosstalk between coagulation, thrombosis, inflammation, and innate immunity. *J Thromb Haemost*. doi: 10.1111/jth.14199

[39] Naito, M., Taguchi, O., Kobayashi, T., Takagi, T., D'Alessandro-Gabazza, C. N., Matsushima, Y., . . . Gabazza, E. C. (2013). Thrombin-activatable fibrinolysis inhibitor protects against acute lung injury by inhibiting the complement system. *Am J Respir Cell Mol Biol, 49*(4), 646-653. doi: 10.1165/rcmb.2012-0454OC

[40] Mehta, P., McAuley, D. F., Brown, M., Sanchez, E., Tattersall, R. S., & Manson, J. J. (2020). COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet (London, England), 395*(10229), 1033-1034. doi: 10.1016/s0140-6736(20)30628-0

[41] Lee, D. C., Sohn, H. A., Park, Z. Y., Oh, S., Kang, Y. K., Lee, K. M., . . . Yeom, Y. I. (2015). A lactate-induced response to hypoxia. *Cell, 161*(3), 595-609. doi: 10.1016/j.cell.2015.03.011

[42] D'Ignazio, L., Bandarra, D., & Rocha, S. (2016). NF-kappaB and HIF crosstalk in immune responses. *FEBS J, 283*(3), 413-424. doi: 10.1111/febs.13578

[43] Messner, C. B., Demichev, V., Wendisch, D., Michalick, L., White, M., Freiwald, A., . . . Ralser, M. (2020). Ultra-High-Throughput Clinical Proteomics Reveals Classifiers of COVID-19 Infection. *Cell Systems, 11*(1), 11-24.e14. doi: https://doi.org/10.1016/j.cels.2020.05.012

[44] D'Alessandro, A., Thomas, T., Dzieciatkowska, M., Hill, R. C., Francis, R. O., Hudson, K. E., . . . Hansen, K. C. (2020). Serum Proteomics in COVID-19 Patients: Altered Coagulation and Complement Status as a Function of IL-6 Level. *Journal of Proteome Research*. doi: 10.1021/acs.jproteome.0c00365

[45] Hui, D. S., Wong, P. C., & Wang, C. (2003). SARS: clinical features and diagnosis. *Respirology, 8 Suppl*(Suppl 1), S20-24. doi: 10.1046/j.1440-1843.2003.00520.x

Table 1 Clinical characteristics and Laboratorial indexes of COVID-19 patients.

Characteristics	Total (N=144)	Non-severe (N=108)	Severe (N=36)	p value
Age, years				
median (IQR)	47 (38-56)	45 (37-54)	55 (48-65)	0.000
Sex, Male	77 (53.5)	57 (52.8)	20(55.6)	0.772
Symptoms on Admission				
Fever, %	104 (72.2)	70(64.8)	34 (94.4)	0.001
Pharyngalgia, %	17 (11.8)	15 (13.9)	2 (5.6)	0.180
Cough, %	65 (45.1)	47 (43.5)	18 (50.0)	0.499
Expectoration, %	26 (18.1)	19 (17.6)	7 (19.4)	0.802
Tiredness, %	16 (11.1)	10 (9.3)	6 (16.7)	0.221
Headache, %	16 (11.1)	9 (8.3)	7 (19.4)	0.066
Medical Treatment				
Oxygen inhalation, %	97 (67.4)	63 (58.3)	34 (94.4)	0.000
Antibiotics, %	26 (18.1)	15 (13.9)	11 (30.6)	0.024
Antivirus, %	144 (100)	108 (100)	36 (100)	1.000
Glucocorticoid, %	37 (25.7)	13 (12.0)	24 (66.7)	0.000
Gamma Immunoglobulin, %	33 (22.9)	7 (6.5)	26 (72.2)	0.000
Laboratorial Indexes				
LDH (ULN) [a]				0.000
Low or normal	115(79.9)	100 (92.6)	15 (41.7)	
High	29 (20.1)	8 (7.4)	21 (58.3)	
ALT (ULN) [b]				0.023
Low or normal	130 (90.3)	101 (93.5)	29 (80.6)	
High	14 (9.7)	7 (6.5)	7 (19.4)	
AST (ULN) [c]				0.000
Low or normal	125(86.8)	101 (93.5)	24 (66.7)	
High	19 (13.2)	7 (6.5)	12 (33.3)	
Total bilirubin (ULN) [d]				0.422
Low or normal	122(84.7)	93(86.1)	29 (80.6)	
High	22 (15.3)	15 (13.9)	7 (19.4)	
Total protein (LLN) [e]				0.477
High or normal	114 (79.2)	87 (80.6)	27 (75.0)	
Low	30 (20.8)	21 (19.4)	9 (25.0)	
Urea (ULN) [f]				0.004
Low or normal	137 (95.1)	106 (98.1)	31 (86.1)	
High	7 (4.9)	2 (1.9)	5 (13.9)	
Creatinine (ULN) [g]				0.029

Low or normal	135 (93.8)	104 (96.3)	31 (86.1)	
High	9 (6.3)	4 (3.7)	5 (13.9)	
Creatine kinase (ULN) [h]				0.000
Low or normal	123 (85.4)	99 (91.7)	24 (66.7)	
High	21 (14.6)	9 (8.3)	12 (33.3)	

Abbreviations: LDH, Lactate dehydrogenase; AST, aspartate aminotransferase; ALT, Alanine aminotransferase.

[a] ULN of LDH for male and female are 285 and 227 U/L, respectively.

[b] ULN of ALT for male and female are 50 and 40 U/L, respectively.

[c] ULN of AST for male and female are 40 and 35 U/L, respectively.

[d] ULN of total bilirubin is 20.5 μmol/L.

[e] LLN of total protein is 65 g/L.

[f] ULN of urea for male and female are 8 and 8.8 U/L, respectively.

[g] ULN of creatinine for male and female are 104 and 84 μmol/L, respectively.

[h] ULN of CK for male and female are 172 and 140 U/L, respectively.

Abbreviations: ULN, upper limit of normal; LLN, lower limit of normal; LDH, Lactate dehydrogenase; AST, aspartate aminotransferase; ALT, Alanine aminotransferase.

556 **Table 2 Medical records of temporal change associated severe patients.**

Patient Num.	HR1	HR2	HR3	HR4	HR5	HR6	HR7	LR1	LR2	LR3	LR4	LR5	LR6	LR7
Sex	F	M	F	M	M	M	M	M	F	M	F	M	M	M
Age	>60	>60	>60	≤60	≤60	≤60	≤60	>60	≤60	>60	≤60	≤60	≤60	≤60
Days for Admission	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Days for Progression to Severe Stage	2	1	4	0*	1	1	3							
Days for Proteomic Sampling	5	2	2	2	2	1	1	2	6	4	5	4	11	1
Days for Metabolomic Sampling	8	2	2	11	3	1	1	2	1	3	1	1	1	1
Basic Diseases	Left breast tumor resection	Benign thyroid tumor resection		Hypertension, gout arthritis, hyperuricemia, chronic renal insufficiency for 14 years	High blood pressure, type 2 diabetes	Chronic viral hepatitis b more than 20 years, hypothyroidism 5 years	Hepatitis b virus infection more than 10 years ago.	Type 2 diabetes, hyperlipidemia						Hepatitis b virus infection 20 years, type 2 diabetes

557 *: Patient HR4 was diagnosed as severe before admission to Taizhou Hospital.

Figure 1 Serum LDH expression levels in the study cohort. A-B) Serum LDH expression levels in different time courses grouped by age and sex. *, $0.01<p\leq0.05$; **, $0.005<p\leq0.01$; ***, $0.001<p\leq0.005$; ****, $p\leq0.001$. C) Serum LDH expression levels of severe and non-severe patients during the admission and discharge stage. D) Expression of serum LDH isoforms during the admission stage. E) Serum LDH expression levels of severe and non-severe patients in time courses at the 3-day interval. F) Receiver operating characteristic (ROC) of the study cohort when setting LDH expression level cutoff as 247 U/L. G) Individual inspection of 14 patients' LDH expression dynamics. Yellow dash line, Patient HR1; Purple dash line, Patient LR4.

Figure 2 Molecular differences between low- and high-risk patients. A) Heatmap of
34 differentially expressed proteins and two differentially expressed metabolites. LR,
low-risk patients. HR, high-risk patients. B) Boxplots of nine selected differentially
expressed proteins and two selected differentially expressed metabolites. C) Protein
network including 12 selected differentially expressed proteins.

Figure 3 Molecular differences between HR (basic) and HR (outliers) patients. A) Heatmap of 38 differentially expressed proteins. LR, low-risk patients. HR, high-risk patients. B) Boxplots of 13 selected differentially expressed proteins. C) Proposed mechanism for serum LDH elevation in COVID-19 patients.