1   Title: Explainable AI enables clinical trial patient selection to retrospectively improve treatment

2   effects in schizophrenia

4   Authors: Monika S. Mellem*, Matt Kollada, Jane Tiller, Thomas Lauritzen

6   Affiliations: BlackThorn Therapeutics, 780 Brannan St., San Francisco, CA 94103

8   * Corresponding Author

Abstract

Background: Heterogeneity among patients' responses to treatment is prevalent in psychiatric disorders. Personalized medicine approaches – which involve parsing patients into subgroups better indicated for a particular treatment – could therefore improve patient outcomes and serve as a powerful tool in patient selection within clinical trials. Machine learning approaches can identify patient subgroups but are often not "explainable" due to the use of complex algorithms that do not mirror clinicians' natural decision-making processes.

Methods: Here we combine two analytical approaches – Personalized Advantage Index and Bayesian Rule Lists – to identify paliperidone-indicated schizophrenia patients in a way that emphasizes model explainability. We apply these approaches retrospectively to randomized, placebo-controlled clinical trial data to identify a paliperidone-indicated subgroup of schizophrenia patients who demonstrate a larger treatment effect (outcome on treatment superior than on placebo) than that of the full randomized sample as assessed with Cohen's d.

Results: Using our combined explainable AI approach to identify a subgroup more responsive to paliperidone than placebo, the treatment effect increased significantly over that of the full sample (p<0.0001 for a one-sample t-test comparing the full sample Cohen's d=0.82 and a generated distribution of subgroup Cohen's d's with mean d=1.22, std d=0.09). In addition, our modeling approach produces simple logical statements (*if-then-else)*, termed a "rule list", to ease interpretability for clinicians. A majority of the rule lists generated from cross-validation found two general psychopathology symptoms, disturbance of volition and uncooperativeness, to predict membership in the paliperidone-indicated subgroup.

Conclusions: These results help to technically validate our explainable AI approach to patient selection for a clinical trial by identifying a subgroup with an improved treatment effect. With

these data, the explainable rule lists also suggest that paliperidone may provide an improved

therapeutic benefit for the treatment of schizophrenia patients with either of the symptoms of

high disturbance of volition or high uncooperativeness.

Trial Registration: clincialtrials.gov identifier: NCT 00083668; registered May 28, 2004

Keywords: machine learning, personalized medicine, explainability, patient selection,

schizophrenia

1. Background

The primary goal in a placebo-controlled clinical trial testing the efficacy of an

experimental medication is to show a treatment effect – that patients randomized to receive the

medication have improved outcomes compared to those receiving placebo. Within psychiatry,

there is heterogeneity in patients' responses, however, with some not responding well or at all

(e.g., Akil et al., 2018; Gillespie et al., 2017) which can weaken the overall response of the

treatment-receiving group compared to placebo. Additionally, the placebo response is robust in

psychiatric disorders (e.g., Weimer et al., 2015) making assessments of treatment efficacy more

difficult. A recently developed method termed Personalized Advantage Index (PAI) has shown

that a certain subgroup of patients may be more responsive to a particular treatment than placebo,

termed "treatment-indicated," and that predictive modeling could lead to personalized medicine

approaches for subtyping treatment-indicated patients (DeRubeis et al., 2014). In particular, this

could also help improve patient selection for clinical trials of that medication to enrich for

patients most likely to show a treatment effect.

Identifying a more responsive subgroup at the onset of a clinical trial is challenging,

however, as it is not necessarily known a priori which baseline patient characteristics lead to

medication responsiveness. Prior work in depression has used the PAI approach to identify the most predictive variables (Webb et al., 2018), but interpretability of the models for clinicians could be further improved as they would require interpretation of regression coefficients and do not suggest clear cutoffs for predictor variables. Here, we additionally used an approach inspired by explainable artificial intelligence (XAI), the Bayesian Rule Lists algorithm (BRL; Letham et al., 2015; Gao et al., 2019), to both help identify the most predictive variables and explain those predictions of treatment-indicated patients with simple *if-then-else* statements that better mirror a clinician's decision-making process by using Boolean criteria with clear cutoffs for predictor variables. The combined analytical approach of PAI and BRL was previously tested in depression and found to retrospectively identify a subgroup with improved treatment effect for the novel antagonist BTRX-246040 (Martin et al., 2019). But it has yet to be tested in other psychiatric populations.

We present this combined approach to patient selection for clinical trials using XAI and validation in schizophrenia patients through a retrospective analysis of a clinical trial. By showing that baseline features alone can be used to identify a subgroup of patients who demonstrate a larger average treatment effect, this approach opens up the possibility of identifying a targeted subgroup of patient prior to randomization to an arm and improving the clinical trial outcome by using this patient subgroup in particular.

2. Methods

2.1 Study

We analyzed data from a 6-week randomized, double-blind, placebo-controlled study evaluating the efficacy of extended-release paliperidone in the treatment of patients with

schizophrenia (clincialtrials.gov identifier: NCT 00083668). This trial was a success, and

paliperidone currently has FDA approval in this population. This study makes use of de-

identified data made available via the YODA Project (https://yoda.yale.edu/; research proposal

number 2019-4080) and was exempt from ethical oversight.

Patients were assessed for eligibility and phenotyped using standard clinical assessments

at baseline and randomized to one of several arms. In this analysis, we used data from the

15mg/day paliperidone arm (n=113) and the placebo arm (n=120). We selected the 15mg/day

arm over arms testing lower doses (3mg or 9mg per day) as it gave the greatest efficacy effect

compared to placebo. The efficacy endpoint was the Positive and Negative Syndrome Scale

(PANSS), a set of 30 questions administered by a trained clinician and scored on a 1-7 ordinal

scale (7 is most severe). The PANSS was administered weekly. After dropping patients with

missing baseline values used as features in the modeling (see below), 95 patients remained in the

treatment arm and 102 in the placebo arm.

2.2 Modeling

Our approach to identify treatment-indicated patients and improve the explainability of

the machine learning algorithm output classifying these patients involved combining two

approaches. First, we used the Personalized Advantage Index (PAI) algorithm (DeRubeis et al.,

2014) to create an index/score that is then used to label a patient as treatment-indicated or rest-

indicated (the rest of the subjects who are not treatment-indicated). This machine learning

approach relies on multiple linear regression which provides some level of explainability through

the coefficients of the predictors but is likely more complex than clinicians' decision making

processes which are closer in form to decision trees or lists. Thus, in a second step, we used the

Bayesian Rule Lists classifier (Letham et al., 2015; Gao et al., 2019) along with the predicted

treatment- and rest-indicated labels from PAI to create a more explainable classifier using

decision lists.

2.2.1 Personalized Advantage Index Modeling and Labeling

In our modeling approach, the PAI algorithm was first used to identify treatment-

indicated patients. The PAI approach predicts actual and counterfactual outcomes (i.e., a

patient's outcome for their assigned arm, drug or placebo, and the non-assigned arm) and

calculates the difference between these two scores (as previously described in DeRubeis et al.,

2014; Webb et al., 2018).

Briefly, we used an Elastic Net regressor (implemented in the python package scikit-

learn), with a grid search for hyperparameter optimization across the range of *alpha* = [0.001,

0.01, 0.1, 1, 10] and *l1_ratio* = [0.1, 0.5, 0.9]. The input features are listed in Table 1 where

"baseline" refers to week 0 of the trial, which precedes treatment arm randomization. The

outcome modeled was the 6-week post-treatment total PANSS score. Scores from patients

missing this 6-week score were replaced with their last observation (n = 80 patients) which was

consistent with the approach used in the original clinical trial analysis. This multiple regression

model then predicts the actual post-treatment PANSS score (on the patient's randomized arm)

and the hypothetical counterfactual score (by substituting the other arm in the regression

equation). PAI then returns a quantitative score for each patient that indicates the difference

between these predicted drug and placebo outcomes with better performance on drug

corresponding to a negative PAI score. A subsequent threshold then creates the two classes of

treatment-indicated and rest-indicated with possible thresholds examined in descending steps of

0.5 (0, -0.5, -1, -1.5,…). We selected the threshold that allocates ~50% of the sample to

treatment-indicated class to maintain a balanced dataset for the BRL classifier. As PAI was

responsible for generating the best possible indication labels for BRL to train on (i.e., generating

"ground truth" labels for BRL), the PAI regression model was trained on all the data. The grid

search of hyperparameters showed that *alpha* = 0.1, *l1_ratio* = 0.1 minimized the R^2.

Model Input Features	**PAI Model Output**	**BRL Model Output**
Demographics • Age • Sex Symptom scales • 30 Baseline individual item PANSS scores • Baseline total PANSS score • Baseline daytime drowsiness and quality of sleep scores from the Sleep Visual Analog Scale (VAS) • Baseline Personal and Social Performance Scale score • Baseline total Schizophrenia Quality of Life Scale score Others • Randomized arm (treatment or placebo; PAI only) • Interactions of randomized arm with the other features (PAI only).	• Predicted actual and counterfactual week 6 PANSS scores • Numerical PAI score (predicted outcome on treatment – predicted outcome on placebo) • Labels: treatment-indicated, rest-indicated created from thresholded PAI score	• Labels: treatment-indicated and rest-indicated

7   ***Table 1:*** *Model inputs and outputs. Several sequential outputs are generated during PAI*

8   *modeling and are listed in order of generation.*

2.2.2 BRL Modeling and Labeling for Additional Explainability

The BRL algorithm was used to create a more explainable model from the initial PAI

treatment-indicated and rest-indicated results. BRL uses sequenced logical rules and Bayesian

inference to make classifications (Letham et al., 2015; Gao et al., 2019). Here, we took in the

same baseline features other than the randomized arm and interactions (Table 1) and classified

patients as treatment- or rest-indicated using the BRL-generated *if-then-else* statements. A

Bayesian rule list is composed of Boolean statements that evaluate if features fit certain criteria

such as "If depression symptom score > 10" and the subsequent classification if the statement is

true – "then, patient is treatment-indicated." These statements are closer to a physician's

decision-making process than are the PAI regression outputs (feature coefficients without clear

cutoffs for feature values). Hyperparameters were set at 3 for the max rule length (number of

Boolean statements combined in an individual rule), 2 for the Bayesian prior hyperparameters of

expected individual rule length and 2 for the expected rule list length (excluding the final base

case). We used a 5-fold cross-validation framework that generated a model (a rules list) on the

80% of training data and used it to classify the patients in the remaining 20% of test data as

treatment- or rest-indicated. Thus five rules lists were generated from the five folds of cross-

validation. We additionally generated a "final" BRL model that could be validated on external

data sets by training the model on the full data set.

2.2.3 Comparing Treatment Effects

After classifying each patient as treatment- and rest-indicated labels using BRL, we

assessed if treatment-indicated patients showed an improved treatment effect compared with the

full sample. A group-level assessment comparing the post-treatment outcomes of patients on the

treatment arm vs. the placebo arm was used to assess a treatment effect. Here, the treatment

effect was assessed on the actual week 6 PANSS total scores of individual patients grouped by

their treatment arm (using Cohen's d as a measure of treatment effect size). After determining

the labels from the BRL outputs, the actual week 6 PANSS total scores for the treatment-

indicated patient subgroup were used to calculate the treatment effect for that subgroup. Then

the treatment effect of the treatment-indicated subgroup was compared with the treatment effect

of the full randomized sample. Our null hypothesis is that patient selection with BRL provides

no improvement of the treatment effect, while our alternative hypothesis is that BRL does

improve the treatment effect relative to that of the full sample. Thus to test this statistically, we

generated a distribution treatment effects by performing BRL 100 times and calculating the

Cohen's d for each BRL-labeled treatment-indicated subgroup. We then compared this

distribution of 100 Cohen's d's with the Cohen's d of the full sample using a two-sided, 1-

sample t-test. Additionally, we assessed the consistency of classification for each subject across

the five rule lists generated from the five folds of cross-validation.

3. Results

Our approach to identify treatment-indicated patients and improve the explainability of

the machine learning algorithm output classifying these patients involved combining two

approaches – PAI and BRL. While the final output and results are the treatment-labeled patient

subgroup and the *if-then-else* rules list from the BRL model, the initial modeling with the PAI

algorithm produced some interim results that we first examined. The PAI regression equation

modeled actual week 6 PANSS scores using actual treatment arm assignment and several

demographic and baseline symptom severity predictor variables (see Table 1). Figure 1 shows

the comparison of measured week 6 PANSS scores with predicted scores (a perfect model would

show all samples sitting on the x=y line). The R^2 of the model shows it explained 58% of the

variance (adjusted R^2 = 0.32). Then, by substituting in the counterfactual randomization arm, the

regression equation was used to make predictions of the week 6 PANSS scores if patients were

receiving the counterfactual treatment. The difference between actual and counterfactual

predictions were used to calculate the PAI scores (predicted score on treatment – predicted score

on placebo) and determine the indication labels to be used by BRL. The distribution of PAI

scores are shown in Figure 2, and a threshold of -9.5 generated two balanced classes: treatment-

indicated (n=100) and rest-indicated (n=97). In this data set, the PAI threshold corresponded to a

30% reduction in average post-treatment PANSS scores for patients in the treatment arm relative

to the scores of patients in the placebo arm.

After determining the indication labels from the thresholded PAI scores, we trained the

BRL classifier on these labels using 5-fold cross-validation. For the training data, the BRL

classifier had an average accuracy of 77.9% (standard deviation = 2.0%), an average Area Under

the ROC Curve (AUC) of 0.83 (std = 0.02), and an average F1 score of 0.77 (std = 0.02) across

the five folds. For the test data, the BRL classifier had an average accuracy of 74.1% (standard

deviation = 5.1%) an average AUC of 0.76 (std = 0.04), and an average F1 score of 0.73 (std =

0.04) across the five folds. Of the 197 patients from the full randomized sample, 87 were labeled

treatment-indicated by the BRL algorithm, and the full confusion matrix of cross-validated labels

are shown in Figure 3. On the full cross-validated test results, the overall accuracy was 74.1%,

the AUC score = 0.74, and the F1 score = 0.73.

Comparison of the two arms (treatment, placebo) for the full sample v. the treatment-

indicated group shows an increase in the Cohen's d between arms from 0.82 to 1.24 (Figure 4).

We also assessed if this large increase in effect size was consistent for the BRL-classified

treatment-indicated subgroup and if it was significantly greater than the full sample effect size.

We generated a distribution of Cohen's d's by performing the same BRL process 100 times and

calculating the treatment effect for the treatment-indicated subgroup each time. We found that

the treatment effects from these 100 iterations of subgrouping were statistically greater than the

full sample treatment effect (BRL-classified treatment-indicated subgroup Cohen's d's mean =

1.22, std = 0.09; two-sided, 1-sample t-test: t=43.2, p<0.0001).

The five rule lists returned by the BRL classifier differ slightly across the five folds but

did identify high disturbance of volition and high uncooperativeness as commonly identifying

baseline features of patients who were more likely to respond on treatment than on placebo.

Table 2 displays the five rule lists.

< PLACE TABLE 2 APPROXIMATELY HERE >

Though the rule lists show some differences across folds, we found that they still

classified patients similarly when applied to the whole data set (not just the test set). Figure 5

shows the number of times that patients were labeled treatment-indicated. For consistent

classifiers, most patients should be labeled either five times or zero times (corresponding to a

rest-indicated patient), and the labeling reflects this well. Additionally, most patients labeled by

the BRL algorithm treatment-indicated four or five times were treatment-indicated according to

the "ground truth" PAI labels. As expected, this was reversed with most patients labeled by the

BRL algorithm treatment-indicated zero times were rest-indicated according to the "ground

truth" PAI labels. Thus, the five rule lists mostly subtype patients similarly though their wording

can differ.

We additionally generated a final BRL model that could be validated on external data sets

(Figure 6). This model generated a single rule list as it is trained on the full data set as opposed

to the cross-validation approach which generated five rules lists across the five folds. High

baseline uncooperativeness and high baseline disturbance of volition each remain predictive of

treatment-indicated subgroup membership. For training and testing on the full data set, accuracy

was 76%, AUC was 0.81, and the F1 score was 0.75.

4. Discussion

With this retrospective analysis, we have technically-validated an approach of using a

combination of machine learning approaches to identify clinically-explainable rules that

effectively subtype paliperidone-indicated schizophrenia patients who show an improved

treatment effect over the full randomized sample. This extends the prior validation that

demonstrated the method's effectiveness in a clinical trial of a novel depression treatment

(Martin et al., 2019).

While this validation was performed on a successful trial where the full sample already

displayed the success of the experimental drug, we demonstrated than the treatment effect can be

further improved with a patient selection approach. Statistically speaking, increasing the effect

size can help decrease the enrollment numbers for patients, thereby possibly decreasing the cost

and time of a clinical trial. Thus, in addition to patient selection that improves the treatment

effect for unsuccessful trials (Martin et al., 2019; Webb et al., 2018), our results suggest that

patient selection could help clinical development even for treatments with stronger effects.

While the primary goal in this study was to validate a patient selection approach, the

methodology also allows us to better understand a potential paliperidone-responsive subtype of

schizophrenia. A majority of the rule lists generated from the cross-validation framework found

two general psychopathology symptoms, disturbance of volition and uncooperativeness, to be

predictive of membership in the paliperidone-indicated subgroup suggesting that paliperidone

may be indicated for the treatment of patients with either of these symptoms. While subtyping

schizophrenia is an active area of research (e.g., Jablensky, 2006; Gillespie et al., 2017), the

higher severity of uncooperativeness and disturbance of volition seen in this paliperidone-

indicated subgroup has not been previously described and should be externally validated. For

this reason, we have included a single BRL rule list as a "final" model which could be tested by

others interested in a paliperidone-indicated subgroup.

This approach is particularly useful in the context of selecting patients for clinical trial

enrollment as the clinical trial outcome is dependent on large effects that are seen for patients in

the treatment arm but not placebo arm. However, the proposed approach with the additional

clinician-friendly explainability of BRL could make it more broadly useable as a clinical

decision support system which are not commonly incorporating machine learning yet (Sutton et

al., 2020; Levy-Fix et al., under review). The framework could be extended to accommodate

multiple classes for indications of multiple treatments. With the proper validation, this could

provide clinicians with a tool to match the best of several possible treatments to a particular

patient.

Some limitations with this approach remain. The PAI model does have some bias in its

predictions as shown by the greater differences in measured and predicted week 6 PANSS scores

in the larger and smaller ranges. This reflects a model that is underfit and could be due to

missing predictor variables or due to using a linear rather than non-linear model. Even with this

weakness, the PAI model still provided adequate predictions to allow the BRL model to find a

treatment-indicated subgroup with improved treatment effect size. Another limitation is the

testing only within a single data set. A more robust approach would be to test the BRL model in

a separate data set to assess generalization of the model and the proposed paliperidone-indicated

schizophrenia phenotype. Some may question using the baseline total PANSS score as a

predictor variable either due to its possible collinearity with other PANSS item scores or that the

resulting use of lower PANSS scores to classify rest-indicated patients (therefore higher PANSS

score is indirectly predictive of treatment-indicated patients) may be reflecting an effect of

regression to the mean for the treatment-indicated patients. Unpublished analyses in our lab did

not find major differences in performance or predictive features whether including or not

including this variable. Additionally, the critical result is not that PANSS scores are reduced for

the treatment-indicated patients, but that they are reduced much more on treatment than on

placebo. Thus selecting patients as rest-indicated based on the baseline total PANSS score and

the implications that has for selecting treatment-indicated patients does not have any bearing on

the improved difference seen between arms of the treatment-indicated patients.

5. Conclusions

These results help to technically validate our explainable AI approach to patient selection for a

clinical trial by identifying a subgroup of schizophrenia with an improved treatment effect.

Importantly, this approach opens up the possibility of identifying a targeted subgroup of patient

prior to randomization to an arm and improving the clinical trial outcome by using this patient

subgroup in particular.

Abbreviations

AI: Artificial Intelligence, AUC: Area Under the ROC Curve, BRL: Bayesian Rule List, FDA: U.S. Food and Drug Administration, PANSS: Positive and Negative Syndrome Scale , PAI: Personalized Advantage Index, PBO: Placebo, ROC: Receiver Operating Characteristic, Rest-ind: Rest-indicated, TRT: Treatment, TRT-ind: treatment-indicated, XAI: Explainable Artificial Intelligence, YODA: Yale University Open Data Access

Declarations

Ethics approval and consent to participate

This study makes use of de-identified data made available via the YODA Project (https://yoda.yale.edu/; research proposal number 2019-4080) and was exempt from ethical oversight.

Consent for publication

Not applicable.

Availability of data and materials

The data that support the findings of this study are available from the YODA Project

(https://yoda.yale.edu/) but restrictions apply to the availability of these data, which were used

under license for the current study, and so are not publicly available.

Competing interests

All authors are current or previous employees of BlackThorn Therapuetics and hold stock or

stock options in BlackThorn Therapeutics.

Funding

Funding was provided by BlackThorn Therapeutics whose employees were involved designing

this post-hoc analysis, performing the analysis, interpreting the results, and writing the

manuscript.

Authors' contributions

JT, MK, MM, and TL helped conceive of the approach to PAI+BRL validation with the YODA

data set and subsequent analysis, and JT advised on methodology with the team who developed

the PAI+BRL approach. MM performed the analysis of the data. MM, MK, and TL interpreted

the results, and wrote or revised the manuscript. All authors read and approved the final

manuscript.

Acknowledgments

We would like to thank Clark Gao, Yuelu Liu, Humberto Gonzalez, and Parvez Ahammad for

their contributions in creating the PAI+BRL methodology that is used here in a different

application, and Kathy Tracy, Rezi Zawadzki, Andrew Jaffe, John Dunlop, and Bill Martin for

further discussions. This study, carried out under YODA Project # 2019-4080, used data

obtained from the Yale University Open Data Access Project, which has an agreement with

JANSSEN RESEARCH & DEVELOPMENT, L.L.C. The interpretation and reporting of

research using this data are solely the responsibility of the authors and does not necessarily

represent the official views of the Yale University Open Data Access Project or JANSSEN

RESEARCH & DEVELOPMENT, L.L.C.

References

Akil H, Gordon J, Hen R, Javitch J, Mayberg H, McEwen B, Meaney MJ, Nestler EJ. Treatment

resistant depression: a multi-scale, systems biology approach. Neuroscience and Biobehavioral

Reviews. 2018; 84: 272-288. doi.org/10.1016/j.neubiorev.2017.08.019

DeRubeis RJ, Cohen ZD, Forand NR, Fournier JC, Gelfand LA, Lorenzo-Luaces L. The

personalized advantage index: translating research on prediction into individualized treatment

recommendations. A demonstration. PLoS One. 2014; 9(1):e83875.

doi:10.1371/journal.pone.0083875

Gao Q, Gonzalez H, Ahammad P. MCA-based rule mining enables interpretable inference in

clinical psychiatry. In Shaban-Nejad A., Michalowski M. (eds) Precision Health and Medicine.

W3PHAI 2019. Studies in Computational Intelligence. Springer, Cham. 2020; vol. 843.

doi.org/10.1007/978-3-030-24409-5_3

Gillespie AL, Samanaite R, Mill J, Egerton A, MacCabe JH. Is treatment-resistant schizophrenia

categorically distinct from treatment-responsive schizophrenia? a systematic review. BMC

Psychiatry. 2017; 17(12):1-14. doi.org/10.1186/s12888-016-1177-y

Jablensky A. Subtyping schizophrenia: implications for genetic research. Mol. Psychiatry. 2006;

11(9):815-836. doi: 10.1038/sj.mp.4001857.

Letham B, Rudin C, McCormick TH, Madigan D. Interpretable classifiers using rules and

Bayesian analysis: building a better stroke prediction model. The Annals of Applied Statistic.*s*

2015; 9(3):1350-1371. doi.org/10.1214/15-AOAS848.

Levy-Fix G, Kuperman GJ, Elhadad N. Machine learning and visualization in clinical decision

support: current state and future directions. 2019; Preprint at arXiv:1906.02664v1

Martin B, Gao Q, Liu Y, Madrid A, Ahammad P, Tiller J. Explainable AI approach reveals

treatment responders in a randomized controlled trial of BTRX-246040, a potent and selective

NOP receptor antagonist. Neuropsychopharmacology, 2019, vol. 44, no. SUPPL 1, pp. 448-448.

Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of

clinical decision support systems: benefits, risks, and strategies for success. NPJ Digital

Medicine. 2020; 3(17). doi.org/10.1038/s41746-020-0221-y

Webb CA, Trivedi MH, Cohen ZD, Dillon DG, Fournier JC, Goer F, et al. Personalized

prediction of antidepressant v. placebo response: evidence from the EMBARC study.

Psychological Medicine. 2018; 1-10. doi.org/10.1017/S0033291718001708

Weimer K, Colloca L, Enck P. Placebo effects in psychiatry: mediators and moderators. Lancet

Psychiatry. 2015; 2(3):246-257. doi:10.1016/S2215-0366(14)00092-3

Figure Legends

Figure 1: *Individual PAI prediction results for actual randomized arms. The plot shows the*

measured total PANSS score at week 6 vs. the averaged predicted total PANSS score at week 6

from the Elastic Net regression model (each dot is an individual patient, n=197). Patients are

colored by their actual randomized arm (paliperidone treatment in blue, placebo in orange), and

as expected the week 6 scores are generally higher for patients receiving placebo. The dashed

line is y=x. Variance explained by the model is 58%. Note that these are not the counterfactual

predictions.

Figure 2: PAI score thresholded graph. A threshold of -9.5 was chosen to create roughly

balanced classes and indicates that membership in the treatment-indicated subgroup required a

predicted treatment arm PANSS score that is 9.5 points less than the predicted placebo arm

PANSS score.

Figure 3: Confusion matrix for cross-validated BRL labels. Actual labels are PAI-derived

labels, and predicted labels are BRL-derived labels.

Figure 4: Comparison of actual post-treatment PANSS scores for the full sample and the

treatment-indicated subgroup. Bars display treatment (TRT) and placebo (PBO) arms for the

full randomized sample (left graph, TRT n=95, PBO n=102) with an illustrative instance of the

BRL-classified treatment-indicated (TRT-indicated) subgroup (right graph, TRT n=41, PBO

n=46). At a Cohen's d of 1.24, the effect size between arms for the treatment-indicated subgroup

is increased more than 50% over the effect size of the full sample (d = 0.82). Error bars are

95% confidence intervals on the mean.

Figure 5: The treatment-indicated (TRT-ind) labeling consistency is seen with a histogram of

BRL-labeled treatment-indicated counts. It reflects the number of times (count) that a patient

was classified as treatment-indicated across all five rule lists from the 5-fold BRL cross-

validation. Most patients are either classified five times or zero times which indicates a higher

level of consistency in patient subtyping across the different rule lists. Additionally, most patients

in the 5-count column were also labeled as TRT-ind by the PAI algorithm (orange portion of the

bar) while most patients in the zero-count column were labeled as Rest-ind by PAI (blue portion

of the bar). Patient numbers corresponding to the orange and blue portions are shown in the

table below the histogram.

Figure 6: *"Final" rule list created by the BRL model when trained on the full data set. The two*

individual item scores for Disturbance of Volition and Uncooperativeness again appear to

classify the treatment-indicated (TRT-ind) subgroup.

Table 2: *List of the five rule lists created by the BRL model across the five folds.*

Fold	Rule List
1

 |
| **2** | |

Table 2 Legend*: The probability shown in parentheses after each rule is estimated from the*

percent of patients who satisfy that rule and were labeled with the given label for the rule by the

PAI "ground-truth" labels, and the confidence intervals were estimated with bootstrapping. The

alphanumeric symbol before each symptom (e.g., P02, G08) refers to the question number from

the PANSS scale. Two individual item scores from the PANSS scale repeatedly were involved in

subtyping the treatment-indicated patients. Both Disturbance of Volition >= 3 and

1   *Uncooperativeness >= 3 appear in multiple rule lists for the treatment-indicated (TRT-ind)*

2   *subgroup and are categorized as General Psychopathology symptoms.*

BRL Count	0	1	2	3	4	5
PAI TRT-ind	19	10	1	1	5	64
PAI Rest-ind	74	7	1	1	2	12