

A Scoping Review of the Experience of Implementing Population Testing for SARS-CoV-2

Clare R Foster, Fiona Campbell, Lindsay Blank, Anna Cantrell, Michelle Black, Andrew C K Lee

School of Health and Related Research, The University of Sheffield, Regent Court, 30 Regent Street, Sheffield S1 4DA, UK

Corresponding author: Clare Foster, clare.foster@sheffield.ac.uk, School of Health and Related Research, The University of Sheffield, Regent Court, 30 Regent Street, Sheffield S1 4DA, UK

Declarations of Interest: None

This research was part-funded by Public Health England.

Abstract

Background: The SARS-CoV-2 pandemic has led to the swift introduction of population testing programmes in many countries across the world, using testing modalities such as drive-through, walk-through, mobile and home visiting programmes. Here, we provide an overview of the literature describing the experience of implementing population testing for SARS-CoV-2.

Methods: We conducted a scoping review using Embase, Medline and the Cochrane Library in addition to a grey literature search. We identified indicators relevant to process, quality and resource outcomes related to each testing modality.

Results: 2,999 titles were identified from the academic literature and the grey literature search, of which 22 were relevant. Most studies were from the USA and the Republic of Korea. Drive-through testing centres were the most common testing modality evaluated and these provided a rapid method of testing whilst minimising resource use.

Conclusions: The evidence base for population testing lacks high quality studies, however, the literature provides evaluations of the advantages and limitations of different testing modalities. There is a need for robust evidence in this area to ensure that testing is deployed in a safe and effective manner in response to the Covid-19 pandemic.

Keywords: mass testing, population testing, SARS-CoV-2, Covid-19

Introduction

In response to the SARS-CoV-2 pandemic, many countries implemented population testing programmes as part of countermeasures to contain the spread of infection and mitigate its health and economic impacts. Population testing provides disease surveillance required to inform broader policy decisions, target resource utilisation and, when twinned with timely case isolation and contact tracing, more effective containment of the virus ¹. Worldwide, population testing programmes are diverse, depending on the population eligible for testing, the technology used to sample and analyse specimens, as well as the timing and frequency of testing.

Population testing seeks to identify people infected with SARS-CoV-2 in a pre-defined group such as health and care workers. By identifying cases of infection through testing, action can be taken to limit infection spread by isolating infected individuals and their contacts during their infectious period. As SARS-CoV-2 may be spread by asymptomatic individuals, including these individuals in testing programmes could help reduce viral transmission. Cumulatively, these actions help control the spread of infection and create conditions that would enable the relatively normal functioning of society.

Modalities used for COVID-19 population testing include drive-in, walk-in, mobile sites, postal testing and home visits. The UK, for example, adopted a testing strategy with 5 pillars, each pillar pertaining to a population subgroup and focusing on either diagnosis, detecting past infection or for surveillance purposes to estimate population prevalence. Testing is co-ordinated centrally and delivered from satellite centres and via postal testing ². South Korea adopted walk-in and drive-through testing ^{3,4}, whilst some areas in Scotland have instituted home-testing to reach more vulnerable groups who cannot access test facilities easily ^{5,6}. There is also increasing political and societal concern of the socioeconomic impact of blunt strategies such as national 'lockdowns' to combat COVID-19 ⁷, and their effect on health inequalities. There may therefore be value in studying the different approaches used worldwide, and to learn from successful programmes from other countries.

What is currently unclear is whether any modality of population testing is more robust and efficacious for containing the virus. There is a need to identify population testing programmes that are more accessible and effective at containing the spread of infection, together with the determinants of success. This can help inform national testing policies as part of the pandemic response efforts to minimise the health, social and economic harms. We conducted a scoping review to describe the volume and type of evidence reporting on the experience of implementing population testing for SARS-CoV-2 in high and upper-middle income countries during the pandemic.

Methods

Scoping review methodology

Scoping reviews aim to rapidly map the key concepts underpinning a research area, by comprehensively summarising evidence in order to inform practice and policy and provide direction for future research ^{8,9}. Scoping reviews use rigorous and transparent literature searching methods but differ from systematic reviews as the quality of included studies are not routinely assessed, nor do they provide a synthesised answer to a particular research question ¹⁰.

This scoping review followed the framework proposed by Arksey and O'Malley ⁸ and refined by Levac et al ¹¹, briefly comprising: identification of the research question, identification of studies, selection of studies, charting of data and collation of results. This review was commissioned by Public Health England who were consulted on the interim outputs of the study. The trial protocol was published on PROSPERO, number CRD42020186506.

Identification of the research question

The purpose of this review was to assess the volume of published literature describing the experience of implementing population testing for Covid-19 and to identify the nature and characteristics of the testing programmes. We sought to elucidate what data was available to assess the outcomes of these testing programmes in terms of processes, populations, quality and/or resource-use. We developed these broad aims in order to generate breadth of coverage and map the literature on this topic so that key concepts and gaps could be identified to inform further practice and policy.

Table 1 outlines the inclusion and exclusion criteria for this review. Studies were included if they described the process of providing or accessing a population testing point for symptomatic and/or asymptomatic individuals for Covid-19, using an antigen or antibody test in any setting, using any testing modality. In order to prioritise research relevant to high income countries such as the UK, we included literature relating to comparable health services from high and upper-middle income countries only.

Studies of laboratory aspects of testing (including diagnostic accuracy), commentaries, opinion pieces and modelling studies were excluded. Studies that described screening where samples were not taken, or that described the testing of passengers at ports or borders, were also considered out of scope and excluded.

Literature identification and selection

A search strategy (see Appendix 1) was developed to retrieve studies that had evaluated or described the process of providing or accessing a testing point for population testing for Covid-19. An information specialist (AC) searched the electronic databases Medline, Embase and the Cochrane Library (Figure 1). Searches were originally conducted in May 2020 and updated in August 2020. The search was limited to studies in English and published between January and August 2020.

Extracted titles and abstracts were screened by at least two reviewers (CF, FC, LB). 250 full text articles were reviewed to clarify whether the article met the inclusion criteria given in Table 1, either because no abstract was available or because it was unclear from the title and abstract alone whether the study met the inclusion criteria. Abstracts were often unavailable due to the large number of commentaries and opinion pieces found by the search strategy. A formal quality appraisal of the evidence was not conducted, given the remit of the scoping review.

In addition, a grey literature search for national and international clinical guidelines was conducted during May 2020. The World Health Organisation (WHO) and the European Centre for Disease Prevention and Control (ECDC) websites were searched, plus websites in the English language from the UK, USA, Canada, Australia, New Zealand, South Korea, China and Taiwan. This search identified 58 potentially relevant guidelines. On further review, only one guideline was relevant to population testing and detailed an approach to drive-through screening implemented in South Korea¹². A further 21 guidelines looked at wider aspects of screening.

Table 1. Inclusion and Exclusion Criteria

Inclusion criteria	Exclusion criteria
<ul style="list-style-type: none">• Any study evaluating or describing the process of providing or accessing a population testing point for Covid-19• Symptomatic and/or asymptomatic individuals• Any setting• Antibody or antigen testing• Any testing modality• High or upper-middle income countries according to World Bank criteria	<ul style="list-style-type: none">• Commentaries, modelling studies or opinion pieces• Studies describing only laboratory aspects of testing, including studies of diagnostic accuracy.• Studies describing only testing at ports or borders• Studies describing only screening in which samples were not taken• Not written in English


Charting of data and collation of results

After the screening was completed, relevant content in the included studies was extracted into a spreadsheet. Data extraction was verified by a second reviewer who checked data extraction from a random sample of four articles. The mode of testing was categorised into one of five different types: drive-through testing, home visiting testing, indoor walk-through centre, outdoor walk-through centre and mobile testing. For outcome data, a thematic framework was used which categorised any quantitative outcomes indicators into one of four groups: process outcomes, participant outcomes, quality outcomes and resource use outcomes.

Results

The database search returned 2,999 results (Figure 1). After automated and manual de-duplication, 2,751 unique references were screened for relevance to the question. On first screening of titles, 250 references were identified as potentially relevant, and on further reading 22 were categorised as relevant.

Figure 1: PRISMA flow diagram



Study Characteristics

A summary of the studies' characteristics is presented in Table 2.

Most of the published literature on this topic is difficult to assign to a study type. Studies were often referred to as brief reports or short communications. The articles typically comprise a description of the testing modality of interest, often with a diagram of the layout of the testing centre and useful operational details followed by an evaluation of its advantages and disadvantages. In some papers, a comparison group was described, however there were no published randomised controlled trials. One article was a qualitative interview study of early experiences of drive-through testing centres¹³.

All the eligible studies described population testing programmes where samples were taken from symptomatic and occasionally asymptomatic individuals. There were no eligible studies of mass testing (defined as regular and/or large-scale testing of individuals from defined populations regardless of symptom status) that used cheaper, faster tests than Q-PCR. For example, Appa et al. described high throughput, community-wide ascertainment of SARS-CoV-2 prevalence using PCR and antibody testing but not rapid methods such as lateral flow tests¹⁴. It was sometimes difficult to categorise articles as 'mass testing' because many studies did not describe the eligible population or tests used in detail. It was also often unclear why asymptomatic individuals were attending for testing.

Many articles describing testing programmes were from the USA (43% of studies) and the Republic of Korea (29% of studies). Most articles described testing programmes in high income countries (95% of articles), with only one originating from an upper-middle income country (Malaysia) ¹⁵.

Testing Modalities

Several different testing modalities were described, which were categorised into five main categories: drive-through, home visiting, mobile testing, indoor walk-through centres and outdoor walk-through centres.

Table 2: Characteristics of included articles

First Author	Country	Mode of Testing
Appa ¹⁴	USA	Drive-through
Brammer ¹³	USA	Drive-through
Chang ¹⁶	Republic of Korea	Drive-through and walk-through
Drees ¹⁷	USA	Drive-through
Flynn ¹⁸	USA	Drive-through and walk-through
Goldberg ¹⁹	USA	Home visiting testing
Halalau ²⁰	USA	Drive-through
Hill ⁵	Scotland	Drive-through and home visiting testing
Kim ²¹	Israel	Drive-through and home visiting testing
Kim ²²	Republic of Korea	Indoor walk-through centre
Kwon ⁴	Republic of Korea	Drive-through
Kwon ²³	Republic of Korea	Outdoor walk-through centre
Lee ¹⁵	Malaysia	Drive-through
Lin ²⁴	Taiwan	Outdoor walk-through centre
Lindholm ²⁵	USA	Drive-through
Manaus ²⁶	Singapore	Indoor walk-through centre
Ministry of Foreign Affairs ¹²	Republic of Korea	Drive-through
Mark [11]	Scotland	Drive-through and home visiting testing
Rivkees ²⁷	USA	Drive-through, walk-through and mobile testing
Seo ²⁸	Republic of Korea	Mobile testing
Shah ²⁹	USA	Drive-through
Ton ³⁰	USA	Drive-through

Sometimes described as off-site COVID-19 testing centres (OSCTCs) ¹³, drive-through testing centres were by far the most common testing modality evaluated (in 72% of articles), in Israel, Malaysia, Korea, Scotland and the USA. This testing modality enabled the use of a vehicle as a self-contained unit, which can reduce the spread of infection. Most were in car parks, stadiums and parks, and one was in an open-air area of a hospital [26]. Some centres enabled individuals without a car to walk in for testing ¹³ in order to increase accessibility. Six articles described drive-through testing in combination with either home visiting testing, mobile testing and/or walk-through testing, enabling a greater proportion of the population to access testing.

Home visiting testing (18% of articles) typically involved a small number of healthcare workers visiting the home of an individual to perform a test. This enabled individuals who are home-bound, frail or have no means of private transportation to access testing without having to use an ambulance, visit a hospital or rely on the assistance of others to access a drive-through site. These schemes were often used as an alternative to local drive-through testing facilities. Home visiting testing took place in Israel, Scotland and the USA.

Two articles described mobile testing. In Korea, testing staff visited workers onsite at their workplaces ²⁸ and in Florida, mobile PCR testing laboratories were used to provide point-of-care testing in different cities ²⁷.

Indoor walk-through centres based in healthcare facilities were used in Singapore and Korea. There were several different designs for walk-through centres, which were located inside hospitals or other healthcare facilities: screening centres ^{22, 26}, negative pressure booths ²² and negative pressure tents ¹⁶. Screening centres permit individuals to access testing inside a building. The Singapore Screening Centre was designed to minimise the movement of patients around the building ²⁶. Patients were assigned a seat number and tagged with a tracker to facilitate contact tracing; staff visited patients in their seats to further reduce contact amongst patients.

Negative pressure booths and tents have been designed to minimise the opportunity for viral spread in an indoor setting. Negative pressure booths ²² were used for sample collection and medical examination procedures in Korea. The booths were inspired by the design of biosafety cabinets and contain a 'glove wall' separating the patient and the medical staff member, who communicate using an interphone. Patients complete registration, questionnaires and payment outside the booths in other sections of the screening centre. Negative pressure booth systems aimed to protect healthcare staff, reduce PPE use and increase throughput compared to other walk-through systems. Negative pressure tents ¹⁶, also located in Korea, were similar to negative pressure booths, but the whole tent is under negative pressure. Staff working in the tent wore full PPE and most of the tent required sterilisation between people tested that took at least 30 minutes.

The final testing modality described was outdoor walk-through centres. These were located outside hospitals in Korea and Taiwan ^{23, 24}. The Korean clinic ²³ screened all patients and visitors to the hospital with the aim of minimising ward closures due to COVID-19 outbreaks. In Taiwan, a 'multifunctional sampling station' was built outside an emergency department, using a 2cm thick clear acrylic board to separate emergency department patients and medical personnel, with inbuilt gloves used to conduct sampling ²⁴.

Populations Tested

Table 3 summarises details of the populations tested and the types of test used. 55% of studies provided information regarding the population eligible to be tested. 41% of articles described accepting both symptomatic and asymptomatic individuals and 14% accepted those with symptoms only. Some testing centres had an algorithm for testing eligibility involving symptoms, epidemiological links, occupational risk factors and/or potential for community exposure. For some, the testing criteria changed over time as the pandemic progressed ^{20, 29}.

Table 3: Populations eligible to be tested and types of testing used

First Author	Population tested	Type of test
Appa ¹⁴	Symptomatic and asymptomatic	Oro-pharyngeal/mid-turbinate swab for PCR testing, finger prick blood test for antibody testing
Brammer ¹³	N/A	N/A
Chang ¹⁶	Symptomatic and asymptomatic	RT-PCR tests of 'specimens'
Drees ¹⁷	Symptomatic only	Not specified
Flynn ¹⁸	Unclear/unspecified	'In-house COVID-19 test' of nasopharyngeal swabs
Goldberg ¹⁹	Symptomatic and/or epidemiological link and/or risk factors and/or potential for community exposure	Nasopharyngeal swab collection for RT-PCR analysis
Halalau ²⁰	Unclear/unspecified	Nasopharyngeal swab collection for RT-PCR analysis
Hill ⁵	Unclear/unspecified	Combined nose and throat swab
Kim ²¹	Symptomatic only	Swab testing
Kim ²²	Symptomatic only	Not specified
Kwon ⁴	Symptomatic and asymptomatic	Naso- and oro-pharyngeal swabs, sputum specimen
Kwon ²³	Symptomatic and/or epidemiological link and/or risk factors and/or potential for community exposure	Pre-testing questionnaire
Lee ¹⁵	Symptomatic and asymptomatic	Temperature measurements, nasopharyngeal and oral swabs
Lin ²⁴	Unclear/unspecified	Nasopharyngeal and oral swab, sputum collection, blood testing for antibodies
Lindholm ²⁵	Symptomatic and/or epidemiological link and/or risk factors and/or potential for community exposure	Screening questionnaire followed by nasopharyngeal swab collection for RT-PCR analysis for those with symptoms or epidemiological link
Manauis ²⁶	Symptomatic and/or epidemiological link and/or risk factors and/or potential for community exposure	Swab testing
Mark [11]	Unclear/unspecified	Not specified
Ministry of Foreign Affairs ¹²	Unclear/unspecified	Upper respiratory tract sample (all), lower respiratory tract sample (only if participant can expectorate sputum alone into a container)
Rivkees ²⁷	Symptomatic and asymptomatic	PCR and antibody testing
Seo ²⁸	Unclear/unspecified	Questionnaire followed by sample collection for RT-PCR testing
Shah ²⁹	Symptomatic and/or epidemiological link and/or risk factors and/or potential for community exposure	Originally naso- and oro-pharyngeal swab, later solely nasopharyngeal swab for RT-PCR testing (following CDC advice)
Ton ³⁰	Unclear/unspecified	Nasal swab

Types of testing used

The most common method for sampling was through nasal or throat swabs. 45% of articles described the use of nasal swabbing, involving either nasal, nasopharyngeal or mid-turbinate sampling. 27% described the use of throat/oropharyngeal swabbing. 27% stated that swabbing was used but did not specify whether these were nasal and/or throat swabs. Less commonly described testing procedures included sputum sampling (14% of articles) and blood sampling for antibody testing (14% of articles) and one included the use of temperature measurements. 9% of publications described the use of pre-screening questionnaires before sampling took place. 18% of articles did not specify the method of testing and 41% of articles described the use of a combination of the above methods.

Outcomes of interest

Outcomes of interest were divided into three categories: process outcomes, quality outcomes and resource use outcomes. No participant outcomes were measured quantitatively in any of the included studies, however the discussion section of many papers contained rich qualitative data describing participant outcomes such as staff and participant safety and wellbeing and service equity.

Process Outcomes

The process outcomes described comprise throughput, duration of test, decontamination time, time to don/doff PPE and waiting time. 77% of articles reported the number of people tested in a specified time period (Table 4). These figures were used to calculate the mean number of people tested each day.¹ Drive-through testing centres tested between 22-539 individuals per day. Indoor walk-through centres tested 9-500 people per day. One outdoor walk-through centre tested 300 people per day. Home visiting testing teams tested 6-15 people per day.

Some studies compared the throughput of different testing modalities. For similar settings, a higher throughput of individuals could be achieved in a drive-through setting compared to walk-through testing¹⁶ or home visiting testing^{5,6}. For the same time period, one indoor walk-through screening centre using negative pressure booths tested more patients than a walk-through centre with no negative pressure booths (>70 people per day compared to 9-10 people per day)²². Multiple booths could be installed and decontamination time between individuals could be reduced to 3-5 minutes from over 30 minutes. It is difficult to compare different studies, as several variables other than the testing modality can affect the number of people tested per day, such as the number of staff present, the procedures used and the number of individuals who could be tested in parallel.

Table 4: Number of people tested per day

Number tested per day (calculated)	Pre-screening (Questionnaire/Temperature only)	Testing (Samples taken)	Ref
Drive-through		22*	6
	107	45	20
		60	12
		192	30
		>100	4
	163	122	25
		200 (max 400)	29
		242	16
		460	14
		539	17
Drive-through and home visiting		2,000 country-wide	21
Drive-through and Walk-through		65 (range 11-127)	18
Indoor Walk-through centres		9-10 (no negative pressure booths) >70 (with negative pressure booths)	22
		41	16
		50-500	26
Home visiting testing		6 (max 11)*	6
		15	5
		15	19
Outdoor Walk-through centre		300	23

¹ *2 studies reported the time period per 'week' but did not state which days of the week were available for testing; a 7-day week was assumed for calculations

The mean duration of a drive-through test was between 3-15 minutes. One study reported a median time per test of 28 minutes (IQR 17-44 minutes)²⁰. Some centres allowed multiple people to be tested per vehicle, whereas others allowed only one person per vehicle. The layout of drive-through testing sites can allow several individuals to be tested at one time, for example one drive- and walk-through centre could test two patients every five minutes¹⁸.

27% of articles compared testing modalities, calculating a range of indicators for process outcomes (Table 5). Drive-through testing was found to be faster than walk-through testing using a negative pressure tent¹⁶ or a screening centre⁴. Testing using an outdoor walk-through centre (2 minutes per test) was faster than using traditional sample collection in a single negative-pressure isolation room (5 minutes per test)²⁴. Home visiting testing (30 minutes per test) was also quicker than transporting patients to hospital for tests with a specialist ambulance (<1 hour)⁶. However, waiting times, defined as the time between arrival at the drive-through centre and testing, was reported to be as high as 7 hours at peak volume²⁰.

Testing in an outdoor walk-through centre dramatically reduced time to don/doff PPE compared to traditional sample collection in a single negative-pressure isolation room (1 minute per patient compared to 21 minutes per patient)²⁴.

Several studies measured the time to disinfect equipment between individuals (decontamination time). The use of drive-through testing eliminated the need for a 30 minute decontamination time between patients in a walk-through centre using a negative pressure tent¹⁶. Drive-through and home visiting testing also required no decontamination time, compared to up to 6 hours of decontamination time needed if a patients was tested in the emergency department⁶.

Decontamination time was much shorter when using an outdoor walk-through centre (10 minutes per patient) compared to a single negative-pressure isolation room (35 minutes per patient). The use of negative pressure booths reduced decontamination time from >30 minutes with no negative pressure booths to 3-5 minutes between patients³¹.

Table 5: Differences in process outcomes when comparing different testing modalities

Ref	Testing Modality	PROCESS OUTCOMES			
		Throughput (mean number tested per day)	Duration of test (mean time per test)	Decontamination time (time to disinfect room between individuals)	Time to don/doff PPE
Chang ¹⁶	Drive-through	242	5-7 minutes	N/A	
	Walk-through (negative pressure tent)	41	30 minutes	At least 30mins between patients	
Hill ⁵	Drive-through	92 patients in 18 hours 15 patients per 3 staff members per day			
	Home visiting testing				
Kwon ⁴	Drive-through		10 minutes		
	Screening centre		30 minutes		
Lin ²⁴	Outdoor walk-through centre (multifunctional sample collection station)		2 minutes	10 minutes per patient	1 minute
	Traditional sample collection (single negative-pressure isolation room)		5 minutes	35 minutes per patient	21 minutes
Mark ⁶	Home visiting testing	79 in 2 weeks	30 minutes + <1 hour travel time	0	
	Drive-through	313 in 2 weeks		0	
	Specialist ambulance and hospital sampling	Not specified	<1h + overnight stay may be required	<6 hours	
Kim ²²	Indoor walk-through screening (negative pressure booths)	>70 per day		3-5 minutes	
	Indoor walk-through screening (no negative pressure booths)	9-10 per day		>30 minutes	

Quality Outcomes

Two different indicators were used to describe quality outcomes: the median time from referral to test and the test turnaround time. Only one study of home visiting testing calculated the median time from referral to test, which was 1 day with a maximum of 3 days⁶. One drive-through testing study calculated the test turnaround time, defined as the time between testing and communication of results. This was found to be 25 hours (IQR 21-29) in-house and 221 hours (IQR 161-269) if outsourced²⁵.

Resource Use Outcomes

Resource use outcomes were measured using cost per patient, use of PPE and impact on hospital closure. Home visiting testing reportedly costed much less (£55 per patient) than the use of a specialist ambulance and hospital sampling (£768 per patient)⁶ (Table 6).

Another study reported that staff in an outdoor walk-through centre used fewer items of PPE than staff working in negative-pressure isolation rooms²⁴. Similarly, drive-through testing can reduce PPE use (96% reduction in mask use, 97% reduction in gown use and 47% reduction in glove use) compared to emergency department based testing³⁰.

Table 6: Resource use outcomes

Ref	Testing Modality	RESOURCE USE OUTCOMES	
		Cost per patient	PPE use
Lin ²⁴	Outdoor walk-through centre (multifunctional sample collection station)		0 PPE items used*
	Traditional sample collection (single negative-pressure isolation room)		24 PPE items used
Mark ⁶	Home visiting testing	£55	
	Drive-through	Not specified	
	Specialist ambulance and hospital sampling	£768	
Ton ³⁰	Drive-through		1,152 masks, 960 gowns/pairs of gloves for 192 patients
	Emergency department testing		42 masks, 24 gowns, 504 pairs of gloves for 192 patients

One study reported the effect of a screening and testing clinic on maintaining the functioning of a tertiary hospital²³. Before the clinic was opened, an average of 36 beds per day were closed due to Covid-19 patients entering the hospital, whereas after the clinic was open and operating well, there was only one closure event (25 beds).

Discussion

This scoping review provides an overview of the literature describing the experience of implementing population testing for SARS-CoV-2 in high and upper-middle income countries. Whilst a range of modalities were reported, the most commonly evaluated were drive-through services using naso- and/oro-pharyngeal swabbing. Drive-through testing provided a rapid and scalable method of testing for COVID-19, reducing the risk of exposure to staff and patients within healthcare settings and minimising PPE use. However, this approach raises questions regarding equity of access for those who do not have access to a private vehicle or are not well enough to drive. The addition of other testing options such as home visiting, mobile testing or walk-through services may help address this issue.

However, the evidence base for population testing lacks robust studies. Many were simply an evaluation of a testing programme with a discussion of its advantages and limitations rather than robust research studies with control groups. The heterogeneous nature of the testing programmes described in the literature makes it difficult to compare between studies. No studies reported quantitative data on participant outcomes.

There is a paucity of published literature on the implementation of mass testing for SARS-CoV-2. There were no studies on mass testing that fit the inclusion criteria of this review, although a few papers have described mass testing of residents in facilities such as care homes and prisons^{32,33}. A recent review by the European Centre for Disease Prevention and Control highlights the need for further studies on the assessment and impact of mass testing³⁴. Indeed, some countries have rushed to adopt mass testing before the benefits, risks and costs of this approach is fully understood³⁵. It is therefore pertinent to draw on the international literature on population testing to inform decision making in order to ensure that testing is deployed in a safe and effective way as part of the overall COVID response³⁶.

A distinction needs to be made between population testing and screening for Covid-19. As Covid-19 is a new disease, it is unsurprising that definitions for screening and testing in this context have not yet been standardised and often appear to be arbitrary. We found the term 'screening' is loosely used in different ways in the literature, including the testing of symptomatic and/or asymptomatic individuals, assessment of risk factors via a questionnaire, and temperature measurement of individuals travelling past a screening post³⁷⁻³⁹.

Our literature search aimed to be comprehensive but timely and expedient. Whilst we used rigorous and transparent search methods, we had to limit our search to articles published in English. Therefore it is possible that some relevant studies have not been included. Although we did not formally assess the quality of included papers, much of the literature was not robust as stated earlier. There was also considerable heterogeneity between studies, which limits our ability to compare outcomes from different articles.

Further exploration is needed of population testing using different SARS-CoV-2 tests as the strengths and limitations of the various SARS-CoV-2 tests could influence the yield, cost-effectiveness and viability of the population testing programmes. Additionally, research into the wider consequences of testing programmes is needed especially on population behaviours as a result of testing. Finally, further study of the cost-effectiveness of population testing compared to other pandemic control measures is also required. As with most public health interventions, there is no 'one-size-fits-all' approach, and whatever population testing approach is adopted it will need to be tailored to the local context and target population.

Acknowledgements

This work was supported by funding from Public Health England.

Supplementary Information

Table 1: Average time per test

Test modality	Mean time per test	Ref
Drive-through	10 min per individual (1/3 shorter than conventional testing)	4
	15min appointments per household scheduled (max 5 per car)	14
	5-8 minutes	15
	28 minutes (median) (IQR 17-44).	20
	5-7 minutes	16
	6 lanes of cars in parallel, 3-5min per car	21
Drive-through and Walk-through	2 patients swabbed every 5 minutes	18
Indoor Walk-through centre	30 minutes	16
	5 minutes	24
Outdoor Walk-through centre	2 minutes	24
Home visiting testing	30 minutes	6

References

1. Walensky RP, Del Rio C. From Mitigation to Containment of the COVID-19 Pandemic: Putting the SARS-CoV-2 Genie Back in the Bottle. *JAMA - Journal of the American Medical Association*. 2020; 323:1889-90.
2. Department of Health and Social Care. Coronavirus (COVID-19): Scaling up our testing programmes 2020.
3. Choi S, Han C, Lee J, Kim SI, Kim IB. Innovative screening tests for COVID-19 in South Korea. *Clin Exp Emerg Med*. 2020; 7:73-7.
4. Kwon KT, Ko J-H, Shin H, Sung M, Kim JY. Drive-Through Screening Center for COVID-19: a Safe and Efficient Screening System against Massive Community Outbreak. *J Korean Med Sci*. 2020; 35.
5. Hill K, Campbell R, Mutch C, Koch O, Mackintosh C. Drive-through testing in COVID-19: experience from NHS Lothian. *Clinical Medicine*. 2020; 20:290-1.
6. Mark K, Steel K, Stevenson J, Evans C, McCormick D, Willocks L, et al. Coronavirus disease (COVID-19) Community Testing Team in Scotland: A 14-day review, 6 to 20 February 2020. *Eurosurveillance*. 2020; 25:2000217.
7. Health Foundation. Will COVID-19 be a watershed moment for health inequalities? 2020.
8. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*. 2005; 8:19-32.
9. Colquhoun HL, Levac D, O'Brien KK, Straus S, Tricco AC, Perrier L, et al. Scoping reviews: time for clarity in definition, methods, and reporting. *J Clin Epidemiol*. 2014; 67:1291-4.
10. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol*. 2018; 18:143.
11. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci*. 2010; 5:69.
12. Republic of Korea Ministry of Foreign Affairs. Standard operating model for "drive-thru" screening clinics. 2020; Available from: http://www.mofa.go.kr/eng/brd/m_22592/view.do?seq=1&srchFr=&srchTo=&p;srchWord=&srchTp=&multi_itm_seq=0&itm_seq_1=0&itm_seq_2=0&company_cd=&company_nm=.
13. Brammer C, Donohue S, Elwell T, Fishbein E, Forschino DA, Horne D, et al. Qualitative review of early experiences of off-site COVID-19 testing centers and associated considerations. *Healthcare*. 2020; 8:100449.
14. Appa A, Chamie G, Sawyer A, Baltzell K, Dippel K, Ribeiro S, et al. SARS-CoV-2 PCR and antibody testing for an entire rural community: methods and feasibility of high-throughput testing procedures. *medRxiv : the preprint server for health sciences*. 2020.
15. Lee E, Mohd Esa NY, Wee TM, Soo CI. Bonuses and pitfalls of a paperless drive-through screening and COVID-19: A field report. *Journal of microbiology, immunology, and infection = Wei mian yu gan ran za zhi*. 2020.
16. Chang MC, Seo W-S, Park D, Hur J. Analysis of SARS-CoV-2 Screening Clinic (Including Drive-Through System) Data at a Single University Hospital in South Korea from 27 January 2020 to 31 March 2020 during the COVID-19 Outbreak. *Healthcare (Basel, Switzerland)*. 2020; 8.

17. Drees ML, Papas MA, Corbo TE, Williams KD, Kurfuerst ST. Identifying community spread of COVID-19 via a free drive-through screening event. *Infection control and hospital epidemiology*. 2020;1-3.
18. Flynn EF, Kuhn E, Shaik M, Tarr E, Scattolini N, Ballantine A. Drive-Through COVID-19 Testing During the 2020 Pandemic: A Safe, Efficient, and Scalable Model for Pediatric Patients and Health Care Workers. *Academic pediatrics*. 2020; 20:753-5.
19. Goldberg SA, Bonacci RA, Carlson LC, Pu CT, Ritchie CS. Home-based Testing for SARS-CoV-2: Leveraging Prehospital Resources for Vulnerable Populations. *The western journal of emergency medicine*. 2020; 21:813-6.
20. Halalau A, Ditkoff J, Hamilton J, Sharrak A, Vanood A, Abbas A, et al. Emergency Center Curbside Screening During the COVID-19 Pandemic: Retrospective Cohort Study. *JMIR public health and surveillance*. 2020; 6:e20040.
21. Kim E. Drawing on Israel's Experience Organizing Volunteers to Operationalize Drive-Through Coronavirus Testing Centers. *Disaster Med Public Health Prep*. 2020:1-3.
22. Kim SI, Lee JY. Walk-Through Screening Center for COVID-19: an Accessible and Efficient Screening System in a Pandemic Situation. *J Korean Med Sci*. 2020; 35.
23. Kwon YS, Park SH, Kim HJ, Lee JY, Hyun MR, Kim HA, et al. Screening Clinic for Coronavirus Disease 2019 to Prevent Intrahospital Spread in Daegu, Korea: a Single-Center Report. *Journal of Korean medical science*. 2020; 35:e246.
24. Lin P-T, Ni T-Y, Chen T-Y, Su C-P, Sun H-F, Chen M-K, et al. Reducing the consumption of personal protective equipment by setting up a multifunctional sampling station in the emergency department to screen for COVID-19 infection in Taiwan. *Environmental health and preventive medicine*. 2020; 25:34.
25. Lindholm DA, Kiley JL, Jansen NK, Hoard RT, Bondaryk MR, Stanley EM, et al. Outcomes of Coronavirus Disease 2019 Drive-Through Screening at an Academic Military Medical Center. *Open forum infectious diseases*. 2020; 7:ofaa306.
26. Manuis CM, Loh M, Kwan J, Chua Mingzhou J, Teo HJ, Teng Kuan Peng D, et al. Bracing for impact: operational upshots from the National Centre for Infectious Diseases Screening Centre (Singapore) during the COVID-19 outbreak. *Journal of the American College of Emergency Physicians open*. 2020.
27. Rivkees SA, Roberson S. The Florida Department of Health STEPS Public Health Approach: The COVID-19 Response Plan and Outcomes Through May 31, 2020. *Public Health Reports*. 2020.
28. Seo E, Mun E, Kim W, Lee C. Fighting the COVID-19 pandemic: onsite mass workplace testing for COVID-19 in the Republic of Korea. *Annals of occupational and environmental medicine*. 2020; 32:e22.
29. Shah A, Challener D, Tande AJ, Mahmood M, O'Horo JC, Berbari E, et al. Drive-Through Testing: A Unique, Efficient Method of Collecting Large Volume of Specimens During the SARS-CoV-2 (COVID-19) Pandemic. *Mayo Clinic proceedings*. 2020; 95:1420-5.
30. Ton AN, Jethwa T, Waters K, Speicher LL, Francis D. COVID-19 drive through testing: An effective strategy for conserving personal protective equipment. *American Journal of Infection Control*. 2020; 48:731-2.
31. Kim E. Comparing the Operational Strategies of South Korea and Israel's Coronavirus Drive-Through Testing Centers and the Implications on Testing Capacity. *Risk management and healthcare policy*. 2020; 13:821-3.

32. Hatfield KM, Reddy SC, Forsberg K, Korhonen L, Garner K, Gulley T, et al. Facility-Wide Testing for SARS-CoV-2 in Nursing Homes - Seven U.S. Jurisdictions, March-June 2020. *MMWR Morbidity and mortality weekly report*. 2020; 69:1095-9.
33. Hagan LM, Williams SP, Spaulding AC, Toblin RL, Figlenski J, Ocampo J, et al. Mass Testing for SARS-CoV-2 in 16 Prisons and Jails - Six Jurisdictions, United States, April-May 2020. *MMWR Morbidity and mortality weekly report*. 2020; 69:1139-43.
34. European Centre for Disease Prevention and Control. Population-wide testing of SARS-CoV-2: country experiences and potential approaches in the EU/EEA and the United Kingdom 2020; Available from: <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-population-wide-testing-country-experiences.pdf>.
35. Gill M, Gray M. Mass testing for covid-19 in the UK. *BMJ*. 2020; 371:m4436.
36. Association of Directors of Public Health and the Faculty of Public Health. Joint Statement from the Association of Directors of Public Health and the Faculty of Public Health on Targeted Community Testing 2020 [cited 2020 04/12/2020]; Available from: <https://www.fph.org.uk/news-events/fph-news/joint-statement-from-the-association-of-directors-of-public-health-and-the-faculty-of-public-health-on-targeted-community-testing/>.
37. Viswanathan M, Kahwati L, Jahn B, Giger K, Dobrescu AI, Hill C, et al. Universal screening for SARS-CoV-2 infection: a rapid review. *Cochrane Database of Systematic Reviews*. 2020.
38. David N, Mash R. Community-based screening and testing for Coronavirus in Cape Town, South Africa: Short report. *African journal of primary health care & family medicine*. 2020; 12:e1-e3.
39. Seddighi H. The Performance of the Iranian Red Crescent by Launching Testing Centers for the Coronavirus Disease. *Disaster medicine and public health preparedness*. 2020:1-2.