Using ICD-10-based Social Determinants of Health Categories to Assess Patients Risk for Acute

Care Utilization

Authors: Peter H. Nguyen, MD[1], James Wang, MD[1], Pamela Garcia-Filion, PhD[1], Deborah

Dominick, LCSW[2], Hamed Abbaszadegan, MD, MBA[1,3], Graciela Gonzalez Hernandez, PhD[4],

Karen O'Connor[4], Shakaib U. Rehman, MD[1,3], Sarada S. Panchanathan, MD, MS[1*]

Affiliation:

1. Department of Biomedical Informatics, College of Medicine, University of Arizona

College of Medicine – Phoenix, 475 N 5[th] St., Phoenix, AZ, United States

2. Department of Social Work, Phoenix VA Healthcare System, 650 E Indian School Rd,

Phoenix, AZ, United States

3. Department of Clinical Informatics, Phoenix VA Healthcare System, 650 E Indian

School Rd, Phoenix, AZ, United States

4. Department of Biostatistics, Epidemiology, and Informatics, Perelman School of

Medicine, University of Pennsylvania, 3400 Civic Center Boulevard, Building 421,

Philadelphia, PA, United States

*Corresponding Author

Key Words: "Social Determinants of Health (SDoH)", "Population Health", "Risk Modeling",

"Electronic Health Records", "Quality Improvement"

ABSTRACT

Background: There has been an increasing recognition of the influence of social, behavioral, economic, and environmental factors on overall patient health. The purpose of this project was to leverage the ICD-10 codes to identify and link social determinants of health (SDoH) to patients with a high probability of utilizing acute care services and to determine if social service intervention reduced care utilization.

Methods: We analyzed retrospective data for active patients at a Department of Veterans Affairs Medical Center (VAMC) from 2015-2017. Eleven categories of SDoH were developed based on existing literature of the social determinants; the relevant ICD-10 codes were divided among these categories. Emergency Room (ER) visits, hospital admissions, and social work visits were determined for each patient in the cohort.

Results: In a cohort of 44,401 patients, the presence of ICD-10 codes within the EHR in the 11 SDoH categories was positively correlated with increased acute care utilization. Veterans with at least one SDoH risk factor were 71% (95%CI: 68% - 75%) more likely to use the ED and 71% (95%CI: 65%-77%) more likely to be admitted to the hospital. Utilization decreased with social service interventions.

Conclusion: This project demonstrates a potentially meaningful method to capture patient social risk profiles through existing EHR data in the form of ICD-10 codes, which can be used to identify the highest risk patients for intervention with the understanding that not all SDoH codes are uniformly used and some SDoHs may not be captured.

INTRODUCTION

Social determinants of Health (SDoH) describe several factors that have become growing areas of interest in healthcare due to their impact on patient health and well-being. The World Health Organization define SDoH as "[t]he conditions in which people are born, grow, live, work, and age"(1). While SDoH are commonly thought of only in terms of socioeconomic factors, the term actually broadly encompasses 1) health-related behaviors, 2) socioeconomic factors, and 3) environmental factors (2). There are a variety of factors that comprise SDoH such as tobacco use, diet and exercise, substance use, education, employment, income, family and social support, community safety, and physical environment safety (3).

In 2007, Dr. Steven Schroeder published an article in the New England Journal of Medicine in which he referenced the impact of social and behavioral factors on health and premature death. The combined contribution of social, environmental, and behavioral factors on a patient's risk for premature death was estimated to be as high as 60%, with genetics contributing 30%, and medical care providing only 10% (4). It is worth noting how low the impact of medical care was estimated to be relative to SDoH.

Ten years after that article, Dr. Sanne Magnan reiterated the importance of SDoH, referencing a model from the University of Wisconsin Population Health Institute's County Health Rankings & Roadmaps program (2). This model was designed to help identify the factors that impact health in terms of quality of life and longevity in order to stimulate discussion and action in the form of potential changes to policies and programs (5). The County Health Rankings model also illustrated the significant impact of the social determinants on health outcomes, but did not include genetics in its factors influencing health outcomes, as the purpose of this specific model was to look at modifiable factors that could be affected by policy changes. It is interesting to note

that, in the County Health Rankings model, modifying health behaviors, social & economic

factors, and physical environment could collectively contribute up to 80% towards improving

health versus only 20% from modifying clinical care issues. This research again strikingly shows

the relatively low impact of direct clinical care, suggesting that SDoH represent a high yield, yet

untapped potential for making significant improvements in patient outcomes in healthcare.

The literature has numerous examples describing the impact of different social determinants on

health outcomes. Braveman and Gottlieb observed that lower educational status and income were

associated with decreased life expectancy at age 25 and increased presence of chronic disease

respectively (6). Yelin et al. noted in the literature that poverty appeared to be associated with

worse systemic lupus erythematous disease damage (7). Joynt et al. suspected in 2012 that 30-

day readmission rates are driven more by a hospital's patient population and the resources of the

community in which it is located than internal factors relating to the hospital itself (8). This

appeared to be supported by a recent study by Hu et al. in 2018 demonstrating that a high area

deprivation index (ADI) score, which reflects how socioeconomically disadvantaged a

geographic region is, was associated with a significantly higher 30-day readmission risk than a

lower ADI score (9).

Despite the mounting evidence regarding the role of SDoH on health outcomes, progress in

integrating their use has been slow. One of the significant barriers in the past has been in

determining a good method to capture the patients' SDoH profile, though this has become easier

in recent years as ICD-10 codes have valid ways to represent many of them to some extent. More

recently, the American Medical Association and United Healthcare have collaborated to plan the

development of up to two dozen new ICD-10 codes to further identify SDoH risk and integrate it

into care (10). The purpose of this project was to leverage the ICD-10 codes to identify and link

SDoH to patients with a high probability of utilizing acute care services and to determine if

social service intervention reduced care utilization.

MATERIALS AND METHODS

Patient population

The data analysis was performed using patient encounter data for an active veteran patient

population from January 2017 to December 2017 for a VAMC in a major metropolitan area. The

patient cohort was comprised of veterans with an active encounter during two timeframes: in

2017 and from 2015 to 2016. Active encounter data from 2017 were compared to encounter data

in the previous two years to examine utilization patterns relative to presence of SDoH and social

services intervention. An additional inclusion criterion for this cohort was availability of a means

test data within the past five years; the means test includes data on income and employment.

Exclusion criteria

Test patients and deceased patients were excluded from the dataset. The Charlson Clinical

Comorbidity Index (CCCI) of co-morbidities was determined for each patient based on

encounter data in 2017 (11). This was used to measure the presence and the degree of major

chronic comorbidities (e.g., heart disease, cancer, etc.). Patients with a CCCI greater than zero

(i.e., presence of co-morbidities) were excluded from the analysis cohort to reduce bias towards

acute care utilization given burden due to major chronic comorbidities.

Data source

Patient encounter data were extracted from the Regional Clinical Data Warehouse for the

VAMC. Microsoft Structured Query Language (SQL) was used for all data querying purposes.

Identifying Social Determinants of Health Risk

A total of 11 SDoH risk factor categories were determined using previous SDoH work by

Schroeder and the Kaiser Family Foundation (3,4). The risk factors were organized by clustering

SDoH that were deemed applicable to the VAMC population and that could be readily extracted

from the EHR. Schroeder identified that social factors, environmental exposures, and behavioral

patterns were modifiable factors influencing health outcomes beyond health care itself. The

Kaiser Family Foundation identified 6 major categories which were further subdivided into 29

minor categories. The category of the Health Care System (including coverage and access to a

primary care physician) was excluded since the entire study was completed with active patients

within a single location within the VA system. The category of food was expanded to include

exercise, and categories that have with significant social and medical impact were added (mental

health challenges and tobacco, alcohol, and substance use). The final SDoH categories used for

this study are listed in Table 1.

Table 1. SDoH categories with sample ICD-10 codes and associated social services.

SDoH Category	Associated Social Services	ICD-10 codes	Example specific codes
Diet and Exercise	• Social Work • Caregiver Support Services	• E66 • Z71.3, Z71.82, Z72.3, Z72.4	E66.3 Overweight Z72.3 Lack of physical exercise
Employment	• Social Work • OEF/OIF Support Group (Military-to-veteran transitions) • Case Management	• Z56	Z56.0 Unemployment, unspecified Z56.2 Threat of job loss
Family/Social Support	• Peer Support	• Z63	Z63.0 Problems in relationship with spouse or partner Z63.31 Absence of family member due to military deployment

Tobacco Use	• Social Work • SARRTP (Substance Abuse Clinic) • Case Management	• F17.2 • Z71.6, Z72.0	F17.213 Nicotine dependence, cigarettes, in withdrawal Z71.6 Tobacco abuse counseling
Education	• Social Work • OEF/OIF Support Group	• Z55	Z55.0 Illiteracy and low-level literacy Z55.3 Underachievement in school
Community Support/Safety	• Peer Support • OEF/OIF Support Group • Community support • Caregiver Support Services	• Z65.1-Z65.3, Z65.8-Z65.9 • Z73.6, Z74, Z75.3-Z75.4	Z65.2 Problems related to release from prison Z75.4 Unavailability and inaccessibility of other helping agencies
Alcohol Use	• Social Work • SARRTP • Case Management	• F10 • Z71.41	F10.20 Uncomplicated Alcohol dependence Z71.41 Alcohol abuse counseling and surveillance of alcoholic
Income	• Social Work • OEF/OIF Support Group • Case Management	• Z59.5-Z59.9	Z59.5 Extreme poverty Z59.7 Insufficient social insurance and welfare support
Mental Health	• Social Work • Case Management	• F20.3, F20.9 • F40.01, F40.10, F40.11, F40.9 • F25, F29, F31 - F39 except F34.0	T14.91 Suicide attempt F20.9 Schizophrenia, unspecified
Substance Use	• Social Work • SARRTP • Case Management	• F19 • F11	F11.23 Opioid dependence with withdrawal F19.931 Other psychoactive substance use, unspecified with withdrawal delirium
Housing Insecurity	• Social Work • HUDVASH (Homeless Support Services) • Case Management	• Z59.0, Z59.1, Z59.4	Z59.0 Homelessness Z59.1 Inadequate housing

When reviewing patient records, one point was assigned per SDoH risk category in which at

least one corresponding ICD-10 code was present. A patient could be assigned a total possible

cumulative risk score of 11 points if they were found to have at least one ICD-

10 corresponding to each category. Multiple codes within the same category were therefore

considered to be one entity for the purposes of this work as the emphasis was placed more on

how many categories of SDoH were relevant rather than the number of codes per category were

present. This would prevent duplication of similar codes from over-estimating the impact

of SDoH due to concepts being associated with each other or overlapping in context (i.e. Z65.2 -

problems related to release from prison vs. Z65.1 - imprisonment and other incarceration).

Identifying Acute Care Utilization

Acute care utilization is defined for the purposes of this project as Emergency Room (ER) visits

and hospital admissions. The number of ER visits for patients in the cohort was queried with

SQL. Hospitalizations were queried by searching for ER disposition fields which corresponded

to hospital admissions from the ER since all hospitalizations at this facility are routed through

the ER. Both the number of ER visits and the number of resulting hospitalizations were separated

by year to correspond to years 2016 and 2017.

Identifying the Impacts of Social Services Utilization

The impact of social services utilization on acute care utilization was examined by ascertaining

social services utilization in 2017 and from 2015 to 2016, and then comparing acute care

utilization rates between patients that received any versus no social services utilization. The total

amount of utilization by social services were calculated and used in this analysis. Patients were

categorized as having either an increase (including newly seen) or decrease (including none) in

social service visits in 2017 compared to the prior time period (2015-2016). The patient cohort

was further narrowed to patients who had at least one SDoH risk factor based on ICD-10 analysis

above and excluded patients who had no acute care services in either 2016 or 2017.

Data Analysis

The t-test function was used in order to calculate statistical significance. Charts were created

with Microsoft Excel, and error bars represent 95% confidence intervals. STATA software was

used to perform logistic regression analysis to examine the relationship between

individual SDoH risk factors, as well as the total number of SDoH risk factors with ER visits and

hospitalizations.

RESULTS

Based on the initial inclusion criteria parameters, an initial cohort of 59,134 patients was

determined. After exclusion criteria stratification, the resulting cohort for January 2017 to

December 2017 was 44,401. Of this cohort, 9,379 patients had ER visits and 2,566 patients had

hospitalizations during this time interval. Of the total cohort of 44,401, the population had an

average age of 58 years +/- 17.4 years and were predominantly male (91%), non-Hispanic

(83.2%), and Caucasian (91.8%). In terms of SDoH risk factors in the cohort, 53.3% of patients

had at least one risk factor. Details about these demographics and number of SDoH risk factors

identified per individual are summarized in Tables 2 and 3, respectively.

Table 2: Summary demographics in the acute care utilization cohort.

Acute Care Utilization Analysis	
Cohort Size	44401
Age (Average/StDev)	58/17.4
Gender	
Male	91.0%
Female	9.0%
Ethnicity	
Hispanic	12.2%
Non-Hispanic	83.2%
Unknown / Unreported Ethnicity	4.6%
Race	
Caucasian	81.8%
Black	8.4%
Asian	0.9%
American Indian / Alaskan	1.9%
Native Hawaiian or Other Pacific	0.8%
Unknown / Unreported Race	6.3%

178 **Table 3. Number of SDoH Risk Factors for patient in acute care utilization cohort**

Number of Risk Factors	N
0	20746
1	14552
2	5344
3	2085
4	894
5	459
6	182
7	91
8	35
9	13

180 Patients who had more SDoH risk factors identified using the ICD-10 categories were likely to

181 have more ER visits and hospitalizations in a given year. Among patients with at least

182 seven SDoH risk factors, the annual ER visit rate was 26 times higher and the annual

183 hospitalization rate was 54 times higher compared to those without a SDoH. The acute care

184 utilization rates by number of SDoH categories is represented in Fig 1.

186
187 **Fig 1:** ER visit (a) and hospitalization (b) rates by number of SDoH risk factors for year 2017 for individuals with a
188 Charlson score of zero.
189

For each SDoH risk factor, the odds ratios (OR) for both ER visits and hospitalization were

greater than one and statistically significant, with ORs for each risk factor ranging from 1.77 to

13.01 for ER visits and 1.86 to 18.44 for hospitalizations (Table 4). Patients who had started to

use a social service or increased their use of social services had decreased rates of ER visits and

hospitalizations, whereas those who had not been seen by social services or decreased their

social service utilization had an increase in their annual rate of acute care visits, regardless of

social service utilized (Fig 2). Those who used more categories of social services were also more

likely to have decreased acute care utilization rates (Fig 3).

Table 4: Logistic regression analysis of the various individual SDoH risk factors, as well as the cumulative total risk score determining their odds ratios on the two outcomes of ER visit or hospital admission.

SDoH Risk Factor	Number of individuals with risk factor	Odds Ratio for ER Visits (95% CI)	Odds Ratio for Hospitalizations (95% CI)
Total	23256	1.71 (1.68, 1.75)	1.71 (1.65, 1.77)
Diet and Exercise	2892	3.19 (2.93, 3.48)	3.12 (2.62, 3.73)
Alcohol Use	1541	3.74 (3.35, 4.18)	6.15 (5.12, 7.39)
Tobacco Use	2016	2.83 (2.56, 3.14)	3.79 (3.13, 4.59)
Drug Use	437	5.74 (4.74, 6.95)	9.16 (6.98, 12.03)
Education	11	13.01 (3.82, 44.6)	18.44 (4.88, 69.61)
Employment	16620	1.77 (1.67, 1.87)	1.86 (1.63, 2.13)
Income	4786	2.27 (2.11, 2.45)	2.62 (2.24, 3.07)
Housing Insecurity	1476	5.41 (4.85, 6.03)	6.87 (5.73, 8.23)
Social Support	213	4.44 (3.36, 5.88)	3.22 (1.82, 5.66)
Community Safety	2635	4.53 (4.16, 4.94)	6.09 (5.21, 7.11)
Mental Health	6875	4.11 (3.86, 4.38)	4.41 (3.85, 5.05)

Fig 2: Average annual change in (a) ER visits and (b) hospitalizations in year 2017 compared to 2016 based on whether patients were newly seen (or had increased visits) or were not seen (or had decreased visits).

Fig 3: The average change in acute care utilization from year 2016 to year 2017 plotted against the number of distinct Social Services that saw the patient between ER visits (a) and hospitalizations (b) from one year to the next.

206
207
208

DISCUSSION

Interest in SDoH is increasing rapidly in healthcare and with good reason. As demonstrated by

our results and consistent with other studies, ER visits and hospitalizations were occurring at

higher rates with increased numbers of SDoH risk factors, which would confirm that there are

more factors outside of currently measured healthcare quality factors that influence outcomes

and other healthcare metrics (1-9). Having any of the 11 SDoH risk factor categories identified

via ICD-10 code in patients without known chronic illnesses was noted to increase the likelihood

of acute care utilization; however, certain risk factors appeared to have a more significant impact

than others.

On one end of the spectrum, drug use-based and housing insecurity-based risk factors, the

categories with higher impacts, had ORs of 5.74 and 5.41, respectively for ER visits and 9.16

and 6.87, respectively for hospitalizations. On the other end of the spectrum, employment-based

risk factors, the category with the lowest impact, had ORs of 1.77 and 1.86, respectively. This

may indicate what types of risk factors to focus on when planning future interventions and may

be helpful in earlier identification of individuals who would be more likely to utilize acute care.

Of note, education-based risk factors had ORs of 13.01 for ER visits and 18.44 for

hospitalizations. However, in the Veteran population, not only is education not generally coded,

all members will have at a minimum a high school diploma or equivalent or they would not have

been allowed to enter active duty. Thus, the education category is likely to be very

underrepresented in this study (only 11 total individuals with this risk factor) and may not be as

reliable in representing SDoH-associated healthcare disparities in the general population.

Social service utilization was found across the board to be positively impactful in reducing acute

care utilization, with case management consistently showing the highest level of impact in both

the ER visit and hospitalization rates. This makes sense as case management has one of the

largest reaches of all the specific social services (seven of 11 SDoH categories impacted) and

involves medical care coordination as well. Utilizing more types of social services also decreased

rates of acute care utilization likely for a similar reason.

Even with increasing interest in integrating SDoH, the most optimal approach to risk screening is

not clear. This project demonstrates that, within the specific healthcare ecosystem of a VA

Medical Center, ICD-10 diagnosis codes corresponding to SDoH risk factors, in addition

to employment and income data already collected as part of routine VA eligibility means

test, can potentially be used to screen for patients for high social risk since ICD-10 has already

been integrated into the EHR. Some efforts have been focused on detecting the presence

of SDoH mentioned in the unstructured portion of the EHR using natural language processing

(NLP) or using machine learning and NLP on a combination of structured and unstructured data

(12–21).

Limitations

While some SDoHs can indeed be ascertained from the EHR, it is important to note that a gap

(not quantified in this study) remains: some SDoHs are known, but not coded either due to a lack

of a correct code or providers not coding for them. For other SDoHs, there is no adequate code:

for example, food insecurity, overcrowded housing, low health literacy or living in an unsafe

neighborhood (important reason why people cannot exercise outside). For these, the only

recourse is extracting relevant mentions of them in the unstructured portions of the record. In

addition, there may be some ICD-10 codes that are missing from this analysis and certain ICD-

10 codes could have been allocated to other categories. Despite the limitations, the results of this

project remain promising in moving healthcare operations forward to integrate SDoH and

mitigate its negative effects.

The optimal method of capturing SDoH remains debatable and some factors may not necessarily

be captured by the method in this project. For example, legal troubles, cultural norms and

barriers, and suspicion about the healthcare system could be perceived as social risk factors,

however they would not neatly fall into the risk factor categories used in this project.

Furthermore, the identification of risk factors in this project is highly dependent on optimal data

entry by clinicians and healthcare employees. If clinicians do not document ICD-10 codes for a

patient with housing insecurity, level of education or issues with family support, this method will

only sub-optimally describe that patient's social risk profile. This limitation should not be

understated, particularly given the current challenge for healthcare workers to accurately

document encounters in the setting of increasing documentation requirements. For instance, the

high OR of acute care utilization for the education SDoH risk factor with a wide confidence

interval may be due to artifact from limited data availability. Also, specific to the VA system,

some data elements such as income level may be more readily available than in other healthcare

systems due to the nature of how VA benefits are determined. In addition, the cohort in this

project was predominantly male, Caucasian, and non-Hispanic and is not representative of the

general patient population in the United States.

Additionally, it should be stated that in the analysis of the utilization of social services, patients

merely having an encounter with a social service provider is being interpreted as a surrogate for

receiving some kind of social service benefit, such as resources, food, or housing. However, it

may be the case that having an encounter with a social service might result in no assistance being

available for a patient. Ultimately, it should be noted that the results of this project identify

correlations, not necessarily causations.

CONCLUSION

Given the increasing importance of the SDoH and their integration into care, the findings from this quality improvement project are important because they represent an efficient method in which SDoH can be captured and linked to health outcomes. Most EHR systems do not yet incorporate SDoH risk screening as part of their data collection, nor do many clinical workflows. The use of the ICD-10 codes does not require any new modification to EHRs. Furthermore, the impact of utilization by the various social services on acute care utilization is encouraging and suggests a means to make SDoH data actionable. Future work will revolve around operationalizing these findings through personalized healthcare and potential use of new modalities such mobile medical applications as well as improved methods of data capture such as NLP.

Acknowledgements

We would like to thank Amit Dahiya, Michael Smith, Don Copp, Pete Fredericks, and Daniel Garcia of the Clinical Informatics Analytics and Applications Department at the Phoenix VA Medical Center for their support with SQL, guidance on clinical data warehouse navigation, and data analysis support. We also acknowledge Dr. Umar Iqbal, Dr. Behnam Vahdati Nia, Dr. Ali Al-Yaqoobi, and Dr. Vikeen Patel from the University of Arizona College of Medicine – Phoenix for their input and support during the execution of this project.

REFERENCES

1. Social determinants of health [Internet]. [cited 2018 Mar 26]. Available from:

   https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-

   ageing/maternal-health/about/social-determinants-of-health--global

2. Magnan S. Social Determinants of Health 101 for Health Care: Five Plus Five. NAM

   Perspectives [Internet]. 2017 Oct 9; Available from: https://nam.edu/social-determinants-

   of-health-101-for-health-care-five-plus-five/

3. Heiman HJ, Artiga S. Beyond Health Care: The Role of Social Determinants in Promoting

   Health and Health Equity. 2015 Nov :11.

4. Schroeder SA. Shattuck Lecture. We can do better--improving the health of the American

   people. N Engl J Med. 2007 Sep 20;357(12):1221–8.

5. Explore Health Rankings | Measures & Data Sources [Internet]. County Health Rankings &

   Roadmaps. [cited 2018 Mar 28]. Available from:

   https://www.countyhealthrankings.org/explore-health-rankings/measures-data-sources

6. Braveman P, Gottlieb L. The social determinants of health: it's time to consider the causes

   of the causes. Public Health Rep. 2014 Feb;129 Suppl 2:19–31.

7. Yelin E, Trupin L, Bunde J, Yazdany J. Poverty, Neighborhoods, Persistent Stress, and

   Systemic Lupus Erythematosus Outcomes: A Qualitative Study of the Patients' Perspective.

   Arthritis Care Res (Hoboken). 2019 Mar;71(3):398–405.

8. Joynt KE, Jha AK. Thirty-day readmissions--truth and consequences. N Engl J Med. 2012

   Apr 12;366(15):1366–9.

9. Hu J, Kind AJH, Nerenz D. Area Deprivation Index Predicts Readmission Risk at an Urban

   Teaching Hospital. Am J Med Qual. 2018 Oct;33(5):493–501.

10. Lagasse J. AMA, UnitedHealthcare team up on billing codes that address social

determinants of health [Internet]. Healthcare Finance News. [cited 2019 Apr 19]. Available

from: https://www.healthcarefinancenews.com/news/ama-unitedhealthcare-team-billing-

codes-address-social-determinants-health

11. Lix LM, Manitoba Centre for Health Policy. Cancer data linkage in Manitoba: expanding

the infrastructure for research. 2016.

12. Erickson J, Abbott K, Susienka L. Automatic address validation and health record review to

identify homeless Social Security disability applicants. J Biomed Inform. 2018;82:41–6.

13. Hatef E, Rouhizadeh M, Tia I, Lasser E, Hill-Briggs F, Marsteller J, et al. Assessing the

Availability of Data on Social and Behavioral Determinants in Structured and Unstructured

Electronic Health Records: A Retrospective Analysis of a Multilevel Health Care System.

JMIR Med Inform. 2019 Aug 2;7(3):e13802.

14. Feller DJ, Bear Don't Walk Iv OJ, Zucker J, Yin MT, Gordon P, Elhadad N. Detecting

Social and Behavioral Determinants of Health with Structured and Free-Text Clinical Data.

Appl Clin Inform. 2020;11(1):172–81.

15. Lindemann EA, Chen ES, Wang Y, Skube SJ, Melton GB. Representation of Social History

Factors Across Age Groups: A Topic Analysis of Free-Text Social Documentation. AMIA

Annu Symp Proc. 2017;2017:1169–78.

16. Bettencourt-Silva JH, Mulligan N, Sbodio M, Segrave-Daly J, Williams R, Lopez V, et al.

Discovering New Social Determinants of Health Concepts from Unstructured Data:

Framework and Evaluation. Stud Health Technol Inform. 2020 Jun 16;270:173–7.

17.   Zhu VJ, Lenert LA, Bunnell BE, Obeid JS, Jefferson M, Halbert CH. Automatically

344   identifying social isolation from clinical narratives for patients with prostate Cancer. BMC

345   Med Inform Decis Mak. 2019 14;19(1):43.

346   18.   Bejan CA, Angiolillo J, Conway D, Nash R, Shirey-Rice JK, Lipworth L, et al. Mining 100

347   million notes to find homelessness and adverse childhood experiences: 2 case studies of

348   rare and severe social determinants of health in electronic health records. J Am Med Inform

349   Assoc. 2018 01;25(1):61–71.

350   19.   Feller DJ, Zucker J, Yin MT, Gordon P, Elhadad N. Using Clinical Notes and Natural

351   Language Processing for Automated HIV Risk Assessment. J Acquir Immune Defic Syndr.

352   2018 Feb 1;77(2):160–6.

353   20.   Dorr D, Bejan CA, Pizzimenti C, Singh S, Storer M, Quinones A. Identifying Patients with

354   Significant Problems Related to Social Determinants of Health with Natural Language

355   Processing. Stud Health Technol Inform. 2019 Aug 21;264:1456–7.

356   21.   Gold R, Cottrell E, Bunce A, Middendorf M, Hollombe C, Cowburn S, et al. Developing

357   Electronic Health Record (EHR) Strategies Related to Health Center Patients' Social

358   Determinants of Health. J Am Board Fam Med. 2017 Aug;30(4):428–47.