Vaginal bacterial load in the second trimester is associated

with early preterm birth recurrence: a nested case-control

study

Authors:
Laura Goodfellow,[1*] Marijn C. Verwijs,[2] Angharad Care,[1] Andrew Sharp,[1] Jelena Ivandic,[1]
Borna Poljak,[3] Devender Roberts,[3] Christina Bronowski,[2] A. Christina Gill,[2] Alistair C. Darby,[4]
Ana Alfirevic,[1] Bertram Muller-Myhsok,[1,5] Zarko Alfirevic,[1†] Janneke H.H.M. van de Wijgert[2,6†]
9
10
Affiliations:
1. Harris Wellbeing Research Centre, Department of Women's and Children's Health
University of Liverpool, United Kingdom
2. Department of Clinical Infection, Microbiology and Immunology, Institute of Infection,
Veterinary, and Ecological Sciences, University of Liverpool, Liverpool, United Kingdom.
3. Liverpool Women's NHS Foundation Trust, Liverpool, United Kingdom.
4. Centre for Genomic Research, University of Liverpool, Liverpool.
5. Department of Translational Research in Psychiatry, Max Planck Institute of Psychiatry,
Munich, Germany.
6. Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht,
Utrecht University, Utrecht, The Netherlands
*Corresponding author. Address: University Department, Liverpool Women's Hospital,
Crown Street, Liverpool, L8 7SS. Tel: 0151 7959557. Fax: 0151 795 9599 email:
l.goodfellow@liverpool.ac.uk
†Contributed equally

26
Running title (60 characters): Vaginal microbiota & recurrent early preterm birth

Word Counts:
Abstract: 249 words (max. 250)
Introduction: 371 words (max. 400)
Text: 3,491 words (max. 3,500)
Tables/figures: 3 tables/1 figure (max. 4)
Appendices: 2

Abstract

<u>Objective</u>: To assess the association between vaginal microbiome (VMB) composition and

recurrent early spontaneous preterm birth (sPTB)/preterm prelabour rupture of membranes

(PPROM).

<u>Design</u>: Nested case-control study.

<u>Setting</u>: UK tertiary referral hospital.

<u>Sample</u>: High-risk women with previous sPTB/PPROM <34^{+0} weeks gestation who had a

recurrence (n=22) or delivered at ≥37^{+0} weeks without PPROM (n=87).

<u>Methods</u>: Vaginal swabs collected between 15-22 weeks gestation were analysed by 16S

rRNA gene sequencing and 16S quantitative PCR.

<u>Main outcome measure</u>: Recurrent early sPTB/PPROM.

<u>Results</u>: 28/109 high-risk women had anaerobic vaginal dysbiosis, with the remainder

dominated by lactobacilli (*L. iners* 36/109, *L. crispatus* 23/109, or other 22/109). VMB type,

diversity, and stability were not associated with recurrence. Women with a recurrence,

compared to those without, had a higher median vaginal bacterial load (8.64 vs. 7.89 $\log_{10}$

cells/μl, adjusted odds ratio (aOR)=1.90, 95% confidence interval (CI)=1.01-3.56, p=0.047)

and estimated *Lactobacillus* concentration (8.59 vs. 7.48 $\log_{10}$ cells/μl, aOR=2.35, CI=1.20-

4.61, p=0.013). A higher recurrence risk was associated with higher median bacterial loads

for each VMB type after stratification, although statistical significance was reached only for

L. iners-domination (aOR=3.44, CI=1.06-11.15, p=0.040). Women with anaerobic dysbiosis or

L. iners-domination had a higher median vaginal bacterial load than women with a VMB

dominated by *L. crispatus* or other lactobacilli (8.54, 7.96, 7.63, and 7.53 $\log_{10}$ cells/μl,

respectively).

<u>Conclusions</u>: Vaginal bacterial load is associated with early sPTB/PPROM recurrence.

Domination by lactobacilli other than *L. iners* may protect women from developing high

bacterial loads. Future PTB studies should quantify vaginal bacteria and yeasts.

<u>Funding</u>: Wellbeing of Women, London, UK

<u>Keywords</u>: vaginal microbiome; *Lactobacillus*; spontaneous preterm birth; preterm

premature rupture of membranes.

<u>Tweetable abstract</u>: Increased vaginal bacterial load in the second trimester may be

associated with recurrent early spontaneous preterm birth.

Introduction

Previous spontaneous preterm birth (sPTB) or preterm prelabour rupture of membranes

(PPROM) is the strongest risk factor for recurrent sPTB/PPROM.[1] This may implicate a

genetic contribution[2] but does not rule out a contribution of persistent microbes and/or

inflammation in the female reproductive tract.[3]

16S ribosomal RNA (16S rRNA) sequencing was first used to assess the relationship between

the bacterial vaginal microbiota (VMB) and PTB in 2014.[4] Since then over 2000 pregnant

women have been assessed in 12 studies (Appendix A, Table A1) . The majority of these

studies suggested a contribution of the VMB composition to PTB, but results have been

inconsistent. The healthy vagina in pregnancy is dominated by *Lactobacillus* species but

contains up to 80 different *Lactobacillus* and other bacterial taxa, and the optimal ways of

summarising and categorising sequencing data to determine clinically relevant associations

with PTB are yet to be determined.[5,6] Studies to date have shown that increased relative

abundances of *L. crispatus, L. gasseri*, and *L. jensenii* were either not associated with PTB or

associated with a reduced PTB risk, and increased relative abundances of bacterial vaginosis

(BV)-associated anaerobes were either not associated with PTB or associated with an

increased PTB risk (Table A1). Results for *L. iners* relative abundance were inconsistent, with

6 studies showing no association, two studies an increased risk, and one study a decreased

risk (Table A1). Importantly, all studies except two[7,8] relied on compositional data, meaning

that an increase in the proportion of one taxon always leads to a decrease in the proportions

of other taxa.[9] Furthermore, an increase in a proportion does not necessarily indicate an

increase in absolute quantity. VMB researchers have only recently begun to semi-quantify

16S rRNA sequencing data by combining sequencing data with a quantitative PCR (qPCR) of

the 16S rRNA gene.[10] The two studies to date that included some quantification reported

opposite results, with one reporting a positive association between overall vaginal bacterial

load and PTB risk,[7] and the other a negative association.[8]

Our aim was to assess associations between VMB compositions and recurrent early

sPTB/PPROM (delivery at <34^{+0} weeks compared to delivery at ≥37^{+0} weeks) among high-risk

pregnant women who had had a previous early sPTB/PPROM, using compositional and semi-

quantitative 16S sequencing data.

Methods

Study design and population

Two cohorts of women at high and low-risk of PTB were enrolled at Liverpool Women's

Hospital, UK, from 1 June 2015 until 31 December 2017. The hospital's patient and public

engagement group helped guide the research and recruitment (Appendix A). High-risk was

defined as a history of sPTB or PPROM at 16^{+0}-33^{+6} weeks gestation, and low-risk as having

had at least one previous term birth (≥37^{+0} weeks) without PPROM and never having had a

PTB or PPROM. In the low-risk cohort, but not in the high-risk cohort, women were excluded

if they had had significant cervical surgery or previous obstetric or medical problems

(Appendix A). For this nested case-control study, we selected 109 women from the high-risk

cohort (N=137) and 145 women from the low-risk cohort (N=223) based on the pregnancy

outcome (sPTB or PPROM 16^{+0}-33^{+6} weeks or birth ≥37^{+0} weeks in the high-risk cohort and

birth ≥39^{+0} weeks in the low-risk cohort) and availability of valid VMB data (Figure 1). Birth

outcomes were classified independently by two obstetricians experienced in PTB

management. When there was a discrepancy between them, the case was reviewed by a

third experienced obstetrician until the team reached consensus. Participants with medically

indicated PTBs, or for whom it was unclear whether a birth was truly spontaneous, were

excluded. We used relevant parts of the CROWN initiative core outcome set for PTB

research[11] (Appendix A).

Clinical procedures

Participants were invited to two study visits at approximately 16 weeks (15^{+1}-18^{+6} weeks)

and 20 weeks (19^{+0}-23^{+0}) gestation. At each visit, a study obstetrician interviewed the

participant; collected high vaginal swabs during a speculum examination for VMB

assessments at the University of Liverpool Center for Genomic Research (HydraFlock

standard tapered swabs, Medical Wire and Equipment, Corsham, UK) and for quantitative

assessment of foetal fibronectin and culture by the local NHS laboratory; determined

cervical length; and did a 3D foetal ultrasound. All other procedures during pregnancy, and

all treatments, were in accordance with UK national guidelines.[12] Preterm birth prevention

therapy (cerclage, vaginal pessary, or vaginal progesterone) was offered if cervical length

was ≤ 25mm or based on clinician and patient preference in case of a large change in cervical

length between measurements. None of the low-risk women required an intervention but

71.6% (78/109) of the high-risk women did (Table 1). Initiation of this treatment was always

after the VMB sampling visits.

VMB laboratory assessments

DNA was extracted from one swab per woman per visit using lysozyme lysis and bead-

beating procedures combined with the Qiagen DNeasy Blood and Tissue kit (Qiagen,

Manchester, UK) (Appendix A).[13] The V3-V4 region of the 16S rRNA gene was amplified and

sequenced on an Illumina HiSeq 2500 instrument (Illumina, San Diego, CA) run in rapid

mode, 2x300bp using a 250PE and 50E kit. The panbacterial 16S rRNA gene copy

concentration per sample was determined using the BactQuant quantitative PCR (qPCR)

assay.[14] Molecular data processing steps are described in Appendix A.

We only used the first set of valid sequencing results for each participant in the analyses

because of the short time period between the two study visits, with one exception. In the

subset of participants with valid sequencing results for both visits (88 high-risk and 129 low-

risk women) we used data from both visits for the VMB stability analyses.

Creation and selection of VMB variables

Existing approaches

We systematically reviewed the published literature to identify VMB variables that were

associated with PTB in at least one study (Appendix A, Table A1). These included: richness

and Simpson diversity (1-D) as continuous variables; two categorical VMB composition

variables with each participant assigned to one category (the community state types (CSTs)

described by Ravel et al[15] and *Lactobacillus* groups (dominant, intermediate, and deplete)

based on *Lactobacillus* relative abundance);[16] stability groups, slightly modified from Romero

et al;[17] and the presence/absence of specific taxa of interest (listed in Table A1).

New approaches

The final three sets of VMB variables are based on our previous work and were applied to

PTB research for the first time.[18–21] The first is a categorical VMB composition variable that

we named 'VMB type'. These VMB types are similar to Ravel's CSTs[15] but with an increased

emphasis on the relative abundance of pathobionts (Appendix A). We defined pathobionts

as bacteria that are considered more pathogenic than BV-anaerobes and that often co-occur

with lactobacilli instead of BV-anaerobes, such as streptococci. Secondly, each non-minority

amplicon sequence variant (ASV; equivalent to a taxon) in each sample was allocated to one

of four 'bacterial groups' based on the published literature (Appendix B): lactobacilli; BV-

anaerobes; pathobionts; and a rest group of 'other bacteria'.[20] Within each sample, read

counts of ASVs belonging to the same bacterial group were summed. This resulted in four

continuous relative abundance variables (one for each bacterial group) per sample, which

sum to 1.0 for each sample. Finally, we converted these bacterial group relative abundances

into estimated concentrations (again four continuous variables, one for each bacterial

group) making use of the BactQuant results (Appendix A). The BactQuant data were also

used to determine estimated concentrations of taxa of interest (listed in Table A1) and the

total bacterial load in each sample.

Statistical analysis

Our primary comparisons were between high-risk women who gave birth ≥37^{+0} weeks

gestation and high-risk women who had a recurrent early sPTB or PPROM <34^{+0} weeks. Low-

risk women who delivered at ≥39^{+0} weeks gestation without PPROM were used as the

normal reference group because we had never before characterised VMB compositions in

pregnant women residing in Liverpool. Characteristics between these three groups were

compared by student's t-test for age, Mann-Whitney U test for other continuous variables,

and Fisher's exact test for binary and categorical variables. The various VMB variables

described above were compared between the two high-risk groups using unadjusted and

adjusted logistic regression, with adjustments for body mass index (BMI; as a quadratic term

due to the bimodal association with PTB), history of cervical surgery (none, single LLETZ, or

multiple LLETZ or knife cone biopsy), and smoking at the time of enrolment (yes/no).

Quartiles of the expected distributions of total vaginal bacterial load and total estimated

Lactobacillus concentration were generated using data from the low-risk reference group,

and high-risk participants were allocated to one of these quartiles. These newly created

categorical variables were also compared between the two high-risk groups of interest by

logistic regression. Finally, in an effort to differentiate between the effects of total vaginal

bacterial load and the types of bacteria that make up this load, women were stratified by

VMB type, and the logistic regression analyses were repeated for each stratum.

Results

We recruited 137 high-risk women and 223 low-risk women. Of 131 high-risk women with

VMB data that passed quality control, 22 had a recurrent early sPTB/PPROM and 87 had

term births $\geq37^{+0}$ weeks without PPROM (Figure 1); the remaining women were excluded

because they delivered between 34^{+0} and 37^{+0} weeks or had a birth initiated by a caregiver.

Of 214 low-risk participants with VMB data that passed quality control, 145 gave birth at

$\geq39^{+0}$ weeks without PPROM and were used as the normal reference group.

The participant characteristics of the three groups were similar (Tables 1 and A2), except for

those that are known risk factors for sPTB/PPROM: a higher proportion of high-risk women

with a recurrence, compared to those who delivered at term, had two or more previous

early sPTB/PPROM events (31.8% vs. 9.2%) and multiple previous LLETZ or knife cone

biopsies (13.6% vs. 1.1%). The median gestational age when the first valid VMB sample was

taken was slightly later in the low-risk group (16^{+5} weeks) than in the high-risk group (16^{+3}

weeks).

VMB compositions and vaginal bacterial loads in high- and low-risk pregnant women

The VMBs of over 70% of pregnant women, high- and low-risk, were dominated by

lactobacilli (defined as ≥75% relative abundance; Tables 2 and A3, heatmap in Figure A1). *L.*

iners domination was most common overall (VMB type Li; 33.0% of high-risk and 22.1% of

low-risk pregnant women), followed by *L. crispatus* domination (VMB type Lcr; 21.1% and

26.9%), and either domination by other lactobacilli (most commonly *L. gasseri* or *L. jensenii*)

or ≥50% bifidobacteria (VMB type Lo+BL; 20.1% and 25.5%). In addition, 12.4% of high-risk

and 12.8% of low-risk women had mild anaerobic dysbiosis (VMB type LA; defined as 25%-

75% lactobacilli with the remainder BV-anaerobes), and 12.8% of high-risk and 13.1% of low-

risk women had severe anaerobic dysbiosis (VMB type BV; ≥75% BV-anaerobes). Pathobionts

and 'other bacteria' were rarely present, and if present, only in low relative abundance or

estimated concentration (Tables 2 and A3).

The VMB of the majority of participants with samples available from both visits between

15^{+0} and 23^{+0} weeks gestation was stable (Table 2). About two thirds of the high-risk (65.9%)

and low-risk women (64.3%) were dominated by the same *Lactobacillus* species at both

visits, and 14.8% of high-risk and 22.5% of low-risk women continued to have some degree

of anaerobic dysbiosis. The remaining women switched *Lactobacillus* species (n=5) or shifted

from lactobacilli-domination to anaerobic dysbiosis (n=12) or vice-versa (n=17).

The median vaginal bacterial load in the low-risk reference cohort was 7.68 $\log_{10}$ cells/µl

(interquartile range (IQR) 6.80-8.35) and in the high-risk cohort 8.07 (IQR 7.28-8.92) $\log_{10}$

cells/µl (Table 2). High-risk women with mild or severe anaerobic dysbiosis (pooled together)

had higher median vaginal bacterial loads than women with domination by *L. iners, L.*

crispatus, or other lactobacilli/bifidobacteria (8.54, 7.79, 7.63, 7.53 $\log_{10}$ cells/µl,

respectively; p-values are <0.05 for all comparisons between dysbiosis and the other groups,

but are not significant for the comparisons between *L. iners*-domination and the other two

lactobacilli groups; Table A4).

Associations with early sPTB/PPROM recurrence – existing approaches

In the high-risk women, we did not identify any associations between recurrence and VMB

richness, Simpson (1-D) diversity, CSTs,[15] categorical *Lactobacillus* groups,[16] categorical

stability groups[17] (Table 2), and presence or relative abundance of common *Lactobacillus*

species (Table A3). *Ureaplasma* species and BVAB TM7-H1 were present in higher

proportions of high-risk women with a recurrence than in high-risk women with a term birth,

and *Dialister* species vice versa, but their relative abundances when present were very low

(Table A3). In contrast, *Bifidobacterium breve* was present in a lower proportion of women

with a recurrence, but its relative abundance when present was also low. The proportions of

low-risk women with these taxa present were similar to those in high-risk women without a

recurrence (Table A3, Figure A2A).

Associations with early sPTB/PPROM recurrence – new approaches

We did not identify any associations between recurrence and VMB type or estimated BV-

anaerobes or pathobionts concentrations (Table 2). However, high-risk women with a

recurrence had a higher overall vaginal bacterial load (8.64 vs. 7.89 $\log_{10}$ cells/µl, adjusted

odds ratio (aOR) 1.90, 95% confidence interval (CI) 1.01-3.56, p=0.047) and a higher total

estimated *Lactobacillus* concentration (8.59 vs. 7.48 $\log_{10}$ cells/µl, aOR 2.35, 95% CI 1.20-

4.61, p=0.013) than high-risk women without a recurrence (Table 2). The aOR was 1.63 (95%

CI 0.93-2.86, p=0.093) for each increase in total vaginal bacterial load quartile, and 2.61 (95%

CI 1.27-5.33, p=<0.001) for each estimated *Lactobacillus* concentration quartile (Table 2).

Table 3 shows estimated concentrations of total vaginal bacteria, the four bacterial groups,

L. cripatus, *L. iners*, and other lactobacilli in high-risk women with and without a recurrence

after stratification by VMB type. A trend towards higher recurrence risk with higher median

bacterial loads was seen for each VMB type after stratification but was only significant in

women with *L. iners*-domination: the median vaginal bacterial loads were 9.03 (IQR 8.02-

9.12) $\log_{10}$ cells/µl in women with a recurrence compared to 7.79 (IQR 6.93-8.65) $\log_{10}$

cells/µl in women with a term birth (aOR 3.44, 95%CI 1.06-11.15, p=0.011). The association

was no longer significant after adjustment for confounders (aOR 2.38, 95%CI 0.71-7.95,

p=0.160). Within each of the other VMB type strata, the median estimated concentrations in

high-risk women who had a term birth were similar to those in low-risk women, and non-

significantly lower than those in high-risk women who had a recurrence.

The majority of high-risk participants with an early sPTB/PPROM recurrence (17/22; 77.3%),

and about half of the high-risk participants without a recurrence (44/87; 50.6%), had at least

one VMB characteristic that was positively associated with recurrence. The following

positively associated VMB characteristics overlapped considerably in both high-risk groups

(Venn diagrams in Figure A2): highest quartile of total bacterial load, highest quartile of

estimated total *Lactobacillus* concentration, highest quartile of estimated *L. iners*

concentration, and presence of BVAB TM7-H1. The only positively associated VMB

characteristic that did not completely overlap with the highest quartile of bacterial and

estimated total *Lactobacillus* concentration was the presence of *Ureaplasma* species (this

was the case in high-risk pregnancies with and without a recurrence).

Discussion

Main Findings

This exploratory study suggests that high vaginal bacterial load in the second trimester is

associated with early sPTB/PPROM recurrence in high-risk pregnant women, irrespective of

the bacterial composition. Our statistical power was limited: only 22 women had an early

sPTB/PPROM recurrence. Therefore, not all associations in line with this conclusion reached

statistical significance, especially after reducing statistical power further by stratification.

However, we believe that this new hypothesis warrants further research for various reasons.

First, the recurrence risk increased for each increase in bacterial and *Lactobacillus*

concentration quartile (on a $\log_{10}$ scale). Second, the findings were consistent for each VMB

type after stratification: higher recurrence in high-risk women with higher median bacterial

loads for all VMB types (reaching significance for *L. iners*-domination), and similar median

bacterial loads in high-risk and low-risk women who had a term birth for all VMB types.

Third, estimated concentrations reached higher levels for anaerobic dysbiosis and *L. iners*-

domination than for domination by other lactobacilli or bifidobacteria. Our findings

therefore do not contradict earlier findings by other groups that indicated increased PTB risk

for anaerobic dysbiosis and high *L. iners* relative abundance.[8,22–24] In fact, the inconsistent

findings related to *L. iners* in the literature might be due to the lack of quantification in most

studies.

Interpretation in light of other evidence

To our knowledge, the potential association between the vaginal bacterial load and PTB has

only been assessed in two previous studies. Freitas et al[7] also showed a higher 16S gene

concentration in the second trimester (approximately 16 weeks gestation) in women who

had an sPTB <37 weeks compared to women with a term birth ≥37 weeks (8.1 vs. 7.8 $\log_{10}$

copies/swab, respectively). Conversely, Elovitz et al[8] reported a similar 16S gene

concentration in women who had a sPTB <37 weeks compared to women with a term birth

≥37 weeks in samples collected before 24 weeks gestation (approximately 8 $\log_{10}$

copies/swab in each group), and a lower concentration (approximately 7.5 vs 7.9 $\log_{10}$

copies/swab) in samples collected at 28 weeks gestation. High vaginal bacterial load may

only be associated with PTB when present prior to 24 weeks gestation, but additional

quantitative studies are needed.

Female genital tract inflammation is considered one potential cause of PTB.[25] BV (a clinical

condition caused by anaerobic dysbiosis) is associated with vaginal inflammation, and is a

very common condition in non-pregnant women.[26] Pregnant women are protected from BV

due to the high oestrogen levels during pregnancy.[27,28] In our study, the prevalence of

anaerobic dysbiosis, and the relative abundances and estimated concentrations of BV-

associated bacteria and pathobionts, were indeed low. This likely explains why we did not

identify any statistically significant associations between the presence or extent of anaerobic

dysbiosis and early sPTB/PPROM recurrence. However, the non-significant trends

consistently suggested increased risk. Another common inflammatory vaginal condition is

vulvovaginal candidiasis, for which pregnant women are at increased risk.[29] In our study,

only one woman (low-risk) reported having been treated for vulvovaginal candidiasis prior to

study enrolment. Future studies should systematically assess vaginal yeasts concentrations

in addition to bacterial concentrations in all participants, and determine their potential

contributions to early sPTB/PPROM recurrence.

In our study, *L. iners* was able to achieve higher bacterial loads than the other lactobacilli,

and was associated with early sPTB/PPROM recurrence. *L. iners* has a genome of just 1.3

Mbp,[30] strikingly smaller than the other common vaginal *Lactobacillus* species, and seems

fully adapted to the vaginal niche.[31] This may explain why it is able to grow to such high

densities in the vagina. *Lactobacillus* species are considered to be non-inflammatory when

present in the vagina,[32] but they might cause inflammation when they travel through the

cervical mucus plug into the upper genital tract. Ascension to the upper genital tract may be

increased when vaginal density is high. Studies have shown that *L. iners* with high expression

of clustered regularly inter-spaced short palindromic repeat (CRISPR) genes are present in

women with BV but not in women without BV.[33] CRISPR genes are the primary bacterial

defence against bacteriophages. The *L. iners* strains associated with high bacterial loads and

recurrent sPTB/PPROM in our study may differ from strains associated with lower bacterial

loads. This requires further research.

Two uncommon taxa (present in fewer than 20% of the women in the low risk group) were

non-significantly associated with early sPTB/PPROM recurrence in our study and were also

associated with PTB in other studies: *Ureaplasma*[7,24] and BVAB TM7-H1[6]. In women with

early sPTB/PPROM recurrence, the presence of *Ureaplasma* species in the vagina did not

completely overlap with having high bacterial loads and lactobacilli concentrations.

Ureaplasma species may therefore play a role in PTB that is independent from overall

354 vaginal bacterial composition and load. Our data confirms that *Bifidobacterium breve* may

355 have a protective effect against early sPTB/PPROM recurrence.[16]

357 *Strengths and Limitations*

358 Our study had some limitations, most notably limited statistical power and insufficient data

359 on vaginal yeasts, but also had strengths. Pregnancy outcomes were very carefully selected

360 and assessed, many VMB composition variables were created covering different aspects of

361 VMB composition, and most of these VMB variables were (semi)quantified. The latter turned

362 out to be crucial, as our most important findings concern bacterial concentrations. The

363 quantification method that we used has recently been validated for non-minority taxa in two

364 different studies.[7,8] We had initially planned to analyse early sPTB and early PPROM as two

365 separate outcomes because the aetiological pathways may differ,[34] and to exclude women

366 who used PTB prevention treatments,[35] but the limited statistical power did not permit this.

367 A final limitation is the lack of VMB data in the third trimester and/or closer to the birth.

369 ## Conclusion

370 Vaginal bacterial load in the second trimester, irrespective of the bacterial composition, was

371 associated with early sPTB/PPROM recurrence. Domination by lactobacilli other than *L. iners*

372 may protect women from developing high bacterial loads. These findings should be

373 confirmed in larger, longitudinal studies that incorporate quantification of vaginal bacteria

374 and yeasts. If they are confirmed, interventions that maintain a non-*iners* lactobacilli/

375 bifidobacteria-dominated VMB may protect women from inflammation-associated PTB.

Acknowledgements

We would like to thank all participants for their enthusiastic involvement in the study, in particular members of the Harris-Wellbeing Patient and Public Engagement group. We would also like to thank the Liverpool Women's Hospital for hosting the research, Tracy Ricketts at the Liverpool Women's Hospital for administrative support, University of Liverpool Centre for Genomic Research staff for 16S sequencing, and Jacques Ravel and Mike Humphrys at the University of Maryland School of Medicine, Institute for Genome Sciences, for the 16S qPCR assays.

Disclosure of interests

All authors have completed the ICMJE uniform disclosure forms and declare: LG, AC, AS, AA, BM-M, AA, ZA and JW received a grant from Wellbeing of Women charity to establish the Harris Wellbeing Research Centre that funded the submitted work. MW, aCG, CB, DR, JI and BP received no support from any organisation for the submitted work. All authors declare no financial relationships with any organizations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Contribution to authorship

JW, AS, DR, BM-M, AC, AA and ZA conceived the study, wrote the protocol and obtained funding. AC, JI, BP and LG contributed to the protocol and recruited participants. AD, JW, aCG, and CB developed the DNA extraction, sequencing, and bioinformatics methods. LG, AC and AS extracted the clinical data. LG performed the laboratory analysis. JW developed the VMB analysis methods, and LG and MV performed the data processing. LG performed the data analysis and wrote the initial draft, and JW, ZA and MV contributed to data interpretation and revised the paper. All authors approved the final manuscript.

Details of Ethics Approval

The study was approved by North West Research Ethics Committee - Liverpool Central, reference 11/NW/0720 on 4 November 2011. The collection of vaginal fluid for this component of the study was approved in amendment 4 on 19 October 2015.

Funding

The prospective cohort study was funded Wellbeing of Women as part of a charitable donation from Lord and Lady Harris to establish the Harris-Wellbeing PTB Research Centre, University of Liverpool. This covered administrative costs, laboratory analysis, salary for AC, study support costs for AC and LG. No additional funding was used. Wellbeing of Women were not involved in the conduct of the research or writing the paper.

References

1. Ferrero DM, Larson J, Jacobsson B, Di Renzo GC, Norman JE, Martin JN, et al. Cross-Country Individual Participant Analysis of 4.1 Million Singleton Births in 5 Countries with Very High Human Development Index Confirms Known Associations but Provides No Biologic Explanation for 2/3 of All Preterm Births. Luo Z-C, editor. PLoS One [Internet]. 2016 Sep 13 [cited 2019 Mar 11];11(9):e0162506. Available from: https://dx.plos.org/10.1371/journal.pone.0162506

2. Zhang G, Srivastava A, Bacelis J, Juodakis J, Jacobsson B, Muglia LJ. Genetic studies of gestational duration and preterm birth. Best Pract Res Clin Obstet Gynaecol [Internet]. 2018 Oct [cited 2019 Mar 11];52:33–47. Available from: http://www.ncbi.nlm.nih.gov/pubmed/30007778

3. Hwang J, Vinturache AE, Jacobsson B, Gyamfi-Bannerman C, Mysorekar IU. Maternal microbiome – A pathway to preterm birth. Semin Fetal Neonatal Med. 2016;21(2):94–9.

4. Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Bieda J, et al. The vaginal microbiota of pregnant women who subsequently have spontaneous preterm labor and delivery and those with a normal delivery at term. Microbiome [Internet]. 2014 Dec 27 [cited 2019 Mar 19];2(1):18. Available from: https://microbiomejournal.biomedcentral.com/articles/10.1186/2049-2618-2-18

5. Peelen MJ, Luef BM, Lamont RF, de Milliano I, Jensen JS, Limpens J, et al. The influence of the vaginal microbiota on preterm birth: A systematic review and recommendations for a minimum dataset for future research. Placenta [Internet]. 2019;79(March):30–9. Available from: https://doi.org/10.1016/j.placenta.2019.03.011

6. Fettweis JM, Serrano MG, Brooks JP, Edwards DJ, Girerd PH, Parikh HI, et al. The vaginal microbiome and preterm birth. Nat Med. 2019;25(6):1012–21.

7. Freitas AC, Bocking A, Hill JE, Money DM. Increased richness and diversity of the vaginal microbiota and spontaneous preterm birth. Microbiome. 2018;6(1):1–15.

8. Elovitz MA, Gajer P, Riis V, Brown AG, Humphrys MS, Holm JB, et al. Cervicovaginal microbiota and local immune response modulate the risk of spontaneous preterm delivery. Nat Commun [Internet]. 2019;10(1):1–8. Available from: http://dx.doi.org/10.1038/s41467-019-09285-9

9. Jian C, Luukkonen P, Yki-Jarvinen H, Salonen A, Korpela K. Quantitative PCR provides a simple and accessible method for quantitative microbiome profiling. bioRxiv [Internet]. 2018;478685. Available from: https://www.biorxiv.org/content/early/2018/11/27/478685

10. Tettamanti Boshier FA, Srinivasan S, Lopez A, Hoffman NG, Proll S, Fredricks DN, et al. Complementing 16S rRNA Gene Amplicon Sequencing with Total Bacterial Load To Infer Absolute Species Concentrations in the Vaginal Microbiome. mSystems. 2020;5(2):1–14.

11. Van't Hooft J, Duffy JMN, Daly M, Williamson PR, Meher S, Thom E, et al. A core outcome set for evaluation of interventions to prevent preterm birth. Obstet Gynecol. 2016;127(1):49–58.

12. National Institute for Health and Care Excellence. Antenatal care for uncomplicated pregnancies. Clin Guidel [Internet]. 2008;(March):1–55. Available from: https://www.nice.org.uk/guidance/cg62/resources/antenatal-care-for-uncomplicated-pregnancies-975564597445

13. Gill C, Van De Wijgert JHHM, Blow F, Darby AC. Evaluation of lysis methods for the extraction of bacterial DNA for analysis of the vaginal microbiota. PLoS One. 2016;11(9):1–16.

14. Liu CM, Aziz M, Kachur S, Hsueh PR, Huang YT, Keim P, et al. BactQuant: an enhanced broad-coverage bacterial quantitative real-time PCR assay. BMC Microbiol. 2012;12.

15. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, et al. Vaginal microbiome of reproductive-age women. Proc Natl Acad Sci U S A. 2011;108(SUPPL. 1):4680–7.

16. Brown RG, Al-Memar M, Marchesi JR, Lee YS, Smith A, Chan D, et al. Establishment of vaginal microbiota composition in early pregnancy and its association with subsequent preterm prelabor rupture of the fetal membranes. Transl Res [Internet]. 2019;1–14. Available from: https://doi.org/10.1016/j.trsl.2018.12.005

17. Romero R, Hassan SS, Gajer P, Tarca AL, Fadrosh DW, Nikita L, et al. The composition and
stability of the vaginal microbiota of normal pregnant women is different from that of non-
pregnant women. Microbiome. 2014;2(1):1–19.
18. van de Wijgert JHHM, Verwijs MC. Lactobacilli-containing vaginal probiotics to cure or prevent
bacterial or fungal vaginal dysbiosis: a systematic review and recommendations for future
trial designs. BJOG An Int J Obstet Gynaecol. 2019;1–13.
19. Verwijs MC, Agaba SK, Darby AC, van de Wijgert JHHM. Impact of Oral Metronidazole
Treatment on the Vaginal Microbiota and Correlates of Treatment Failure. Am J Obstet
Gynecol [Internet]. 2019;(August). Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0002937819310075
20. van de Wijgert JHHM, Verwijs MC, Gill AC, Borgdorff H, van der Veer C, Mayaud P.
Pathobionts in the Vaginal Microbiota: Individual Participant Data Meta-Analysis of Three
Sequencing Studies. Front Cell Infect Microbiol. 2020;10(April).
21. van de Wijgert JHHM, Verwijs MC, Agaba SK, Bronowski C, Mwambarangwe L, Uwineza M, et
al. Intermittent Lactobacilli-containing Vaginal Probiotic or Metronidazole Use to Prevent
Bacterial Vaginosis Recurrence: A Pilot Study Incorporating Microscopy and Sequencing. Sci
Rep [Internet]. 2020;10(1):1–15. Available from: http://dx.doi.org/10.1038/s41598-020-
60671-6
22. Kindinger LM, Bennett PR, Lee YS, Marchesi JR, Smith A, Cacciatore S, et al. The interaction
between vaginal microbiota, cervical length, and vaginal progesterone treatment for preterm
birth risk. Microbiome [Internet]. 2017;5(1):1–14. Available from:
http://dx.doi.org/10.1186/s40168-016-0223-9
23. Callahan BJ, DiGiulio DB, Goltsman DSA, Sun CL, Costello EK, Jeganathan P, et al. Replication
and refinement of a vaginal microbial signature of preterm birth in two racially distinct
cohorts of US women. Proc Natl Acad Sci U S A [Internet]. 2017;114(37):9966–71. Available
from:
http://www.ncbi.nlm.nih.gov/pubmed/28847941%0Ahttp://www.pubmedcentral.nih.gov/art
iclerender.fcgi?artid=PMC5604014
24. DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, et al. Temporal
and spatial variation of the human microbiota during pregnancy. Proc Natl Acad Sci U S A.
2015;112(35):11060–5.
25. Romero R, Espinoza J, Gonçalves LF, Kusanovic JP, Friel L, Hassan S. The Role of Inflammation
and Infection in Preterm Birth. Semin Reprod Med. 2007;25(01):21–39.
26. Kenyon C, Colebunders R, Crucitti T. The global epidemiology of bacterial vaginosis: A
systematic review. Am J Obstet Gynecol [Internet]. 2013;209(6):505–23. Available from:
http://dx.doi.org/10.1016/j.ajog.2013.05.006
27. Cruickshank R, Sharman A. The Biology of the Vagina in the Human Subject. BJOG An Int J
Obstet Gynaecol [Internet]. 1934 Apr 1;41(2):208–26. Available from:
https://doi.org/10.1111/j.1471-0528.1934.tb08759.x
28. Hay P. Bacterial Vaginosis as a Mixed Infection. In: Brogden K, Guthmiller J, editors.
Polymicrobial Diseases [Internet]. Washington (DC); 2002. Available from:
https://www.ncbi.nlm.nih.gov/books/NBK2495
29. Fidel PL, Sobel JD. Immunopathogenesis of recurrent vulvovaginal candidiasis. Clin Microbiol
Rev. 1996;9(3):335–48.
30. Macklaim JM, Gloor GB, Anukam KC, Cribby S, Reid G. At the crossroads of vaginal health and
disease, the genome sequence of Lactobacillus iners AB-1. Proc Natl Acad Sci U S A.
2011;108(SUPPL. 1):4688–95.
31. Van Der Veer C, Hertzberger RY, Bruisten SM, Tytgat HLP, Swanenburg J, De Kat Angelino-Bart
A, et al. Comparative genomics of human Lactobacillus crispatus isolates reveals genes for
glycosylation and glycogen degradation: Implications for in vivo dominance of the vaginal
microbiota. Microbiome. 2019;7(1):1–14.
32. Könönen E. Anaerobic Cocci and Anaerobic Gram-Positive Nonsporulating Bacilli. In: Bennett

J, Dolin R, Blaser M, editors. Mandell, Douglas, and Bennett's Principles and Practice of
Infectious Diseases. 8th ed. Chicago: Elsevier Inc.; 2015. p. 2781-2786.e2.

33.   Macklaim JM, Fernandes AD, Di Bella JM, Hammond JA, Reid G, Gloor GB. Comparative meta-
RNA-seq of the vaginal microbiota and differential expression by Lactobacillus iners in health
and dysbiosis. Microbiome. 2013;1(1):1–11.

34.   Capece A, Vasieva O, Meher S, Alfirevic Z, Alfirevic A. Pathway analysis of genetic factors
associated with spontaneous preterm birth and pre-labor preterm rupture of membranes.
PLoS One. 2014;9(9).

35.   Peelen MJ, Luef BM, Lamont RF, de Milliano I, Jensen JS, Limpens J, et al. The influence of the
vaginal microbiota on preterm birth: A systematic review and recommendations for a
minimum dataset for future research. Placenta [Internet]. 2019;79(December 2018):30–9.
Available from: https://doi.org/10.1016/j.placenta.2019.03.011

Tables, Figures, and Supplementary Materials Caption List

Table 1: Participant characteristics by pregnancy risk group and outcome

Table 2: Associations between VMB variables and recurrent early sPTB/PPROM in high-risk women

Table 3: Associations between estimated median concentrations of vaginal bacteria and early recurrent sPTB/PPROM stratified by VMB type

Figure 1: Participant flow

Appendix A: Supplementary methods, figures, and tables

Appendix B: Assignment of amplicon sequence variants to bacterial groups

Table 1: Participant characteristics by pregnancy risk group and outcome

		Low-risk pregnancy (LR) Term birth n=145		High-risk pregnancy (HR) Term birth n=87		Early sPTB or PPROM n=22		P* value LR vs HR	P* value HR term vs HR preterm
Age (years)	mean (SD)	31.1	(4.5)	30.0	(4.5)	31.4	(5.4)	0.145	0.211
BMI (kg/m2)¶	median (IQR)	24.5	(22-29)	25	(22-28)	27.5	(22-35)	0.619	0.139
Current smoker¶	number (%)	15	(10.6)	15	(17.4)	5	(23.8)	0.096	0.537
Ethnicity (%)	White	139	(95.9)	81	(93.1)	22	(100.0)	0.540	0.732
	Black	3	(2.1)	4	(4.6)	0	0.0		
	Asian	0	0.0	0	0.0	0	0.0		
	Other	2	(1.4)	0	0.0	0	0.0		
	Not recorded	1	(0.1)	2	(2.3)	0	0.0		
Parity (%)	0	0	0.00	9	(10.34)	1	(4.55)	0.000	0.017
	1	102	(70.34)	43	(49.43)	10	(45.45)		
	2	36	(24.83)	27	(31.03)	4	(18.18)		
	≥ 3	7	(4.83)	8	(9.20)	7	(31.82)		
Number of previous PPROM or sPTB 16+0-33+6 weeks (%)	1			79	(90.8)	15	(68.2)	na	0.012
	≥2			8	(9.2)	7	(31.8)		
Previous cervical surgery	Nil significant	131	(90.3)	79	(90.8)	17	(77.3)	na	0.033
	Single LLETZ	14	(9.7)	7	(8.0)	2	(9.1)		
	Multiple LLETZ or knife cone			1	(1.1)	3	(13.6)		
Gestational age at first valid VMB sample (weeks)	median and IQR	16+5	(16+2-17+1)	16+2	(16+0-16+6)	16+3	(15+6-17+3)	0.000	0.794
Cervical length¶ (mm) (median and IQR)	16 weeks	41	(36-47.1)	35.5	(31-42)	36	(26.5-43.5)	0.000	0.578
	20 weeks	41	(37-45)	35	(30-40)	30	(26-37)	0.000	0.097
qfFN¶ (ng/ml) (median and IQR)	16 weeks	7	(5-16)	7	(5-22)	9	(6-23.5)	0.766	0.410
	20 weeks	7	(5-12)	7	(5-14)	7	(6-12)	0.274	0.914
Preterm birth prevention treatment used after study visit	None	139	(100.0)	61	(70.1)	17	(77.3)	na	0.097
	Cervical cerclage			4	(4.6)	2	(9.1)		
	Progesterone			2	(2.3)	1	(4.5)		
	Arabin pessary			20	(23.0)	2	(9.1)		
Antimicrobial in pregnancy prior to sampling (%)	None	115	(79.3)	65	(74.7)	17	(77.3)	0.038	0.895
	Metronidazole or clindamycin	0	0.0	4	(4.6)	1	(4.5)		
	Other antibiotic	21	(14.5)	9	(10.3)	3	(13.6)		
	Clotrimazole	1	(0.7)	0	0.0	0	0.0		
	Not recorded/ unsure	8	(5.5)	9	(10.3)	1	(4.5)		
Gestational age at PPROM (weeks)	median and range					31+1	18+0-33+6	na	na
Gestational age at birth (weeks)	median and range	40+1	(39+0-41+6)	38+6	37+0-41+5	31+5	18+0-35+5	na	na
Birthweight¶ (g)	mean (SD)	3594	(439)	3234	(489)	1778	(673)		

Abbreviations: BMI, body mass index; HR, high-risk pregnancy; LR, low-risk pregnancy; na, not applicable; PPROM, preterm prelabour rupture of membranes; qfFN, quantitative foetal fibronectin level in cervicovaginal fluid; sPTB, spontaneous preterm birth.

* P values by Student's t-test for age, Mann-Whitney U test for BMI, cervical length, qfFN and gestational age at sampling, and Fisher's exact test for remainder of variables.

¶Missing values: BMI 1 each for LR and HR term; smoking 3 for LR, and 1 each for HR term and early preterm; birthweight 2 each for HR term and HR preterm. Cervical length at 16 weeks: HR term n=80, HR preterm n=20, LR n=136. Cervical length at 20 weeks: HR term n=79, HR preterm n=14, LR n=135. qfFN at 16 weeks: HR term n=77, HR preterm n=20, LR n=142. qfFN at 20 weeks: HR term n=75, HR preterm n=19, LR n=136.

⌂Parity was defined as number of previous pregnancies with live births or stillbirths ≥24^{+0} weeks gestation. Eligibility for the HR cohort included ≥1 sPTB or PPROM at 16^{+0} to 33^{+6} weeks gestation. Therefore, all HR women with parity=0 had at least one previous late miscarriage. Antimicrobial usage was only assessed in the pregnancy prior to sampling. Clotrimazole is available without medical review as an 'over the counter' medication in the UK, and so usage after the study visit was not possible to assess within our study design.

Table 2: Associations between VMB variables and recurrent early sPTB/PPROM in high-risk women

Basis for assessment			Low-risk pregnancy (LR) Birth≥39 weeks n=145			High-risk pregnancy (HR) All high-risk pregnancies n=109			Birth ≥37 weeks n=87			sPTB or PPROM <34 weeks n=22			p value* difference between HR≥37 vs. <34 weeks	Adjusted¶ logistic regression model of HR≥37 vs <34 weeks (n=106)		
			Median (IQR)	n	(%)	Median (IQR)	n	%	Median (IQR)	n	%	Median (IQR)	n	%		OR	95% CI	P
Relative abundance data (16S rRNA sequencing)	Global assessments	Richness (n)	7 (4-14)			7 (4-15)			7 (4-15)			8 (5-14)			0.35	0.99	(0.93-1.06)	0.785
		Simpson diversity (1-D)	0.38 (0.07-0.53)			0.30 (0.06-0.53)			0.39 (0.08-0.54)			0.19 (0.05-0.45)			0.60	0.29	(0.04-2.18)	0.232
	VMB type⌑	Lcr		39	(26.9)		23	(21.1)		17	(19.5)		6	(27.3)		Comparator		
		Li		32	(22.1)		36	(33.0)		28	(32.2)		8	(36.4)		0.59	(0.15-2.34)	0.455
		Lo+BL		37	(25.5)		22	(20.2)		17	(19.5)		5	(22.7)	0.696	0.81	(0.17-3.79)	0.793
		LA		18	(12.4)		14	(12.8)		13	(14.9)		1	(4.5)		0.16	(0.01-1.98)	0.154
		BV		19	(13.1)		14	(12.8)		12	(13.8)		2	(9.1)		0.48	(0.07-3.24)	0.451
	Community State Type	I (L. crispatus)		43	(29.7)		23	(21.1)		17	(19.5)		6	(27.3)		Comparator		
		II (L. gasseri)		16	(11.0)		11	(10.1)		9	(10.3)		2	(9.1)		0.38	(0.04-3.95)	0.421
		III (L. iners)		31	(21.4)		33	(30.3)		26	(29.9)		7	(31.8)	0.848	0.55	(0.13-2.24)	0.402
		IV (mixed non Lactos)		35	(24.1)		29	(26.6)		25	(28.7)		4	(18.2)		0.43	(0.09-2.01)	0.287
		V (L. jensenii)		20	(13.8)		13	(11.9)		10	(11.5)		3	(13.6)		0.94	(0.17-5.12)	0.944
	Group according to Lactobacillus relative abundance	Lactobacillus dominance (>75%)		105	(72.4)		80	(73.4)		61	(70.1)		19	(86.4)		Comparator		
		Intermediate (25-74.9%)		14	(9.7)		6	(5.5)		5	(5.7)		1	(4.5)	0.277	0.91	(0.09-9.36)	0.940
		Lactobacillus deplete (<25%)		26	(17.9)		23	(21.1)		21	(24.1)		2	(9.1)		0.29	(0.05-1.53)	0.140
	Stability: change in VMB between 16 and 20 weeks gestation⌑	Same Lactobacilli		83	(64.3)		58	(65.9)		46	(64.8)		12	(70.6)		Comparator		
		Different Lactobacilli		3	(2.3)		2	(2.3)		1	(1.4)		1	(5.9)		4.40	(0.19-104)	0.359
		Remain non-LD	n=129	29	(22.5)	n=88	13	(14.8)	n=71	12	(16.9)	n=17	1	(5.9)	0.579	0.42	(0.04-4.11)	0.457
		Non-LD to LD		7	(5.4)		10	(11.4)		8	(11.3)		2	(11.8)		0.96	(0.16-5.90)	0.965
		LD to non-LD		7	(5.4)		5	(5.7)		4	(5.6)		1	(5.9)		0.41	(0.02-7.21)	0.543
Relative abundance combined with BactQuant, producing estimated concentration in log10 cells/µl	Global assessment	Bacterial load	7.68 (6.80-8.35)	145	(100.0)	8.07 (7.28-8.92)	109	(100.0)	7.89 (7.22-8.80)	87	(100.0)	8.64 (7.47-9.08)	22	(100.0)	0.035	1.90	(1.01-3.56)	0.047
	Bacterial group	Total Lactobacillus	7.31 (6.42-8.01)	143	(98.6)	7.38 (6.76-8.17)	107	(98.2)	7.48 (6.87-8.45)	85	(97.7)	8.59 (7.47-8.87)	22	(100.0)	0.005	2.35	(1.20-4.61)	0.013
		Total BV associated	5.41 (1.91-7.03)	109	(75.2)	5.49 (0-7.65)	81	(74.3)	5.20 (0-7.65)	62	(71.3)	6.65 (5.09-8.36)	19	(86.4)	0.102	1.15	(0.96-1.38)	0.126
		Total pathobionts	0.00 (0-0)	34	(23.4)	0.00 (0-0)	24	(22.0)	0.00 (0-0)	20	(23.0)	0.00 (0-0)	4	(18.2)	0.596	0.93	(0.712-1.22)	0.596
		Total other bacteria	0.00 (0-4.24)	50	(34.5)	0.00 (0-4.93)	45	(41.3)	0.00 (0-4.84)	35	(40.2)	0.00 (0-5.12)	10	(45.5)	0.671	1.04	(0.86-1.25)	0.698
	Type of Lactobacillus	L. crispatus	5.88 (4.41-6.83)	125	(86.2)	6.22 (4.44-6.95)	89	(81.6)	6.17 (4.08-6.91)	71	(81.6)	6.32 (5.79-7.76)	18	(81.8)	0.319	1.02	(0.83-1.25)	0.843
		L. iners	4.65 (0-7.14)	82	(56.6)	5.47 (0-7.33)	76	(69.7)	6.55 (0-7.98)	63	(72.4)	7.42 (0-8.81)	13	(59.1)	0.684	0.97	(0.84-1.12)	0.644
		Other Lactobacilli	5.49 (3.77-6.91)	125	(86.2)	5.94 (4.24-7.07)	99	(90.8)	5.81 (4.21-7.09)	78	(89.7)	5.95 (5.34-7.61)	21	(95.5)	0.172	1.14	(0.88-1.47)	0.311
	Allocation of high risk participants to quartiles of low risk		Concentration															
	Quartile of total bacterial load (n, %)	1	<6.80	37	(25.5)		12	(11.0)		11	(12.6)		1	(4.5)				
		2	≥6.80-<7.68	36	(24.8)		34	(31.2)		28	(32.2)		6	(27.3)	0.093	1.63	(0.93-2.86)	0.087
		3	≥7.68-<8.35	35	(24.1)		18	(16.5)		17	(19.5)		1	(4.5)				
		4	≥8.35	37	(25.5)		45	(41.3)		31	(35.6)		14	(63.6)				
	Quartile of total Lactobacillus (n, %)	1	<6.42	36	(24.8)		13	(11.9)		12	(13.8)		1	(4.5)				
		2	≥6.42->7.31	36	(24.8)		25	(22.9)		24	(27.6)		1	(4.5)	0.018	2.61	(1.27-5.33)	0.009
		3	≥7.31->8.01	36	(24.8)		29	(26.6)		23	(26.4)		6	(27.3)				
		4	≥8.01	37	(25.5)		42	(38.5)		28	(32.2)		14	(63.6)				

Abbreviations: HR, high-risk pregnancy; LR, low-risk pregnancy; PPROM, preterm prelabour rupture of membranes; sPTB, spontaneous preterm birth; VMB, vaginal microbiota.
*P values by Mann-Whitney U test for continuous variables and Fisher's exact test for remainder. ¶Logistic regression adjusted for body mass index as quadratic term, history of cervical surgery (none/single LLETZ/multiple LLETZ and/or knife cone biopsy) and smoking (yes/no at time of visit). Adjusted analyses excluded two HR women with missing smoking data, one with missing BMI data, and variable numbers because of collinearity within the regression. ⌑VMB type definitions: Li=L. iners-dominated (≥75% lactobacilli with L. iners the most common); Lcr=L. crispatus-dominated (≥75% lactobacilli with L. crispatus the most common); Lo+BL=other lactobacilli- or Bifidobacterium dominated (Lo+BL; either≥75% lactobacilli with L. jensenii or L. gasseri the most common, or ≥50% Bifidobacterium); lactobacilli and anaerobes (LA; 25%- 75% lactobacilli); and BV=mixture of BV-anaerobes (<25% lactobacilli). ⌑VMB changes between 16 and 20 weeks were only assessed in the subset of participants with valid VMB assessments at both time points: LR n=129; n=71 HR term n=71, and HR preterm n=17.

Table 3: Associations between estimated median concentrations of vaginal bacteria and early recurrent sPTB/PPROM stratified by VMB type

VMB type¶			Low risk pregnancy (LR) Birth ≥39 weeks n=145 Median (IQR)	High risk pregnancy (HR) All high-risk pregnancies n=109 Median (IQR)	Birth ≥37 weeks n=87 Median (IQR)	sPTB or PPROM <34 weeks (early preterm) n=22 Median (IQR)	Unadjusted OR	95% CI	P	Adjusted* OR	95% CI	P
LCr		n	39	23	17	6	23			18		
	Bacterial load		7.75 (6.44-8.46)	7.63 (7.14-8.63)	7.55 (7.14-8.32)	8.20 (7.42-9.06)	1.67	(0.57-4.89)	0.349	1.35	(0.41-4.40)	0.623
	Bacterial group	Total Lactobacillus	7.73 (6.42-8.46)	7.63 (7.14-8.59)	7.55 (7.14-8.25)	8.20 (7.41-8.86)	1.64	(0.55-4.85)	0.375	1.32	(0.40-4.30)	0.649
		Total BV associated	4.74 (0-6.29)	4.62 (0-6.07)	4.29 (0-5.36)	5.24 (0-6.71)	1.09	(0.79-1.50)	0.601	0.87	(0.55-1.36)	0.535
		Total pathobionts	0.00 (0-0)	0.00 (0-0)	0.00 (0-0)	0.00 (0-0)	0.93	(0.49-1.75)	0.825	1.07	(0.48-2.39)	0.866
		Total other bacteria	0.00 (0-3.19)	0.00 (0-0)	0.00 (0-0)	0.00 (0-0)	1.01	(0.59-1.71)	0.980	0.97	(0.56-1.68)	0.904
	Type of Lactobacillus	L. crispatus	7.72 (6.41-8.45)	7.63 (7.12-8.37)	7.55 (7.12-8.22)	8.19 (7.41-8.80)	1.66	(0.55-5.02)	0.365	1.31	(0.40-4.27)	0.658
		L. iners	0.00 (0-4.23)	0.00 (0-4.81)	0.00 (0-5.32)	0.00 (0-0)	0.71	(0.41-1.21)	0.204	0.84	(0.49-1.42)	0.514
		Other Lactobacilli	5.09 (3.39-5.94)	5.34 (3.02-7.16)	4.64 (3.02-6.14)	6.20 (5.34-7.39)	1.17	(0.77-1.76)	0.457	1.03	(0.93-1.80)	0.930
Li		n	32	36	28	8	36			34		
	Bacterial load		7.64 (6.94-8.25)	7.79 (6.93-8.65)	7.79 (6.93-8.65)	9.03 (8.02-9.12)	3.44	(1.06-11.15)	**0.040**	2.38	(0.71-7.95)	0.160
	Bacterial group	Total Lactobacillus	7.61 (6.90-8.25)	7.96 (7.23-8.83)	7.79 (6.93-8.60)	8.91 (8.02-9.10)	3.40	(1.04-11.11)	**0.042**	2.37	(0.70-8.04)	0.165
		Total BV associated	4.55 (0-6.37)	5.16 (0-6.60)	4.75 (0-5.79)	6.77 (5.33-7.73)	1.45	(0.98-2.15)	0.061	1.57	(0.95-2.59)	0.079
		Total pathobionts	0.00 (0-0)	0.00 (0-0)	0.00 (0-0)	0.00 (0-2.31)	1.21	(0.84-1.74)	0.307	1.15	(0.69-1.91)	0.590
		Total other bacteria	0.00 (0-3.61)	0.00 (0-4.26)	0.00 (0-3.38)	4.07 (0-5.69)	1.39	(1.00-1.94)	0.050	1.43	(0.95-2.16)	0.085
	Type of Lactobacillus	L. crispatus	5.89 (3.75-6.60)	6.19 (4.45-6.84)	6.02 (3.83-6.78)	6.38 (6.12-7.79)	1.26	(0.82-1.94)	0.295	1.09	(0.69-1.72)	0.695
		L. iners	7.52 (6.90-8.12)	7.93 (7.13-8.79)	7.73 (6.93-8.59)	8.82 (8.02-9.07)	3.35	(1.03-10.86)	**0.044**	2.33	(0.70-7.76)	0.169
		Other Lactobacilli	4.97 (0-6.75)	5.59 (4.08-6.59)	5.40 (4.41-7.30)	5.81 (4.08-6.59)	1.20	(0.78-1.85)	0.398	1.06	(0.69-1.63)	0.782
Lo or BL		n	37	22	17	5	22			19		
	Bacterial load		7.12 (6.32-7.69)	7.53 (7.24-8.64)	7.48 (7.21-8.73)	7.64 (7.46-8.61)	1.36	(0.48-3.86)	0.563	3.47	(0.61-19.7)	0.161
	Bacterial group	Total Lactobacillus	7.12 (6.08-7.69)	7.45 (7.10-8.61)	7.36 (6.94-7.89)	7.61 (7.42-8.61)	1.68	(0.54-5.29)	0.371	5.96	(0.65-54.5)	0.114
		Total BV associated	4.64 (0-5.59)	4.29 (0-6.19)	0.00 (0-4.94)	6.36 (6.31-6.58)	1.84	(0.97-3.50)	0.063	model failed		
		Total pathobionts	0.00 (0-2.98)	0.00 (0-4.11)	0.00 (0-4.11)	0.00 (0-0)	0.87	(0.53-1.41)	0.560	0.66	(0.23-1.87)	0.430
		Total other bacteria	0.00 (0-5.01)	1.13 (0-5.93)	0.00 (0-5.95)	5.09 (0-5.12)	1.05	(0.76-1.44)	0.769	1.21	(0.77-1.91)	0.410
	Type of Lactobacillus	L. crispatus	5.17 (4.00-5.88)	5.55 (4.44-6.54)	5.52 (5.26-6.54)	5.59 (0-6.16)	0.85	(0.59-1.21)	0.362	0.83	(0.53-1.29)	0.410
		L. iners	0.00 (0-6.41)	0.00 (0-7.09)	4.42 (0-7.09)	0.00 (0-0)	0.84	(0.61-1.16)	0.297	0.87	(0.58-1.30)	0.497
		Other Lactobacilli	6.99 (5.97-7.51)	7.46 (6.84-8.51)	7.24 (6.93-7.84)	7.61 (7.42-8.30)	1.87	(0.58-6.05)	0.293	8.97	(0.66-121)	0.099
LA or BV		n	37	28	25	3	28			27		
	Bacterial load		7.93 (7.44-8.52)	8.54 (8.11-9.22)	8.31 (8.07-9.16)	9.15 (9.11-9.28)	9.46	(0.28-318)	0.210	12.42	(0.31-501)	0.182
	Bacterial group	Total Lactobacillus	6.98 (6.30-7.45)	7.38 (6.63-8.37)	7.25 (6.56-8.19)	8.26 (8.15-8.86)	4.57	(0.54-38.5)	0.162	5.49	(0.17-63.1)	0.172
		Total BV associated	7.93 (7.19-8.51)	8.41 (8.07-9.19)	8.18 (8.07-9.15)	9.05 (8.83-9.24)	6.58	(0.33-133)	0.219	9.93	(0.42-237)	0.156
		Total pathobionts	0.00 (0-0)	0.00 (0-3.92)	0.00 (0-4.49)	0.00 (0-0)	model failed			model failed		
		Total other bacteria	0.00 (0-4.40)	4.57 (0-6.07)	4.59 (0-5.96)	0.00 (0-6.30)	0.83	(0.54-1.28)	0.402	0.70	(0.30-1.62)	0.407
	Type of Lactobacillus	L. crispatus	4.98 (0-5.93)	5.21 (0-6.41)	4.88 (0-6.60)	5.79 (0-5.90)	1.04	(0.69-1.55)	0.859	0.84	(0.48-1.46)	0.537
		L. iners	5.65 (0-7.11)	6.68 (4.80-8.21)	6.56 (4.74-7.98)	8.26 (8.15-8.86)	4.74	(0.49-45.6)	0.179	9.82	(0.35-274)	0.179
		Other Lactobacilli	4.79 (2.37-5.84)	5.32 (4.12-6.04)	5.34 (4.01-6.06)	5.20 (5.10-5.44)	1.09	(0.64-1.85)	0.756	1.03	(0.55-1.93)	0.916

Abbreviations: 95% CI, 95% confidence interval; HR, high-risk pregnancy; IQR, interquartile range; LR, low-risk pregnancy; OR, odds ratio; PPROM, preterm prelabour rupture of membranes; sPTB, spontaneous preterm birth; VMB, vaginal microbiota. *Logistic regression adjusted for body mass index as quadratic term, history of cervical surgery (none/single LLETZ/multiple LLETZ and/or knife cone biopsy) and smoking (yes/no at time of visit). Adjusted analyses excluded two HR women with missing smoking data, one with missing BMI data, and variable numbers because of collinearity within the regression. Participants were included in these analyses even if a taxa was not present in that sample (with concentration=0). ¶VMB type definitions: Li=*L. iners*-dominated (≥75% lactobacilli with *L. iners* the most common); Lcr=*L. crispatus*-dominated (≥75% lactobacilli with *L. crispatus* the most common); Lo+BL=other lactobacilli- or *Bifidobacterium* dominated (Lo+BL; either≥75% lactobacilli with *L. jensenii* or *L. gasseri* the most common, or ≥50% *Bifidobacterium*); lactobacilli and anaerobes (LA; 25%- 75% lactobacilli); and BV=mixture of BV-anaerobes (<25% lactobacilli).