Sleep disorders predict the one-year onset, persistence, but not remission of psychotic experiences in 10-11 year old children: a longitudinal analysis of the ABCD cohort data

Short title:

Sleep disorders and psychotic experiences in children

Authors:

Sarah Reeve[1], Vaughan Bell[1,2]

Affiliations:

[1] Department of Clinical, Educational, and Health Psychology, University College London, UK

[2] South London and Maudsley NHS Foundation Trust, London, UK

Corresponding author:

sarah.reeve.18@ucl.ac.uk

Abstract 258

Word count: 3862

Abstract

Sleep problems have been reliably associated with psychotic experiences in adults and have been suggested as target for intervention. However, the relationship between sleep disorder and psychotic experiences in children has not been extensively studied despite the potential for guiding intervention. The Adolescent Brain Cognitive Development (ABCD) dataset, containing baseline and one-year follow-up data of over 11,000 10-11 year olds, was utilised to investigate this relationship. More specifically, a set of pre-registered multi-level regression models were applied to test whether a) baseline sleep disorder predicts baseline psychotic experiences cross-sectionally; b) baseline sleep disorder predicts psychotic experiences one year later; c) the persistence of sleep disorder predicts the persistence psychotic experiences at one year; d) the remission of sleep disorder predicts the remission of psychotic experiences. After controlling for potential confounders, sleep disorder was associated with psychotic experiences cross-sectionally (OR=1.40, 95% CI 1.20-1.63), at one-year follow-up (OR=1.32, 95% CI 1.11-1.57), and the persistence of sleep disorder predicted the persistence of psychotic experiences (OR=1.72, 95% CI 1.44-2.04). However, remission of sleep problems did not predict remission of psychotic experiences (OR=1.041, 95% CI 0.80-1.35). In all models where an association was found, sleep was one of the two strongest predictors of psychotic experiences (with stimulant medication being the other). The results indicate that sleep problems in children are common and strongly associated with psychotic experiences but the lack of co-remission raises questions about the mechanism of association. However given existing evidence in adults, further investigation and interest in sleep as a preventative mental health intervention in this age group is warranted.

Introduction

Sleep disorders are hypothesised to be a significant causal factor in the development and maintenance of psychosis (Reeve et al., 2015; Waite et al., 2019). Experimental research supports that sleep disruption increases psychotic experiences (Reeve, Emsley, et al., 2018), and that treating sleep problems reduces psychotic experiences in adults (Freeman et al., 2017). While the relationship between sleep and psychotic symptoms is thought to be bidirectional, longitudinal studies have indicated that sleep problems predict later psychotic symptoms to a greater extent than vice versa (Hennig & Lincoln, 2017; Reeve, Nickless, et al., 2018). These and other findings have raised the possibility of targeting sleep problems as a preventative, early intervention in mental health (Freeman et al., 2017). This proposal is further supported by sleep problems acting as a generic contributor to a range of mental health problems in addition to (or perhaps as part of) their relationship with psychosis (Freeman et al., 2020), including during childhood and adolescence (Gradisar et al., 2020).

This raises the question of how early this relationship between sleep and psychosis exists. It is known that psychotic-like experiences are roughly 10x more common in children than they are in adults (reported by 66% of 7-8 year old children compared to 5.8-7.2% of adults; (Laurens et al., 2012; Linscott & van Os, 2013; McGrath et al., 2015). Clearly not all childhood psychotic-like experiences are pathological (Maijer et al., 2019), nevertheless persistence, distress, and amount of different psychotic-like experiences reported in childhood is associated with higher risk of developing adult psychotic disorder (Dominguez et al., 2011; Fisher et al., 2013) and a predictor of poor physical and mental health more generally later in life (Davies et al., 2018; Kelleher et al., 2012; Trotta et al., 2020). The age at which psychotic-like experiences are reported is also positively related to risk of later psychotic disorder. Reporting psychotic-like experiences at age 10-11 is associated with a 5x increased likelihood of developing a later psychotic disorder (Fisher et al., 2013), whereas reporting these experiences by ages 14-17 is associated with a 10x increased

likelihood (Dominguez et al., 2011). This set of findings have been encompassed under a 'proneness to persistence' model of psychosis, with these non-clinical psychotic experiences existing on a continuum with psychotic disorders (Linscott & van Os, 2013).

The relationship between sleep problems and psychotic experiences in children is less well-researched than in adults (Reeve et al., 2015). However, many studies have demonstrated that children or adolescents with sleep problems (including nightmares, insomnia, or oversleeping) are more likely to report psychotic experiences (Jeppesen et al., 2014; Koopman-Verhoeff et al., 2018; Y. J. Lee et al., 2012). Furthermore, longitudinal observational studies support that insomnia is specifically linked to paranoia (Bird et al., 2017; Hennig & Lincoln, 2017), with Hennig and colleagues additionally showing that the relationship seems primarily from insomnia to paranoia rather than vice versa. Several large population cohort studies have reported that sleep problems during childhood predict psychotic experiences in teenage years (Fisher et al., 2014; Morales-Muñoz et al., 2020; Thompson et al., 2015). Furthermore, as in adults, sleep problems in children are linked to a wide range of other negative mental and physical health outcomes (Carskadon & Barker, 2020; Dong et al., 2019; Gregory & Sadeh, 2012; Zhang et al., 2017).

Despite this evidence indicating that sleep problems have an important role in the onset of psychotic experiences during childhood and adolescence, it is less clear whether sleep problems contribute to persistence of psychotic experiences in this age group. This is particularly important given that persistence is a particularly strong predictor of poor outcome. The effect of remission of sleep problems on psychotic experiences is also unknown – but if improvements in sleep are associated with a reduction in psychotic experiences, this would provide evidence suggesting sleep interventions to reduce psychotic experiences could be viable.

The current study

We used the Adolescent Brain and Cognitive Development study cohort data (release 2.0.1) to investigate the relationship between sleep and psychotic experiences in children aged 9-11 to address a set of pre-registered hypotheses:

1. Do sleep disorders and psychotic experiences co-occur in childhood?
2. Do sleep disorders predict later onset of psychotic experiences?
3. Do sleep disorders predict persistence of psychotic experiences?
4. Does remission of sleep disorders predict remission of psychotic experiences?

These were examined use a pre-registered set of linear and logistic regression models applied to the baseline and one-year follow up data using a pre-registered analysis plan with all analysis available as open code: https://osf.io/8ks72/

Method

Recruitment

The Adolescent Brain and Cognitive Development (ABCD; https://abcdstudy.org/) 2.0.1 release study data was used in the current analysis (Feldstein Ewing & Luciana, 2018; Volkow et al., 2018). The ABCD is an ongoing longitudinal cohort study aimed at recruiting a representative sample of US children. At the initial stage, all families of children aged 9-10 in geographic catchment area of study sites across the USA were contacted via schools with information about the study. Families that volunteered were then contacted by the research team and screened for inclusion. Participants were purposively recruited to fulfil specific representativeness requirements to match sociodemographic factors in the US, for example utilising targets for each of five major race/ethnicity classifications (White, African-American, Hispanic, Asian All Other). The 2.0.1 release includes baseline data on 11,873 individuals, and one year follow-up from 4,951 of those participants. Full details of the ABCD study design are

available in a special issue of Developmental Cognitive Neuroscience (Feldstein Ewing &
Luciana, 2018).

Measures

Psychotic experiences

Children's psychotic experiences were assessed using the Prodromal Questionnaire – Brief Child
version (PQBC). This measure was adapted for and validated within the current ABCD dataset
showing high reliability (Cronbach's alpha = 0.863 for total score and 0.873 for distress subscale;
(Karcher et al., 2018). The PQBC is a self-report questionnaire assessing presence and distress
associated with psychotic like symptoms in children. For each psychotic-like experience the child
is asked if they experience it (yes/no), and if they do, how much it distresses/bothers them on a
pictographic 1 to 5 scale showing a human cartoon figure in various levels of distress. This
questionnaire yields two outcome variables – the sum of symptoms endorsed (range 0-21) and
the sum of distress reported (range 0-126).

For the current study, both continuous and dichotomous outcomes of the sum of symptoms
were used. For the dichotomous outcome the symptom score was transformed to a categorical
0-1 variable where 1 indicates presence of at least one psychotic symptom and 0 indicates no
psychotic symptoms present. When applied to baseline and follow up this was then used to
derive variables indicating 'new onset' of psychotic symptoms (i.e. 0 at baseline, 1 at follow up)
'persistence' of psychotic symptoms (i.e. 1 at both time points), and 'remission' (1 at baseline, 0
at follow up).

Sleep disorders

Presence of sleep disorders was assessed using the Sleep Disorder Scale for Children (SDSC),
which is a parent-reported questionnaire assessing the presence of a range of sleep disorder
symptoms in children (Bruni et al., 1996). It is composed of 26 Likert items assessing the

frequency of various disturbances over the past 6 months on a 1 to 5 Likert scale (1 = never, 5 = always/daily experiencing a particular issue). All items can be summated for a total measure of sleep disturbance, for which a cut-off point at 39 has sensitivity of 0.89 and specificity of 0.74, correctly identifying 73.4% of a control group and 89.1% of sleep disordered participants.

This cut-off was used to categorise participants according to absence or presence of disturbed sleep. The categorical score at baseline and follow up was then used to derive variables 'persistence' of sleep disturbance (i.e. present at both time points), and 'remission' (present at baseline, absent at follow up).

Potential confounders

Variables were chosen to address a range of possible confounders to the hypothesised relationship between sleep and psychosis. Gender (male/female), ethnicity (white/black/Hispanic/Asian/other) were reported within the basic demographic questionnaires of the study. Ethnicity was re-coded in to white/non-white for the purposes of all analyses. Socioeconomic status was indexed by using the sum score of seven yes/no items in the parent demographic survey questions relating to experiences of family hardship (e.g. 'in the past 12 months has there been a time when you and your immediate family needed food but couldn't afford to buy it or couldn't afford to go out to get it?'), with higher scores on this indicating lower socioeconomic status. Neighbourhood deprivation was assessed using the area deprivation index of the home address, which provides a national percentile where higher values indicate higher levels of deprivation. Child IQ was assessed using the WISC-V matrix reasoning subscale score (range 1-19), with higher values indicating higher IQ. Family conflict was measured using the 9-item family conflict subscale of the family experiences, each item reported by parents as true or false (e.g. 'We fight a lot in our family'), with higher values indicating higher levels of conflict. Lastly prescribed medication fields were searched for any stimulant medications (e.g.

Methylphenidate) including their trade names (e.g. Ritalin) with their presence/absence coded accordingly.

Analysis

The pre-registration document outlining the hypotheses and analysis plan, and the analysis code used in this study are available online at the following link: https://osf.io/8ks72/. R Studio version 3.6.2 was used for all analyses, Supplementary Materials 1 contains a list of packages and version numbers.

For all hypotheses, a set of planned regression analyses were pre-specified. The pre-registration was completed before the ABCD 2.0 data release, i.e. before the one-year follow-up data was made available. In each case, the regression model was first estimated with only the key explanatory (sleep) and dependent (psychosis) variables, and if a significant association was found, the process was repeated with potential confounders included in order to test robustness of the hypothesised association between sleep and psychosis.

We tested the following four hypotheses:

1. Do sleep disorders and psychotic experiences co-occur in childhood?

Tested cross-sectionally at baseline with a) a linear regression to test continuous association between sleep symptoms and psychotic experiences b) a logistic regression to test if sleep symptoms (continuous) predicted presence of psychotic experiences (dichotomous) and c) a logistic regression to test if presence of sleep disorder (dichotomous) predicted presence of psychotic experiences (dichotomous).

2. Do sleep disorders predict later onset of psychotic experiences?

We tested whether sleep disorders at baseline predict psychotic experiences at follow-up. This was investigated by logistic regression models testing if the presence of sleep disorder a) at

baseline and b) at both baseline and follow-up predicted psychotic experiences at one year follow up.

3. Do sleep disorders predict persistence of psychotic experiences?

We tested whether sleep disorders at baseline predict the persistence of psychotic experiences from baseline to follow-up (i.e. psychotic experiences being present at both time points). This was investigated by logistic regression models testing if the presence or persistence of sleep disorder predicted persistence of psychotic experiences.

4. Does remission of sleep disorders predict remission of psychotic experiences?

Finally, we tested whether the remission of sleep disorders (i.e. present at baseline, absent at follow up) also predicted remission of psychotic experiences using logistic regression.

Alterations to the pre-registered statistical plan

Alterations to our analysis plan were made according to subsequently published guidance on analysis of the ABCD cohort (Heeringa & Berglund, 2020), namely introducing multi-level clustering by site and family. The planned unweighted results are reported here, but population weighted demographic and descriptive variables and weighted analyses (where population weights have been converted to scaling weights to allow their inclusion in the multi-level regression model using the 'Method A' and 'Method B' algorithm proposed by (Carle, 2009)), are available in supplementary material.

Results

Demographic and descriptive results

Table 1 displays descriptive statistics of the study. Most descriptive variables remained relatively consistent between baseline and follow-up, however the follow-up group has a preponderance of

White children (59.5% compared to 52.1%) and fewer Black children (9.3% at follow up compared to 15.0% at baseline).

Table 1 about here

Sleep disorders were present in the cohort at similar proportions at baseline and follow up (30.4% and 28.4% respectively). However, this masks some turnover, with roughly a third of the sleep disorder cases at follow-up representing 'new onset' cases (9.8%) to replace a similar proportion that remitted over the year (10.0%). For psychotic experiences there is a clear decline (61.5% of children reporting at least one at baseline, versus 46.8% at follow-up), a result of the high remittance rate (24.6%) compared to new onset (9.6%). This is also reflected in the reduced average PQ-BC score at follow-up versus baseline.

Nevertheless, where sleep disorders and psychotic experiences were present the majority were persistent across the study period: 65.4% of sleep disorder cases and 78.7% of psychotic experience cases at follow up had been present at baseline. This is supported by cross-tabulation of sleep disorders and psychotic experiences which shows relatively low numbers of cases where both sleep disorders and psychotic experiences are new onset (1.1%) and where they have both remitted (2.5%). Out of cases where both sleep disorder and psychotic experiences were present at follow up, the vast majority were 'persistent' cases, where both difficulties had been present at baseline (joint persisting group represents 91.5% of joint onset and persistent cases).

RQ1: Do sleep disorders and psychotic experiences co-occur in childhood?

The regression results associated with this research question can be found in Table 2. They illustrate that sleep disorders and psychotic experiences are significantly and strongly associated within the baseline time point. Sleep disorder symptoms above cut off is associated with an OR of 1.50 (95% CI 1.30-1.74) of at least one psychotic experience being present. The OR remains

significant but reduces slightly (to 1.40, 95% CI 1.20-1.63) once the control variables are added (RQ1f). In this final model, sleep disorder cut off is the largest predictor of current psychotic experiences out of all the parameters tested.

Table 2 about here

RQ2: Do sleep disorders predict onset of psychotic experiences?

The results of the logistic regression analyses on this research question are shown in Table 3. Sleep disorders were found to be strongly associated with reporting psychotic experiences in all models. Sleep disorder symptoms at baseline were found to predict psychotic experiences one year later (OR=1.62, 95% CI 1.41-1.86), even once sleep disorder symptoms at the later time point were controlled for (OR=1.41 95% CI 1.20-1.65). This association remained after control variables were added (OR=1.32, 95% CI 1.11-1.57). The only stronger predictor than sleep disorder in the final model (RQ2d) was stimulant medication (OR=1.61, 95% CI 1.22-2.10).

Table 3 about here

RQ3: Do sleep disorders predict persistence of psychotic experiences?

The results of the logistic regression analyses relating to persistence of psychotic experiences are presented in Table 4. Presence of sleep disorder symptoms at baseline (OR=1.71, 95% CI 1.48-1.97) and persistence of sleep disorder symptoms (OR=1.89, 95% CI 1.60-2.23) were found to highly significantly predict persistence of psychotic experiences at 12 months. These effects remained strong even once control variables were added (ORs of 1.56 and 1.72 for baseline and persisting sleep disorder symptoms respectively). Persistence of sleep disorder symptoms was the strongest overall predictor of persistence of psychotic experiences in the corrected model (RQ3d), with the next largest effect being stimulant medication (OR=1.46, 95% CI 1.12-1.91).

Table 4 about here

RQ4: Do sleep disorders predict remission of psychotic experiences?

Remission of sleep disorder symptoms was not a significant predictor of remission of psychotic experiences (OR=1.041, p=0.766, 95% CI 0.80, 1.35) in the uncontrolled model, therefore the analysis with control variables was not carried out.

Differences arising from weighted analysis

The results using the population weighted values are reported in the supplementary material. The pattern of all results remained substantially the same except for non-white ethnicity which became a significant predictor of onset of psychotic experiences at 12 months (alongside all other factors). See supplementary material for full details.

Discussion

We report that sleep disorder symptoms are strong and significant predictors of both the co-occurrence and persistence of psychotic experiences in 9-11 year old children, even after controlling for a range of other variables including socioeconomic status, ethnicity, IQ, and stimulant medication. However, the remission of sleep disorders did not predict remission of psychotic experiences, and we discuss possible reasons for this negative association below.

Psychotic experiences were common in our study – reported by over two-thirds of children at baseline. This is similar to previous studies in this age group (Laurens et al., 2012), and the relatively high rate of remission versus new onset by the one year follow up is also in keeping with existing developmental models of psychotic experiences. Repeating these analyses through adolescence – where persistence of psychotic experiences over time increasingly predicts risk of later psychotic disorder – will be relevant to further specify what factors are most associated with onset, persistence, and remission of psychotic experiences.

Sleep disorder symptoms were also relatively common in the study group, with just under a third of children having significant sleep issues. Guidelines for parents and clinicians on child sleep difficulties tend to advise that many difficulties ameliorate naturally over time, which is contrary to our findings that around two-thirds do not (at least over the one-year observation period reported here). Previous studies have also found a similar rate of persistence (Fricke-Oerkermann et al., 2007). There is evidence of increasing rates of child and adolescent sleep problems over time (Matricciani et al., 2012) suggesting the need for additional support for parents and health services to address childhood sleep difficulties.

Besides our primary focus on the influence of sleep on psychotic experiences, our findings support the association of a wide range of epidemiological factors (male gender, socioeconomic status, neighbourhood deprivation) with psychotic experiences in this age group. Lower socioeconomic status of parents and increased neighbourhood deprivation were consistent predictors in every model as supported by previous research (Solmi, Lewis, Zammit, & Kirkbride, 2020). Non-white ethnicity was not found to be a significant predictor in the unweighted models but was significant in some analyses once the models were weighted (see Supplementary materials), supporting the longstanding robust association reported elsewhere e.g. (Jongsma et al., 2020). The results therefore support the importance of broader scale social interventions as preventative of mental health issues, and further investigation of the mechanisms between, for example, experiences of deprivation and the development of psychotic experiences.

Lastly, the strong relationship between stimulant medication and psychotic experiences is in keeping with previous literature e.g. (Shyu et al., 2015). However, the full range of reasons for the extent of this association are not discernible from the data reported here. Given the well-established association between dopamine agonist stimulant medication and psychotic experiences (Tost, Alam, & Meyer-Lindenberg, 2010), it is likely that a proportion of this

association can be accounted for as an adverse effect of the medication. However, confounding by indication is likely also to play a role and children who may fulfil criteria for prescribing may also have shared risk factors for psychotic experiences that manifest through other causal pathways (Hollis et al., 2019).

We also note the lack of a relationship between the remission of sleep problems and the remission of psychotic-experiences in this study. As this study was observational, it is not possible to confirm a causal relationship although we suggest several hypotheses. The first is that sleep problems may cause an increase and persistence in psychotic symptoms but through altering a mediating mechanism that remains when sleep problems resolve. In adult studies, the role of affective dysregulation has been highlighted as a potential mediating factor (Akram et al., 2020) with some evidence that this may also be apparent in children (Jeppesen et al., 2014) and we suggest this is a plausible candidate for such a mechanism. Alternatively, sleep problems may cause sensitisation of the mechanisms that generate psychotic symptoms but more rapidly than desensitisation occurs when sleep problems resolve, and although recovery may occur, it may need more than the year interval measured with this data. We note here that 'mechanism' may be neurocognitive (for example, arousal-based increases in aberrant salience) or social (for example, sleep disorder causing behaviour difficulties, exclusion or victimisation, subsequently impacting on psychotic experiences). Nevertheless, this does raise a question about the potential efficacy of sleep improvement intervention in reducing psychotic-like symptoms in preadolescent children, given than an association between improved sleep and reduced psychotic experiences would be expected if there were a simple causal association. However, we note that this study only tested remission of sleep problems as observed in the study, and it may be that a sleep intervention would address some of the shared risk factors for both poor sleep and psychotic-like experiences leading to an improvement in psychotic-like experiences not apparent in naturalistic remission.

Limitations

We note that the observational nature of this study limits the causal conclusions we can draw as we cannot rule out that any changes or associations observed are due to unobserved factors. Another limitation of this being an observational study is that the application of planned analyses is reliant on the data that emerges. This may have limited our ability to reliably test some aspects of the relationship. For example, the particularly low proportion of cases (2.71%) where psychotic experiences and sleep disorders remitted from baseline possibly limited the ability of the remission models to identify any effect (as indicated by the wide confidence intervals for these models). However, given that prior research has suggested a large proportion of psychotic experiences and sleep problems in this age group naturally remit, the lack of such an effect observed in this cohort deserves further investigation. The slight demographic shift between baseline and follow-up group is also worth considering as this may have reduced the accuracy of our estimates (even post-weighting) due to the follow-up group being less representative of the study sample. Confirmation of these findings once the full one-year dataset is released – and with later phases of ABCD data - would be worthwhile.

We have been limited in variables available by the measures used in the ABCD study, which have their own benefits and limitations. For example, while self-reported scales for psychotic experiences in adolescence are recommended, false positives are known to occur (Kelleher et al., 2011; K.-W. Lee et al., 2016) and therefore these associations need to be additionally explored with alternative measures. The sleep measure was parent-reported and parents may vary in their detection of sleep disorders in their children, and some sleep disorders may be more detectable by parents than others (e.g. sleep-walking versus insomnia).

In conclusion, this study indicates that sleep problems in late childhood/early adolescence are common, and are associated with increased likelihood of reporting psychotic experiences both at the time and at one year follow up. Sleep problems were also strongly associated with persistence

of psychotic experiences from baseline to one year – this is of particular importance given that persistence of psychotic experiences in this period is associated with raised risk of later psychotic disorder. These associations remained strong even while controlling for a broad range of confounding variables, including ethnicity, neighbourhood deprivation, family conflict, and prescription of stimulant medication. In conclusion, this study suggests further attention towards sleep problems in this period both as a plausible contributing factor to later mental health problems, but also as a possible preventative treatment target.

Acknowledgements

Data used in the preparation of this article were obtained from the Adolescent Brain Cognitive Development[SM] (ABCD) Study (https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal study designed to recruit more than 10,000 children age 9-10 and follow them over 10 years into early adulthood. The ABCD Study® is supported by the National Institutes of Health and additional federal partners under award numbers U01DA041022, U01DA041028, U01DA041048, U01DA041089, U01DA041106, U01DA041117, U01DA041120, U01DA041134, U01DA041148, U01DA041156, U01DA041174, U24DA041123, U24DA041147, U01DA041093, and U01DA041025. A full list of supporters is available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a complete listing of the study investigators can be found at https://abcdstudy.org/Consortium_Members.pdf. ABCD consortium investigators designed and implemented the study and/or provided data but did not necessarily participate in analysis or writing of this report. This manuscript reflects the views of the authors and may not reflect the opinions or views of the NIH or ABCD consortium investigators.

The ABCD data repository grows and changes over time. The ABCD data used in this report came from doi: 10.15154/1521349. DOIs can be found at https://doi.org

Table 1: Demographic and descriptive variables

Descriptive statistics	Baseline n = 11830	12-month follow-up n = 4910
Age – mean (SD)	9y10m (7.4m)	11y0m (7.6m)
Gender – n male (%)	6162 (52.1%)	2565 (52.2%)
Ethnicity – n(%)		
White	6161 (52.1%)	2923 (59.5%)
Black	1769 (15.0%)	457 (9.3%)
Hispanic	2391 (20.2%)	930 (18.9%)
Asian	250 (2.1%)	115 (2.3%%)
Other	1238 (10.5%)	485 (9.9%)
Socioeconomic status scale – mean (SD)	0.47 (1.1)	0.38 (1.0)
Child IQ Scaled Score – mean (SD)	9.86 (3.0)	10.14(2.9)
Neighbourhood deprivation percentile – mean (SD)	39.22 (27.3)	36.23 (25.3)
Family conflict scale – mean (SD)	2.54 (2.0)	2.47 (1.9)
Stimulant medication prescribed – n (%)	722 (6.1%)	322 (5.6%)
PQ-BC total – mean (SD)	2.63 (3.6)	1.71 (3.0)
SDSC total– mean (SD)	36.53 (8.2)	36.35 (7.9)

Derived variable counts

Sleep disorders present *(≥39 cut off on SDSC)* – n (%)	3602 (30.4%)	1392 (28.4%)
Sleep disorders onset *(absent at baseline, present at follow up)* – n(%)		482 (9.8%)
Sleep disorders persist *(present at both baseline and follow up)* – n (%)		909 (18.5%)
Sleep disorders remit *(present at baseline, absent at follow up)* – n (%)		490 (10.0%)
Psychotic experiences present *(≥1 on PQ-BC total)* – n (%)	7278 (61.5%)	2298 (46.8%)
Psychotic experiences onset – n (%)		473 (9.6%)
Psychotic experiences persist – n (%)		1808 (36.8%)
Psychotic experiences remit – n (%)		1208 (24.6%)

Cross tabulation (sleep disorders vs psychotic experiences)

Onset (absent at baseline, present at follow up)		Psychotic experience onset No	Yes
Sleep disorder onset	No	4008 (81.6%)	420 (8.6%)
	Yes	429 (8.7%)	53 (1.1%)

Persistence (present at baseline and follow up)		Psychotic experience persisting No	Yes
Sleep disorder persisting	No	2621 (53.3%)	1379 (28.1%)
	Yes	480 (9.8%)	430 (8.8%)

Remission (present at baseline, absent at follow up)		Psychotic experience remitting No	Yes
Sleep disorder remission	No	3336 (67.9%)	1084 (22.1%)
	Yes	365 (7.4%)	125 (2.5%)

SDSC: Sleep Disorder Scale for Children, PQBC: Prodromal Questionnaire – Brief Child version ; SD = standard deviation

Table 2: Regression analyses of co-occurrence of sleep disorders and psychotic experiences at baseline (T0)

Model (outcome)	Parameters	B	95% CI	p-value	df	AIC
Model 1a (PQBC total T0) [a]	SDSC total T0	0.048[a]	0.04, 0.06	<0.001	4863	25850

Model (outcome)	Parameters	Odds	95% CI	p-value	df	AIC
Model 1b (PQBC cut-off T0) [b]	SDSC total T0	1.028	1.02, 1.04	<0.001	4904	6320
Model 1c (PQBC cut-off T0) [b]	SDSC cut off T0	1.503	1.30, 1.74	<0.001	4904	6330

Model (outcome)	Parameters	B	95% CI	p-value	df	AIC
Model 1d (PQBC total T0) [a]	SDSC total T0	0.037	0.02, 0.05	<0.001	4566	24131
	Gender	-0.289	-0.49, -0.09	0.004		
	Ethnicity	0.039	-0.04, 0.12	0.347		
	Socioeconomic status	0.183	0.08, 0.29	0.001		
	Neighbourhood deprivation	0.011	0.01, 0.02	<0.001		
	IQ	-0.062	-0.10, -0.03	<0.001		
	Family conflict	-0.003	-0.06, 0.05	0.927		
	Stimulant medication	0.603	0.21, 1.00	0.003		

Model (outcome)	Parameters	Odds	95% CI	p-value	df	AIC
Model 1e (PQBC cut-off T0) [b]	SDSC total T0	1.022	1.01, 1.03	<0.001	4592	5860
	Gender	0.829	0.73, 0.94	0.005		
	Ethnicity	1.003	0.95, 1.06	0.897		
	Socioeconomic status	1.124	1.04, 1.21	0.003		
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001		
	IQ	0.937	0.92, 0.96	<0.001		
	Family conflict	0.998	0.96, 1.03	0.913		
	Stimulant medication	1.247	0.94, 1.65	0.121		
Model 1f (PQBC cut off T0) [b]	SDSC cut off T0	1.401	1.20, 1.63	<0.001	4592	5863
	Gender	0.828	0.73, 0.94	0.004		
	Ethnicity	1.006	0.95, 1.06	0.826		
	Socioeconomic status	1.131	1.05, 1.22	0.001		
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001		
	IQ	0.937	0.92, 0.96	<0.001		
	Family conflict	1.003	0.97, 1.04	0.890		
	Stimulant medication	1.264	0.96, 1.67	0.098		

[a] = linear regression; [b] = logistic regression; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; B = standardised beta; CI = confidence intervals; df = degrees of freedom; AIC = Akaike information criterion.

Table 3: Regression analyses of whether sleep disorders at baseline predict onset of psychotic experiences at 12 months

Model (outcome)	Parameters	Odds	95% CI	p-value	df	AIC
Model 2a (PQBC cut off T1)	SDSC cut off T0	1.621	1.41, 1.86	<0.001	4904	6548
Model 2b (PQBC cut off T1)	SDSC cut off T0	1.407	1.20, 1.65	<0.001	4903	6538
	SDSC cut off T1	1.325	1.13, 1.56	0.001		
Model 2c (PQBC cut off T1)	SDSC cut off T0	1.496	1.29, 1.74	<0.001	4592	6048
	Gender	0.799	0.70, 0.91	0.001		
	Ethnicity	1.031	0.98, 1.09	0.257		
	Socioeconomic status	1.171	1.09, 1.26	<0.001		
	Neighbourhood deprivation	1.008	1.00, 1.01	<0.001		
	IQ	0.959	0.94, 0.98	<0.001		
	Family conflict	0.986	0.95, 1.02	0.430		
	Stimulant medication	1.649	1.26, 2.16	<0.001		
Model 2d (PQBC cut off T1)	SDSC cut off T0	1.320	1.11, 1.57	0.001	4591	6041
	SDSC cut off T1	1.294	1.09, 1.53	0.003		
	Gender	0.794	0.70, 0.91	0.001		
	Ethnicity	1.030	0.98, 1.09	0.267		
	Socioeconomic status	1.163	1.08, 1.25	<0.001		
	Neighbourhood deprivation	1.008	1.00, 1.01	<0.001		
	IQ	0.959	0.94, 0.98	<0.001		
	Family conflict	0.983	0.95, 1.02	0.356		
	Stimulant medication	1.605	1.22, 2.10	<0.001		

SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; df = degrees of freedom; AIC = Akaike information criterion

Table 4: Regression analyses of whether sleep disorders at baseline predict persistence of psychotic experiences at 12 months

Model (outcome)	Parameters	Odds	95% CI	p-value	df	AIC
Model 3a (PQBC persist)	SDSC cut off T0	1.707	1.48, 1.97	<0.001	4904	6192
Model 3b (PQBC persist)	SDSC persist	1.891	1.60, 2.23	<0.001	4904	6188
Model 3c (PQBC persist)	SDSC cut off T0	1.562	1.34, 1.82	<0.001	4592	5710
	Gender	0.798	0.70, 0.91	0.001		
	Ethnicity	0.999	0.95, 1.05	0.957		
	Socioeconomic status	1.163	1.08, 1.25	<0.001		
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001		
	IQ	0.953	0.93, 0.98	<0.001		
	Family conflict	0.989	0.95, 1.03	0.553		
	Stimulant medication	1.471	1.13, 1.92	0.005		
Model 3d (PQBC persist)	SDSC persist	1.716	1.44, 2.04	<0.001	4592	5706
	Gender	0.796	0.70, 0.91	0.001		
	Ethnicity	0.998	0.95, 1.05	0.943		
	Socioeconomic status	1.160	1.08, 1.24	<0.001		
	Neighbourhood deprivation	1.009	1.01, 1.01	<0.001		
	IQ	0.952	0.93, 0.98	<0.001		
	Family conflict	0.991	0.96, 1.03	0.627		
	Stimulant medication	1.460	1.12, 1.91	0.005		

persist = above cut off at T0 and T1; SDSC = Sleep Disorder Scale for Children; PQBC = Prodromal Questionnaire – Brief Child version; CI = confidence intervals; df = degrees of freedom; AIC = Akaike information criterion.

References

Akram, U., Gardani, M., Irvine, K., Allen, S., Ypsilanti, A., Lazuras, L., Drabble, J., Stevenson, J. C., & Akram, A. (2020). Emotion dysregulation mediates the relationship between nightmares and psychotic experiences: Results from a student population. *Npj Schizophrenia*, *6*(1), 1–7. https://doi.org/10.1038/s41537-020-0103-y

Bird, J. C., Waite, F., Rowsell, E., Fergusson, E. C., & Freeman, D. (2017). Cognitive, affective, and social factors maintaining paranoia in adolescents with mental health problems: A longitudinal study. *Psychiatry Research*, *257*(July), 34–39. https://doi.org/10.1016/j.psychres.2017.07.023

Bruni, O., Ottaviano, S., Guidetti, V., Romoli, M., Innocenzi, M., Cortesi, F., & Giannotti, F. (1996). The Sleep Disturbance Scale for Children (SDSC) construction and validation of an instrument to evaluate sleep disturbances in childhood and adolescence. *Journal of Sleep Research*, *5*(4), 251–261. https://doi.org/10.1111/j.1365-2869.1996.00251.x

Carle, A. C. (2009). Fitting multilevel models in complex survey data with design weights: Recommendations. *BMC Medical Research Methodology*, *9*(1), 49. https://doi.org/10.1186/1471-2288-9-49

Carskadon, M. A., & Barker, D. H. (2020). Editorial Perspective: Adolescents' fragile sleep—shining light on a time of risk to mental health. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, *61*(10), 1058–1060. https://doi.org/10.1111/jcpp.13275

Davies, J., Sullivan, S., & Zammit, S. (2018). Adverse life outcomes associated with adolescent psychotic experiences and depressive symptoms. *Social Psychiatry and Psychiatric Epidemiology*, *53*(5), 497–507. https://doi.org/10.1007/s00127-018-1496-z

Dominguez, M. D. G., Wichers, M., Lieb, R., Wittchen, H.-U., & van Os, J. (2011). Evidence that onset of clinical psychosis is an outcome of progressively more persistent subclinical psychotic experiences: An 8-year cohort study. *Schizophrenia Bulletin*, *37*(1), 84–93. https://doi.org/10.1093/schbul/sbp022

Dong, L., Gumport, N. B., Martinez, A. J., & Harvey, A. G. (2019). Is improving sleep and circadian problems in adolescence a pathway to improved health? A mediation analysis. *Journal of Consulting and Clinical Psychology*, *87*(9), 757–771. https://doi.org/10.1037/ccp0000423

Feldstein Ewing, S., & Luciana, M. (2018). The Adolescent Brain Cognitive Development (ABCD) Consortium: Rationale, Aims, and Assessment Strategy [Special Issue]. *Developmental Cognitive Neuroscience*, *32*, 1–164.

Fisher, H., Caspi, A., & Poulton, R. (2013). Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: A birth cohort study. *Psychological Medicine*, *43*(10), 2077–2086. https://doi.org/10.1017/S0033291712003091

Fisher, H., Lereya, S. T., Thompson, A., Lewis, G., Zammit, S., & Wolke, D. (2014). Childhood parasomnias and psychotic experiences at age 12 years in a United Kingdom birth cohort. *Sleep*, *37*(3), 475–482. https://doi.org/10.5665/sleep.3478

Freeman, D., Sheaves, B., Goodwin, G. M., Yu, L. M., Nickless, A., Harrison, P. J., Emsley, R., Luik, A. I., Foster, R. G., Wadekar, V., Hinds, C., Gumley, A., Jones, R., Lightman, S., Jones, S., Bentall, R., Kinderman, P., Rowse, G., Brugha, T., … Espie, C. A. (2017). The effects of improving sleep on mental health (OASIS): A randomised controlled trial with mediation analysis. *Lancet Psychiatry*, *4*(10), 749–758. https://doi.org/10.1016/S2215-0366(17)30328-0

Freeman, D., Sheaves, B., Waite, F., Harvey, A. G., & Harrison, P. J. (2020). Sleep disturbance and psychiatric disorders. *The Lancet Psychiatry*, *7*(7), 628–637. https://doi.org/10.1016/S2215-0366(20)30136-X

Fricke-Oerkermann, L., Plück, J., Schredl, M., Heinz, K., Mitschke, A., Wiater, A., & Lehmkuhl, G. (2007). Prevalence and Course of Sleep Problems in Childhood. *Sleep*, *30*(10), 1371–1377. https://doi.org/10.1093/sleep/30.10.1371

Gradisar, M., Gregory, A. M., & Tikotzky, L. (2020). Is sleep the red flag to psychopathology's bull? Journal of Child Psychology and Psychiatry and Allied Disciplines, 61(10), 1055–1057. https://doi.org/10.1111/jcpp.13331

Gregory, A. M., & Sadeh, A. (2012). Sleep, emotional and behavioral difficulties in children and adolescents. *Sleep Medicine Reviews*, *16*(2), 129–136. https://doi.org/10.1016/j.smrv.2011.03.007

Heeringa, S., & Berglund, P. (2020). A Guide for Population-based Analysis of the Adolescent Brain Cognitive Development (ABCD) Study Baseline Data. *BioRxiv*, 2020.02.10.942011. https://doi.org/10.1101/2020.02.10.942011

Hennig, T., & Lincoln, T. M. (2017). Sleeping Paranoia Away? An Actigraphy and Experience-Sampling Study with Adolescents. *Child Psychiatry & Human Development*, 1–10. https://doi.org/10.1007/s10578-017-0729-9

Hollis, C., Chen, Q., Chang, Z., Quinn, P. D., Viktorin, A., Lichtenstein, P., … Larsson, H. (2019). Methylphenidate and the risk of psychosis in adolescents and young adults: a population-based cohort study. *The Lancet Psychiatry*, *6*(8), 651–658. https://doi.org/10.1016/S2215-0366(19)30189-0

Jeppesen, P., Clemmensen, L., Munkholm, A., Rimvall, M. K., Rask, C. U., Jørgensen, T., Larsen, J. T., Petersen, L., van Os, J., & Skovgaard, A. M. (2014). Psychotic experiences co-occur with sleep problems, negative affect and mental disorders in preadolescence. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, *56*(5), 558–565. https://doi.org/10.1111/jcpp.12319

Jongsma, H. E., Gayer-Anderson, C., Tarricone, I., Velthorst, E., Van Der Ven, E., Quattrone, D., … Kirkbride, J. B. (2020). Social disadvantage, linguistic distance, ethnic minority status and first-episode psychosis: Results from the EU-GEI case-control study. Psychological Medicine. https://doi.org/10.1017/S003329172000029X

Karcher, N. R., Barch, D. M., Avenevoli, S., Savill, M., Huber, R. S., Simon, T. J., Leckliter, I. N., Sher, K. J., & Loewy, R. L. (2018). Assessment of the prodromal questionnaire-brief child version for measurement of self-reported psychoticlike experiences in childhood. *JAMA Psychiatry*, *75*(8), 853–861. https://doi.org/10.1001/jamapsychiatry.2018.1334

Kelleher, I., Harley, M., Murtagh, A., & Cannon, M. (2011). Are Screening Instruments Valid for Psychotic-Like Experiences? A Validation Study of Screening Questions for Psychotic-Like Experiences Using In-Depth Clinical Interview. *Schizophrenia Bulletin*, *37*(2), 362–369. https://doi.org/10.1093/schbul/sbp057

Kelleher, I., Keeley, H., Corcoran, P., Lynch, F., Fitzpatrick, C., Devlin, N., Molloy, C., Roddy, S., Clarke, M. C., Harley, M., Arseneault, L., Wasserman, C., Carli, V., Sarchiapone, M., Hoven, C., Wasserman, D., & Cannon, M. (2012). Clinicopathological significance of psychotic experiences in non-psychotic young people: Evidence from four population-based studies. *The British Journal of Psychiatry*, *201*(1), 26–32. https://doi.org/10.1192/bjp.bp.111.101543

Koopman-Verhoeff, M. E., Bolhuis, K., Cecil, C. A. M., Kocevska, D., Hudziak, J. J., Hillegers, M. H. J., Mileva-Seitz, V. R., Reiss, I. K., Duijts, L., Verhulst, F., Luijk, M. P. C. M., & Tiemeier, H. (2018, December 14). During day and night: Childhood psychotic experiences and objective and subjective sleep problems. *Schizophrenia Research*, *206*, 127–134. https://doi.org/10.1016/j.schres.2018.12.002

Laurens, K. R., Hobbs, M. J., Sunderland, M., Green, M. J., & Mould, G. L. (2012). Psychotic-like experiences in a community sample of 8000 children aged 9 to 11 years: An item response theory analysis. *Psychological Medicine*, *42*(7), 1495–1506. https://doi.org/10.1017/S0033291711002108

Lee, K.-W., Chan, K.-W., Chang, W.-C., Lee, E. H.-M., Hui, C. L.-M., & Chen, E. Y.-H. (2016). A systematic review on definitions and assessments of psychotic-like experiences. *Early Intervention in Psychiatry*, *10*(1), 3–16. https://doi.org/10.1111/eip.12228

Lee, Y. J., Cho, S.-J., Cho, I. H., Jang, J. H., & Kim, S. J. (2012). The relationship between psychotic-like experiences and sleep disturbances in adolescents. *Sleep Medicine*, *13*(8), 1021–1027. https://doi.org/10.1016/j.sleep.2012.06.002

Linscott, R. J., & van Os, J. (2013). An updated and conservative systematic review and meta-analysis of epidemiological evidence on psychotic experiences in children and adults: On the pathway from proneness to persistence to dimensional expression across mental disorders. *Psychological Medicine*, *43*(6), 1133–1149. https://doi.org/10.1017/S0033291712001626

Maijer, K., Hayward, M., Fernyhough, C., Calkins, M. E., Debbané, M., Jardri, R., Kelleher, I., Raballo, A., Rammou, A., Scott, J. G., Shinn, A. K., Steenhuis, L. A., Wolf, D. H., & Bartels-Velthuis, A. A. (2019). Hallucinations in Children and Adolescents: An Updated Review and Practical Recommendations for Clinicians. *Schizophrenia Bulletin*, *45*(Supplement_1), S43–S55. https://doi.org/10.1093/schbul/sby119

Matricciani, L., Olds, T., & Petkov, J. (2012). In search of lost sleep: Secular trends in the sleep time of school-aged children and adolescents. *Sleep Medicine Reviews*, *16*(3), 203–211. https://doi.org/10.1016/j.smrv.2011.03.005

McGrath, J. J., Saha, S., Al-Hamzawi, A., Alonso, J., Bromet, E. J., Bruffaerts, R., Caldas-de-Almeida, J. M., Chiu, W. T., de Jonge, P., Fayyad, J., Florescu, S., Gureje, O., Haro, J. M., Hu, C., Kovess-

Masfety, V., Lepine, J. P., Lim, C. C. W., Mora, M. E. M., Navarro-Mateu, F., … Kessler, R. C. (2015). Psychotic Experiences in the General Population. *JAMA Psychiatry*, *72*(7), 697. https://doi.org/10.1001/jamapsychiatry.2015.0575

Morales-Muñoz, I., Broome, M. R., & Marwaha, S. (2020). Association of Parent-Reported Sleep Problems in Early Childhood With Psychotic and Borderline Personality Disorder Symptoms in Adolescence. *JAMA Psychiatry*. https://doi.org/10.1001/jamapsychiatry.2020.1875

Reeve, S., Emsley, R., Sheaves, B., & Freeman, D. (2018). Disrupting Sleep: The Effects of Sleep Loss on Psychotic Experiences Tested in an Experimental Study With Mediation Analysis. *Schizophrenia Bulletin*, *44*(3), 662–671. https://doi.org/10.1093/schbul/sbx103

Reeve, S., Nickless, A., Sheaves, B., & Freeman, D. (2018). Insomnia, negative affect, and psychotic experiences: Modelling pathways over time in a clinical observational study. *Psychiatry Research*, *269*, 673–680. https://doi.org/10.1016/j.psychres.2018.08.090

Reeve, S., Sheaves, B., & Freeman, D. (2015). The role of sleep dysfunction in the occurrence of delusions and hallucinations: A systematic review. *Clinical Psychology Review*, *42*, 96–115. https://doi.org/10.1016/j.cpr.2015.09.001

Shyu, Y.-C., Yuan, S.-S., Lee, S.-Y., Yang, C.-J., Yang, K.-C., Lee, T.-L., & Wang, L.-J. (2015). Attention-deficit/hyperactivity disorder, methylphenidate use and the risk of developing schizophrenia spectrum disorders: A nationwide population-based study in Taiwan. *Schizophrenia Research*, *168*(1–2), 161–167. https://doi.org/10.1016/J.SCHRES.2015.08.033

Solmi, F., Lewis, G., Zammit, S., & Kirkbride, J. B. (2020). Neighborhood Characteristics at Birth and Positive and Negative Psychotic Symptoms in Adolescence: Findings from the ALSPAC Birth Cohort. *Schizophrenia Bulletin*, *46*(3), 581–591. https://doi.org/10.1093/schbul/sbz049

Thompson, A., Lereya, S. T., Lewis, G., Zammit, S., Fisher, H. L., & Wolke, D. (2015). Childhood sleep disturbance and risk of psychotic experiences at 18: UK birth cohort. *British Journal of Psychiatry*, *207*(1), 23–29. https://doi.org/10.1192/bjp.bp.113.144089

Tost, H., Alam, T., & Meyer-Lindenberg, A. (2010). Dopamine and psychosis: Theory, pathomechanisms and intermediate phenotypes. *Neuroscience & Biobehavioral Reviews*, *34*(5), 689–700. https://doi.org/10.1016/j.neubiorev.2009.06.005

Trotta, A., Arseneault, L., Caspi, A., Moffitt, T. E., Danese, A., Pariante, C., & Fisher, H. L. (2020).

    Mental Health and Functional Outcomes in Young Adulthood of Children With Psychotic

    Symptoms: A Longitudinal Cohort Study. *Schizophrenia Bulletin*, *46*(2), 261–271.

    https://doi.org/10.1093/schbul/sbz069

Volkow, N. D., Koob, G. F., Croyle, R. T., Bianchi, D. W., Gordon, J. A., Koroshetz, W. J., Pérez-Stable,

    E. J., Riley, W. T., Bloch, M. H., Conway, K., Deeds, B. G., Dowling, G. J., Grant, S., Howlett,

    K. D., Matochik, J. A., Morgan, G. D., Murray, M. M., Noronha, A., Spong, C. Y., … Weiss, S.

    R. B. (2018). The conception of the ABCD study: From substance use to a broad NIH

    collaboration. *Developmental Cognitive Neuroscience*, *32*, 4–7.

    https://doi.org/10.1016/j.dcn.2017.10.002

Waite, F., Sheaves, B., Isham, L., Reeve, S., & Freeman, D. (2019). Sleep and schizophrenia: From

    epiphenomenon to treatable causal target. *Schizophrenia Research*.

    https://doi.org/10.1016/J.SCHRES.2019.11.014

Zhang, J., Paksarian, D., Lamers, F., Hickie, I. B., He, J., & Merikangas, K. R. (2017). Sleep Patterns and

    Mental Health Correlates in US Adolescents. *Journal of Pediatrics*, *182*, 137–143.

    https://doi.org/10.1016/j.jpeds.2016.11.007