Modelling the COVID-19 Fatality Rate in England and its Regions

Nigel Saunders

Thermotech Ltd., Surrey Technology Centre, The Surrey Research Park,
Guildford, Surrey GU2 7YG, U.K.

Abstract

A model to account for the fatality rate in England and its regions is proposed. It follows the clear observation that, rather than two connected waves, there have been many waves of infections and fatalities in the regions of England of various magnitudes, usually overlapping. The waves are self-limiting, in that clear peaks are seen, particularly in reported positive test rates. The present model considers fatalities as the data reported are more reliable than positive test rates, particularly so during the first wave when so little testing was done.

The model considers the observed waves are essentially similar in form and can be modelled using a single wave form, whose final state is only dependent on its peak height and start date. The basic wave form was modelled using the observed fatality rates for London, which unlike the other regions, exhibited almost completely as a single wave in the "first wave". Its form matches rather well with the "Do Nothing" model reported by Imperial College on 16th March 2020, but reduced substantially from its expected peak.

There are, essentially, only two adjustable parameters used in the model, the start date of the relevant wave and its height. The modelled fatalities for each wave are summed per day and a cumulative curve is matched to that reported. The minimal number of adjustable parameters, alongside the fact that the waves invariably overlap, provides highly stringent conditions on the fitting process.

Results are presented for each region for both the "first" and "second' waves. High levels of accuracy are obtained with R^2 values approaching 100% against the ideal fit for both waves. It can also be seen there are fundamental differences between the underlying behaviour of the "first" and "second" waves.

1. Introduction

While there are daily reports of the progress of COVID-19 infections in the UK, it seems little work has been done in trying to understand how the virus has actually spread and grown in the UK. While regional R rates are regularly presented, infection or fatality rates over the whole period for the regions are rarely, if ever presented. However, by doing so, a far more detailed view of how the infection has spread and grown is revealed. Fig.1 below shows reported population adjusted infection rates for the regions of England from the beginning of the pandemic, up to 1st January 2021 [1]. A 7-day average of cases by specimen date (*averaged at midweek*) is used and the last 5 days not used due to likely incompleteness as advised by the GOV.uk site.

---

1. UK Government website for data and insights on Coronavirus (COVID-19) https://coronavirus.data.gov.uk/

Fig.1 Reported population adjusted infection rates for the regions of England from the beginning of the pandemic, up to 1st January 2021 [1] using a 7-day average of cases by specimen date

Examination of Fig.1 provides clear light on how the virus has spread and grown across England. (Note, the variability in the curves at the end of December appears due to under recording around Christmas day.)

- There have been many waves of infection in the regions of various magnitudes, many major, but with some small but distinctly observable.

- In the "first wave", with the exception of London, which exhibited as an essentially singular wave, multiple overlapping waves can be seen in the other regions, exhibiting as multiple peaks within a broad "first" regional wave.

- After the first lockdown, infections continued to fall in a quite continuous fashion. There were some minor waves but, effectively, rates plateaued, showing only a slight rise until week 35, after which rates increased in all regions nationwide.

- Since week 35 there appears to be multiple overlapping, but distinct waves in all of the regions. While the lower rate regions exhibit these waves quite clearly, it is also clear they exist in the high-rate regions and can be seen as inflections in the slope of the test data, which occur as an increasingly dominant wave adds to the infection curve.

- During the second week of October, with the exception of the North West which peaked in the second half of September, there was a fall in infection rates for all regions, with the Northern and Midland regions exhibiting particularly sharp falls. Subsequently, around the end of November/beginning of December, infection rates rose in all regions, but most sharply in London, the South East and the East of England.

2. Development of the model

From Fig.1, London is the only region that exhibited as mainly a single wave during the "first wave", which is rather symmetrical in form, and can be used a basis for developing a model consisting of multiple waves. Rather than using positive test data, the model has been developed taking as its basis fatality rates. The data is far more reliably reported over the whole period and the reported test data also does not accurately reflect the true total number of infections due to asymptomatic cases.

Fig.2 Fatality rates reported for London including those from the present model and using rate adjusted ICU and ICS models.

Fig.2 shows reported fatalities by date of death [1], as a 7-day average for London. The wave is quite symmetrical in nature until the 29th April, at which point a further wave becomes apparent, adding to the first, before diminishing to almost zero. The model used here fits the curve up until then, before itself diminishing to zero (Appendix 1). Fig.2 also includes two further curves taken after the Imperial College report of 16th March 2020 [2].

(i) A curve based on the "Do Nothing" scenario (hereafter called the Imperial College Unconstrained (ICU) model), scaled to match the peak height of the reported curve for London and date adjusted.

(ii) A curve based on the scenario of Schools and Universities Closure, Case Isolation and Population-Wide Social Distancing (hereafter called the Imperial College Suppression (ICS) model). The reported curve is provided in terms of critical care beds occupied and, based on the assumption it is directly proportional to subsequent fatalities, scaled to match the peak height for London and date adjusted.

The scaled ICU curve matches the reported data for the first wave rather well, though is a little wider, while the ICS curve matches the reported data almost exactly to the peak height before markedly diverging.

The fitted curve can now be used in a multiple wave model which requires three sets of input. The first is the number of waves, which is usually clear to see or becomes required during the fitting process. Otherwise, the fitting process requires only two parameters (i) the start date of the relevant wave and (ii) its height. The calculated fatalities for each wave are then summated and the cumulative curve matched to the reported data. The model was implemented here using spreadsheets but should suit a computer programme utilising a least squares methodology to minimise the error.

2. N.M. Ferguson et al., Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-19 mortality and healthcare demand, (Imperial College COVID-19 Response Team, 16 March 2020)

3. Results

3.1 "First Wave"

Figs. 3-11(a&b) show results for the fitted curves to reported data alongside plots for the divergence from the "ideal fit" in terms of the value of R^2. The data is given as fatalities, reported by date of death [1], as a 7-day average.

Considering the simplicity of the model, it matches the general behaviour of fatality rates very accurately, but there are some spikes in fatalities which cannot be matched. It may be, straightforwardly, that the model is unable to account for minor variations in behaviour or due to clusters of fatalities in settings where fatality rates would be higher than for the overall population average.

Fig. 12 shows the sum of all the waves of the regions compared to fatality data reported for England [1] as a 7-day average, while Fig.13 compares the raw data as reported by date of death for completeness.

Fig.3 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for London in the "first wave"

Fig.4 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the East of England in the "first wave"

Fig.5 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the East Midlands in the "first wave

Fig.6 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the North East in the "first wave"

Fig.7 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the North West in the "first wave"

Fig.8 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the South East in the "first wave"

Fig.9 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the South West in the "first wave"

Fig.10 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the West Midlands in the "first wave"

Fig.11 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for Yorkshire & Humber in the "first wave"

Fig. 12 Comparison between the sum of all calculated waves of the regions compared to reported fatalities by date of death (7-day average) for England during the "first wave"

Fig. 13 Comparison between the sum of all calculated waves of the regions compared to reported fatalities by date of death for England during the "first wave"

3.2 "Second Wave"

Figs. 14-22 (a&b) show, similarly to the "first wave", results for the fitted curves to reported data [1] alongside plots for the divergence from the "ideal fit" in terms of the value of R^2. As for Fig.1 the last 5 reported days are not considered in the reported data. However, it is noted that there is significant delay in reporting, with fatality results still being reported well after the 5-day delay accounted for here. The reported and subsequently fitted rates for the later dates, therefore, have significant uncertainty. Fig.23 show the sum of all the waves of the regions compared to fatality data reported for England [1] as a 7-day average, while Fig.24 compares the raw data as reported per day for completeness.

All regions show at least 5 waves and, with the exception of the Midlands regions, a further 6th wave is becoming apparent in the other regions. It should be considered highly likely that a further wave will manifest itself in the Midlands regions during early 2021.

A common feature for all regions is the periodic nature of the waves. The model cannot explain why that should be the case but, a periodic nature is also seen quite clearly in the infection curves and further analysis is required to understand why it has arisen.

Fig.14 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for London in the "second wave"

Fig.15 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the East of England in the "second wave"

Fig.16 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the East Midlands in the "second wave"

Fig.17 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the North East in the "second wave"

Fig.18 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the North West in the "second wave"

Fig.19 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the South East in the "second wave"

Fig.20 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the South West in the "second wave"

Fig.21 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the West Midlands in the "second wave"

Fig.22 (a) Comparison between calculated and reported fatalities by date of death (7-day average) and (b) divergence from the "ideal fit" for the West Midlands in the "second wave"

Fig. 23 Comparison between the sum of all calculated waves of the regions compared to reported fatalities by date of death (7-day average) for England during the "second wave"

Fig. 24 Comparison between the sum of all calculated waves of the regions compared to reported fatalities by date of death for England during the "second wave"

4. Discussion

The accuracy of the fitted curves to reported fatality rates, with R^2 values approaching 100% against the ideal, strongly supports the validity of the current model and it reveals detail that would otherwise remain hidden or considered purely speculative. For example, in the "first wave" some regions show smoothly falling fatality rates with time, while others clearly show a multiple wave format. The strength of the model is it accounts for both types of behaviour in a unified approach.

It also accounts for the observed "exponential" rises in infection/fatality rates often quoted during the "second wave" which, when extrapolated into the future, substantially overestimated subsequent reported rates. Such apparent rises can happen when multiple waves, each increasing in magnitude with time, are added to each other and appear to be one continuous wave, e.g., Yorkshire & Humber.

Fig. 25 Extrapolation of the fatality curve for England in the "second wave" showing overestimation of future fatality rates by assuming exponential growth

However, when the subsequent wave/waves either do not increase in size or diminish, it can lead to large errors in extrapolation. Fig.25 shows reported fatality rates as a 7-day average for England [1] plotted on an exponential y-axis. Extrapolation, using data for the first 2-3 weeks of October, leads to a prediction of close to 2,000 fatalities per day by the end of November, while the real value had peaked around 400 before then.

By its very nature, the method used here is an evidence-based approach and quite unlike the more theoretical approaches adopted by groups such as in Imperial College. It has the advantage that it uses pre-existing evidence to develop an approach that can quantitatively match reported behaviour. As such, it is not within the scope of the present paper to undertake any detailed discussion on the differences between the two approaches.

However, it is noted that the success of the present model in accounting for fatality rates so accurately lies in the fact that it considers each region separately. In effect, it models England as a set of closed systems that allows regional variability of fatality rates to be taken into account. The ability to do so becomes particularly important for the "second wave". In a more fundamental model, which includes transmission between the regions, it could be considered as a loosely coupled model with each region having its own individual characteristics yet allowing transmission between regions.

The behaviour of the "first" and "second" waves is very different. The "first wave" consisted of a large initial spike in fatalities in all regions and was followed by a series of significantly sized secondary waves which diminished with time. London, which is the most highly interconnected of the regions, exhibited almost completely as a single wave unlike the other regions. It therefore seems reasonable to assume that the secondary waves in the other regions was due to the virus spreading into previously unaffected areas.

The "second wave" has behaved in a fundamentally different way. It began after a long period of very low fatality rates and in all regions simultaneously. Unlike the "first wave", the "second wave" is characterised by a series of periodic waves in all regions, including London which is now exhibiting as a series of periodic waves rather than a predominantly single wave. It is also characterised by a substantial regional variation in fatality rates over time.

Fatality rates per head of population were initially highest in the Northern and Midlands regions and peaked during November. On the other hand, in London, the South East and the East of England rates were initially quite low. They then rose sharply at the end of November and still had not peaked by the end of December. The South West behaved quite differently to the other regions, in that fatality rates remained low per head of population and peaked in November.

The fatality rates for England, therefore, reflect regional behaviour, with a peak exhibiting in rates at the end of November, when the Northern and Midlands regions peaked. Rates fell for a short time, before the sharp rise in fatality rates in London, the South East and the East of England caused an increase during December. The appearance of two waves, therefore, arises due to the regional variations in England, rather than something that reflects the general behaviour of the population of England.

Conclusions

The current paper makes no claims to provide a theoretical explanation for how COVID-19 has transmitted in England and its regions. Its purpose is to see if it is possible to quantitatively model the clear observation from measured infection/fatality rates that there have been many waves of infection and subsequent fatalities in the regions of England. The fact that the model can provide such exact correlation with observed fatality rates suggests there would be merit in developing a theoretical approach to describe how such a multiple wave form has arisen, particularly so for the "second wave".

It further provides a clear and quantitative explanation as to why exponential extrapolations which have been made during the "second wave" have invariably failed by such large margins and accounts for fatality curves which quite clearly exhibit multiple waves as well as those that appear quite continuous.

About the author

The author is a researcher and scientific modeller in the physical sciences and has written this paper as a contribution to the further understanding of the spread of COVID-19.

Appendix 1. Fitted curve for London Fatalities per day
based on a 7-day average of reported data [1]

The curve below is used as a "master wave". It is based on the reported 7-day average for London and, fitted to the reported data using a combination of fitting to a polynomial function and manual adjustment to provide a smoothly changing curve.

While the fitted curve matches the reported fatalities for London very well, it would clearly be preferable for a more meaningful wave function to be developed to match the behaviour of London. It would also be of interest to examine other highly interconnected regions/cities, particularly if reasonably isolated from other populated areas, to see if there are common features in their behaviour during the "first wave" of the pandemic.

Date	Value	Date	Value
09/03/2020	0.00	14/04/2020	154.83
10/03/2020	1.10	15/04/2020	145.55
11/03/2020	2.46	16/04/2020	136.55
12/03/2020	4.20	17/04/2020	127.83
13/03/2020	6.41	18/04/2020	119.38
14/03/2020	9.16	19/04/2020	111.22
15/03/2020	12.55	20/04/2020	103.33
16/03/2020	16.64	21/04/2020	95.73
17/03/2020	21.52	22/04/2020	88.41
18/03/2020	27.26	23/04/2020	81.38
19/03/2020	33.95	24/04/2020	74.63
20/03/2020	41.66	25/04/2020	68.17
21/03/2020	50.47	26/04/2020	61.99
22/03/2020	60.46	27/04/2020	56.11
23/03/2020	71.71	28/04/2020	50.51
24/03/2020	84.31	29/04/2020	45.21
25/03/2020	98.32	30/04/2020	40.20
26/03/2020	111.25	01/05/2020	35.49
27/03/2020	123.76	02/05/2020	31.07
28/03/2020	136.62	03/05/2020	26.95
29/03/2020	149.36	04/05/2020	23.12
30/03/2020	161.51	05/05/2020	19.60
31/03/2020	172.61	06/05/2020	16.37
01/04/2020	182.17	07/05/2020	13.45
02/04/2020	191.00	08/05/2020	10.83
03/04/2020	198.00	09/05/2020	8.51
04/04/2020	204.00	10/05/2020	6.50
05/04/2020	206.50	11/05/2020	4.79
06/04/2020	207.60	12/05/2020	3.39
07/04/2020	207.00	13/05/2020	2.30
08/04/2020	204.50	14/05/2020	1.52
09/04/2020	199.00	15/05/2020	1.05
10/04/2020	191.50	16/05/2020	0.60
11/04/2020	184.31	17/05/2020	0.22
12/04/2020	174.21	18/05/2020	0.00
13/04/2020	164.38		