SARS-CoV-2 B.1.1.7 escape from mRNA vaccine-elicited neutralizing antibodies

Dami A. Collier[1,2,3]*, Anna De Marco[4]* , Isabella A.T.M. Ferreira*[1,2] Bo Meng[1,2]*, Rawlings
Datir *[1,2,3], Alexandra C. Walls[5], Steven A. Kemp S[1,2,3], Jessica Bassi[4], Dora Pinto[4], Chiara
Silacci Fregni[4], Siro Bianchi[4], M. Alejandra Tortorici[5], John Bowen[5], Katja Culap[4], Stefano
Jaconi[4], Elisabetta Cameroni[4], Gyorgy Snell[6], Matteo S. Pizzuto[4], Alessandra Franzetti
Pellanda[7], Christian Garzoni[7], Agostino Riva[8], The CITIID-NIHR BioResource COVID-19
Collaboration[9], Anne Elmer[10], Nathalie Kingston[10], Barbara Graves [10], Laura E McCoy[3],
Kenneth GC Smith [1,2], John R. Bradley [2,10], Lourdes Ceron-Gutierrez L[11], Gabriela Barcenas-
Morales [11,12],   The COVID-19 Genomics UK (COG-UK) consortium[13] , William Harvey[14],
Herbert W. Virgin[6], Antonio Lanzavecchia[4], Luca Piccoli[4], Rainer Doffinger[11], Mark Wills[2] ,
David Veesler[5], Davide Corti[4]*, Ravindra K. Gupta[1,2, 15]*

*Equal contribution

[1]Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), Cambridge, UK.
[2]Department of Medicine, University of Cambridge, Cambridge, UK.
[3]Division of Infection and Immunity, University College London, London, UK.
[4]Humabs Biomed SA, a subsidiary of Vir Biotechnology, 6500 Bellinzona, Switzerland.
[5]Department of Biochemistry, University of Washington, Seattle, WA 98195, USA
[6]Vir Biotechnology, San Francisco, CA 94158, USA.
[7]Clinic of Internal Medicine and Infectious Diseases, Clinica Luganese Moncucco, 6900 Lugano,
Switzerland
[8]Division of Infectious Diseases, ASST Fatebenefratelli Sacco, Luigi Sacco Hospital, University of
Milan, 20157 Milan, Italy
[9] The CITIID-NIHR BioResource COVID-19 Collaboration, see appendix 1 for author list
[10] NIHR Cambridge Clinical Research Facility, Cambridge, UK.
[11]Department of Clinical Biochemistry and Immunology, Addenbrookes Hospital, UK
[12] Laboratorio de Inmunologia, S-Cuautitlán, UNAM, Mexico
[13]https://www.cogconsortium.uk. Full list of consortium names and affiliations are in Appendix 2.
[14] Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow,
Glasgow, UK
[15] Africa Health Research Institute, Durban, South Africa

Correspondence: dcorti@vir.bio, rkg20@cam.ac.uk

Key words: SARS-CoV-2; COVID-19; antibody, vaccine, neutralising antibodies;
mutation;

Abstract

SARS-CoV-2 transmission is uncontrolled in many parts of the world, compounded in some areas by higher transmission potential of the B1.1.7 variant now seen in 50 countries. It is unclear whether responses to SARS-CoV-2 vaccines based on the prototypic strain will be impacted by mutations found in B.1.1.7. Here we assessed immune responses following vaccination with mRNA-based vaccine BNT162b2. We measured neutralising antibody responses following a single immunization using pseudoviruses expressing the wild-type Spike protein or the 8 mutations found in the B.1.1.7 Spike protein. The vaccine sera exhibited a broad range of neutralizing titres against the wild-type pseudoviruses that were modestly reduced against B.1.1.7 variant. This reduction was also evident in sera from some convalescent patients. Decreased B.1.1.7 neutralization was also observed with monoclonal antibodies targeting the N-terminal domain (9 out of 10), the Receptor Binding Motif (RBM) (5 outof 29), but not in neutralizing mAbs binding outside the RBM. Introduction of the E484K mutation in a B.1.1.7 background to reflect newly emerging viruses in the UK led to a more substantial loss of neutralizing activity by vaccine-elicited antibodies over that conferred by the B.1.1.7 mutations alone. Further work is needed to establish the impact of these observations on protective vaccine efficacy in the context of the evolving B.1.1.7 lineage.

Introduction

The outbreak of a pneumonia of unknown cause in Wuhan, China in December 2019, culminated in a global pandemic due to a novel viral pathogen, now known to be SARS-COV-2[1]. The unprecedented scientific response to this global challenge has led to the rapid development of vaccines aimed at preventing SARS-COV-2 infection and transmission. Continued viral evolution led to the emergence and selection of SARS-CoV-2 variants with enhanced infectivity/transmissibility[2,3] [4,5] and ability to circumvent drug[6] and immune control[7,8].

SARS-CoV-2 vaccines have recently been licensed that target the Spike (S) protein, either using mRNA or adenovirus vector technology with protection rates over a few months ranging from 62 to 95%[9-11]. The BNT162b2 vaccine encodes the full-length trimerised S protein of SARS CoV-2 and is formulated in lipid nanoparticles to optimise delivery to cells[12]. Other vaccines include the Moderna mRNA-1273 vaccine, which is also a lipid nanoparticle formulated S glycoprotein[13] and the Oxford-AstraZeneca ChAdOx1 nCoV-19 vaccine (AZD1222) which is a replication-deficient chimpanzee adenoviral vector ChAdOx1, containing the S glycoprotein[14]. The duration of immunity conferred by these vaccines is as yet unknown. These vaccines were designed against the Wuhan-1 isolate discovered in 2019. Concerns have been raised as to whether these vaccines will be effective against new SARS-CoV-2 variants, such as B.1.1.7 (N501Y.V1), B.1.351 (N501Y.V2) and P1 (N501Y.V2) that

originated in the UK, South Africa, and Brazil and are now being detected all over the
world[15-17].

The phase I/II studies of the Pfizer-BioNTech BNT162b2 vaccine determined the
immunogenicity of different dosing regimens. The geometric mean concentration (GMC) of
RBD-binding IgG 21 days after the first dose of 30 µg of the BNT162b2 vaccine, which is the
dose approved in the UK, was higher than the GMC of a panel of convalescent plasma (1,536
vs 602 U/ml). Nevertheless, the corresponding neutralisation geometric mean titre (GMT) was
3-fold lower than a panel of convalescent plasma (29 vs 94)[12], but substantially increased after
boost immunization[18]. In older adults mean GMT was only 12 in a preliminary analysis of 12
participants[19] and increased to 109 after the second dose.

In this study, we assess antibody responses induced three weeks after vaccination with
the first dose of BNT162b2 following the rollout in the UK. In addition, by using a panel of
human neutralizing monoclonal antibodies (mAbs) we show that the B.1.1.7 variant can
escape neutralization mediated by most NTD-specific antibodies tested and by a fraction of
RBM-specific antibodies. We also show that the recent appearance of the E484K mutation in
B.1.1.7 isolates from the UK, similarly to the B.1.351 and P1 isolates, results in incremental
loss of neutralization by BNT162b2 mRNA-elicited antibodies.

Results

Twenty-four participants had received the BNT162b2 mRNA vaccine three weeks prior to
blood draw for serum and peripheral blood monocnulear cells (PBMC) collection. Median age
was 82 years (IQR 64-85) and 30% were female (**Supplementary Table 1**). Serum IgG titres
to N protein, S and the S RBD were assayed by particle based flow cytometry on a Luminex
analyser (**Fig. 1a**). These data showed Spike and RBD antibody titres much higher than in
healthy controls, similar to both convalescent plasma units used for therapeutic purposes as
well as to serum from recovered individuals. The raised N titres relative to control could be
the result of non specific cross reactivity that is increased following vaccination. However, the
antibody response was heteorgenorus with almost 100-fold variation in IgG titres to S and
Spike RBD across the vaccinated participants.

Using lentiviral pseudotyping we generated wild type S proteins on the surface of enveloped
virions in order to measure neutralisation activity of vaccine-elicited sera. This system has
been shown to give results correlating with replication competent authentic virus[20,21]. The

vaccine sera exhibited a range of inhibitory dilutions giving 50% neutralisation (ID50) (**Fig.**
1c-d). The GMT against wild type (WT) pseudovirus was 24. Seven out of 23 participants
exhibited no appreciable neutralisation against the WT pseudotyped virus. There was
reasonable correlation between full length S IgG titres and serum neutralisation titres
(**Extended Data Fig. 1a**). A broad range of T cell responses was measured by IFN gamma
FLUOROSPOT against SARS-CoV-2 peptides in vaccinees. These cell responses did not
correlate with IgG S antibody titres, but there was some correlation with serum neutralisation
against WT virus (**Extended Data Fig. 1b-c**).

We next generated mutated pseudoviruses carrying S protein with mutations N501Y,
A570D and the H69/V70 deletion. We observed no reduction in the ability of sera from
vaccinees to inhibit either WT or mutant virus (**Extended Data Fig. 2**). In fact, we observed
that vaccine sera displayed higher neutralizing activity against the N501Y, A570D and
H69/V70 deletion mutant relative to WT virus. Next, we then tested a panel of sera from 11
recovered individuals and found that these sera also neutralised both wild type and the
mutated viruses similarly (**Extended Data Fig. 3**). The findings for vaccine sera may be
related to a potential allosteric effect of ΔH69/V70 that might enhance neutralization by
antibodies targeting cryptic sites on the S, such as the RBD site II (also named as class 4 site).

We then generated the full set of eight mutations in the S protein present in B.1.1.7
variant (**Fig. 1b** and **Table 1**), ΔH69/V70, Δ144, N501Y and A570D in the S_1 subunit and
P681H, T716I, S982A and D1118H in the S_2 subunit. All constructs also contained D614G.
We found that among 15 individuals with neutralisation activity three weeks after receiving a
single dose of the the BNT162b2 mRNA vaccine, 10 showed evidence of reduction in
efficacy of antibodies against the B.1.1.7 mutant (**Fig. 1c-d**). The highest fold change was
approximately 6. Amongst sera from 7 recovered individuals, only 2 demonstrated reduced
potency against B.1.1.7 (**Fig. 1e-h** and **Extended Data Fig. 4**).

B.1.1.7 variant escapes from NTD- and RBM-specific mAb-mediated neutralization.
To investigate the role of the full set of mutations in NTD, RBD and S2 present in the B.1.1.7
variant, we tested 61 mAbs isolated from 15 individuals that recovered from WT SARS-
CoV-2 infection with an *in-vitro* pseudotyped neutralization assay using VeroE6 target cells
expressing TMPRSS2 (**Extended Data Table 1**). We found that 20 out of 61 (32.8%) mAbs
showed a greater than 2-fold loss of neutralizing activity of B.1.1.7 variant compared to WT

SARS-CoV-2 (**Fig. 2a,b** and **Extended Data Fig. 5**). Remarkably, the B.1.1.7 mutant virus

was found to fully escape neutralization by 7 out of 10 NTD-targeting mAbs (70%), and

partial escape from additional 2 mAbs (20%) (**Fig. 2c**). We previously showed that the

deletion of residue 144 abrogates binding by 4 out of 6 NTD-specific mAbs tested, possibly

accounting for viral neutralization escape by most NTD-specific antibodies[22]. Of the 29

RBM-targeting mAbs, 5 (17.2%) showed more then 100-fold decrease in B.1.1.7

neutralization, and additional 6 mAbs (20.7%) had a partial 2-to-10-fold reduction (**Fig. 2d**).

Finally, all RBD-specific non-RBM-targeting mAbs tested fully retained B.1.1.7 neutralizing

activity (**Fig. 2e**).

To address the role of B.1.1.7 N501Y mutation in the neutralization escape from

RBM-specific antibodies, we tested the binding of 51 RBD-specific mAbs to WT and N501Y

mutant RBD by biolayer interferometry (**Fig. 2f** and **Extended Data Fig. 6**). The 5 RBM-

specific mAbs that failed to neutralize B.1.1.7 variant (**Fig. 2d**) showed a complete loss of

binding to N501Y RBD mutant (**Fig. 2g, h**), demonstrating a critical role for this mutation as

an escape mechanism for RBM-targeting mAbs.

The decreased neutralizing activity of the immune sera from vaccinees and

convalescent patients against B.1.1.7, but not against Δ69/70-501Y-570D mutant (**Fig. 1** and

Extended Data Fig. 2), could be the result of a loss of neutralizing activity of both RBD- and

NTD-targeting antibodies, and suggests that the key mutations driving polyclonal escape are

Δ144 and N501Y.

SARS-CoV-2 B.1.1.7 binds human ACE2 with higher affinity than WT

SARS-CoV-2 and SARS-CoV enter host cells through binding of the S glycoprotein to

ACE2[1,23]. Previous studies showed that the binding affinity of SARS-CoV for human ACE2

correlated with the rate of viral replication in distinct species, transmissibility and disease

severity [24-26]. To understand the potential contribution of receptor interaction to infectivity,

we set out to evaluate the influence of the B.1.1.7 RBD substitution N501Y on receptor

engagement. We used biolayer interferometry to study binding kinetics and affinity of the

purified human ACE2 ectodomain (residues 1-615) to immobilized biotinylated SARS-CoV-

2 B.1.1.7 or WT RBDs. We found that ACE2 bound to the B.1.1.7 RBD with an affinity of

22 nM compared to 133 nM for the WT RBD (**Extended Data Fig. 7**), in agreement with our

previous deep-mutational scanning measurements using dimeric ACE2[27]. Although ACE2

bound with comparable on-rates to both RBDs, the observed dissociation rate constant was

slower for B.1.1.7 than for the WT RBD (**Table 2**). Enhanced binding of the B.1.1.7 RBD to

human ACE2 resulting from the N501Y mutation might participate to the efficient ongoing
transmission of this newly emerged SARS-CoV-2 lineage, and possibly reduced opportunity
for antibody binding.

Further loss of neutralization of vaccine-elicited antibodies against a B.1.1.7 Spike
carrying E484K mutation

The E484K substitution (**Fig 3a**) has been identified as antigenically important, being
reported as an escape mutation for several monoclonal antibodies including C121, C144,
REGN10933 and REGN10934. E484K is also known to be present in the B.1.351 (501Y.V2)
and the P.1 (501Y.V3) lineages in combination with amino acid replacements at N501 and
K417. As of 1st Feb 2021, fourteen English and two Welsh B.1.1.7 sequences from viral
isolates contained the E484K substitution (**Fig. 3b**), The number of B.1.1.7 sequences has
been increasing since the start of December 2020 (**Fig. 3c**). Phylogenetic analysis suggests
that there were two independent acquisitions in English and Welsh sequences (**Fig. 3b**).

We generated pseudoviruses bearing B.1.1.7 Spike mutations with or without additional
E484K and tested these against vaccine sera. We observed a significant loss of neutralising
activity compared with WT and with B.1.1.7; mean fold change for the E484K B.1.1.7 Spike
was   9.6   compared   to   2.4   for   B.1.1.7   alone   (p<0.05)   (**Fig.   4**).

Discussion

Serum neutralizing activity is a correlate of protection for other respiratory viruses, including influenza[28] and respiratory syncytial virus where prohylaxis with monoclonal antibodies has been used in at-risk groups[29,30]. Neutralising antibody titres appeared to be highly correlated with vaccine protection against SARS-CoV-2 rechallenge in non-human primates, and importantly, there was no correlation between T cell responses (as measured by ELISPOT) and protection[31]. Moreover, passive transfer of purified polyclonal IgGs from convalescent macaques protected naïve macaques against subsequent SARS-CoV-2 challenge[32]. Coupled with multiple reports of re-infection, there has therefore been significant attention placed on virus neutralisation.

This study reports on the neutralisation escape from sera collected after one dose of the BNT162b2 vaccine. The participants of this study had a median age of over 80, in line with the targeting of this age group in the initial rollout of the vaccination campaign in the UK. Participants showed similar neutralising activity against wild type pseudovirus as in the phase I/II study, with geometric mean titres of 24 and 29, respectively[12]. The three mutations in S (N501Y, A570D, ΔH69/V70) did not appear to impact neutralisation in a pseudovirus assay. However, we demonstrated that a pseudovirus bearing S protein with the full set of mutations present in the B.1.1.7 variant (i.e., ΔH69/V70, Δ144, N501Y, A570D, P681H, T716I, S982A, D1118H) did result in small reduction in neutralisation by sera from vaccinees. A reduction in neutralization titres from mRNA-elicited antibodies in volunteers who received two doses (using both mRNA-1273 and BNT162b2 vaccines) was also observed by Wang et al.[33] using pseudoviruses carrying the N501Y mutation. Another study reported on a modest and not significant (average 1.2 fold) reduction of neutralization against the B.1.1.7 variant in sera from individuals vaccinated with two doses of mRNA-1273[34]. The level of neutralizing antibodies observed in both of these studies was approximately one log higher than the one observed in the cohort of vaccinees described in this study. This may be related to the older age of the individuals from this study as well as to the fact that these were exposed to only one dose of the BNT162b2 vaccine. In general, it is expected that the effect of mutations on the neutralization by polyclonal serum antibodies might be more prominent on low-titre in contrast to high-titre sera. It is important to study virus neutralisation at lower serum neutralisation titres because decline in neutralisation titres over time is expected to occur following vaccination.

The reduced neutralizing activity observed with polyclonal antibodies elicited by
mRNA vaccines observed in this study and in Wang et al.[33] is further supported by the loss of
neutralizing activity observed with human mAbs directed to both RBD and, to a major extent,
to NTD. In the study by Wang et al., 6 out 17 RDB-specific mAbs isolated from mRNA-1273
vaccinated individuals showed more than 100-fold neutralization loss against N501Y mutant,
a finding that is consistent with the loss of neutralization by 5 out 29 RBM-specific mAbs
described in this study.

Multiple variants, including the 501Y.V2 and B.1.1.7 lineages, harbor multiple
mutations as well as deletions in NTD, most of which are located in a site of vulnerability that
is targeted by all known NTD-specific neutralizing antibodies[22]. The role of NTD-specific
neutralizing antibodies might be under-estimated, in part by the use of neutralization assays
based on target cells over-expressing ACE2 receptor. NTD-specific mAbs were suggested to
interfere with viral entry based on other accessory receptors, such as DC-SIGN and L-SIGN[35],
and their neutralization potency was found to be dependent on different in vitro culture
conditions[22]. The observation that 9 out of 10 NTD-specific neutralizing antibodies failed to
show a complete or near-complete loss of sneutralizing activity against B.1.1.7 indicates that
this new variant may have evolved also to escape from this class of antibodies, that may have
a yet unrecognized role in protective immunity. Wibmer et al.[36] have also recently reported
the loss of neutralization of 501Y.V2 by the NTD-specific mAb 4A8, likely driven by the
R246I mutation. This result is in line with the lack of neutralization of B.1.1.7 by 4A8 mAb
observed in this study, likely caused by Δ144 due to loss of binding[22]. Finally, the role of
NTD mutations (in particular, L18F, Δ242-244 and R246I) was further supported by the
marked loss of neutralization observed by Wibmer et al.[36] against 501Y.V2 compared to the
chimeric virus carrying only the RBD mutations K417N, E484K and N501Y. Taken together,
the presence of multiple escape mutations in NTD is supportive of the hypothesis that this
region of the Spike, in addition to RBM, is also under immune pressure.

E484K has been shown to impact neutralisation by monoclonal antibodies or
convalescent sera, especially in combination with N501Y and K417N[16,37-39]. Worryingly, we
have shown that there are multiple B.1.1.7 sequences in the UK bearing E484K with early
evidence of transmission as well as independent aquisitions. We measured further reduction
neutralisation titers by vaccine sera when E484K was present alongside the B.1.1.7 S
mutations. Consistent with our findings, Wu and co-authors[34] have shown that variants

carrying the E484K mutation resulted in 3-to-6 fold reduction in neutralization by sera from

mRNA-1273 vaccinated individuals.

Evidence for the importance role of NTD deletions in combination with E484K in immune

escape is provided by Andreano *et al*. who describe the emergence of Δ140 in virus co-

incubated with potently neutralizing convalescent plasma, causing a 4-fold reduction in

neutralization titre. This Δ140 mutant subsequently acquired E484K which resulted in a

further 4-fold drop in neutralization titre indicating a two residue change across NTD and

RBD represents an effective pathway of escape that can dramatically inhibit the polyclonal

response.

Our study was limited by relatively small sample size. Although the Spike pseudotyping

system has been shown to faithfully represent full length infectious virus, there may be

determinants outside the S that influence escape from antibody neutralization either directly

or indirectly in a live replication competent system. On the other hand live virus systems

allow replication and therefore mutation to occur, and rigorous sequencing at multiple steps is

needed.

Amidst high transmission in many parts of the world, vaccines are a key part of a long term

strategy to bring SARS-CoV-2 under control. Our data suggest that vaccine escape will be

inevitable in the future, and should be mitigated by designing next generation vaccines with

mutated S sequences and using alternative viral antigens. Currently however, vaccines are

likely to contribute controlling SARS-CoV-2 infections in the short term.

Acknowledgements

We would like to thank Cambridge University Hospitals NHS Trust Occupational Health

Department. We would also like to thank the NIHR Cambridge Clinical Research Facility

and staff at CUH and. We would like to thank Eleanor Lim and Georgina Okecha. We thank

Dr James Voss for the kind gift of HeLa cells stably expressing ACE2. RKG is supported by a

Wellcome Trust Senior Fellowship in Clinical Science (WT108082AIA). LEM is supported

by a Medical Research Council Career Development Award (MR/R008698/1). SAK is

supported by the Bill and Melinda Gates Foundation via PANGEA grant: OPP1175094. DAC

is supported by a Wellcome Trust Clinical PhD Research Fellowship. KGCS is the recipient

of a Wellcome Investigator Award (200871/Z/16/Z). This research was supported by the
National Institute for Health Research (NIHR) Cambridge Biomedical Research Centre, the
Cambridge Clinical Trials Unit (CCTU), and the NIHR BioResource. The views expressed
are those of the authors and not necessarily those of the NIHR or the Department of Health
and Social Care. JAGB is supported by the Medical Research Council (MC_UP_1201/16).
IATM is funded by a SANTHE award.

Author contributions

Conceived study: D.C., RKG, DAC. Designed study and experiments: RKG, DAC, LEM, JB,
MW, JT, LCG, GBM, RD, BG, NK, AE, M.P., D.V., L.P., A.D.M, J.B., D.C. Performed
experiments: BM, DAC, RD, IATMF, LCG, GBM. Interpreted data: RKG, DAC, BM, RD,
IATMF, LEM, JB, KGCS. A.D.M., J.B. and C.S.F. carried out pseudovirus neutralization
assays. D.P. produced pseudoviruses. M.S.P., L.P., D.V. and D.C. designed the experiments.
A.C.W., N.S. and S.J. expressed and purified the proteins. K.C., S.J. and E.C. sequenced and
expressed antibodies. E.C. and K.C. performed mutagenesis for mutant expression plasmids.
A.C.W. M.A.T., J.E.B., and S.B. performed binding assays. A.R., A.F.P and C.G contributed
to donor's recruitment and sample collection related to mAbs isolation. H.W.V., G.S., A.L.,
D.V., L.P. and D.C. analyzed the data and prepared the manuscript with input from all
authors.

Competing interests

A.D.M., J.B., D.P., C.S.F., S.B., K.C., N.S., E.C., G.S., S.J., A.L., H.W.V., M.S.P., L.P. and
D.C. are employees of Vir Biotechnology and may hold shares in Vir Biotechnology. H.W.V.
is a founder of PierianDx and Casma Therapeutics. Neither company provided funding for
this work or is performing related work. D.V. is a consultant for Vir Biotechnology Inc. The
Veesler laboratory has received a sponsored research agreement from Vir Biotechnology Inc.
The remaining authors declare that the research was conducted in the absence of any
commercial or financial relationships that could be construed as a potential conflict of
interest. RKG has received consulting fees from UMOVIS Lab, Gilead and ViiV.

MATERIALS AND METHODS

Participant recruitment and ethics

Participants who had received the first dose of vaccine and individuals with COVID-19 were

consented into the Covid-19 cohort of the NIHR Bioresource. The study was approved by the

East of England – Cambridge Central Research Ethics Committee (17/EE/0025).

SARS-CoV-2 serology by multiplex particle-based flow cytometry (Luminex):

Recombinant SARS-CoV-2 N, S and RBD were covalently coupled to distinct carboxylated

bead sets (Luminex; Netherlands) to form a 3-plex and analyzed as previously described

(Xiong et al. 2020). Specific binding was reported as mean fluorescence intensities (MFI).

Recombinant expression of SARS-CoV-2-specific mAbs.

Human mAbs were isolated from plasma cells or memory B cells of SARS-CoV-2 immune

donors, as previously described [40-42]. Recombinant antibodies were expressed in ExpiCHO

cells at 37°C and 8% CO_2. Cells were transfected using ExpiFectamine. Transfected cells

were supplemented 1 day after transfection with ExpiCHO Feed and ExpiFectamine CHO

Enhancer. Cell culture supernatant was collected eight days after transfection and filtered

through a 0.2 µm filter. Recombinant antibodies were affinity purified on an ÄKTA xpress

FPLC device using 5 mL HiTrap™ MabSelect™ PrismA columns followed by buffer

exchange to Histidine buffer (20 mM Histidine, 8% sucrose, pH 6) using HiPrep 26/10

desalting columns

Generation of S mutants

Amino acid substitutions were introduced into the D614G pCDNA_SARS-CoV-2_S plasmid

as previously described[43] using the QuikChange Lightening Site-Directed Mutagenesis kit,

following the manufacturer's instructions (Agilent Technologies, Inc., Santa Clara, CA).

Sequences were checked by Sanger sequencing.

Preparation of B.1.1.7 SARS-CoV-2 S glycoprotein-encoding-plasmid used to produce

SARS-CoV-2-MLV based on overlap extension PCR. Briefly, a modification of the overlap

extension PCR protocol[44] was used to introduce the nine mutations of the B.1.1.7 lineage in

the SARS-CoV-2 S gene. In a first step, 9 DNA fragments with overlap sequences were

amplified by PCR from a plasmid (phCMV1, Genlantis) encoding the full-length SARS-

CoV-2 S gene (BetaCoV/Wuhan-Hu-1/2019, accession number mn908947). The

mutations (del-69/70, del-144, N501Y, A570D, D614G, P681H, S982A, T716I and

D1118H) were introduced by amplification with primers with similar Tm. Deletion of the C-

terminal 21 amino acids was introduced to increase surface expression of the recombinant

S[45]. Next, 3 contiguous overlapping fragments were fused by a first overlap PCR (step

2) using the utmost external primers of each set, resulting in 3 larger fragments with

overlapping sequences. A final overlap PCR (step 3) was performed on the 3 large

fragments using the utmost external primers to amplify the full-length S gene and the

flanking sequences including the restriction sites KpnI and NotI. This fragment was digested

and cloned into the expression plasmid phCMV1. For all PCR reactions the Q5 Hot Start

High fidelity DNA polymerase was used (New England Biolabs Inc.), according to the

manufacturer's instructions and adapting the elongation time to the size of the amplicon.

After each PCR step the amplified regions were separated on agarose gel and purified

using Illustra GFX™ PCR DNA and Gel Band Purification Kit (Merck KGaA).

Pseudotype virus preparation

Viral vectors were prepared by transfection of 293T cells by using Fugene HD transfection

reagent (Promega). 293T cells were transfected with a mixture of 11ul of Fugene HD, 1µg of

pCDNAΔ19Spike-HA, 1ug of p8.91 HIV-1 gag-pol expression vector[46,47], and 1.5µg of

pCSFLW (expressing the firefly luciferase reporter gene with the HIV-1 packaging signal).

Viral supernatant was collected at 48 and 72h after transfection, filtered through 0.45um filter

and stored at -80°C. The 50% tissue culture infectious dose (TCID$_{50}$) of SARS-CoV-2

pseudovirus was determined using Steady-Glo Luciferase assay system (Promega).

Serum/plasma pseudotype neutralization assay

Spike pseudotype assays have been shown to have similar characteristics as neutralisation

testing using fully infectious wild type SARS-CoV-2[20]. Virus neutralisation assays were

performed on 293T cell transiently transfected with ACE2 and TMPRSS2 using SARS-CoV-

2 Spike pseudotyped virus expressing luciferase[48]. Pseudotyped virus was incubated with

serial dilution of heat inactivated human serum samples or sera from vaccinees in duplicate

for 1h at 37°C. Virus and cell only controls were also included. Then, freshly trypsinized

293T ACE2/TMPRSS2 expressing cells were added to each well. Following 48h incubation

in a 5% CO_2 environment at 37°C, luminescence was measured using the Steady-Glo

Luciferase assay system (Promega).

IFNγ FLUOROSPOT assays

Frozen PBMCs were rapidly thawed, and the freezing medium was diluted into 10ml of

TexMACS media (Miltenyi Biotech), centrifuged and resuspended in 10ml of fresh media

with 10U/ml DNase (Benzonase, Merck-Millipore via Sigma-Aldrich), PBMCs were

incubated at 37°C for 1h, followed by centrifugation and resuspension in fresh media

supplemented with 5% Human AB serum (Sigma Aldrich) before being counted. PBMCs

were stained with 2ul of each antibody: anti-CD3-fluorescein isothiocyanate (FITC), clone

UCHT1; anti-CD4-phycoerythrin (PE), clone RPA-T4; anti-CD8a-peridinin-chlorophyll

protein - cyanine 5.5 (PerCP Cy5.5), clone RPA-8a (all BioLegend, London, UK),

LIVE/DEAD Fixable Far Red Dead Cell Stain Kit (Thermo Fisher Scientific). PBMC

phenotyping was performed on the BD Accuri C6 flow cytometer. Data were analysed with

FlowJo v10 (Becton Dickinson, Wokingham, UK). 1.5 to 2.5 x 105 PBMCs were incubated

in pre-coated Fluorospot plates (Human IFNγ FLUOROSPOT (Mabtech AB, Nacka Strand,

Sweden)) in triplicate with peptide mixes specific for Spike, Nucleocapsid and Membrane

proteins of SARS-CoV-2 (final peptide concentration 1µg/ml/peptide, Miltenyi Biotech) and

an unstimulated and positive control mix (containing anti-CD3 (Mabtech AB),

Staphylococcus Enterotoxin B (SEB), Phytohaemagglutinin (PHA) (all Sigma Aldrich)) at

37ºC in a humidified CO2 atmosphere for 48 hours. The cells and medium were decanted

from the plate and the assay developed following the manufacturer's instructions. Developed

plates were read using an AID iSpot reader (Oxford Biosystems, Oxford, UK) and counted

using AID EliSpot v7 software (Autoimmun Diagnostika GmbH, Strasberg, Germany). All

data were then corrected for background cytokine production and expressed as SFU/Million

PBMC or CD3 T cells.

406
MAbs pseudovirus neutralization assay

MLV-based SARS-CoV-2 S-glycoprotein-pseudotyped viruses were prepared as previously

described (Pinto et al., 2020). HEK293T/17cells were cotransfected with a WT or B.1.1.7

SARS-CoV-2 Spike glycoprotein-encoding-plasmid, an MLV Gag-Pol packaging construct

and the MLV transfer vector encoding a luciferase reporter using X-tremeGENE HP

transfection reagent (Roche) according to the manufacturer's instructions. Cells were cultured

for 72 h at 37°C with 5% CO_2 before harvesting the supernatant. VeroE6 stably expressing

human TMPRSS2 were cultured in DMEM containing 10% FBS, 1% penicillin–

streptomycin, 8 µg/mL puromycin and plated into 96-well plates for 16–24 h. Pseudovirus

with serial dilution of mAbs was incubated for 1 h at 37°C and then added to the wells after

washing 2 times with DMEM. After 2–3 h DMEM containing 20% FBS and 2% penicillin–

streptomycin was added to the cells. Following 48-72 h of infection, Bio-Glo (Promega) was

added to the cells and incubated in the dark for 15 min before reading luminescence with

Synergy H1 microplate reader (BioTek). Measurements were done in duplicate and relative

luciferase units were converted to percent neutralization and plotted with a non-linear

regression model to determine IC50 values using GraphPad PRISM software (version 9.0.0).

Antibody binding measurements using bio-layer interferometry (BLI)

MAbs were diluted to 3 µg/ml in kinetic buffer (PBS supplemented with 0.01% BSA) and

immobilized on Protein A Biosensors (FortéBio). Antibody-coated biosensors were

incubated for 3 min with a solution containing 5 µg /ml of WT or N50Y SARS-CoV-2

RBD in kinetic buffer, followed by a 3-min dissociation step. Change in molecules bound to

the biosensors caused a shift in the interference pattern that was recorded in real time using

an Octet RED96 system (FortéBio). The binding response over time was used to calculate

the area under the curve (AUC) using GraphPad PRISM software (version 9.0.0).

Production of SARS-CoV-2 and B.1.1.7 receptor binding domains and human ACE2

The SARS-CoV-2 RBD (BEI NR-52422) construct was synthesized by GenScript into

CMVR with an N-terminal mu-phosphatase signal peptide and a C-terminal octa-histidine tag

(GHHHHHHHH) and an avi tag. The boundaries of the construct are N-$_{328}$RFPN$_{331}$ and

$_{528}$KKST$_{531}$-C[49]. The B.1.1.7 RBD was synthesized by GenScript into pCMVR with the same

boundaries and construct details with a mutation at N501Y. These plasmids were transiently

transfected into Expi293F cells using Expi293F expression medium (Life Technologies) at

37°C 8% CO_2 rotating at 150 rpm. The cultures were transfected using PEI cultivated for

5 days. Supernatants were clarified by centrifugation (10 min at 4000xg) prior to loading onto

a nickel-NTA column (GE). Purified protein was biotinylated overnight using BirA (Avidity)

prior to SEC into PBS. Human ACE2-Fc (residues 1-615 with a C-terminal thrombin

cleavage site and human Fc tag) were synthesized by Twist. Clarified supernatants were

affinity purified using a Protein A column (GE LifeSciences) directly neutralized and buffer

exchanged. The Fc tag was removed by thrombin cleavage in a reaction mixture containing

3 mg of recombinant ACE2-FC ectodomain and 10 µg of thrombin in 20 mM Tris-HCl

pH8.0, 150 mM NaCl and 2.5 mM $CaCl_2$.The reaction mixture was incubated at 25°C

overnight and re-loaded on a Protein A column to remove uncleaved protein and the Fc tag.

The cleaved protein was further purified by gel filtration using a Superdex 200 column

10/300 GL (GE Life Sciences) equilibrated in PBS.

Protein affinity measurement using bio-layer interferometry

Biotinylated RBD (either WT or N501Y) were immobilized at 5 ng/uL in undiluted 10X
Kinetics Buffer (Pall) to SA sensors until a load level of 1.1nm, A dilution series of either
monomeric ACE2 or Fab in undiluted kinetics buffer starting at 1000-50nM was used for
300-600 seconds to determine protein-protein affinity. The data were baseline subtracted and
the plots fitted using the Pall FortéBio/Sartorius analysis software (version 12.0). Data were
plotted in Prism.

462　**Supplementary Table 1:** Vaccinee Participant ages

ID	Age
1	60-65
2	80-85
3	40-45
4	85-90
5	25-30
6	35-40
7	80-85
8	80-85
9	55-60
10	85-90
11	80-85
12	80-85
13	80-85
14	85-90
15	75-80
16	80-85
17	80-85
18	85-90
19	85-90
20	80-85
22	55-60
25	60-65

470 **Table 2. Kinetic analysis of human ACE2 binding to SARS-CoV-2 B.1.1.7 and Wuhan-1**
471 **RBDs by biolayer interferometry.** Values reported represent the global fit to the data shown
472 in Extended Data Fig. 7.

		SARS-CoV-2 RBD WT	SARS-CoV-2 RBD B.1.1.7
K_D (nM)		133	22
k_{on} (M^{-1}.s^{-1})	hACE2	$1.3*10^5$	$1.4*10^5$
k_{off} (s^{-1})		$1.8*10^{-2}$	$3*10^{-3}$

Extended Data Table 1. Neutralization, V gene usage and other properties of tested mAbs.

mAb	Domain (site)	VH usage (% id.)	Source (DSO)	IC50 WT (ng/ml)	IC50 B.1.1.7 (ng/ml)	ACE2 blocking	SARS-CoV	Escape residues	Ref.
4A8	NTD (i)	1-24	N/A	38	-	Neg.	-	S12P; C136Y; Y144del; H146Y; K147T; R246A	[50]
S2L26	NTD (i)	1-24 (97.2)	Hosp. (52)	70	-	Neg.	-	N/A	[22]
S2L50	NTD (i)	4-59 (95.4)	Hosp. (52)	264	50	Neg.	-	N/A	[22]
S2M28	NTD (i)	3-33 (97.6)	Hosp. (46)	295	12'207	Neg.	-	P9S/Q; S12P; C15F/R; L18P; Y28C; A123T; C136Y; G142D; Y144del; K147Q/T; R246G; P251L; G252C	[22]
S2X107	NTD (i)	4-38-2 (97)	Sympt. (75)	388	-	Neg.	-	N/A	[22]
S2X124	NTD (i)	3-30 (99)	Sympt. (75)	221	-	Neg.	-	N/A	[22]
S2X158	NTD (i)	1-24 (96.3)	Sympt. (75)	56	-	Neg.	-	N/A	[22]
S2X28	NTD (i)	3-30 (97.9)	Sympt. (48)	1'399	-	Neg.	-	P9S; S12P; C15W; L18P; C136G/Y; F140S; L141S; G142C/D; Y144C/N; K147T/Q/E; R158G; L244S; R246G	[22]
S2X303	NTD (i)	2-5 (95.9)	Sympt. (125)	69	-	Neg.	-	N/A	[22]
S2X333	NTD (i)	3-33 (96.5)	Sympt. (125)	66	-	Neg.	-	P9L; S12P; C15S/Y; L18P; C136G/Y; F140C; G142D; K147T	[22]
S2D106	RBD (I/RBM)	1-69 (97.2)	Hosp. (98)	27	20	Strong	-	N/A	[8]
S2D19	RBD (I/RBM)	4-31 (99.7)	Hosp. (49)	128	75'200	Moderate	-	N/A	[8]
S2D32	RBD (I/RBM)	3-49 (98.3)	Hosp. (49)	26	11	Strong	-	N/A	[8]
S2D65	RBD (I/RBM)	3-9 (96.9)	Hosp. (49)	24	12	Weak	-	N/A	[8]
S2D8	RBD (I/RBM)	3-23 (96.5)	Hosp. (49)	27	58'644	Strong	-	N/A	[8]
S2D97	RBD (I/RBM)	2-5 (96.9)	Hosp. (98)	20	17	Weak	-	N/A	[8]
S2E11	RBD (I/RBM)	4-61 (98.3)	Hosp. (51)	27	16	Weak	-	N/A	[8]
S2E12	RBD (I/RBM)	1-58 (97.6)	Hosp. (51)	27	31	Strong	-	G476S (3x)	[8,51]
S2E13	RBD (I/RBM)	1-18 (96.2)	Hosp. (51)	34	77	Strong	-	N/A	[8]
S2E16	RBD (I/RBM)	3-30 (98.3)	Hosp. (51)	36	38	Strong	-	N/A	[8]
S2E23	RBD (I/RBM)	3-64 (96.9)	Hosp. (51)	139	180	Strong	-	N/A	[8]
S2H14	RBD (I/RBM)	3-15 (100)	Sympt. (17)	460	64'463	Weak	-	N/A	[8,52]
S2H19	RBD (I/RBM)	3-15 (98.6)	Sympt. (45)	239	-	Weak	-	N/A	[8]
S2H58	RBD (I/RBM)	1-2 (97.9)	Sympt. (45)	27	14	Strong	-	N/A	[8]
S2H7	RBD (I/RBM)	3-66 (98.3)	Sympt. (17)	492	573	Weak	-	N/A	[8]
S2H70	RBD (I/RBM)	1-2 (99)	Sympt. (45)	147	65	Weak	-	N/A	[8]
S2H71	RBD (I/RBM)	2-5 (99)	Sympt. (45)	36	9	Moderate	-	N/A	[8]
S2M11	RBD (I/RBM)	1-2 (96.5)	Hosp. (46)	11	4	Weak	-	Y449N; L455F; E484K; E484Q; F490L; F490S; S494P	[8,51]
S2N12	RBD	4-39 (97.6)	Hosp. (51)	76	40	Strong	-	N/A	[8]

	(I/RBM)								
S2N22	RBD (I/RBM)	3-23 (96.5)	Hosp. (51)	32	21	Strong	-	N/A	[8]
S2N28	RBD (I/RBM)	3-30 (97.2)	Hosp. (51)	72	21	Strong	-	N/A	[8]
S2X128	RBD (I/RBM)	1-69-2 (97.6)	Sympt. (75)	50	112	Strong	-	N/A	[8]
S2X16	RBD (I/RBM)	1-69 (97.6)	Sympt. (48)	45	103	Strong	-	N/A	[8]
S2X192	RBD (I/RBM)	1-69 (96.9)	Sympt. (75)	326	-	Weak	-	N/A	[8]
S2X227	RBD (I/RBM)	1-46 (97.9)	Sympt. (75)	26	14	Strong	-	N/A	
S2X246	RBD (I/RBM)	3-48 (96.2)	Sympt. (75)	35	30	Strong	-	N/A	
S2X30	RBD (I/RBM)	1-69 (97.9)	Sympt. (48)	32	53	Strong	-	N/A	[8]
S2X324	RBD (I/RBM)	2-5 (97.3)	Sympt. (125)	8	23	Strong	-	N/A	
S2X58	RBD (I/RBM)	1-46 (99)	Sympt. (48)	32	47	Strong	-	N/A	[8]
S2H90	RBD (II)	4-61 (96.6)	Sympt. (81)	77	32	Strong	+	N/A	[8]
S2H94	RBD (II)	3-23 (93.4)	Sympt. (81)	123	144	Strong	+	N/A	[8]
S2H97	RBD (V)	5-51 (98.3)	Sympt. (81)	513	248	Weak	+	N/A	
S2K15	RBD (II)	2-26 (99.3)	Sympt. (87)	361	235	0	+	N/A	
S2K21	RBD (II)	3-33 (96.2)	Sympt. (118)	201	189	0	+	N/A	
S2K30	RBD (II)	1-2 (97.2)	Sympt. (87)	185	134	0	+	N/A	
S2K63v2	RBD (II)	3-30-3 (95.6)	Sympt. (118)	144	215	0	+	N/A	
S2L17	RBD (?)	5-10-1 (98.3)	Hosp. (51)	313	127	Moderate	+	N/A	[8]
S2L37	RBD (II)	3-13 (98.3)	Hosp. (51)	14'370	824	Moderate	+	N/A	
S2L49	RBD (?)	3-30 (97.9)	Hosp. (51)	24	32	Neg.	+	N/A	[8]
S2X259	RBD (IIa)	1-69 (94.1)	Sympt. (75)	145	91	Moderate	+	N/A	
S2X305	RBD (?)	1-2 (95.1)	Sympt. (125)	34	21	Strong	-	N/A	
S2X35	RBD (IIa)	1-18 (98.6)	Sympt. (48)	140	143	Strong	+	N/A	[52]
S2X450	RBD (?)	2-26 (96.9)	Sympt. (271)	368	198	Strong	+	N/A	
S2X475	RBD (?)	3-21 (93.8)	Sympt. (271)	1'431	851	Strong	+	N/A	
S2X607	RBD (?)	3-66 (95.4)	Sympt. (271)	41	23	Strong	-	N/A	
S2X608	RBD (?)	1-33 (93.2)	Sympt. (271)	21	35	Strong	-	N/A	
S2X609	RBD (?)	1-69 (93.8)	Sympt. (271)	47	35	Strong	-	N/A	
S2X613	RBD (?)	1-2 (91.7)	Sympt. (271)	28	19	Strong	-	N/A	
S2X615	RBD (?)	3-11 (94.8)	Sympt. (271)	23	17	Strong	-	N/A	
S2X619	RBD (?)	1-69 (92.7)	Sympt. (271)	36	60	Strong	-	N/A	
S2X620	RBD (?)	3-53 (95.1)	Sympt. (271)	34	45	Strong	-	N/A	

id., identity. DSO, days after symptom onset. * as described in Piccoli et al and McCallum et al. N/A, not available; -, not neutralizing

References

1 Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable
bat origin. *Nature* **579**, 270-273, doi:10.1038/s41586-020-2012-7 (2020).
2 Davies, N. G. *et al.* Estimated transmissibility and severity of novel SARS-CoV-2
Variant of Concern 202012/01 in England. *medRxiv*, 2020.2012.2024.20248822,
doi:10.1101/2020.12.24.20248822 (2020).
3 Volz, E. *et al.* Transmission of SARS-CoV-2 Lineage B.1.1.7 in England: Insights from
linking epidemiological and genetic data. *medRxiv*, 2020.2012.2030.20249034,
doi:10.1101/2020.12.30.20249034 (2021).
4 Korber, B. *et al.* Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G
Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819,
doi:10.1016/j.cell.2020.06.043 (2020).
5 Yurkovetskiy, L. *et al.* Structural and Functional Analysis of the D614G SARS-CoV-2
Spike Protein Variant. *Cell* **183**, 739-751 e738, doi:10.1016/j.cell.2020.09.032 (2020).
6 Martinot, M. *et al.* Remdesivir failure with SARS-CoV-2 RNA-dependent RNA-
polymerase mutation in a B-cell immunodeficient patient with protracted Covid-19.
Clin Infect Dis, doi:10.1093/cid/ciaa1474 (2020).
7 Kemp, S. *et al.* Neutralising antibodies in Spike mediated SARS-CoV-2 adaptation.
medRxiv, 2020.2012.2005.20241927, doi:10.1101/2020.12.05.20241927 (2020).
8 Thomson, E. C. *et al.* The circulating SARS-CoV-2 spike variant N439K maintains
fitness while evading antibody-mediated immunity. *bioRxiv*, 1-49,
doi:papers3://publication/doi/10.1101/2020.11.04.355842 (2020).
9 Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N Engl
J Med*, doi:10.1056/NEJMoa2035389 (2020).
10 Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N
Engl J Med* **383**, 2603-2615, doi:10.1056/NEJMoa2034577 (2020).
11 Voysey, M. *et al.* Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222)
against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil,
South Africa, and the UK. *Lancet* **397**, 99-111, doi:10.1016/S0140-6736(20)32661-1
(2021).
12 Mulligan, M. J. *et al.* Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults.
Nature **586**, 589-593, doi:10.1038/s41586-020-2639-4 (2020).
13 Corbett, K. S. *et al.* SARS-CoV-2 mRNA Vaccine Development Enabled by Prototype
Pathogen Preparedness. *bioRxiv*, 2020.2006.2011.145920,
doi:10.1101/2020.06.11.145920 (2020).
14 Folegatti, P. M. *et al.* Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine
against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised
controlled trial. *Lancet* **396**, 467-478, doi:10.1016/S0140-6736(20)31604-4 (2020).
15 Kemp, S. A. *et al.* Recurrent emergence and transmission of a SARS-CoV-2 Spike
deletion ΔH69/V70. *bioRxiv*, 2020.2012.2014.422555,
doi:10.1101/2020.12.14.422555 (2020).
16 Tegally, H. *et al.* Emergence and rapid spread of a new severe acute respiratory
syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations
in South Africa. *medRxiv*, 2020.2012.2021.20248640,
doi:10.1101/2020.12.21.20248640 (2020).

17   Faria, N. R. *et al.* *Genomic characterisation of an emergent SARS-CoV-2 lineage in*
Manaus: preliminary findings, <https://virological.org/t/genomic-characterisation-
of-an-emergent-sars-cov-2-lineage-in-manaus-preliminary-findings/586> (2021).
18   Jackson, L. A. *et al.* An mRNA Vaccine against SARS-CoV-2 - Preliminary Report. *N*
Engl J Med **383**, 1920-1931, doi:10.1056/NEJMoa2022483 (2020).
19   Walsh, E. E. *et al.* Safety and Immunogenicity of Two RNA-Based Covid-19 Vaccine
Candidates. *New England Journal of Medicine* **383**, 2439-2450,
doi:10.1056/NEJMoa2027906 (2020).
20   Schmidt, F. *et al.* Measuring SARS-CoV-2 neutralizing antibody activity using
pseudotyped and chimeric viruses. 2020.2006.2008.140871,
doi:10.1101/2020.06.08.140871 %J bioRxiv (2020).
21   Brouwer, P. J. M. *et al.* Potent neutralizing antibodies from COVID-19 patients define
multiple targets of vulnerability. *Science* **369**, 643-650, doi:10.1126/science.abc5902
(2020).
22   McCallum, M. *et al.* N-terminal domain antigenic mapping reveals a site of
vulnerability for SARS-CoV-2. *bioRxiv*, doi:10.1101/2021.01.14.426475 (2021).
23   Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS- CoV-2 Spike
Glycoprotein. *Cell* **181**, 281-292.e286,
doi:papers3://publication/doi/10.1016/j.cell.2020.02.058 (2020).
24   Guan, Y. *et al.* Isolation and characterization of viruses related to the SARS
coronavirus from animals in southern China. *Science* **302**, 276-278,
doi:10.1126/science.1087139 (2003).
25   Li, W. *et al.* Efficient replication of severe acute respiratory syndrome coronavirus in
mouse cells is limited by murine angiotensin-converting enzyme 2. *J Virol* **78**, 11429-
11433, doi:10.1128/JVI.78.20.11429-11433.2004 (2004).
26   Li, W. *et al.* Receptor and viral determinants of SARS-coronavirus adaptation to
human ACE2. *EMBO J* **24**, 1634-1643, doi:10.1038/sj.emboj.7600640 (2005).
27   Starr, T. N. *et al.* Deep Mutational Scanning of SARS-CoV-2 Receptor Binding Domain
Reveals Constraints on Folding and ACE2 Binding. *Cell* **182**, 1295-1310 e1220,
doi:10.1016/j.cell.2020.08.012 (2020).
28   Verschoor, C. P. *et al.* Microneutralization assay titres correlate with protection
against seasonal influenza H1N1 and H3N2 in children. *PloS one* **10**, e0131531,
doi:10.1371/journal.pone.0131531 (2015).
29   Kulkarni, P. S., Hurwitz, J. L., Simoes, E. A. F. & Piedra, P. A. Establishing Correlates of
Protection for Vaccine Development: Considerations for the Respiratory Syncytial
Virus Vaccine Field. *Viral Immunol* **31**, 195-203, doi:10.1089/vim.2017.0147 (2018).
30   Goddard, N. L., Cooke, M. C., Gupta, R. K. & Nguyen-Van-Tam, J. S. Timing of
monoclonal antibody for seasonal RSV prophylaxis in the United Kingdom. *Epidemiol*
Infect **135**, 159-162, doi:10.1017/S0950268806006601 (2007).
31   Mercado, N. B. *et al.* Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus
macaques. *Nature* **586**, 583-588, doi:10.1038/s41586-020-2607-z (2020).
32   McMahan, K. *et al.* Correlates of protection against SARS-CoV-2 in rhesus macaques.
Nature, doi:10.1038/s41586-020-03041-6 (2020).
33   Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating
variants. *bioRxiv*, doi:10.1101/2021.01.15.426911 (2021).

34   Wu, K. *et al.* mRNA-1273 vaccine induces neutralizing antibodies against spike
mutants from global SARS-CoV-2 variants. *bioXriv*, doi:10.1101/2021.01.25.427948
(2021).
35   Soh, W. T. *et al.* The N-terminal domain of spike glycoprotein mediates SARS-CoV-2
infection by associating with L-SIGN and DC-SIGN. 1-30,
doi:papers3://publication/doi/10.1101/2020.11.05.369264 (2020).
36   Wibmer, C. K. *et al.* SARS-CoV-2 501Y.V2 escapes neutralization by South African
COVID-19 donor plasma. *bioRxiv*, doi:10.1101/2021.01.18.427166 (2021).
37   Greaney, A. J. *et al.* Comprehensive mapping of mutations to the SARS-CoV-2
receptor-binding domain that affect recognition by polyclonal human serum
antibodies. *bioRxiv*, 2020.2012.2031.425021, doi:10.1101/2020.12.31.425021
(2021).
38   Greaney, A. J. *et al.* Complete mapping of mutations to the SARS-CoV-2 spike
receptor-binding domain that escape antibody recognition. *Cell Host & Microbe*
(2020).
39   Weisblum, Y. *et al.* Escape from neutralizing antibodies by SARS-CoV-2 spike protein
variants. *Elife* **9**, e61312, doi:10.7554/eLife.61312 (2020).
40   Corti, D. *et al.* A neutralizing antibody selected from plasma cells that binds to group
1 and group 2 influenza A hemagglutinins. *Science* **333**, 850-856,
doi:10.1126/science.1205669 (2011).
41   Pinto, D. *et al.* Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295, doi:10.1038/s41586-020-2349-y (2020).
42   Tortorici, M. A. *et al.* Ultrapotent human antibodies protect against SARS-CoV-2
challenge via multiple mechanisms. *Science*, doi:10.1126/science.abe3354 (2020).
43   Gregson, J. *et al.* HIV-1 viral load is elevated in individuals with reverse transcriptase
mutation M184V/I during virological failure of first line antiretroviral therapy and is
associated with compensatory mutation L74I. *Journal of Infectious Diseases* (2019).
44   Forloni, M., Liu, A. Y. & Wajapeyee, N. Creating Insertions or Deletions Using Overlap
Extension Polymerase Chain Reaction (PCR) Mutagenesis. *Cold Spring Harb Protoc*
2018, doi:10.1101/pdb.prot097758 (2018).
45   Case, J. B. *et al.* Neutralizing Antibody and Soluble ACE2 Inhibition of a Replication-
Competent VSV-SARS-CoV-2 and a Clinical Isolate of SARS-CoV-2. *Cell Host Microbe*
28, 475-485 e475, doi:10.1016/j.chom.2020.06.021 (2020).
46   Naldini, L., Blomer, U., Gage, F. H., Trono, D. & Verma, I. M. Efficient transfer,
integration, and sustained long-term expression of the transgene in adult rat brains
injected with a lentiviral vector. *Proc Natl Acad Sci U S A* **93**, 11382-11388,
doi:10.1073/pnas.93.21.11382 (1996).
47   Gupta, R. K. *et al.* Full length HIV-1 gag determines protease inhibitor susceptibility
within in vitro assays. *AIDS* **24**, 1651 (2010).
48   Mlcochova, P. *et al.* Combined point of care nucleic acid and antibody testing for
SARS-CoV-2 following emergence of D614G Spike Variant. *Cell Rep Med*, 100099,
doi:10.1016/j.xcrm.2020.100099 (2020).
49   Walls, A. C. *et al.* Elicitation of Potent Neutralizing Antibody Responses by Designed
Protein Nanoparticle Vaccines for SARS-CoV-2. *Cell* **183**, 1367-1382 e1317,
doi:10.1016/j.cell.2020.10.043 (2020).

50   Chi, X. *et al.* A neutralizing human antibody binds to the N-terminal domain of the
619   Spike protein of SARS-CoV-2. *Science*, eabc6952-6913,
620   doi:papers3://publication/doi/10.1126/science.abc6952 (2020).
621   51   Tortorici, M. A. *et al.* Ultrapotent human antibodies protect against SARS-CoV-2
622   challenge via multiple mechanisms. *Science* **4**, eabe3354-3316,
623   doi:papers3://publication/doi/10.1126/science.abe3354 (2020).
624   52   Piccoli, L. *et al.* Mapping neutralizing and immunodominant sites on the SARS-CoV-2
625   spike receptor-binding domain by structure-guided high-resolution serology. *Cell*, 1-
626   55, doi:papers3://publication/doi/10.1016/j.cell.2020.09.037 (2020).
627

Figure 1

Figure 1. Neutralisation by sera against wild type and B.1.1.7 Spike mutant SARS-CoV-2 pseudotyped viruses. a, Serum IgG responses against N protein, Spike and the Spike Receptor Binding Domain (RBD) from 23 vaccinated participants (green), 20 recovered COVID-19 cases (red), 3 convalescent plasma units and 20 healthy controls (grey) as measured by a flow cytometry based Luminex assay. MFI, mean fluorescence intensity. **b,** Spike in open conformation with a single erect RBD (PDB: 6ZGG) in trimer axis vertical view with the locations of mutated residues highlighted in red spheres and labelled on the monomer with erect RBD. Vaccine (**c-d**) and convalescent sera (**e-f**) against WT and B.1.1.7 Spike mutant with N501Y, A570D, ΔH69/V70, Δ144/145, P681H, T716I, S982A and D1118H. **g-h,** neutralisation curves for serum from two convalescent individuals with reduced susceptibility to B.1.1.7 Spike mutant, means of technical replicates are plotted with error bars representing standard error of mean. Data are representative of 2 independent experiments.

Figure 2

Figure 2. Neutralisation and binding by a panel of NTD- and RBD-specific mAbs against WT and B.1.1.7 SARS-CoV-2 viruses. a, Neutralisation of WT (black) and B.1.1.7 (orange) Spike pseudotyped SARS-CoV-2-MLV by 3 selected mAbs (S2E12, S2X333 and S2H14) from one representative experiment. Shown is the mean ± s.d. of 2 technical replicates. **b,** Neutralisation of WT and B.1.1.7 SARS-CoV-2-MLV by 61 mAbs targeting NTD (n=10), RBM (n=29) and non-RBM sites in the RBD (n=22). Shown are the mean IC50 values (ng/ml) of n=2 independent experiments. **c-e,** Neutralisation shown as mean IC50 values (upper panel) and mean fold change of B.1.1.7 relative to WT (lower panel) of NTD (c), RBM (d) and non-RBM (e) mAbs. Lower panel shows IC50 values from 2 independent experiments. **f-h,** Kinetics of binding of mAbs to WT (black) and N501Y (blue) RBD as measured by bio-layer interferometry (BLI). Shown in (f) are the 6 RBM-targeting mAbs with reduced binding to N501Y RBD and 2 representative mAbs (RBM-targeting S2E12 and non-RBM-targeting S2X35). Area under the curve (AUC) (g) and AUC fold change (h) of 51 mAbs tested against WT and N501Y RBD. mAbs with a >1.3 AUC fold change shown in blue.

Figure 3. E484K appearing in background of B.1.1.7 with evidence of transmission a.
Representation of Spike RBM:ACE2 interface (PDB: 6M0J) with residues E484, N501 and K417
highlighted as spheres coloured by element **b.** Maximum likelihood phylogeny of a subset of
sequences from the United Kingdom bearing the E484K mutation (green) and lineage B.1.1.7
(blue). As of 1st Feb 2021, 16 sequences from the B.1.1.7 lineage (two unrelated sequences in
Wales, and one cluster of 14 in England) have acquired the E484K mutation (red). c. Sequence
accumulation over time in GISAID for sequences with B.1.1.7 and E484K.

Figure 4

Figure 4. Neutralisation potency of vaccine sera against pseudovirus virus bearing Spike mutations in the B1.1.7 lineage with and without E484K in the receptor binding domain (all In Spike D614G background). **a,** Neutralisation curves for five vaccinated individuals. Data points represent mean of technical replicates with standard error and are representative of two independent experiments **b, c.** Indicated is 50% neutralisation titre for each virus expressed as fold change relative to WT. WT (n=24), B.1.1.7 (n=24), B.1.1.7 +E484K (n=12). In c only data for participants with FC from three viruses are presented. Data points represent means of two independent experiments.

Extended Data Figure 1

Extended Data Figure 1: Immune responses three weeks after first dose of Pfizer SARS-CoV-2 vaccine BNT162b2 a, Relationship between serum IgG responses as measured by flow cytometry and serum neutralisation ID50. **b,** Relationship between serum neutralisation ID50 and T cell responses against SARS-CoV-2 by IFN gamma ELISPOT. SFU: spot forming units. **c**, Relationship between serum IgG responses and T cell responses.

Extended Data Fig. 2

Extended data Fig 2. Neutralisation by vaccine sera against wild type and mutant SARS-CoV-2 pseudotyped viruses: (**a-b**) WT and Spike mutant with N501Y, A570D, ΔH69/V70. Data are representative of 2 independent experiments.

Extended Data Fig. 3

Extended Data Fig. 3. Neutralisation potency of convalescent sera against pseudovirus virus bearing three Spike mutations (N501Y, A570D, ΔH69/V70) present in B.1.1.7 versus wild type (all In Spike D614G background). Indicated is serum $\log_{10}$ inverse dilution against % neutralisation. Where a curve is shifted to the right this indicates the virus is less sensitive to the neutralising antibodies in the serum. Data points represent means of technical replicates and error bars represent standard error of the mean.

Extended Data Fig. 4

Extended Data Fig. 4. Neutralisation potency of convalescent sera against wild type and B.1.1.7 Spike mutant SARS-CoV-2 pseudoviruses. Indicated is serum log_{10} inverse dilution against % neutralisation. Where a curve is shifted to the right this indicates the virus is less sensitive to the neutralising antibodies in the serum. Data points represent means of technical replicates and error bars represent standard error of the mean.

Extended Data Fig. 5

Extended Data Fig. 5. Neutralisation of WT and B.1.1.7 SARS-CoV-2 of a panel of 58 mAbs. **a-c,** Neutralisation of WT (black) and B.1.1.7 (orange) SARS-CoV-2-MLV by 9 NTD-targeting (a), 27 RBM-targeting (b) and 22 non-RBM-targeting (c) mAbs.

Extended Data Fig. 6

Extended Data Fig. 6. Kinetics of binding to WT and N501Y SARS-CoV-2 RBD of 43 RBD-specific mAbs. a-b, Binding to WT (black) and N501Y (blue) RBD by 22 RBM-targeting (a) and 21 non-RBM-targeting (b) mAbs. An antibody of irrelevant specificity was included as negative control.

Extended Data Fig. 7

Extended Data Fig. 7. Binding of human ACE2 to SARS-CoV-2 B.1.1.7 and WT RBDs. a-b. BLI binding analysis of the human ACE2 ectodomain (residues 1-615) to immobilized SARS-CoV-2 WT RBD (a) and B.1.1.7 RBD (b). Black lines correspond to a global fit of the data using a 1:1 binding model.