Age-dependent heterogeneity of lymph node metastases and survival: A

population-based study.

Michael Behring[1], Sumit Agarwal[1], Prachi Bajpai[1], Amr Elkholy[1], Hyung-Gyoon Kim[1], Shi

Wei[1, 2], Sadeep Shrestha[2, 3], Upender Manne[1, 2*]

Institutional Affiliation: [1]Department of Pathology; [2]O'Neal Comprehensive Cancer Center,

[3]Department of Epidemiology, University of Alabama at Birmingham; 1720 Second Avenue

South, Birmingham, AL 35294.

*Correspondence:

Upender Manne, MS, PhD; Professor of Pathology, University of Alabama at Birmingham, AL

35294, USA. upendermanne@uabmc.edu

Abstract

Background: For several cancers, including those of the breast, young age at diagnosis is associated with an adverse prognosis. Although this effect is often attributed to heritable mutations such as BRCA1/2, the relationship between pathologic features, young age of onset, and prognosis for breast cancer remains unclear. In the present study, we highlight links between age of onset and lymph node metastasis (NM) in US women with breast cancer.

Methods: Case listings from Surveillance, Epidemiology, and End Result (SEER) 18 registry data for women with breast cancer, which include information on race, were used. NM and its associated outcomes were evaluated for a subset of women with receptor subtype information and then compared against a larger, pre-subtype validation set of data from the same registry. Age of diagnosis was a 5-category variable; under 40 years, 40-49 years, 50-59 years, 60-69 years and 70+ years. Univariate and adjusted multivariate survival models were applied to both sets of data.

Results: As determined with adjusted logistic regression models, women under 40 years old at diagnosis had 1.55 times the odds of NM as women 60-69 years of age. The odds of NM for (HR = hormone receptor) HR+/HER2+, HR-/HER2+, and triple-negative breast cancer subtypes were significantly lower than those for HR+/HER2-. In subtype-stratified adjusted models, age of diagnosis had a consistent trend of decreasing odds of NM by age category, most noticeable for HR+ subtypes of luminal A and B. Univariate 5-year survival by age was worst for women under 40 years, with NM attributable for 49% of the hazard of death from cancer in adjusted multivariate models.

33 **Conclusions:** Lymph node metastasis is age-dependent, yet not all molecular subtypes are

34 clearly affected by this relationship. For <40-yr-old women, NM is a major cause for shorter

35 survival. When stratified by subtype, the strongest associations were in HR+ groups, suggesting

36 a possible hormonal connection between young age of breast cancer onset and NM.

37 **Keywords:** Breast cancer, Nodal metastasis, Early age of diagnosis, SEER cancer registry

Background:

In 2020, an estimated 276,480 new cases of breast cancer were diagnosed in the United States (https://seer.cancer.gov/statfacts/html/breast.html). For women with breast cancer, the infiltration of tumor cells into surrounding lymph nodes is associated with a poor prognosis. Lymph node metastasis (NM), a means for the regional and distant spread of tumor cells, has a considerable influence upon treatment options and patient survival. Approximately 60% of all newly diagnosed cases of breast cancer are localized (non-metastatic). However, one third of the patients with localized cancers will eventually develop metastatic disease (1). Of all new cases of breast cancer, another third have regional NM at the time of diagnosis (1). Lymph nodes usually represent the first site of metastasis of breast cancer, and they initiate the process of metastasis of the disease.

Breast cancer is distinctive among highly prevalent cancers in that women with a young age of onset often have a more aggressive form of the disease. Although younger women are eligible for more intensive therapy, they nevertheless have, relative to older patients, worse survival and a higher recurrence rate (2,3). Although links between molecular/receptor subtypes and disease progression (metastasis in particular) are commonplace (4–6), there has been little research into the durability of these relationships across age groups. Estrogen receptor-positive status has an unclear prognostic influence across ages (7–9). However, *ERBB2*/*HER2* receptor status seems to be more frequent and is associated with a lower survival of younger women (10). A few negatively associated prognostic variables for early onset breast cancer are identified (11–13), yet the connection between patient characteristics and metastasis has not been fully assessed. In the present study, we used nation-wide data from Surveillance Epidemiology and End Result

(SEER) cancer registries to gain a better understanding of the relationship between lymph node

metastasis, receptor subtype, and age-dependent patient characteristics of breast cancer.

Materials and Methods

Study population data

The present analysis involved data from the SEER 18 SEER*Stat8.3.8 database case listings.

The data in SEER 18 represents 27.8% of the total US population and 18 cancer registries (14).

After excluding patients with non-ductal or lobular tumor histology, male gender, missing

information on lymph node metastasis, and follow up of less than 6 months, our final sample

consisted of 717,331 women diagnosed with breast cancer from 1975-2017. Since SEER did not

start recording information regarding the HER2 receptor subtype until 2010, we used a subset of

the overall data that offered complete receptor subtype analysis. The receptor subtype set of data

is made up of 223,986 cases of breast cancer with first and only primary tumor for patients

diagnosed for the ~ 7 year period of 2010-2017.

Study design

The present study used both case-control and follow-up designs. Cases were defined as breast

cancer patients with NM at the time of diagnosis. Controls were patients with no NM at

diagnosis. Data for all eligible patients were used, and no matching of controls to cases was

done. A response variable of lymph node metastasis was defined as a binary outcome using

AJCC 7[th] edition tumor , nodes , and metastases (TNM) staging (15). This large data set with NM

information is consistent with AJCC 6[th] edition TNM staging (16). All nodes (N) values of N0

(including N0 (i-) and N0 (mol-)) were considered NM negative, and any other N values (N1-

N4) were considered positive. Included were patient demographic measures of age and ethnicity

as well as tumor differentiation and staging information. Breast cancer subtype was based on

receptor status. The "HR" abbreviation for hormone receptor represents both estrogen (ER) and

progesterone (PR) receptor status. HER2 indicates human epidermal growth factor 2 receptor

status. Borderline information on HER2 status was excluded from analysis of SEER*Stat data

queries (17). Age of onset was considered as both a continuous and a categorical variable.

Categories of age were based on preceding literature (8,18): under 40 years, 40-49 years, 50-59

years, 60-69 years, and 70+ years. Additional survival analysis was accomplished with time to

death from cancer as an outcome (Tables 1a and 1b).

Statistical analyses

In depicting the univariate relationship between NM, receptor subtype, and individual covariates,

we applied chi-square tests for categorical, and t-tests for continuous p-values. Logistic

regression models were constructed for each variable to obtain univariate odds of NM and 95%

confidence intervals. To examine whether the effect of age upon odds of NM was modified by

receptor subtype, we performed 4 separate, adjusted logistic regression analyses stratified by

subtype. Logistic regression modeling was also used to adjust for potential confounding

variables, both for full data set and in receptor-based subtype subset analyses. Kaplan-Meier log-

rank tests and Cox proportional hazard models were used to estimate the effect of age upon

survival and NM. We measured 5-year survival for the similar age categories used in the

analyses of Alteri et al. (19). For 5 separate age category-stratified survival models, we

calculated the proportion of hazard of death that was attributable to NM (attributable fraction

analysis) by comparing a counterfactual survival function (excluding NM from baseline) to a

factual survival function, including NM (20,21). In Cox models, variables adjusted for were NM,

race, size of tumor, and receptor subtype. All statistical analyses utilized R version 3.6.2 (2019-

12-12) (22).

Results

Overall study population

Associations with NM were consistent across overall (1975-2017) and subset (2010—2017)

analyses and showed that, at diagnosis, 32-36% of women had nodal metastasis of breast cancer.

In follow up, women with NM made up a larger proportion of breast cancer deaths, were more

African American, had tumors of larger size at diagnosis, and had more distant metastases at

diagnosis (Table 1a and 1b). Additional information unique to the test data set also showed

higher grade/more poorly differentiated tumors for NM cases. Women with nodal metastases

were also younger than controls (non-NM), with an average age at diagnosis of 57 years (61

years for controls). Lastly, a larger proportion of women had HR+/HER2+, triple-negative, and

HR-/HER2+ subtype cancers than controls. See Tables 1a and 1b for a description of patient

variables across nodal metastasis outcomes. In a subset analysis (data not shown) of women

under 40 years in 1975-2017 data, no effect modification by age was found for the association

between race, grade, and NM.

Odds of nodal metastasis

The adjusted odds of NM for common variables in both datasets were available for race, age,

tumor size, and tumor grade. The relationship between age and the adjusted odds of NM had

similar estimates for both sets of data. Among all age groups, women under 40 had the highest

adjusted odds of NM (1.55 in subset and 1.74 in overall data). Both sets of data confirmed tumor

size as being strongly associated with NM. Relative to Whites, African American women had

higher adjusted odds of NM (1.13 in subset and 1.23 in overall data). Higher tumor grades were

also positively associated with NM, with the effect increasing parallel to the loss of

differentiation. See Tables 2a and 2b for details of univariate and adjusted odds of NM.

Receptor subtype

In receptor-stratified data, adjusted estimates of NM by subtype showed that TNBC cancers had

lower odds of nodal metastasis relative to the HER2-/HR+ receptor subtype (OR 0.74, 95% CI

0.70-0.78) (Table 2b). There was, however, no significant increase in the odds of having NM for

either HER2+ subtypes of HER2+/ER+ or HER2+/ER- (Table 2a). There was a young-to-old

gradient in the odds of NM, which is most apparent in for HR+ subtypes of tumors, HER2-/ER+

and lHER2+/ER+ (Figure 1). In HR-/HER2+subtypes, age at diagnosis had a similar trend in

odds estimates per age group, but showed no significant difference between ages <40-59. In

addition, triple-negative subtypes showed a slightly higher association with NM for those

younger than 60 but with no age gradient (Figure 1).

Survival analysis

For both subset and overall data, Kaplan-Meier analyses consistently showed women under 40 as

sharing the worst survival outcomes with women aged 70 years and above. Although women 70

years and older had low survival, women under 40 years old began follow-up with survival

similarly high with other age categories (40-69 years), but there was a decline starting at ~15

months which surpassed 70-year-olds at ~22 months. Women under 40 years of age had the

lowest probability of 5-year survival at 0.87, with age groups from 40-69 having similar 5-year

survivals at approximately 0.91-0.93, with the value for the oldest group of women slightly

decreasing to 0.89. As determined with adjusted Cox analyses, in both sets of data, women

younger than 40 also had a higher fraction of hazard of death from cancer (~50% vs. ~37% at

start) due to NM than women 70 years and above (Figure 2).

Discussion

In adult females younger than 40, there is considerable debate about whether breast cancer

should be considered as a distinct disease. Women under 40 have consistently poor survival

outcomes across various study populations (12,13,23–26) yet the reasons for this are not fully

explained. The present study showed that the odds of NM at diagnosis have a gradient

relationship to patient age of onset. This age-decreasing slope is most prominent for hormone

receptor-positive subtypes. Findings from survival analysis confirmed that women under 40 have

a hazard of death from breast cancer equal to or greater than the next-poorest group, women 70

years or older. Furthermore, the hazard of death attributable to NM was higher in women under

40 years, suggesting that NM has a stronger causal role in breast cancer mortality for younger

women as compared to late age of diagnosis survival factors.

Previous studies have suggested that a higher incidence of the HR-/HER2+ subtype for young

women with breast cancer partially explains their more aggressive disease (23–25). Indeed,

although our analysis showed a higher proportion of women under 40 years old having the HR-

/HER2+ subtype, there was no significant difference in the odds of NM and age among the HR-

/HER2+, and HR+ subtypes. However, a study, using a subset of the SEER data shows that,

regardless of patients age, the HR+/HER2-breast cancer subtype has a higher rate of lymph node

involvement at diagnosis than the triple-negative subtype (11). From this study, NM at diagnosis

does not appear to be related to HR-/HER2+ aggressiveness in young age of diagnosis.

Furthermore, this observation may be of importance in considering whether prior research

examining the link between the HR-/HER2+ subtype and nodal metastasis was adequately

controlled for age at diagnosis.

The level of generalizability of the SEER 2010-2017 receptor-only subset of data may be

questioned as a liability for analysis. We evaluated how representative subset data was using the

complete, 42-year data. For all the variables common between datasets, both measures of

association and measures of effect were consistent across data. Furthermore, we examined

whether there was a time trend in proportion of NM positive patients at diagnosis in relation to

age group. Interestingly, we found that, for women <40 years, the proportion of NM at diagnosis

vs. non-NM remained stable across time at approximately 50%. In women aged 70+, there was a

marked split of NM proportion from 50% starting in 1980, decreasing to ~25% by 2008 (Figure

3) .

The concordance of IHC-based receptor subtype with intrinsic molecular subtype has been

shown to vary by subtype. Luminal B (HR+/HER2+) is the worst performing subtype in

measures of both concordance and accuracy across several studies (27,28), suggesting that the

relationship between young age of onset, NM, and interplay between estrogen/progesterone

hormone receptors and HER2 should be studied further with a thorough consideration of

molecular subtype classification. In contrast, previous research has shown the concordance and

accuracy between both luminal A (HR+/HER2-) and triple-negative (HR-/HER2-) to be the

highest among subtypes (29), suggesting that the results found in our study for these subtypes are

likely not due to misclassification.

Lastly, there is also a question of consistency in measurement of variables over time. Perhaps the

phenomenon of aggressive disease for younger women with breast cancer is influenced by

changes in screening and diagnostic practices. From the standpoint of the association between

age of diagnosis and NM, we included year of diagnosis as a predictor variable for both sets used

for analysis. There was no confounding effect of the year of diagnosis upon the association

between NM and age of diagnosis.

Involvement of lymph nodes is a key component in decisions for postoperative therapy,

particularly radiation therapy, because clinicians evaluate need for lymph node radiation to

minimize toxicity of treatment. Therefore, our results showing that the higher incidence of NM

in young (<40 years) HR+ breast cancer groups (HR+/HER2- and HR+/HER2+) is clinically

relevant. This relationship between NM and age of diagnosis is clear in women with luminal

tumors, and may explain observations from previous studies linking the luminal subtype, young

age of diagnosis, and poor prognosis (8). Thus, our findings may aid in identifying aggressive

disease in young women with luminal disease.

Conclusions

Our results suggest that a predisposition towards more severe breast cancer for women with

younger age of diagnosis is driven in part by NM. This relationship between NM and age of

diagnosis for women with luminal tumors is strong and may be related to a poor prognosis.

These findings also have implications in identifying high-risk, young HR+ groups (HR+/HER2-

and HR+/HER2+) of breast cancer patients for aggressive therapy.

List of abbreviations

NM, nodal metastasis; SEER, Surveillance, Epidemiology, and End Results; TNM, tumor, node,

and metastasis; HR, hormone receptor; ER, estrogen receptor; PR, progesterone receptor;

ERBB2/HER2, human epidermal growth factor receptor 2; TNBC, triple-negative breast cancer.

DECLARATIONS:

Ethics approval and consent to participate: This study used publicly available data, thus these

factors are not applicable.

Consent for publication: Not applicable

Availability of data and materials: The datasets used and/or analyzed during the current study

are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests

Funding: The funding sources for the research are reported in the Acknowledgement section

below. The NCI training grant to Dr. Michael Behring is provided to support his salary, and the

NCI/NIH grant to Dr. Manne has covered a portion of his effort on this study, particularly to

analyze the data based on the race/ethnicity of patients. However, the funding bodies did have

any role in the design of the study or in collection, analysis, and interpretation of data or in

writing the manuscript.

Authors' contributions: UM and MB conceptualized the study; MB, SA, PB, AE, and H-GK

collected the data. MB analyzed the data, and MB, SW, SS, and UM interpreted the data

regarding breast cancer. MB was a major contributor for this study; all authors contributed to

writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements: This study was supported, in part, by grant 5U54CA118948 (NCI/NIH) and

institutional/departmental impact funds to UM, and by CA047888 (NCI/NIH), the UAB Cancer

Prevention and Control Training Program to MB. We thank Dr. Donald Hill for his help in

editing the manuscript.

Authors' information: MB, MPH, PhD-Postdoctoral Fellow of the Cancer Control and

Prevention Program; PB, PhD-Researcher V; AE, MS- Graduate Student; H-GK, PhD -

Researcher V; SW, MD, PhD- breast diagnostic pathologist, Professor of Pathology, and member

of the O'Neal Comprehensive Cancer Center; SS, PhD, cancer epidemiologist, Professor of

Epidemiology; and UM, MS, PhD, cancer translational researcher, Professor of Pathology, and

member of the O'Neal Comprehensive Cancer Center.

Table 1a. Clinical and demographic characteristics of US women with breast cancer by lymph node status (SEER, 2010-2017)

Lymph node metastasis Characteristics	negative (N=151827)	positive (N=72159)
Breast tumor by receptor subtype		
- HR-/HER2- (triple-negative)	18418 (12.1%)	9189 (12.7%)
- HR-/HER2+	6819 (4.5%)	4927 (6.8%)
- HR+/HER2-	110069 (72.5%)	47318 (65.6%)
- HR+/HER2+	16521 (10.9%)	10725 (14.9%)
Age (years)	60.5 ± 12.9	56.9 ± 13.3
Age categories		
- <40	7342 (4.8%)	6702 (9.3%)
- 40-49	25117 (16.5%)	15678 (21.7%)
- 50-59	38185 (25.2%)	19930 (27.6%)
- 60-69	42631 (28.1%)	16746 (23.2%)
- 70+	38552 (25.4%)	13103 (18.2%)
Cause of death		
- alive	142734 (94.0%)	60886 (84.4%)
- breast	4014 (2.6%)	8767 (12.1%)
- other	5079 (3.3%)	2506 (3.5%)
Race		
- White	120218 (79.2%)	54391 (75.4%)
- American Indian/Alaska Native	925 (0.6%)	512 (0.7%)
- Asian or Pacific Islander	14083 (9.3%)	6811 (9.4%)
- African American	15499 (10.2%)	10011 (13.9%)
- Unknown	1102 (0.7%)	434 (0.6%)
Metastasis (AJCC M 7[th] ed. 2010)		
- No distant metastasis	138047 (98.8%)	60336 (90.3%)
- Distant metastasis	1737 (1.2%)	6485 (9.7%)
Tumor size (AJCC T 7[th] ed.2010)		
- T1 (<2 cm)	98954 (71.7%)	22125 (33.8%)
- T2 (2 cm-5 cm)	33586 (24.3%)	29815 (45.6%)
- T3 (>5 cm)	3788 (2.7%)	7631 (11.7%)
- T4 (extension into chest wall/skin)	1727 (1.3%)	5798 (8.9%)
Grade		
- Well differentiated; Grade I	38066 (25.1%)	7535 (10.4%)
- Moderately differentiated; Grade II	63272 (41.7%)	29342 (40.7%)
- Poorly differentiated; Grade III	45972 (30.3%)	32305 (44.8%)
- Undifferentiated; anaplastic; Grade IV	340 (0.2%)	244 (0.3%)
- Unknown	4177 (2.8%)	2733 (3.8%)

Table 1b. Clinical and demographic characteristics of US women with breast cancer by lymph node status (SEER 1975-2017)

Lymph node metastasis Characteristics	negative (N=459656)	positive (N=257675)
Age (years)	60.7 ± 13.5	57.3 ± 13.8
Age categories		
- <40	24485 (5.3%)	24526 (9.5%)
- 40-49	80917 (17.6%)	57697 (22.4%)
- 50-59	110958 (24.1%)	67060 (26.0%)
- 60-69	112844 (24.5%)	54294 (21.1%)
- 70+	130452 (28.4%)	54098 (21.0%)
Cause of death		
- alive	332223 (72.3%)	141349 (54.9%)
- breast	36964 (8.0%)	77168 (29.9%)
- other	90469 (19.7%)	39158 (15.2%)
Race		
- White	378616 (82.4%)	204948 (79.5%)
- American Indian/Alaska Native	2316 (0.5%)	1514 (0.6%)
- Asian or Pacific Islander	35863 (7.8%)	19267 (7.5%)
- African American	40497 (8.8%)	30963 (12.0%)
- Unknown	2364 (0.5%)	983 (0.4%)
Grade		
- Well differentiated; Grade I	94056 (20.5%)	20760 (8.1%)
- Moderately differentiated; Grade II	174711 (38.0%)	87336 (33.9%)
- Poorly differentiated; Grade III	131972 (28.7%)	105851 (41.1%)
- Undifferentiated; anaplastic; Grade IV	4847 (1.1%)	4105 (1.6%)
- Unknown	54070 (11.8%)	39623 (15.4%)

Table 2a. Odds of nodal metastasis, unadjusted and adjusted models (SEER, 2010-2017)

Odds of lymph node metastasis	Unadjusted		Adjusted	
Characteristics	**OR**	**95% CI**	**OR**	**95% CI**
Breast tumor by receptor subtype				
- HR-/HER2- (triple negative)	1.16	1.13-1.19	0.60	0.58-0.62
- HR-/HER2+	1.68	1.62-1.75	0.87	0.83-0.91
- HR+/HER2-	1	ref	1	ref
- HR+/HER2+	1.51	1.47-1.55	0.94	0.91-0.97
Age categories				
- <40	2.32	2.24-2.41	1.55	1.49-1.62
- 40-49	1.59	1.55-1.63	1.37	1.33-1.41
- 50-59	1.33	1.30-1.36	1.21	1.18-1.25
- 60-69	1	ref	1	ref
- 70+	0.87	0.84-0.89	0.80	0.77-0.82
Race				
- White	1	ref	1	ref
- American Indian/Alaska Native	1.22	1.10-1.36	1.08	0.95-1.23
- Asian or Pacific Islander	1.07	1.04-1.10	0.92	0.88-0.95
- African American	1.43	1.39-1.47	1.13	1.09-1.17
- Unknown	0.85	0.74-0.98	0.87	0.76-0.99
Metastasis (AJCC M 7[th] ed. 2010)				
- No distant metastasis	1	ref	1	ref
- Distant metastasis	8.54	8.10-9.02	3.68	3.46-3.91
Tumor size (AJCC T 7[th] ed. 2010)				
- T1 (<2 cm)	1	ref	1	ref
- T2 (2 cm-5 cm)	3.97	3.89-4.06	3.40	3.32-3.48
- T3 (>5 cm)	9.01	8.64-9.39	7.00	6.71-7.32
- T4 (extension into chest wall/skin)	15.02	14.21-15.88	9.91	9.34-10.51
Grade				
- Well differentiated; Grade I	1	ref	1	ref
- Moderately differentiated; Grade II	2.34	2.28-2.41	1.70	1.65-1.76
- Poorly differentiated; Grade III	3.55	3.45-3.65	2.00	1.94-2.08
- Undifferentiated; anaplastic; Grade IV	3.63	3.07-4.29	1.97	1.62-2.38
- Unknown	3.31	3.13-3.49	1.62	1.52-1.74

Table 2b. Odds of nodal metastasis, unadjusted and adjusted models (SEER 1975-2017)

Odds of lymph node metastasis	Unadjusted		Adjusted	
Characteristics	**OR**	**95% CI**	**OR**	**95% CI**
Age categories				
- <40	2.08	2.04-2.12	1.74	1.71-1.78
- 40-49	1.48	1.46-1.50	1.39	1.37-1.41
- 50-59	1.26	1.24-1.27	1.22	1.20-1.23
- 60-69	1	ref	1.00	ref
- 70+	0.86	0.85-0.87	0.87	0.85-0.89
Race				
- White	1	ref	1	ref
- American Indian/Alaska Native	1.21	1.13-1.28	1.14	1.06-1.21
- Asian or Pacific Islander	0.99	0.97-1.01	0.92	0.91-0.94
- African American	1.41	1.39-1.43	1.23	1.21-1.25
- Unknown	0.77	0.71-0.83	0.76	0.70-0.81
Grade				
- Well differentiated; Grade I	1	ref	1	ref
- Moderately differentiated; Grade II	2.26	2.23-2.30	2.20	2.17-2.24
- Poorly differentiated; Grade III	3.63	3.57-3.70	3.31	3.18-3.57
- Undifferentiated; anaplastic; Grade IV	3.84	3.67-4.01	3.53	3.38-3.69
- Unknown	3.32	3.25-3.39	3.23	3.17-3.30

Figure 1. Comparison of adjusted (M stage, T stage, grade) odds of nodal metastasis by age, stratified by receptor subtype, SEER 2010-2017 (ref. group age 60-69 years). Adjusted for race and tumor size.

Figure 2. Univariate survival and NM-attributable fraction from adjusted models by selected age groups. A) Kaplan-Meier plot of survival by age group, SEER 2010-2017; B) Fraction of hazard of death due to NM for women under 40 years, and for women 70 years or more, SEER 2010-2017

Figure 3. Lymph node metastasis (NM) as a proportion of total cases by month from 1975-2017,

for ages under 40 years (top) and 70 years or older (bottom). NM in red, non-NM in blue.

Bibliography

1.   Howlader N, Noone AM, Krapcho M, Garshell J, Miller D, Altekruse SF, Kosary CL, Yu

M, Ruhl J, Tatalovich Z, Mariotto A, Lewis DR, Chen HS, Feuer EJ CK (eds). SEER

Cancer Statistics Review, 1975-2011 [Internet]. National Cancer Institute. 2013. Available

from: http://seer.cancer.gov/csr/1975_2011/

2.   Narod S a. Breast cancer in young women. Nat Rev Clin Oncol. 2012;9(8):460–70.

3.   Rick Alteri, MD; Cammie Barnes, MBA; Adriane Burke, MPH; Ted Gansler, MD, MPH;

Susan Gapstur, PhD; Mia Gaudet, PhD; Joan Kramer, MD; Lisa A Newman, MD, MPH;

Dearell Niemeyer, MPH; Cheri Richards, MS; Carolyn Runowicz, MD; Debbie Saslow,

PhD; Scott Simpson M. Breast Cancer Facts & Figures 2013-2014. Am Cancer Soc.

2014;38.

4.   Perou CM, Sørlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular

portraits of human breast tumours. Nature. 2000;406:747–52.

5.   Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, et al. Repeated

observation of breast tumor subtypes in independent gene expression data sets. Proc Natl

Acad Sci U S A. 2003;100:8418–23.

6.   Sotiriou C, Neo S-Y, McShane LM, Korn EL, Long PM, Jazaeri A, et al. Breast cancer

classification and prognosis based on gene expression profiles from a population-based

study. Proc Natl Acad Sci U S A. 2003;100:10393–8.

7.   Partridge AH, Gelber S, Piccart-Gebhart MJ, Focant F, Scullion M, Holmes E, et al.

Effect of age on breast cancer outcomes in women with human epidermal growth factor

receptor 2-positive breast cancer: results from a herceptin adjuvant trial. J Clin Oncol.

2013;31(21):2692–8.

8.   Partridge AH, Hughes ME, Warner ET, Ottesen RA, Wong Y-N, Edge SB, et al. Subtype-

Dependent Relationship Between Young Age at Diagnosis and Breast Cancer Survival. J

Clin Oncol. 2016;34(27).

9.   Azim H a, Michiels S, Bedard PL, Singhal SK, Criscitiello C, Ignatiadis M, et al.

Elucidating prognosis and biology of breast cancer arising in young women using gene

expression profiling. Clin Cancer Res. 2012;18(5):1341–51.

10.   Keegan THM, DeRouen MC, Press DJ, Kurian AW, Clarke C a. Occurrence of breast

cancer subtypes in adolescent and young adult women. Breast Cancer Res.

2012;14(2):R55.

11.   Fabiano V, Mandó P, Rizzo M, Ponce C, Coló F, Loza M, et al. Breast Cancer in Young

Women Presents With More Aggressive Pathologic Characteristics: Retrospective

Analysis From an Argentine National Database. JCO Glob Oncol. 2020 Sep;(6):639–46.

12.   Shannon C, Smith IE. Breast cancer in adolescents and young women. Eur J Cancer. 2003

Dec;39(18):2632–42.

13.   Yazdani-Charati R, Hajian-Tilaki K, Sharbatdaran M. Comparison of pathologic

characteristics of breast cancer in younger and older women. Casp J Intern Med.

2019;10(1):42–7.

14.   National Cancer Institute. SEER*Stat Database: NPCR and SEER Incidence - U.S. Cancer

Statistics Public Use Database, Nov 2017 submission (2001-2015). 2017.

15.   Edge SB, Compton CC. The American Joint Committee on Cancer: the 7th edition of the

AJCC cancer staging manual and the future of TNM. Vol. 17, Annals of surgical

oncology. 2010. p. 1471–4.

16.   American Joint Committee on Cancer. AJCC Cancer Staging Handbook. Cancer. 2010. 52

p.

17.   NCI. Breast Subtype (2010+) [Internet]. SEER Documentation. Available from:

https://seer.cancer.gov/seerstat/databases/ssf/breast-subtype.html

18.   Anders CK, Johnson R, Litton J, Phillips M, Bleyer A. Breast Cancer Before Age 40

Years. Semin Oncol. 2009 Jun;36(3):237–49.

19.   Rick Alteri, MD; Cammie Barnes, MBA; Adriane Burke, MPH; Ted Gansler, MD, MPH;

Susan Gapstur, PhD; Mia Gaudet, PhD; Joan Kramer, MD; Lisa A Newman, MD, MPH;

Dearell Niemeyer, MPH; Cheri Richards, MS; Carolyn Runowicz, MD; Debbie Saslow,

PhD; Scott Simpson M. Breast Cancer Facts & Figures 2013-2014. Am Cancer Soc.

2014;38.

20.   Sjölander A, Vansteelandt S. Doubly robust estimation of attributable fractions in survival

analysis. Stat Methods Med Res. 2017;26(2):948–69.

21.   Chen L, Lin DY, Zeng D. Attributable fraction functions for censored event times.

Biometrika. 2010;97(3):713–26.

22.   R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing. Vienna, Austria; 2015.

23.   Agrup M, Stal O, Olsen K, Wingren S. C-erbB-2 overexpression and survival in early

onset breast cancer. Breast Cancer Res Treat. 2000;63(1):23–9.

24.   Hartley MC, McKinley BP, Rogers EA, Kalbaugh CA, Messich HS, Blackhurst DW, et al.

Differential expression of prognostic factors and effect on survival in young (< or =40)

breast cancer patients: a case-control study. Am Surg. 2006;72(12):1185–9.

25.   Akiyama T, Sudo C, Ogawara H, Toyoshima K, Yamamoto T. The product of the human

c-erbB-2 gene: a 185-kilodalton glycoprotein with tyrosine kinase activity. Science.

1986;232(4758):1644–6.

26.   Mattes MD, Bhatia JK, Metzger D, Ashamalla H, Katsoulakis E. Breast Cancer Subtype

as a Predictor of Lymph Node Metastasis according to the SEER Registry. J Breast

Cancer. 2015;18(2):143.

27.   Ades F, Zardavas D, Bozovic-Spasojevic I, Pugliano L, Fumagalli D, De Azambuja E, et

al. Luminal B breast cancer: Molecular characterization, clinical management, and future

perspectives. J Clin Oncol. 2014;32(25):2794–803.

28.   Nielsen TO, Parker JS, Leung S, Voduc D, Ebbert M, Vickery T, et al. A comparison of

PAM50 intrinsic subtyping with immunohistochemistry and clinical prognostic factors in

tamoxifen-treated estrogen receptor-positive breast cancer. Clin Cancer Res.

2010;16(21):5222–32.

29.   Jamshidi N, Yamamoto S, Gornbein J, Kuo MD. Receptor-based surrogate subtypes and

discrepancies with breast cancer intrinsic subtypes: Implications for image biomarker

development. Radiology. 2018;289(1):210–7.

Figure Legends

Figure 1. Comparison of adjusted (M stage, T stage, grade) odds of nodal metastasis by age, stratified by receptor subtype, SEER 2010-2017 (ref. group age 60-69 years). Adjusted for race and tumor size.

Figure 2. Univariate survival and NM-attributable fraction from adjusted models by selected age groups. A) Kaplan-Meier plot of survival by age group, SEER 2010-2017; B) Fraction of hazard of death due to NM for women under 40 years, and for women 70 years or more, SEER 2010-2017.

Figure 3. Lymph node metastasis (NM) as a proportion of total cases by month from 1975-2017, for ages under 40 years (top) and 70 years or older (bottom). NM in red, non-NM in blue.