Review Protocol

Oral ulcers of COVID-19 patients: a scoping review protocol

Author(s)

Abanoub RIAD,[1,2,†] Julien ISSA,[3,†] Veronika CHUCHMOVA,[2] Esraa GOMAA,[4] Andrea POKORNA,[1,5] Jitka

KLUGAROVA,[1,2] Miloslav KLUGAR[1,2]

Affiliation(s)

[1] Czech National Centre for Evidence-Based Healthcare and Knowledge Translation (Cochrane Czech

Republic, Czech EBHC: JBI Centre of Excellence, Masaryk University GRADE Centre), Institute of

Biostatistics and Analyses, Faculty of Medicine, Masaryk University, Brno, Czech Republic

[2] Department of Public Health, Faculty of Medicine, Masaryk University, Brno, Czech Republic

[3] Department of Esthetic and Restorative Dentistry, Faculty of Dentistry, Saint Joseph University,

Beirut, Lebanon

[4] Department of Oral Medicine and Oral Radiology, Faculty of Dentistry, Egyptian Russian University,

Cairo, Egypt

[5] Department of Nursing and Midwifery, Faculty of Medicine, Masaryk University, Brno, Czech

Republic

Corresponding Author

Dr. Abanoub RIAD, DDS

P.O.: Czech National Centre for Evidence-Based Healthcare and Knowledge Translation (Cochrane

Czech Republic, Czech EBHC: JBI Centre of Excellence, Masaryk University GRADE Centre), Institute

of Biostatistics and Analyses, Faculty of Medicine, Masaryk University, Kamenice 5, 625 00, Brno,

Czech Republic

Email: abanoub.riad@med.muni.cz

[†] These authors contributed equally to the work.

Abstract

Objective: This scoping review aims to systematically identify the types, characteristics, and possible
pathophysiologic etiologies of the oral ulcers that emerge in COVID-19 patients.

Introduction: The oral cavity is a vulnerable niche for the most diverse microbial ecosystem in the
human body; therefore, it presents a wide array of mucocutaneous complications that could indicate
various acute and chronic conditions. The COVID-19-related oral conditions, including oral ulcers,
had been widely debated as direct manifestations or indirect complications of the SARS-CoV-2
infection. According to a preliminary search of PROSPERO, MEDLINE, the Cochrane Database of
Systematic Reviews and the *JBI Evidence Synthesis*, there is no published nor registered scoping
review concerned with the oral ulcers of COVID-19 patients.

Inclusion criteria: The review will include studies included COVID-19 patients whose infection had
been confirmed by RT-PCR testing regardless of infection severity and clinical course. Only the
studies that reported COVID-19 patients with oral ulcers.

Methods: A three-phase search strategy will be carried out: an initial limited search, a full electronic
search, and hand search using the reference lists of all included records. The main bibliographic
databases of published literature will include MEDLINE (PubMed), Cochrane COVID-19 Study
Register, EMBASE via Ovid, and Scopus. All identified records will be managed using EndNote 9.2,
and the titles and abstracts will be screened against the inclusion criteria before the full text of all
potentially relevant studies will be examined. The data will be presented in tabular form, rating
maps, and narrative summary.

Registration: This protocol had been pre-registered in Open Science Framework (OSF) Registries.[1]

Keywords: Aphthous Stomatitis, COVID-19, Oral Manifestations, Oral Ulcer

Abstract word count: 244

Total manuscript word count: 1532

Introduction

The outbreak of the coronavirus disease (COVID-19) had required sensitive case definitions to be
developed according to the dynamically growing epidemiologic evidence.[2] The COVID-19-related
extrapulmonary manifestations including the cutaneous lesions had been frequently reported since
the beginning of this pandemic; however, it took several months until these lesions were recognized
by the World Health Organization (WHO) within the clinical case definitions of COVID-19.[2,3]
According to the review of Daneshgaran *et al.* 2020, the vesicular rashes were suggested to be used
for early diagnosis of COVID-19; while acral lesions were useful for epidemiological uses, and
vascular rashes for the prognosis of the disease severity.[4]

The oral cavity is subjected to numerous environmental factors that make it vulnerable to
mucocutaneous complications. This vulnerability is supported by the fact that the oral cavity is a
locus of the most diverse microbial ecosystem in the human body; therefore, it can harbour
accessible lesions that could be diagnostic for an array of systemic conditions.[5] Ulceration of the
oral epithelium, underlying connective tissues, or both may last for two weeks or more, and it can be
triggered by chronic conditions including drug-induced oral ulcers, lichen planus, pemphigus vulgaris,
mucous membrane pemphigoid, lupus erythematosus, Reiter's syndrome, tuberculosis, eosinophilic
ulcers, precancerous ulcers, and mycoses-related ulcers.[6] Additionally, oral ulcers may be of acute
nature and last for less than two weeks, e.g. traumatic ulcers, recurrent aphthous stomatitis (RAS),
Bechet's disease, allergic reactions, erythema multiform, necrotizing sialometaplasia, hematologic
disorders-related ulcers, and bacterial and viral infections. The clinical presentation of some viral
infections like herpes simplex virus, varicella-zoster virus, coxsackievirus, cytomegalovirus,
Epstein-Barr virus, measles virus, and human immunodeficiency virus (HIV) includes typically oral
ulcers and vesicles which may aid in their timely diagnosis and management.[7]

The oral symptoms of COVID-19 had been first reported in Lyon by Chaux-Bodard *et al.* 2020 as an
irregular tongue ulcer which evolved concurrently with acral erythema preceding the COVID-19
respiratory symptoms thus suggesting a promising epidemiological and clinical significance of the
oral lesions as inaugural symptoms.[8] In the following weeks, hundreds of individual cases with
different oral lesions were reported in multiple countries to explain the nature and the aetiology of
these lesions, which were equally distributed in females vs males.[9–11] The COVID-19-related oral
symptoms could be depicted as drug reactions, viral enanthem, stress-induced lesions, bacterial or
fungal co-infections, or inflammatory reaction.[9,11] Heretofore, the pathophysiological hypotheses of

COVID-19-related oral symptoms are supported by collateral observations that require further
investigation; for instance, the hypothesis of viral enanthema is supported by the high expression of
angiotensin-converting enzyme-2 (ACE-2) receptors in oral epithelium which serve as the primary
entry of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2); also the inflammatory
response hypothesis is supported by the increasing number of cases of the pediatric multisystem
inflammatory syndrome temporally associated with SARS-CoV-2 (PMIS-TS).[9,12] However, the onset
of oral lesions had considerably varied across the reported cases due to lack of use of reference
timepoints, it may indicate the variety of pathophysiologic etiologies and the natures of these
lesions.[9]

The COVID-19-related oral ulcers had been reported in critically-ill,[13,14] moderate,[15] mild[15–17] and
asymptomatic cases[16]. Aphthous-like lesions were reported without necrosis and hemorrhagic
crusts in younger patients, while in older patients with severe infection, the aphthous-like lesions
had been necrotic and hemorrhagic.[11] In a recent review of aphthous stomatitis in COVID-19
patients, the aphthous lesions were slightly predominant in females and people below 40 years old,
and their onset ranged between 0 to 10 days since the emergence of the respiratory symptoms.[15]
The tongue was the most common location for ulcerative and erosive lesions of non-specific origin
which appeared 3 days before the onset of respiratory symptoms in one of the cases.[11] Herpetiform
lesions were painful and emerged before, during, or after the respiratory symptoms.[11]

This scoping review's primary objectives are to evaluate the breadth and depth of the evidence on
oral lesions of COVID-19 patients and to evaluate their proposed pathophysiological courses and
epidemiological and clinical significance. A preliminary search of PROSPERO, MEDLINE, the Cochrane
Database of Systematic Reviews and the *JBI Evidence Synthesis* was conducted on January 20[th], 2020,
and no current or underway scoping reviews or systematic reviews on the topic were identified.

Review question(s)

What are the possible etiologies, types, and characteristics of the oral ulcers that emerge in
COVID-19 patients?

Inclusion criteria

Participants

This review will consider studies that include COVID-19 patients whose infection had been confirmed

by means of RT-PCR.

Concept

This review will consider studies that reported any type of oral ulcers for example: solitary vs.
multiple, acute vs. recurrent, painful vs. painless, necrotic vs. non-necrotic, hemorrhagic vs.
non-hemorrhagic, or contagious vs. non-contagious.

Context

This review will consider studies that reported COVID-19 patients with mild, moderate, severe and
critic clinical courses according to the Australian guidelines for clinical care of COVID-19 patients.[18]
The patients hospitalized either at intensive care or intermediate care (which provides more
intensive monitoring and patient management than the general ward but less than is offered in the
intensive care),[19] and the patients who underwent home isolation will be included in this review.

Types of sources

This scoping review will consider descriptive as well as analytical epidemiological studies.
Furthermore, systematic reviews, narrative reviews and correspondence articles will be considered
for inclusion in the proposed scoping review.

Methods

The proposed scoping review will be conducted in accordance with the JBI methodology for scoping
reviews.[20]

Search strategy

The search strategy will locate both published and unpublished primary studies, reviews, and
correspondence articles. A three-phase search strategy will be carried out. The first phase, which has
been already completed, was an initial limited search of MEDLINE (PubMed) and Google Scholar to
identify relevant articles, followed by an analysis of the text words contained in the titles and
abstracts of retrieved records, in addition to the index terms used to describe these records. A full
search strategy for MEDLINE (PubMed) is stipulated in <u>Appendix 1</u>. Secondly, the search strategy will
be developed according to the previous phase using all identified keywords and index terms, and it
will be customized for each included information source. Finally, the bibliographical reference lists of

all the included records will be screened for additional records.

Only studies published in English will be included. All the studies published since January 1st, 2020
will be included as the first cluster of COVID-19 cases was reported in Wuhan, China on December
30th, 2019.

Information sources

The databases to be searched include: MEDLINE (PubMed), Cochrane COVID-19 Study Register,
EMBASE via Ovid, and Scopus. The search for unpublished studies and gray literature will include:
medRxiv, ClinialTrials.gov, and the WHO's International Clinical Trials Registry Platform (ICTRP)
through the Cochrane COVID-19 Study Register [Clinicians working with COVID-19 patients].

Study selection

Following the search, all identified records will be collated and uploaded into EndNote 9.2 (Clarivate
Analytics, PA, USA) and duplicates removed. Following a pilot test, titles and abstracts will then be
screened by two independent reviewers (A.R. and J.I.) against the inclusion criteria. The potentially
relevant records will be retrieved in full and their citation details imported into the JBI System for
the Unified Management, Assessment and Review of Information (JBI SUMARI; JBI, Adelaide,
Australia).[21] The full text of the selected records will be assessed in detail against the inclusion
criteria by two independent reviewers (A.R. and J.I.). Reasons for exclusion of full-text records that
do not meet the inclusion criteria will be recorded and reported in the scoping review. Any
disagreements that arise between the reviewers at each stage of the selection process will be
resolved through discussion or with a third reviewer (M.K.). The search results will be reported in full
in the final scoping review and presented in a Preferred Reporting Items for Systematic Reviews and
Meta-analyses for Scoping Reviews (PRISMA-ScR) flow diagram.[22]

Data extraction

Data will be extracted from records included in the scoping review by two independent reviewers
(A.R. and J.I.) using a draft charting form developed by the reviewers. The team will trial the charting
form to ensure that all relevant results are extracted, and the table will be revise and amended
when required during the data charting process. Modifications will be detailed in the full scoping
review. Any disagreements that arise between the reviewers will be resolved through discussion or

with a third reviewer. The primary studies' authors will be contacted to request missing or additional

data, where required. The data extracted will include specific details such as author(s), year of

publication, country (where the cases were reported), study design, study population (age and

gender), medical history, COVID-19 symptoms, ulcer characteristics (color, size, number, location,

onset, and duration) and treatment. A draft extraction tool is provided (see Appendix 2).

Data analysis and presentation

The extracted data will be presented in tabular form according to the objectives of this scoping

review. Additionally, rating maps will be used to display the frequency of reported cases. The

tabulated results will be accompanied by a narrative summary demonstrating how the results relate

to the scoping review's objectives.

Ethical consideration

The ethical committee of Masaryk University – Faculty of Medicine has exempted this study from

getting approval due to the fact that it uses open-access data.

Conflicts of interest

The authors declare no conflict of interest.

Funding statement

Masaryk University supported the work of A.R. and V.C. by the grants MUNI/A/1608/2020 and

MUNI/IGA/1543/2020. The work of A.R., A.P. J.K. and M.K. was supported by the INTER-EXCELLENCE

grant number LTC20031 – "Towards an International Network for Evidence-based Research in

Clinical Health Research in the Czech Republic".

Data availability statement

The data that support the findings of this study are openly available in Open Science Framework at

https://doi.org/10.17605/OSF.IO/JWAU4

References

1.   Riad A, Issa J, Chuchmova V, Gomaa E, Pokorna A, Klugarova J, et al. Oral ulcers of COVID-19

patients: a scoping review protocol [Internet]. Open Science Framework (OSF) Registeries.

2021 [cited 2021 Jan 22]. Available from: https://osf.io/jwau4

2.   World Health Organization (WHO). Coronavirus disease (COVID-19) [Internet]. [cited 2021 Jan
20]. Available from:
https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19

3.   Wollina U, Karadağ AS, Rowland-Payne C, Chiriac A, Lotti T. Cutaneous signs in COVID-19
patients: A review [Internet]. Vol. 33, Dermatologic Therapy. Blackwell Publishing Inc.; 2020
[cited 2021 Jan 20]. Available from: /pmc/articles/PMC7273098/?report=abstract

4.   Daneshgaran G, Dubin DP, Gould DJ. Cutaneous Manifestations of COVID-19: An
Evidence-Based Review [Internet]. Vol. 21, American Journal of Clinical Dermatology. Adis;
2020 [cited 2021 Jan 20]. p. 627–39. Available from:
https://doi.org/10.1007/s40257-020-00558-4

5.   Chi AC, Neville BW, Krayer JW, Gonsalves WC. Oral Manifestations of Systemic Disease
[Internet]. Vol. 82, American Family Physician. 2010 Dec [cited 2021 Jan 20]. Available from:
www.aafp.org/afpAmericanFamilyPhysician

6.   Muñoz-Corcuera M, Esparza-Gómez G, González-Moles MA, Bascones-Martínez A. Oral
ulcers: clinical aspects. A tool for dermatologists. Part II. Chronic ulcers. Clin Exp Dermatol
[Internet]. 2009 Jun 1 [cited 2021 Jan 20];34(4):456–61. Available from:
http://doi.wiley.com/10.1111/j.1365-2230.2009.03219.x

7.   Muñoz-Corcuera M, Esparza-Gómez G, González-Moles MA, Bascones-Martínez A. Oral
ulcers: clinical aspects. A tool for dermatologists. Part I. Acute ulcers. Clin Exp Dermatol
[Internet]. 2009 Apr 1 [cited 2021 Jan 20];34(3):289–94. Available from:
http://doi.wiley.com/10.1111/j.1365-2230.2009.03220.x

8.   Chaux-Bodard AG, Deneuve S, Desoutter A. Oral manifestation of Covid-19 as an inaugural
symptom? [Internet]. Vol. 26, Journal of Oral Medicine and Oral Surgery. EDP Sciences; 2020
[cited 2020 Nov 25]. p. 18. Available from: https://doi.org/10.1051/mbcb/2020011

9.   Riad A, Klugar M, Krsek M. COVID-19 Related Oral Manifestations, Early Disease Features?
Oral Dis [Internet]. 2020 Jun 30 [cited 2020 Jul 13];odi.13516. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/odi.13516

10.   Eghbali Zarch R, Hosseinzadeh P. COVID-19 from the perspective of dentists: A case report
and brief review of more than 170 cases. Dermatol Ther [Internet]. 2021 Jan [cited 2021 Jan

220   21]; Available from: https://onlinelibrary.wiley.com/doi/10.1111/dth.14717

221 11. Iranmanesh B, Amiri R, Zartab H, Aflatoonian M. Oral manifestations of COVID-19 disease: A
222   review article. Dermatol Ther [Internet]. 2020 Nov 25 [cited 2020 Dec 8];dth.14578. Available
223   from: https://onlinelibrary.wiley.com/doi/10.1111/dth.14578

224 12. Riad A, Boccuzzi M, Sagiroglu D, Klugar M, Krsek M. Pediatric multisystem inflammatory
225   syndrome temporally associated with SARS-COV-2: Oral manifestations and implications. Int J
226   Paediatr Dent [Internet]. 2020 Aug 21 [cited 2020 Aug 30];ipd.12694. Available from:
227   https://onlinelibrary.wiley.com/doi/abs/10.1111/ipd.12694

228 13. Amorim dos Santos J, Normando AGC, Carvalho da Silva RL, De Paula RM, Cembranel AC,
229   Santos-Silva AR, et al. Oral mucosal lesions in a COVID-19 patient: New signs or secondary
230   manifestations? [Internet]. Vol. 97, International Journal of Infectious Diseases. Elsevier B.V.;
231   2020 [cited 2020 Dec 10]. p. 326–8. Available from:
232   https://pubmed.ncbi.nlm.nih.gov/32526392/

233 14. Brandão TB, Gueiros LA, Melo TS, Prado-Ribeiro AC, Nesrallah ACFA, Prado GVB, et al. Oral
234   lesions in patients with SARS-CoV-2 infection: could the oral cavity be a target organ? Oral
235   Surg Oral Med Oral Pathol Oral Radiol [Internet]. 2020 Aug 18 [cited 2020 Dec 12]; Available
236   from: /pmc/articles/PMC7434495/?report=abstract

237 15. Riad A, Kassem I, Stanek J, Badrah M, Klugarova J, Klugar M. Aphthous Stomatitis in COVID-19
238   Patients: Case-series and Literature Review. Dermatol Ther [Internet]. 2021 Jan 3 [cited 2021
239   Jan 3]; Available from: https://onlinelibrary.wiley.com/doi/10.1111/dth.14735

240 16. Riad A, Kassem I, Hockova B, Badrah M, Klugar M. Tongue ulcers associated with SARS-CoV-2
241   infection: A case series. Oral Dis [Internet]. 2020 Sep 25 [cited 2021 Jan 13];odi.13635.
242   Available from: https://onlinelibrary.wiley.com/doi/10.1111/odi.13635

243 17. Al-Khanati NM, Riad A, Sahloul ME, Klugar M. Aphthous-like stomatitis of COVID-19 patients.
244   Brazilian J Oral Sci [Internet]. 2020 Nov 4 [cited 2020 Dec 4];19:e201354–e201354. Available
245   from: http://dx.doi.org/10.20396/bjos.v19i0.8661354

246 18. NHMRC NH and MRC. Australian Guidelines for the clinical care of people with COVID-19
247   [Internet]. Australian Clinical Practice Guidelines. 2020 [cited 2020 Sep 17]. Available from:
248   https://www.clinicalguidelines.gov.au/register/australian-guidelines-clinical-care-people-covi

d-19

19.   Vincent JL, Rubenfeld GD. Does intermediate care improve patient outcomes or reduce costs?
Crit Care [Internet]. 2015 Mar 2 [cited 2021 Jan 23];19(1):89. Available from:
http://ccforum.com/content/19/1/89

20.   Peters M, Godfrey C, McInerney P, Baldini Soares C, Khalil H, Parker D. JBI Manual for
Evidence Synthesis [Internet]. JBI Manual for Evidence Synthesis. JBI; 2020 [cited 2021 Jan
21]. Available from: https://wiki.jbi.global/display/MANUAL

21.   Munn Z, Aromataris E, Tufanaru C, Stern C, Porritt K, Farrow J, et al. The development of
software to support multiple systematic review types: The Joanna Briggs Institute System for
the Unified Management, Assessment and Review of Information (JBI SUMARI). Int J Evid
Based Healthc [Internet]. 2019 Mar 1 [cited 2021 Jan 22];17(1):36–43. Available from:
https://pubmed.ncbi.nlm.nih.gov/30239357/

22.   Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for
scoping reviews (PRISMA-ScR): Checklist and explanation [Internet]. Vol. 169, Annals of
Internal Medicine. American College of Physicians; 2018 [cited 2021 Jan 22]. p. 467–73.
Available from: https://pubmed.ncbi.nlm.nih.gov/30178033/

Appendix I: Search strategy

MEDLINE (PubMed)

Search conducted on January 22ⁿᵈ, 2021, from (01.01.2020) to present.

Search	Query	Records retrieved
#1	COVID-19[all] OR SARS-CoV-2[all] OR 2019-nCOV[all] OR severe acute respiratory syndrome[all] OR coronavirus[all] OR corona virus[all]	119,920
#2	Ulcer*[all] OR aphthous[all] OR herpe*[all] OR blister*[all] OR eruption*[all] OR erosion*[all] OR lesion*[all] OR vesicle*[all] OR stomatitis[all]	1,472,960
#3	Oral[all] OR mouth[all] OR tongue[all] OR lip*[all] OR labial[all] OR buccal[all] OR perioral[all] OR palate[all] OR pharyn*[all]	1,271,073
#4	#1 AND #2 AND #3	219
Limited to the date range from (01.01.2020) to present		126

268 ## Appendix II: Data extraction instrument

Author, Date, Location	Number of patients	Gender	Age	Medical anamnesis	COVID-19 symptoms	COVID-19 treatment	Ulcer characteristics				Ulcer Treatment
							Location	Number / Size	Necrosis / Hemorrhage	Onset / Duration	