Genome-wide meta-analysis of pneumonia suggests a role for mucin biology and provides novel drug repurposing opportunities

William R. Reay[1,2], Michael P. Geaghan[1,2], 23andMe Research Team[3†], Murray J. Cairns[1,2*]

[1]School of Biomedical Sciences and Pharmacy, Faculty of Health and Medicine, The University of Newcastle, Callaghan, NSW, 2308, Australia

[2]Hunter Medical Research Institute, Newcastle, NSW, 2305, Australia

[3]23andMe Inc., Sunnyvale, CA, 94086, United States of America

[†]A list of authors and their affiliations appears at the end of the paper

[*]To whom correspondence should be addressed:

Professor Murray Cairns, Medical Sciences Building, University Drive, Callaghan, NSW 2308, Australia

Email: Murray.cairns@newcastle.edu.au, Phone: +61 02 4921 8670

ABSTRACT

Pneumonia remains one of the leading causes of death worldwide, particularly amongst the elderly and young children. We performed a genome-wide meta-analysis of lifetime pneumonia diagnosis (N=266,277), that encompassed the largest collection of cases published to date. Genome-wide significant associations with pneumonia were uncovered for the first time beyond the major histocompatibility complex region, with three novel loci, including a signal fine-mapped to a cluster of mucin genes. Moreover, we demonstrated evidence of a polygenic effect of common and low frequency pneumonia associated variation impacting several other mucin genes and *O*-glycosylation, further suggesting a role for these processes in pneumonia pathophysiology. The pneumonia GWAS was then leveraged to identify drug repurposing opportunities, including evidence that supports the use of lipid modifying agents in the prevention and treatment of the disorder. We also propose how polygenic risk could be utilised for precision drug repurposing through pneumonia risk scores constructed using variants mapped to pathways with known drug targets. In summary, we provide novel insights into the genetic architecture of pneumonia susceptibility, with future study warranted to functionally interrogate novel association signals and evaluate the suitability of the compounds prioritised by this study as repositioning candidates.

INTRODUCTION

Pneumonia is characterised as an acute infection of the lung, with fluid filled alveoli and resultant restriction of oxygen intake being a key hallmark of its pathophysiology. There are a number of mechanisms known to cause pneumonia, however, bacterial or viral infection are the most common aetiologies[1]. Pharmacological intervention in pneumonia treatment is largely dependent on the infection source – for instance, bacterial induced pneumonia is treated with antibiotics. Yearly mortality rates worldwide from pneumonia remain high, even in the developed world where access to antibiotics and routine hospital care is usually unrestricted[2,3]. This necessitates a greater understanding of the mechanisms involved in pneumonia susceptibility and pathogenesis, which could be leveraged to identify novel treatments and inform the repositioning of existing drugs.

There has been considerable work undertaken to identify host factors which influence the onset and clinical course of pneumonia. Twin-based estimates of pneumonia heritability are still lacking, however, the heritability of death due to infections disease has been estimated as high as 40%, although further study is required [4]. There have also been few studies which have used modern statistical genetics approaches to test for the existence of risk-increasing or protective alleles associated with pneumonia with sufficient power for surpassing genome-wide significance. Previously, a genome-wide association study of lifetime self-reported pneumonia diagnosis was published using participants obtained by 23andMe Inc. that identified a significant signal in the major histocompatibility complex (MHC) region on chromosome six[5]. We sought to increase statistical power to detect association signals by performing a genome-wide meta-analysis of self-reported pneumonia in the 23andMe cohort with SNP effects on a clinically ascertained pneumonia phenotype from the FinnGen consortium. The genetic architecture of pneumonia was further interrogated to identify novel risk genes and salient biological pathways, along with an estimate of genetic correlation with clinically significant phenotypes. These data were then considered in light of drug repurposing and provided support to a number of plausible repositioning opportunities.

MATERIALS AND METHODS

Genome-wide meta-analysis of pneumonia

The genome-wide meta-analysis was performed using two primary study cohorts from 23andMe Inc. and FinnGen (release 3), respectively, with full details of these cohorts and the meta-analysis procedure detailed in the supplementary methods. Summary statistics for a self-reported pneumonia phenotype were obtained from 23andMe as outlined by Tian *et al.*[5]. This self-reported phenotype was derived from an online survey of 23andMe customers about their medical history. In the final GWAS after quality control (QC), there were 40600 cases and 90039 controls. In addition, summary statistics for pneumonia were downloaded from the third release of the FinnGen database which combines genotype data from Finnish biobanks and digital health record data from Finnish health registries. The pneumonia phenotype chosen was *All pneumoniae* (J10 pneumonia), for which 15771 cases and 119867 controls were available for GWAS after QC.

The 23andMe and FinnGen summary statistics were meta-analysed using an inverse-variance weighted model with fixed effects as implemented by METAL version March 2011[6]. Firstly, we meta-analysed common variants, defined as sites with allele frequency > 1% in both the 23andMe and FinnGen cohorts. Variants were retained if they were available in both summary statistics and had an imputation quality that exceeded a minimum of 0.3 or a mean of 0.5 for variants not physically genotyped, resulting in 6888413 sites with an effect size estimate from the meta-analysis and a total sample size of 266277 individuals. Imputed rare variants available in both studies were subjected to a stricter filtering threshold for imputation quality such that only variants with a minimum imputation quality > 0.5 or a mean value > 0.7 were subjected to meta-analysis, with 834366 low frequency variants considered. In both instances, we further tested for heterogeneity between the contributing studies using Cochran's Q test. Genome-wide summary statistics from the IVW meta-analysis were processed using the FUMA v1.3.6 (Functional Mapping and Annotation of Genome-Wide Association Studies) platform[7]. Genome-wide significant variants were characterised using the traditional $P < 5 \times 10^{-8}$ threshold, whilst suggestive significance was defined using a more lenient threshold of $P < 1 \times 10^{-5}$. We utilised the default settings for defining independent significant SNPs ($r^2 \leq 0.6$) and lead SNPs ($r^2 \leq 0.1$). The reference panel population for LD estimation was the UK biobank release 2b 10k White British panel, with LD blocks within 250 kb of each other merged into a single locus. We examined the effect of conditioning on two smoking GWAS via the

multi-trait-based conditional & joint analysis (mtCOJO) framework implemented in GCTA v

1.93.2 beta (Supplementary Methods)[8,9]. For the *MUC5AC* lead SNP, we additionally

performed a phenome-wide association study using the IEUGWAS database version 3.7.0

(https://gwas.mrcieu.ac.uk/), reporting SNPs using a conventional phenome-wide significance

threshold of $P < 1 \times 10^{-5}$. Given the most significant association in this database was a GWAS

of adult-onset asthma [10], we tested whether the association of SNPs proximal to *MUC5AC* was

driven by the same underlying causal variant, assuming a single causal variant, via the *coloc*

colocalisation methodology implemented in version 4 of the package [11]. We also sought to

replicate our results in two UK biobank (UKBB) pneumonia GWAS, specifically, a self-

reported pneumonia phenotype performed in the automated GWAS pipeline by the MRC IEU

group (ukb-b-4533, https://gwas.mrcieu.ac.uk/datasets/ukb-b-4533/), as well as a phecode

ICD-10 UKBB GWAS performed in an automated series of GWAS by the authors of the

SAIGE methodology (https://pheweb.org/UKB-SAIGE/pheno/480) [12].

Estimation of SNP-based heritability

SNP based heritability was computed using LD score regression (LDSR) with 1000 genomes

phase 3 LD scores and weights[13]. We converted the heritability estimate to the liability scale

assuming the population prevalence of pneumonia as that of pneumonia in the FinnGen dataset

(12.61%), as well as a more conservative estimate based on ICD-10 diagnosed pneumonia in

the UK biobank (UKBB) sample (3.20% - Supplementary Methods).

Finemapping genome-wide significant loci

We finemapped the three-novel genome-wide significant loci outside of the MHC region by

using a method which leverages asymptotic Bayes' factors (ABF) to estimate credible sets

under the assumption of a single causal variant[14]. Specifically, we utilised Wakefield's method

to approximate ABFs assuming a prior variance of 0.2^2, which reflects the belief that the

confidence intervals of estimated variant effect sizes expressed as odds ratios ranging from

around 0.68 to 1.48. Given that the posterior probability for causality of each variant is

proportional to its Bayes' factor, these can be summed until a prespecified probability (ρ) is

reached, thus, constituting a ρ set of putative causal variants. In this study, we derived 95%

credible sets. A single causal variant was assumed such that we did not have to account for LD

between variants, which has been demonstrated to be problematic in finemapping studies

which prespecify more than one causal variant using references external to the GWAS like the

1000 genomes project panel[15].

Gene-based and gene-set association

Common variant (MAF > 0.01) SNP-wise *P* values were aggregated at gene-level using MAGMA v1.07b[16], as described in the supplementary methods. The Bonferroni threshold for genic association was $P < 2.68 \times 10^{-6}$, accounting for the number of genes tested. Moreover, gene-based *P* values were leveraged for gene-set association using 1379 hallmark and canonical gene-sets from the Molecular Signatures Database (MSigDB)[17]. Rare variants (MAF < 0.01) were also aggregated at gene-level by leveraging the properties of the Cauchy distribution (Supplementary Methods). Code for the Cauchy combination test was obtained from (https://github.com/yaowuliu/ACAT) and outlined by Liu and Xie[18,19]. The MAGMA approach for common variants accounts for dependency between *P* values by estimating their covariance as a function of pairwise LD in a population sample – however, there are methodological challenges with this approach for rare variants and likely much larger samples would be required for accurate estimation of dependency between rare variants, if any exists[20,21]. Therefore, we employed Cauchy transformation to combine *P* values as it guards against type I error inflation due to potential unknown covariance between rare variants (Supplementary Methods). In addition, we constructed a model for rare variant gene-set association analogous to the MAGMA approach for common variants that leverages gene-based *Z* values (probit transformation of *P*). The same collection of pathways from MSigDB were considered, with genic *Z* values regressed against a binary indicator of set membership (β_S), covaried for logarithmically transformed gene-length, and rare variant count per gene. A one-sided test was performed for β_S, such that the null hypothesis is $\beta_S = 0$ and the alternative $\beta_S > 0$. Only gene-sets with rare variants overlapping at least 5 genes were retained.

Transcriptome-wide association studies of pneumonia

A transcriptome-wide association study (TWAS) of pneumonia was performed using the FUSION package[22]. We utilised GTEx v7 SNP weights from three tissues that would be plausibly involved in the pathophysiology of pneumonia (whole blood, lung, and spleen). We corrected for the number of *cis*-heritable genes outside the MHC region for which a TWAS *Z* could be calculated in each tissue (Supplementary Methods). For transcriptome-wide significant genes, we tested whether the expression and pneumonia-associated signal displayed statistical colocalisation as encompassed by the SNP weights with the *coloc* package as implemented by FUSION[11]. In addition, we probabilistically finemapped transcriptome-wide significant regions using the FOCUS approach to derive a credible set of putative causal genes, as described previously[23]. We utilised the default Bernoulli prior ($p = 1 \times 10^{-3}$) and chi-square

171   prior variance ($n\sigma^2 = 40$) to approximate Bayes' factors for each gene, and thus, derive the

172   posterior inclusion probabilities (*PIP*) for each gene to be causal given its observed TWAS *Z*.

Genetic correlation and causal inference

We estimated genetic correlation between pneumonia and 180 high quality, European ancestry

GWAS using LDSR as implemented by the LDhub application[24]. For Bonferroni significant

genetic correlation estimates, we constructed a latent causal variable (LCV) model using the

most significant trait from each LDhub phenotypic category and pneumonia to evaluate

evidence for genetic causality between traits, as outlined extensively elsewhere[25–27]. A strong

estimate of the posterior genetic causality proportion (GCP) was defined as significantly

different from zero (one sided *t*-test) and an absolute GCP estimate > 0.6. Weak GCP estimates

close to zero for genetically correlated traits imply that their relationship is potentially mediated

by horizontal pleiotropy, whereby there are shared pathways, but the two traits do not likely

exhibit vertical pleiotropy by acting within the same pathway. We additionally evaluated

evidence for a causal relationship between HDL cholesterol and pneumonia by constructing a

multivariable Mendelian randomisation (MR) model using the TwoSampleMR package[28]. This

multivariable model leveraged genetic instrumental variables (IV) from three highly

biologically interconnected lipid traits (LDL, HDL, and triglycerides) and estimated the effects

of these IVs on the outcome conditioned on their association with the other two lipid classes[29].

Genetically informed drug repurposing

We implemented three strategies to propose drug repurposing candidates: i) single loci drug-

gene matching, ii) genetic correlation and/or evidence of a putative causal relationship between

a biochemical trait that could be targeted by an approved drug, and iii) precision drug

repurposing using the polygenic scoring orientated *pharmagenic enrichment score* (PES)

approach. Full details of these analyses are described in the supplementary methods. We

utilised a panel of 50 biochemical GWAS performed by the Neale lab from the UK biobank

which had high or medium confidence estimates of SNP heritability that were significantly

different from zero (http://www.nealelab.is/uk-biobank) and estimated genetic correlation and

the posterior mean GCP for trait pairs that survived Bonferroni correction. We sought to

replicate the results of the multivariable MR for the three lipid classes (LDL, HDL, and

Triglycerides) using the UK biobank GWAS. MR was then performed to evaluate further

evidence for a causal effect between gamma-glutamyltransferase (GGT) and pneumonia, as

well as triglycerides and pneumonia (univariable estimate). Specifically, we defined independent, non-palindromic genome-wide significant variants as IVs and constructed four MR models with differing underlying assumptions (two inverse-variance weighted estimators with fixed or multiplicative random effects, weighted median estimator, weighted mode estimator, and MR-Egger)[30–33]. A series of sensitivity analyses to evaluate statistical evidence for confounding pleiotropy was then undertaken as outlined in the supplementary methods[32–35].

The PES framework is based on the postulation that an enrichment of genetic risk within a biological pathway with known drug targets may be an impetus to repurpose a drug which modulates that pathway for individuals who carry the high genetic load mapped to the pathway, as described elsewhere[36,37]. Specifically, we identify druggable pathways with an enrichment of common variant associations relative to the rest of the genes tested and construct pathway-based risk scores for these gene-sets (Supplementary Methods). We utilised the UK biobank (UKBB) cohort to test the association between a pneumonia PES and pneumonia phenotypes recorded for these participants[38,39]. These analyses are described in detail in the supplementary methods. Briefly, we retained 336,896 unrelated white British ancestry participants and 13,568,914 autosomal variants that survived a series quality control steps, including, imputation quality filtering (INFO > 0.8), $MAF > 1 \times 10^{-4}$, call rate > 0.98, and filtering strong deviations from the Hardy-Weinberg equilibrium. Self-reported pneumonia diagnosis and ICD-10 codes from hospital inpatient records were used to construct the pneumonia phenotype (Supplementary Methods). There were 10,540 individuals from the genotyped subset of the cohort included in the PES calculation with a primary or secondary diagnosis using the ICD-10 primary or secondary diagnosis codes relevant to pneumonia. In the strict phenotype definition, we defined cases as those satisfying ICD-10 criteria, and controls as all those who did not have one of those codes recorded along with any individual who self-reported pneumonia without a pneumonia ICD-10 code ($N_{Controls} = 320,213$). Individuals who self-reported pneumonia but were not assigned a relevant ICD-10 code were excluded from the study cohort in this strict configuration. In the broad-phenotype definition, pneumonia cases were individuals with a relevant ICD-10 code or a self-reported lifetime pneumonia diagnosis (N = 15,138). In other words, the strict definition only included individuals with a pneumonia ICD-10 code. The PES was then constructed using common, autosomal variants outside of the MHC region mapped to genes in that pathway with PRsice2 assuming an additive model, with further details provided in the supplementary methods[40]. The *P* value threshold of including

variants in the PES was the same as what was used to identify the gene-set. A genome wide

PRS for pneumonia susceptibility was also constructed in an analogous fashion.

We explored the phenotypic relevance of pneumonia PES profiles using a random subset of

the UKBB (N ~ 10,000) for which a multiplex assay was performed to quantity

immunoglobulin G (IgG) antibody response to a series of antigens for infectious agents

selected for study, as outlined elsewhere[41]. The two phenotypes of interest here were a binary

indicator of *seropositivity* for 14 infections with seroprevalence > 5% in our genotyped subset

of the cohort, and amongst seropositive individuals, a continuous measure of antibody response

(mean fluorescence intensity) to each antigen for that infection – termed *seroreactivity*. The

correlation between PES or PRS and seroreactivity was assessed by linear regression, whilst

logistic regression was utilised for seropositivity. Both models were covaried for sex, age, age^2,

ten SNP derived principal components, genotyping batch, and two QC metrics related to the

antibody assay (Supplementary Methods). A heatmap of the regression *t* statistics was

constructed using the ComplexHeatmap package[42]

RESULTS

Novel common and rare variant loci associated with pneumonia

We performed a genome-wide meta-analysis of pneumonia using common and rare (MAF <

0.01) overlapping variants from 23andMe and FinnGen release three, with 6,888,413 and

834,366 common and low frequency sites tested, respectively. We estimated the SNP based

heritability as approximately 3.24% on the liability scale (Fig. 1b), using the incidence of

pneumonia in the FinnGen cohorts as the population prevalence (12.67%), although we

acknowledge the population prevalence of pneumonia is difficult to quantify. As a result, we

re-estimated h^2 using a more conservative population prevalence value based on phenotype

data from the UK biobank (3.20%), resulting in a lower estimate of $h^2_{SNP} = 0.0213$. The point

estimate of SNP-based h^2 was higher in the 23andMe cohort. $h^2_{SNP} = 0.054$, although the

estimate was more precise in the meta-analysis than in 23andMe and FinnGen alone: $Z_{Meta} =$

9.26, $Z_{23andMe} = 7.44$, and $Z_{FinnGen} = 4.12$. In line with previous comparisons between self-

reported and clinically ascertained phenotypes, the heritability estimate was lower in FinnGen

than 23andMe. There was some evidence of test statistic inflation when visualised as a *QQ* plot

(Supplementary Figure 1), however, the proportion of the polygenic signal in the meta-analysis

attributed to model misspecification and/or confounding was around 11% (LDSR ratio =

0.1145), whilst the mean χ^2 was large enough to estimate heritability (1.11).

There were four common genomic loci that surpassed the conventional genome-wide

significance threshold ($P < 5$ x 10^{-8}, Table 1, Fig. 1a,c, Supplementary Fig. 2-5). The effect

sizes of these common variant signals were small in accordance with expectation, with each

SNP increasing or decreasing the odds of pneumonia by around 5%. Unsurprisingly, the most

significant signal spanned the major histocompatibility complex (MHC) region, which has

been previously published as associated with pneumonia in the 23andMe cohort[5]. Due to the

complexity of this region, we define the MHC signal as a single locus, with the minor allele of

the lead common SNP associated with a small reduction in the odds of pneumonia (OR = 0.94

[95% CI: 0.92, 0.96], $P = 6.44$ x 10^{-13}). The most significant novel common signal in this study

was a region located on chromosome 11, with the lead SNP (rs28624253) located upstream of

the *MUC5AC* gene that encodes a mucin protein, with three other genes that encode a mucin

protein within 400 kilobases of the lead SNP *MUC6*, *MUC2*, and *MUC5B*. Importantly,

rs28624253 was similarly associated in the 23andMe ($P = 2.65$ x 10^{-6}) and FinnGen ($P = 3.42$

x 10^{-6}) cohorts and there was no appreciable evidence for differences in population structure

between the input GWAS driving this signal. Mucins are heavily glycosylated proteins that

play a number of important roles, particularly in relation to the maintenance of mucosal

barriers[43] . Mucin genes are known the exhibit somewhat pervasive genomic complexity and

evidence of heterogeneity between populations, to ensure that this signal is not just an artefact

of this, we performed a phenome-wide association study of the lead SNP and found that this

variant was associated with only relevant phenotypes to pneumonia. Specifically, it was linked

to adult-onset asthma, self-reported regular cough and mucus, and eosinophil count at a

conventional phenome-wide significance threshold ($P < 1$ x 10^{-5}). Given rs28624253 was

associated with adult-onset asthma at a more stringent level of genome-wide significance, we

tested whether the mucin signal observed for pneumonia and adult-onset asthma were driven

by the same underlying causal variant and found strong evidence to support this hypothesis

(posterior probability > 90%). We caution that this assumes the existence of a single causal

variant, which may be unrealistic given the complexity of this region. As we visualise in

supplementary figure 6, if one utilises a more conservative prior probability of a shared causal

variant than there is some evidence that there is a different underlying causal variant but that

the locus is still associated with both traits. It should also be noted that the odds increasing

allele for pneumonia (*G*) is perhaps counterintuitively associated with decreased risk of asthma,

which suggests a multifaceted biological mechanism which may be related to the role of mucins
in the airway.
308
The third most significant locus encompassed several genes including *SAMD8*, *DUSP13*, and
VDAC2, whilst locus four was largely intergenic, with some variants overlapping long-
noncoding RNAs with limited annotation information (Supplementary Text). There were also
two loci that almost surpassed genome-wide significance; specifically, a locus physically
mapped to the interleukin-6 receptor region (*IL6R* – lead SNP: rs12730036, $P = 5.61$ x 10^{-8}),
and a region on chromosome 12 where the lead SNP is mapped to an intron of *TNFRSF1A*,
which encodes a tumour necrosis factor receptor (lead SNP: rs1800693, $P = 5.5$ x 10^{-8}). We
integrated functional genomic and annotation data to prioritise candidate genes from the novel
genome-wide significant loci in this study, as described in the supplementary text. We found
that genes in the non-MHC loci implicated by at least two lines of evidence were enriched in
phenotypically relevant pathways such as mucus layer, lung fibrosis, and *O*-linked
glycosylation of mucins (Supplementary Figure 5, Supplementary Text). .Interestingly, despite
the meta-analysis combining a self-reported phenotype with clinically ascertained pneumonia
diagnoses, only the lead SNP in the MHC locus demonstrated any significant heterogeneity in
the effect sizes between the two cohorts (Cochran's Q, $P_{Het} = 0.04$, Table 1, Supplementary
Text, Supplementary Figure 7).

Figure 1. Genome-wide meta-analysis of pneumonia susceptibility. (a) Manhattan

plot of common variant GWAS for pneumonia, as is usual practice, each point is the -

$\log_{10} P$ value of a variant for association with pneumonia, with the red dotted line

indicative of genome-wide significance ($P < 5 \times 10^{-8}$). Lead SNPs are highlighted and

labelled on the plot, except for the MHC locus which we denote as "MHC" due to its

complexity. **(b)** Estimates of SNP-based heritability (h^2) on the liability scale for the

23andMe and FinnGen cohorts individually, as well as the using the inverse-variance

weighted effects meta-analysis of the two cohorts. The error bars represent the standard

error of h^2. **(c)** Region plots for the three-novel genome-wide significant loci outside of

the MHC region, the LD for each variant with the lead SNP estimates from the 1000

genomes phase III European reference set was utilised to colour the points.

Table one: Lead SNPs within common genome-wide significant loci associated with

pneumonia

Lead SNP	Locus	EA/NEA	EAF (NFE)	EAF (FIN)	OR	95% CI	P_{GWAS}	P_{Het}
rs9268966	MHC	G/A	0.26	0.33	0.94	0.92, 0.96	6.44e^{-13}	0.04
rs28624253	chr11:1110395-1232702	G/A	0.37	0.41	1.05	1.04, 1.07	5.27e^{-11}	0.44
rs10762653	chr10:76815686-76993015	G/A	0.16	0.18	1.06	1.04, 1.08	2.70e^{-9}	0.61
rs264954	chr2:104056454-104380545	T/C	0.46	0.45	0.96	0.95, 0.98	4.44e^{-8}	0.53

Common (MAF > 0.01) lead SNPs were defined as independent SNPs ($r^2 < 0.1$) within each genomic locus.

The effect allele (EA) and non-effect allele (NEA) was reported for this table such that the effect allele was the

minor allele. The effect allele frequency (EAF) is denoted in gnomAD v2.1.1 for non-Finish Europeans (NFE)

and Finns (FIN). All odds ratio and their respective confidence intervals were calculated relative to the EA.

Heterogeneity of effect between the 23andMe and FinnGen cohorts was tested using Cochran's Q, with the P

value of that test reported here (P_{Het}). Due to the complexity of locus 1 (MHC), we report only a single common

SNP for this locus. Locus coordinates in in hg19 assembly.

The novel genome-wide significant loci outside the MHC region were finemapped to estimate

a 95% credible set of plausible causal variants assuming a single causal variant in each region

(Supplementary Tables 1-3). The variants encompassed by the 95% credible set for the

rs28624253 locus were all proximally upstream/downstream of *MUC5AC*, or within the gene itself, supporting the relevance of this mucin gene for that association signal. The second most significant novel pneumonia associated locus discovered in this study (rs10762653 lead SNP) had a smaller credible set, with the lead SNP having a considerably high posterior probability than the remaining credible set SNPs (*PP* = 0.676). Interestingly, the lead SNP for this locus has low estimated LD with other proximal genome-wide significant SNPs (Fig 1c), suggesting the existence of several causal variants that cannot be accounted for by this method. Finally, the intergenic region spanned by the third novel genome-wide significant locus yielded a large credible set of over 200 variants and, as a result, further functional interrogation is required to mechanistically interpret this locus.

Smoking status was not included as a covariate in the respective GWAS meta-analysed, and thus, we sought to investigate whether genetic variants associated with smoking may confound our findings in this GWAS. Specifically, we genetically conditioned common variant associations on their effect size from a GWAS of smoking initiation and smoking heaviness using mtCOJO[8]. The effect sizes of the lead SNPs from the novel genome-wide significant loci were not greatly attenuated after conditioning on either of the smoking phenotypes and remained genome wide significant, with the exception of the locus tagged by rs264954 (Conditioned on smoking initiation: $P = 7.15 \times 10^{-7}$; Conditioned on smoking heaviness: $P = 1.05 \times 10^{-7}$). Furthermore, there was a slight reduction in the SNP heritability estimate on the liability scale, although this only amounted to a less than 0.5% difference after conditioning on either smoking phenotype – $h^2_{\text{Conditioned on smoking initiation}} = 3.08\%$, $h^2_{\text{Conditioned on smoking heaviness}} = 2.95\%$.

We also uncovered a genome-wide significant association between a rare intergenic variant in the MHC region and pneumonia - rs11962863, OR = 1.59 [95% CI: 1.44, 1.74], $P = 1.15 \times 10^{-9}$. This relatively large effect allele, however, did display statistically significant heterogeneity in its effect between the two cohorts ($P = 8.20 \times 10^{-5}$). This locus is considerably rarer in the Finnish population (AF = 5.8×10^{-4}) than non-Finnish Europeans in gnomAD (AF = 2.9×10^{-3}), which may account for its larger effect size in the FinnGen cohort. Due to the complexity of recombination and linkage in the MHC locus, the functional consequences of this variant remains difficult to interpret at an individual level without considering the local genomic context of affected individuals, such as HLA type. We also detected six additional regions with

rare variants that surpassed suggestive significance for association with pneumonia ($P < 1$ x 10^{-5}, Supplementary Table 4).

We sought to replicate our genome-wide significant and suggestively associated common loci using two GWAS from the independent UK Biobank cohort – specifically, we utilised two automated GWAS that encompassed a self-reported pneumonia phenotype (N_{Case} = 6572, $N_{Controls}$ = 456,361) and ICD-10 derived pneumonia diagnoses (N_{Case} = 10,059, $N_{Controls}$ = 398,538). We investigated both phenotyping approaches given our GWAS was a meta-analysis of self-reported and clinically ascertained data. In the self-reported pneumonia UKBB GWAS, we found that no SNPs replicated at genome-wide significance, however, the *MUC5AC* lead SNP was nominally associated in the same direction (β = 0.001, *SE* = 2.6 x 10^{-3}, P = 0.022), with the MHC lead SNP also was also nominally significant (P = 0.03) and the remaining two lead SNPs demonstrated no significant evidence of replication. The ICD-10 phenotype GWAS in the UKBB did not replicate any of our non-MHC genome-wide significant SNPs at even nominal significance, although MHC SNPs were found to be nominally significant. It should be noted that a limitation of both GWAS is that they focused only on either the self-reported or clinically ascertained phenotype in the UKBB, meaning some controls plausibly would have had pneumonia, and thus, decreasing power. Moreover, the effective sample sizes (N_{eff}) of these UKBB GWAS were markedly smaller than ours (177749 in the current discovery meta-analysis versus 25915 and 39246, respectively).

We also considered two very recent smaller sample-size pneumonia GWAS without publicly available summary statistics to see if we could replicate their findings. Firstly, Chen *et al.* performed a GWAS of pneumonia susceptibility and severity in the Vanderbilt University Biobank (BioVU, N_{Case} = 8889, $N_{Controls}$ = 60,767, N_{eff} = 31019) , European ancestry cohort)[44]. They found that a genome-wide significant common signal in Europeans associated with pneumonia severity, with the lead SNP rs10786398 nominally associated in our meta-analysis: β = -0.029, *SE* = 0.001, P = 2.5 x 10^{-4}, whilst we were unable to replicate the significant rare-variant association signal from that study as the variant was not available in our analyses. Moreover, a meta-analysis of a smaller previous FinnGen release and ICD-10 (N_{eff} = 94584) derived pneumonia in the UKBB found two genome-wide significant index SNPs in the 15q15.1 region that were directionally consistent in our analyses, although not statistically significant, with a trend observed for rs76474922: β = 0.025, *SE* = 0.01, P = 0.08 [45]. The SNP-

based heritability estimate from that study also closely mirrored ours (3.3% on the liability

scale), supporting the reliability of this study's estimate in a larger sample.

Gene and gene-set association further supports a role for mucin biology in pneumonia

We performed gene and gene-set association to investigate associations with pneumonia

beyond univariable SNP-phenotype relationships (Supplementary Tables 5,6). Six genes

outside of the MHC region were significant in the meta-analysis after the application of

multiple-testing correction (*MUC5AC*, *MUC5B*, *DUPD1*, *SAMD8*, *DUSP13*, and *TOX*), all of

which were within genome-wide significant loci except for the *TOX* gene ($P = 1.26$ x 10^{-6}),

which plays a role in T cell persistence during response to a pathogen[46,47]. Gene-set association

revealed a single gene-set for which its member genes were enriched with pneumonia

associated common variation relative to all other genes tested: *Termination of O-glycan*

biosynthesis – $\beta = 0.89$, $SE = 0.21$, $P = 1.05$ x 10^{-5}, $q = 0.01$. The strongest gene-based signal

in this pathway was accounted for by two mucin genes that span part of the genome-wide

significant loci on chromosome 11 (*MUC5AC, MUC2*), whilst there were four other mucin

genes within this set that displayed a nominal gene-based association ($P < 0.05$) outside that

region (*MUC15*, *MUC16*, *MUC12*, and *MUC17*). The signal from this gene-set remained

relatively robust upon using a more conservative definition of the genic boundaries for SNP to

gene annotation during gene-based association: $\beta = 0.66$, $SE = 0.19$, $P = 3.04$ x 10^{-4}. Rare

variants were then subjected to gene-based association by leveraging the properties of the

Cauchy distribution, such that covariance between P values did not need to be estimated. No

genes surpassed Bonferroni correction, considering genes with at least two rare variants. The

most significant gene found upon testing all genic rare variants was *ZNF19*, $P = 6.3$ x 10^{-5},

whilst *FREM1* was the top gene ($P = 8.16$ x 10^{-4}) considering variants annotated as exonic only

(Supplementary Tables 7,8). Furthermore, we utilised the gene-based results to construct a

competitive rare variant gene-set association model and tested the same gene-sets as in the

common variant analyses (Supplementary Tables 9,10). While there were no gene-sets that

survived multiple-testing correction utilising all genic rare variants or those with only exonic

annotated sites, we found some support for B lymphocyte antigen response in the rare variant

architecture of pneumonia, as the most significant association from the rare variant model was

with the *B cell antigen receptor* pathway - $\beta = 0.59$, $SE = 0.19$, $P = 7.7$ x 10^{-4}. Interestingly,

there was also a nominal rare variant signal amongst pathways related to carbohydrate

metabolism (*Metabolism of carbohydrates* – $P = 6.22$ x 10^{-3}, *Glycosaminoglycan degradation*

– $P = 0.02$) and glycosylation (*N-glycan biosynthesis* – $P = 0.01$), although these results must

be interpreted cautiously given, they do not survive multiple testing correction.

A transcriptome-wide association study was then undertaken to identify genes for which

genetically predicted expression was correlated with pneumonia susceptibility. We selected

SNP weights from three tissues which are plausibly biologically relevant to pneumonia

pathophysiology: lung, whole blood, and spleen. After applying Bonferroni correction to the

number of tests within each tissue outside the MHC individually, significant correlation was

observed between decreased predicted expression of *VDAC2* and pneumonia ($Z = -4.39$, $P =$

1.15×10^{-5}, Supplementary Table 11) using spleen tissue as the prediction model. The

association between *VDAC2* and pneumonia was also supported by SNP weights from the lung

and whole blood, although these did not surpass the Bonferroni threshold: $P_{\text{Lung}} = 5.52 \times 10^{-4}$,

$P_{\text{Whole blood}} = 9.97 \times 10^{-4}$. We tested a related, but distinct hypothesis by assessing evidence for

colocalisation between the GWAS signal spanning *VDAC2* and the SNP weights from the three

tissues utilised to construct the model of genetically predicted *VDAC2* expression.

Colocalisation assumes a biologically conservative parameter of a single shared causal variant

that underlies the relationship between *VDAC2* expression and pneumonia. We observed

heterogeneity between the three tissues, with moderately strong evidence of colocalisation

between *VDAC2* expression and SNP weights from whole blood ($PP_{\text{H4}} = 0.848$), however,

using lung and spleen expression SNP weights there was no strong evidence ($PP > 0.8$) for any

of the five colocalisation hypotheses. In lung, there was evidence for an association between

VDAC2 lung expression SNP weights and pneumonia ($PP_{\text{H3}} = 0.568$, $PP_{\text{H4}} = 0.363$), although

we were not able to clearly determine whether there were two independent SNPs driving the

association (H3) or a shared variant (H4). Interestingly, there was moderate support in the

model leveraging the spleen SNP expression weights that this signal was only associated with

pneumonia and not *VDAC2* expression ($PP_{\text{H2}} = 0.643$), which could be driven by the spleen

model of genetically regulated expression being somewhat less predictive ($R^2 = 0.038$, best

linear unbiased prediction), than in blood ($R^2 = 0.1$, LASSO) and lung ($R^2 = 0.083$, elastic net),

respectively. Probabilistic finemapping of the marginal TWAS Z scores in the *VDAC2* region

was then undertaken using a multi-tissue panel to assess evidence for whether *VDAC2* is the

causal gene in this region. This locus was dense with genes, with 26 unique genes within the

90% credible set of causal genes. There was moderate evidence that *VDAC2* was the most

probable causal gene at this locus given it had the largest absolute TWAS Z, the highest *PIP*

for an individual model from the adrenal gland ($PIP = 0.279$), and a relatively large cumulative

PIP for all 17 *VDAC2* models derived a variety of tissues that were finemapped ($PIP_{\text{Cumulative}}$

= 0.683). Further investigation is thus needed to refine this locus.

There were a number of other genes that trended towards surviving multiple-testing correction

in spleen and the other two tissues – including, *PLD4* in spleen and *STPG1* in lung. We

subjected genes that trended towards multiple testing correction (two orders of magnitude

above the Bonferroni threshold) to gene-set overrepresentation analysis to identify gene

ontologies enriched for these genes, with *immune system process* and *immune response*

overrepresented after multiple-testing correction, supporting the biological relevance of this

signal (Supplementary Table 12). For instance, downregulation of tumour necrosis factor

receptor gene *TNFRS19* in lung trended towards correlation with pneumonia, $Z = -3.44$, $P =$

5.78×10^{-4}.

Pneumonia displays genetic correlation with clinically significant phenotypes

We estimated genetic correlation between pneumonia and 180 GWAS using LDSR, with a

significant non-zero estimate of genetic correlation obtained for twenty phenotypes after the

application of multiple testing correction ($P < 2.78 \times 10^{-4}$, Supplementary Table 13, Fig. 2).

Interestingly, the most significant genetic correlation was found between pneumonia and

insomnia ($r_g = 0.496$, $SE = 0.075$, $P = 3.55 \times 10^{-11}$), which supports previous observational data

that insomnia and reduced sleep duration increased the risk of developing pneumonia[48,49]. In

addition, we uncovered significant genetic correlation with other clinically interesting

phenotypes including forced vital capacity ($r_g = -0.215$, $SE = 0.0361$, $P = 2.62 \times 10^{-9}$), obesity

($r_g = 0.3023$, $SE = 0.0466$, $P = 9.15 \times 10^{-11}$), and HDL cholesterol ($r_g = -0.2876$, $SE = 0.0664$,

$P = 1.48 \times 10^{-5}$).

Figure 2. Estimates of genetic correlation between pneumonia and a panel of

GWAS that survive multiple testing correction. Each panel of the forest plot is the

estimate of genetic correlation by LD score regression (+/- its standard error, denoted

by the error bars). The dotted lines represent a genetic correlation of zero. Panels were

grouped by the phenotypic category of the trait subjected to LDSR.

A latent causal variable (LCV) model was then constructed for the most significantly

correlated trait-pairs from each phenotypic category that comprised the 180 GWAS tested for

genetic correlation. The LCV approach leverages the bivariate effect size distribution of

SNPs in two GWAS and their LD scores to estimate a genetic causality proportion (GCP),

such that, evidence of partial genetic causality can be distinguished from genetic correlation.

We found strong evidence ($|\widehat{GCP}| > 0.6$) for partial genetic causality of cigarettes per day and

HDL on pneumonia. Both posterior mean GCP estimates were significantly different from

zero, however, the estimate of HDL $\rightarrow$ pneumonia ($\widehat{GCP} = 0.758$, $SE = 0.159$, $P = 2.20 \times 10^{-11}$)

was more precise than that of cigarettes per day $\rightarrow$ pneumonia ($\widehat{GCP} = 0.713$, $SE = 0.225$,

$P = 3.51 \times 10^{-3}$). The magnitude of the potential causal relationship between HDL and

pneumonia was further investigated using mendelian randomisation (MR). Given the

biological overlap between the genetic architecture of HDL and other lipid classes, we

constructed a multivariable MR model that conditioned HDL instrumental variables on their

association with LDL cholesterol and triglycerides, obtaining the SNP-exposure estimates

from the Willer *et al.* global lipids genetics consortium paper , as has been outlined elsewhere [29,50]. There was no evidence of a causal effect of HDL ($P = 0.65$) or LDL ($P = 0.47$) conditioned on the remaining lipid classes, although there was nominal evidence of a risk increasing effect of triglycerides on pneumonia – $\beta = 0.058$, $SE = 0.028$, $P = 0.035$. These data highlight the complexities of distinguishing between confounding pleiotropy and evidence for causal relationships, with further work needed to resolve whether the directionally disproportionate variant effect sizes for HDL → pneumonia captured by the LCV model represent a true evidence for a causal relationship or whether other factors like triglycerides may explain this relationship. In addition, the effect of other confounders like BMI on these relationships cannot also be ruled out, although BMI did not show evidence of a causal effect in the LCV model.

Opportunities for drug repurposing by leveraging the genetic architecture of pneumonia

We sought to interrogate the pneumonia GWAS to propose novel drug repurposing candidates that could be useful to treat patients diagnosed with pneumonia more effectively. Firstly, we utilised a very liberal approach which identified genes that were targeted by approved drugs outside the MHC region physically mapped to loci associated with pneumonia at a minimum of a suggestive significance threshold ($P < 1 \times 10^{-5}$). There were five such genes that displayed a high confidence interaction with an approved pharmacological agent with a known mechanism of action – *IL6R*, *SCNN1A*, *ATP2B1*, *ERBB2*, and *STAT5B*). For instance, Tocilizumab is a monoclonal antibody that targets *IL6R* which has been suggested as a repurposing opportunity to use for severe illness following SARS-CoV2 infection, although results from randomised controlled trials have been mixed in terms of efficacy[51]. We examined these genes in the TWAS analyses, however, only *IL6R* was significantly *cis*-heritable in one of the three tissues we utilised. Interestingly, there was a trend towards a correlation between downregulation of *IL6R* and increased odds of pneumonia, which would not support the use of anti-IL-6 receptor agents like tocilizumab – $Z = -3.148$, $P = 1.64 \times 10^{-3}$. In contrast, the lead SNP and odds increasing allele of the *IL6R* locus in this GWAS has been associated with increased *IL6R* levels in a protein quantitative trait loci study (pQTL), which would support the efficacy of tocilizumab. We caution that the pQTL signal is in high LD with a missense variant (rs2228145), and thus, antigen binding affinity may be altered to create an artefactual pQTL association. These antigen-binding affinity related effects require further investigation, particularly in light of the phenomenon of the non-synonymous rs2228415 C allele displaying correlation with increased protein abundance from the pQTL study[52] but with decreased

expression via RNAseq derived eQTL estimates from whole blood by GTEx, along with
decreased CRP levels in the UK biobank, a well-characterised biomarker of IL-6 receptor (IL-
6R) inhibition. Previous functional analyses of the rs2228145 non-synonymous allele have
demonstrated that it likely impairs IL6 signalling, with increased expression of soluble
circulating IL-6R but downregulation of the membrane bound isoform[53]. As a result, we
conclude that based on the genetic association alone that the anti-inflammatory effect of IL-6R
blockade may have a risk-increasing impact on pneumonia, although further work is required
to evaluate tocilizumab as a potential repurposing opportunity, particularly as its efficacy
would likely be dependent on its temporal application in the clinical course of the disease.

A panel of 50 metabolites and blood cell count phenotypes from the UK biobank (UKBB) with
moderate to high confidence SNP-based heritability estimates were then tested for genetic
correlation with pneumonia (Supplementary Table 14). The concept underlying this is that if
there is evidence of a relationship between a biochemical trait and pneumonia, then a drug
which modulates that trait in a risk-decreasing direction could be clinically useful, and thus,
repurposed for pneumonia. We found 13 biochemical traits from the UKBB panel that were
correlated with pneumonia after multiple-testing correction (Supplementary Table 15). The
most significant correlation was a positive genetic correlation with triglycerides, followed by
a positive correlation with glycaeted haemoglobin (HbA1c), which is interesting given
previous evidence that hyperglycaemia has a deleterious effect on lung function[27,54]. The
correlation between HbA1c and pneumonia may not be necessarily driven by glycaemic
biology, particularly as HbA1c is strongly influenced by haematological factors, although we
did observe weak evidence for a positive correlation between glucose and pneumonia that does
not survive multiple testing correction ($P = 0.01$, Supplementary Table 15). It should be noted
that this UKBB sample is a larger sample size than the triglyceride GWAS subjected to LDSR
in the broad LDhub GWAS panel from earlier in the manuscript, and thus, the estimate is more
significant in this analysis. In addition, the HDL GWAS from the UKBB was not included due
to anomalous estimates of large standard error when considering its heritability estimate
(Supplementary Methods). There was only very weak evidence from the LCV model for a
potential causal influence of triglycerides on pneumonia ($\widehat{GCP} = 0.465$, $SE = 0.226$, $P = 0.052$,
Figure 3a), although this broadly supports the results of the lipid multivariable MR described
earlier in the manuscript that provided nominal evidence of a deleterious impact of increased
triglycerides. We attempted to replicate the earlier multivariable MR results using the LDL,

HDL, and triglyceride GWAS from the UK biobank as the exposure traits rather than the Willer

et al. global lipids genetics consortium GWAS, with an analogous result that demonstrated an

odds-increasing effect of triglycerides but a non-significant effect of LDL or HDL cholesterol

(Supplementary Table 16). We followed this up by estimating a univariable estimate of

triglycerides → pneumonia, with a 5.4% mean increase in the odds of pneumonia per standard

deviation increase in triglycerides across the five MR methods utilised (Supplementary

Methods, Supplementary Table 17). The causal estimate was only statistically significant in

the case of the two IVW estimators: OR = 1.056 [95% CI: 1.01, 1.104], P = 0.017 (IVW

multiplicative random effects), however, the remaining models had relatively consistent point

estimates of the effect of triglycerides on pneumonia (Figure 3b). There was some nominal

evidence of heterogeneity amongst IV exposure-outcome estimates (Q = 212.51, df = 168, P =

0.01), however, the Egger intercept was not significantly different from – thus, we observed

no direct statistical evidence of confounding pleiotropy. As a result, these data suggest a

potential repurposing opportunity for drugs prescribed for hypertriglyceridemia, such as statins

and fibrates, with the caveat that the MR and LCV models from this study provide only

relatively weak to moderate support, particularly as the mean posterior GCP estimate was low,

and thus, genetic correlation could contaminate the MR estimate.

The remaining LCV models for biochemical traits genetically correlated with pneumonia did

not indicate any strong evidence of partial genetic causality, with the exception of a putative

effect of gamma-glutamyltransferase (GGT) on pneumonia that approximately reached the

threshold for a strong point estimate of the GCP - $\widehat{GCP}$ = 0.59, SE = 0.212, P = 1.08 x 10^{-4}.

GGT is an enzyme that is commonly characterised as a biomarker of liver dysfunction, with

some evidence of an immunological role for this enzyme, as well as its involvement in alveolar

gas exchange. We further investigated this putative causal relationship by leveraging 207

independent SNPs associated with GGT (P < 5 x 10^{-8}) in the UK biobank as IVs – approximated

variance explained by IVs = 6.47%, F-statistic = 114.91. The five MR models implemented in

this study yielded a roughly similar effect size per SD increase in GGT that increased the odds

of pneumonia (mean pneumonia OR per SD GGT increase = 1.055, Supplementary Table 18).

However, the estimates were not statistically significant, with the exception of the weighted

median model (OR = 1.087 [95% CI: 1.01, 1.17], P = 0.025). There was evidence of

heterogeneity amongst the IV exposure-outcome effects, which may be indicative of

confounding pleiotropy (Q = 301.87, df = 206, P = 1.501 x 10^{-5}), although given the large

number of IVs observed heterogeneity is not a surprising phenomenon. Importantly, there was

no statistical evidence of pleiotropy from the intercept of the MR egger model. We caution that

statistical approaches to assess pleiotropy do not and cannot rule out a confounding influence

on the MR estimate, and detailed biological annotation of the IVs would be warranted to

investigate this further. In summary, there may be a causal relationship between GGT and

increased odds of pneumonia which could support inhibition of this enzyme as a treatment

target, with several GGT inhibitors under active development and explored for use in

respiratory illness[55,56].

Furthermore, we compared the MR estimate of the effect of a standard deviation increase in

GGT and triglyceride concentration on the odds of pneumonia to an observational estimate

from the UK biobank sample (Supplementary Methods). We found that the observational

association between a standard deviation increase in triglyceride concentration and pneumonia

was analogous to the univariable and multivariable MR estimates (OR = 1.059 [95% CI: 1.042,

1.076], $P = 2$ x 10^{-12}), whilst the observed association between GGT and pneumonia was

stronger than the MR estimate, with each standard deviation associated with a 13.54% [95%

CI: 12.43%, 14.56%] increase in the odds of pneumonia amongst UK biobank participants.

Interestingly, these associations were also consistent amongst individuals in the UK biobank

with relatively lower risk of pneumonia, that is, non-smoking females aged 45 or younger at

time of assessment – triglycerides: OR = 1.335 [95% CI: 1.13, 1.555], $P = 2.67$ x 10^{-4}, GGT:

OR = 1.354 [95% CI: 1.161, 1.553], $P = 2.75$ x 10^{-5}. These variables are extremely

heterogeneous and there are many potential confounders of the observed effect sizes, however,

it supports the inferred relationship from the LCV and MR models.

701
702 **a**
712 **b**

Figure 3. Investigating the potential utility of modulating biochemical traits as drug repurposing opportunities for pneumonia. (**a**) Latent causal variable models constructed between genetically correlated trait pairs after Bonferroni correction. Each point represents the genetic causality proportion, with the *y* axis denoting the precision of the GCP estimate, that is, its *Z* score. The genetic correlation estimate between the two traits was utilised to shade the points, with the larger points also indicative of a larger GCP *Z* score. A positive GCP estimate is indicative of partial genetic causality of the biochemical trait → pneumonia, whilst a negative estimate represents the converse. (**b**) A Mendelian randomisation (MR) analysis of the effect of genetically proxied gamma glutamyl-transferase (GGT) and triglycerides on the odds of pneumonia. The scatter plot visualises the effect of each instrumental variable SNP on GGT or triglycerides verses its effect on pneumonia, with the regression trend line the MR estimate from each of the five models implemented. Similarly, the forest plot indicates the pneumonia odds ratio for each of the MR models, with the error bar indicative of the 95% confidence intervals. The MR models were as follows: mode = weighted mode estimator, median = weighted median estimator, IVW-FE = inverse-variance weighted estimator with fixed effects, IVW-MRE = inverse-variance weighted estimator with multiplicative random effects, Egger = MR-Egger regression.

Precision drug repurposing to treat pneumonia

We implemented the *pharmagenic enrichment score* (PES) approach to identify drug repurposing candidates that could be targeted more precisely based on genetic risk[27,37]. Briefly, the PES is a genetic risk score specifically within a biological pathway that is targeted by approved drugs. The concept underlying the PES is that individuals with elevated genetic risk within a particular druggable set of genes may benefit from a pharmacological agent that modulates the pathway in question. Firstly, we identified five druggable pathways that displayed an enrichment of the common variant genetic architecture of pneumonia at one of four *P*-value thresholds for the inclusion of variants in the model (FDR < 0.05, Table 2, Supplementary Table 19, Supplementary Methods). These included two complement-related pathways, *p53 signalling*, and *bile acid metabolism*.

Table 2. Candidate druggable gene-sets that could be utilised to calculate *pharmagenic enrichment scores*

PES gene-set	P_T[1]	P	Example drug/nutraceutical
P53 signalling pathway	0.05	2.13×10^{-8}	Fostamatinib
Lectin induced complement pathway	0.05	2.98×10^{-8}	Human immunoglobulin
Complement pathway	0.05	6.81×10^{-8}	Zinc
Bile acid metabolism	0.005	1×10^{-6}	Atorvastatin
RIG-I-like receptor signalling pathway	0.005	1.58×10^{-5}	Etanercept

[1]P_T = P value threshold for variant inclusion in the model

There were a number of diverse compounds that targeted these pathways, with compounds identified as repurposing candidates through testing whether there was a statistically significant overrepresentation of their targets in the gene-set, along with high confidence single drug-gene interactions (Supplementary Tables 20,21). For instance, targets of the micronutrient zinc were overrepresented amongst genes in the *Complement pathway* gene-set, supporting previous evidence that zinc can modulate complement activation [57,58].

Characteristics of pneumonia pharmagenic enrichment scores

We investigated the properties of these five scores in the UKBB cohort. Interestingly, there were no large correlations (all $r < 0.07$) between the PES and a genome-wide polygenic risk score for pneumonia, which supports our hypothesis that pathway-based risk scores may provide novel biological insights that are not encompassed by PRS constructed from variants throughout the genome. Both the genome wide PRS and the PES profiles were not significantly associated with pneumonia diagnosis (self-reported/clinically ascertained or clinically ascertained only, Supplementary Tables 22,23, Supplementary Methods). This is perhaps not surprising given the low SNP heritability of pneumonia and the heterogeneity of the phenotype – however, we believe this does not preclude the relevance of PES at an individual level. For instance, given the putative relationship between hypertriglyceridemia and increased odds of pneumonia, as supported by this study, we tested the relationship between the pneumonia *Bile acid metabolism* PES and measured triglycerides in the UKBB. There was a small but significant positive correlation between the PES and triglyceride concentration – $\beta = 0.01$, *SE* = 0.002, $P = 1.69 \times 10^{-4}$, which was significant even after adjusting for statin use or using triglyceride values as the outcome variable winsorized at three standard deviations above the

mean to guard against an excessive influence of outliers (β = 0.01, *SE* = 0.002, *P* = 1.43 x 10

^{4}). This relationship with triglycerides was not seen using genome wide PRS (β = 1.57 x 10^{-4},

SE = 0.002, *P* = 0.93).

Associations between pneumonia pharmagenic enrichment scores and host susceptibility to

infection

The relationship between PES, as a function of host genetic susceptibility to pneumonia, and

antibody response to infection was then assessed in a subset of the UKBB with these data

available and passing our genotyping QC thresholds (N = 6443). We implemented these

analyses given the host-immune response is directly relevant to pneumonia and if the PES

displays distinct correlations with antibody response compared to genome-wide pneumonia

PRS, it would emphasise the potential biological utility of the PES framework. Firstly, we

tested the association between each PES and antibody response to 28 antigens amongst

individuals with detectable levels of antibodies for 14 infections (seroreactivity, Fig. 4c,

Supplementary Table 24). The strongest relationship between a PES profile and seroreactivity

was a positive correlation between the *Complement pathway* PES and IgG response to the

major capsid protein VP1 of the BK polyomavirus, with each SD increase in the PES associated

with a 0.05 (0.01) SD increase in antibody response, *P* = 4.1 x 10^{-4}, which trended towards

surviving multiple testing correction (*q* = 0.07). There was no association between genome

wide PRS and IgG mediated response to this antigen. There were other nominally significant

correlations observed (Fig. 4a), with most of these correlations positive and potentially

indicative of an increased immune response. In addition, susceptibility to infection

(seropositivity) was also investigated (Supplementary Table 25). For instance, a nominal

association uncovered between *RIG-I-like receptor signalling pathway* and decreased odds of

herpes simplex virus-1 infection (OR = 0.93 per SD in score [95% CI: 0.87, 0.98], *P* = 4 x 10

^{3}), whilst *Complement pathway* PES was nominally associated with increased odds of positive

H. pylori serostatus - OR = 1.07 per SD in score [95% CI: 1.01, 1.12], *P* = 0.02.

a

b

c

Figure 4. The relationship between host genetics pneumonia pathway based *pharmagenic enrichment scores* **and antibody response**. (**a**) Nominally significant associations between pneumonia PES and seropositivity as a binary variable. The forest plot denotes the odds ratio of each infection serostatus (error bars represent 95% confidence intervals) per SD increase in the PES tested. The infections abbreviated are as follows: MCV = Merkel Cell Polyomavirus, JCV = Human Polyomavirus JCV, HSV = herpes simplex virus 1, HSV2 = herpes simplex virus 2, *H. pylori* = Helicobacter pylori. (**b**) IgG response (mean fluorescence intensity) to the major capsid protein VP1

847 of the BK polyomavirus in the 10th percentile (1st decile, denoted 1), 50th percentile (5th

848 decile, denoted 5), and 90th percentile (10th decile, denoted 10) of the complement

849 pathway PES amongst the genotyped subset of the UKBB subjected to antibody

850 screening included in our analyses. (**c**) Heatmap of the regression t statistic

851 (beta/standard error) for the correlation between each PES and PRS with respective IgG

852 response to antigens amongst individuals seropositive for that infection. Hierarchical

853 clustering was applied to the rows and columns using Pearson's correlation distance.

855 **DISCUSSION**

856 In this study, we uncovered the first significant association signals for lifetime pneumonia

857 susceptibility outside of the MHC region. Interestingly, there was also a novel low frequency

858 variant in the MHC itself which reached genome-wide significant that confers a relatively large

859 (~ 59%) increase in the odds of pneumonia, reflecting the immense heterogeneity spanned

860 within this region. Further analyses of the MHC signal are warranted, particularly to

861 deconvolve specific HLA types that may contribute to pneumonia susceptibility and

862 progression. Each of the loci beyond the MHC identified this study were relatively complex,

863 although we were able to derive a 95% credible set for the most significant non-MHC locus on

864 chromosome 11 that implicated the mucin gene *MUC5AC*. It should be noted that our fine-

865 mapping approach is relatively biologically naïve as it assumes a single causal variant, and

866 thus, the involvement of other genes remains unclear. *MUC5AC* is an interesting candidate

867 given that it has been previously implicated in the pathogenesis of respiratory illness and the

868 role of mucins in physical defence against pathogens via mucociliary clearance[59]. This heavily

869 glycosylated protein is lowly expressed in normal respiratory epithelium, however, is

870 upregulated upon in response to perturbagens, such as viral infection[60,61]. We posit that

871 upregulation of *MUC5AC* may be deleterious in the context of pneumonia given recent

872 evidence that this protein can enhance airway inflammation induced by viral infection[62],

873 although dissection of the mechanisms of variants in this locus are warranted, particularly given

874 the apparent discordant relationship of this signal we observed on the odds on asthma.

875 Interestingly, there are some preliminary data that suggests *MUC5AC* is upregulated in the

876 airway mucus of patients with severe COVID-19, although these studies were conducted using

877 small sample sizes[63,64]. There was also evidence of a more expansive polygenic signal amongst

878 mucin and beta-galactoside/N-acetylgalactosaminide genes in the *termination of O-glycan*

879 *biosynthesis pathway*, whereby sialic acid residues conjugated to mucins can terminate O-

880 glycan biosynthesis[65]. Interestingly, therapies specifically targeting mucin-linked O-

glycosylation are now under active development, including a recently proposed hexosamine

analog that demonstrated potent inhibition of O-glycan biosynthesis and downregulation of

neutrophil infiltration in rodents[66].

Drug repurposing is an attractive downstream application for GWAS, and we demonstrate its

potential utility for pneumonia through three distinct methods. Causal inference between

biochemical traits and pneumonia may provide repositioning opportunities along with a greater

understanding of pathological mechanisms in the disorder. For instance, we reveal evidence to

suggest a potential protective effect of HDL cholesterol and a deleterious impact of elevated

triglycerides on the odds of pneumonia. These data support observational data that lower

baseline HDL and elevated triglycerides have risk-increasing properties for pneumonia[67–69], .

We emphasise that our data only provided weak to moderate support for a causal influence of

lipid abundance on pneumonia, and replicated, well-powered randomised controlled trials are

needed to definitively assess the suitability of lipid-modifying agents like statins. A key

limitation of proposing drug repurposing candidates for the phenotype as a singular entity is

that it ignores the inherent heterogeneity of pneumonia onset and clinical course. It has been

shown previously that individual genes supported by GWAS significantly increase the

likelihood of a candidate drug being successfully approved from the phase I stage [70], however,

the relevance of more expansive association signals relating to biological networks for

precision medicine remains unclear As a result, we sought to implement an approach which

seeks to target drug repositioning opportunities to those with elevated genetic risk within

pathways relevant to the compound (PES, one of the prioritised pathways for the construction

of a PES was the *Bile acid metabolism* gene-set, further supporting the relevance of compounds

that modulate cholesterol. In the UKBB cohort, we demonstrated that these scores were distinct

from genome-wide PRS, and thus, may offer biological insights that were missed by using a

genome-wide score. For example, the *Bile acid metabolism* PES was positively correlated with

triglyceride levels, whilst this was not observed for PRS. We caution that one cannot draw a

causal inference from this relationship. In addition, the putative relationship between PES and

IgG response to antigens is biologically relevant both in terms of increased and decreased

antibody response. One can conceptualise the phenotype of pneumonia as having a contribution

from increased likelihood of infection, but also an aberrant inflammatory response once

infected with a pathogen. The distinct signal observed with these IgG phenotypes for the

pathway-based PES compared to genome-wide PRS, therefore, further highlights the potential

utility of the PES framework. Further work is required to evaluate the suitability of compounds

that modulate the PES pathways of interest and to categorise drug repurposing candidates based

on their suitability for prevention and/or treatment of pneumonia. The key advantage of the

PES framework is that only individuals with relevant genetic background in a pathway of

interest would be prioritised for the respective repurposing candidate, which would be useful

given the polygenic nature of complex disorders. Detailed discussion of the strengths and

limitations of the PES methodology have been featured in previous publications[27,37].

There are a number of important limitations that should be considered in light of the pneumonia

GWAS itself and our drug repurposing analyses. Firstly, this GWAS was conducted using

samples from European ancestry as large, diverse, genotyped cohorts with pneumonia status

information are not yet available. It will be critical to translate findings related to host-genetic

influences on pneumonia that future efforts strive to collect trans-ancestral data, particularly

due to concerns about the portability of European GWAS signals and the advantages in

finemapping afforded by including multiple ancestries[71]. The SNP heritability for pneumonia

derived in this study was also relatively low, and it remains unclear how heterogeneity amongst

the phenotype definition of pneumonia may contribute to this. In other words, given that

pneumonia is caused by a variety of factors and may go undiagnosed in some individuals,

detailed phenotyping data would potentially assist in resolving the genetic architecture of this

disorder. For example, a GWAS on susceptibility verses pneumonia severity will likely reveal

different biological insights. This could also be aided by stratified analyses by age, given

pneumonia is more pervasive in the elderly. The putative drug repurposing candidates

suggested in this study must also be viewed in light of the low heritability of pneumonia and

need for clinical validation. Despite these challenges, we believe that further resolving the host-

genetic architecture of pneumonia will be invaluable to public health efforts to more effectively

prevent and manage the illness. In summary, we revealed novel genome-wide significant loci

associated with life-time pneumonia susceptibility beyond the MHC region. These data

provided some support for the potential utility of triglycerides and GGT as treatment targets

for pneumonia, however, randomised controlled trials are now required to establish the efficacy

of such interventions. Moreover, the *pharmagenic enrichment score* approach may provide a

precision medicine-based intervention for drug repurposing for pneumonia prophylaxis and

treatment given an individual's composition of genetic risk. The properties of these scores and

the prospects of integrating them with other clinical metrics warrants further research.

DECLARATION OF INTERESTS

W.R.R and M.J.C have filed a patent related to the use of the pharmagenic enrichment score framework in complex disorders, the remaining authors declare no competing financial interests.

ACKNOWLEDGEMENTS

We wish to acknowledge the participants and investigators of FinnGen study and the 23andMe Inc. study from which these data were derived for the GWAS meta-analysis. In addition, his research has been conducted using the UK Biobank Resource under the application 58432. This study was supported by an NHMRC project grant (1147644). M.J.C. is supported by an NHMRC Senior Research Fellowship (1121474).

CONSORTIA

The 23andMe Research Team

Michelle Agee[3], Babak Alipanahi[3], Robert K. Bell[3], Katarzyna Bryc[3], Sarah L. Elson[3], Pierre Fontanillas[3], Nicholas A. Furlotte[3], Barry Hicks[3], David A. Hinds[3], Karen E. Huber[3], Ethan M. Jewett[3], Yunxuan Jiang[3], Aaron Kleinman[3], Keng-Han Lin[3], Nadia K. Litterman[3], Jennifer C. McCreight[3], Matthew H. McIntyre[3], Kimberly F. McManus[3], Joanna L. Mountain[3], Elizabeth S. Noblin[3], Carrie A. M Northover[3], Steven J. Pitts[3], G. David Poznik[3], J. Fah Sathirapongsasuti[3], Janie F. Shelton[3], Suyash Shringarpure[3], Chao Tian[3], Joyce Y. Tung[3], Vladimir Vacic[3], Xin Wang[3] & Catherine H. Wilson[3].

AUTHOR INFORMATION

School of Biomedical Sciences and Pharmacy, Faculty of Health and Medicine, The University of Newcastle, Callaghan, NSW, 2308, Australia

William R. Reay, Michael P. Geaghan, and Murray J. Cairns

Hunter Medical Research Institute, Newcastle, NSW, 2305, Australia

William R. Reay, Michael P. Geaghan, and Murray J. Cairns

23andMe Inc., Sunnyvale, CA, 94086, United States of America

The 23andMe Research Team

WEB RESOURCES

DGIdb v4.2.0 - https://www.dgidb.org/

FAVOR - http://favor.genohub.org/

FOCUS version July 24 2019 - https://github.com/bogdanlab/focus

FUMA v1.3.6 - https://fuma.ctglab.nl/

FUSION version May 29 2020 - https://github.com/gusevlab/fusion_twas

LCV version March 15 2019 - https://github.com/lukejoconnor/LCV

LDSR v1.0.1 - https://github.com/bulik/ldsc

MAGMA v.1.07b - https://ctg.cncr.nl/software/magma

METAL version March 2011 - https://genome.sph.umich.edu/wiki/METAL_Quick_Start

MR-PRESSO v1 - https://github.com/rondolab/MR-PRESSO/commits/master

mtCOJO (GCTA version 1.93.2 beta mac) -

https://cnsgenomics.com/software/gcta/#mtCOJO

PLINK2 v.2.00a3LM - https://www.cog-genomics.org/plink/2.0/

PRSice-2 v.2.3.3 (linux) - https://www.prsice.info/

TwoSampleMR v.0.5.5 - https://github.com/MRCIEU/TwoSampleMR

VEP GRCh37 release 101 August 2020 -

https://grch37.ensembl.org/Homo_sapiens/Tools/VEP?db=core

WebGestaltR v.4.0.2 - https://github.com/bzhanglab/WebGestaltR

1002

1003

DATA AND CODE AVAILABILITY

All data in this study are publicly available, summary statistics from 23andMe Inc. can be

obtained upon application to the company. As per 23andMe data sharing policies, we are

only able to release 10,000 SNPs from our meta-analysis, which we have made available on

GitHub

(https://github.com/Williamreay/Pneumonia_meta_GWAS_drug_repurposing/tree/master/Su

mmary_statistics). The full GWAS summary statistics for the 23andMe discovery data set

will be made available through 23andMe to qualified researchers under an agreement with

23andMe that protects the privacy of the 23andMe participants.

Researchers wishing to recapitulate our meta-analysis can apply for access for the 23andMe

subset of the study (https://research.23andme.com/dataset-access/), and then meta-analyse

with FinnGen release 3 summary statistics as described in our manuscript. Code utilised in

this study is available also on GitHub -

https://github.com/Williamreay/Pneumonia_meta_GWAS_drug_repurposing.

REFERENCES

1. Mackenzie, G. (2016). The definition and classification of pneumonia. Pneumonia (Nathan) *8*, 14.

2. Restrepo, M.I., Faverio, P., and Anzueto, A. (2013). Long-term prognosis in community-acquired pneumonia. Curr Opin Infect Dis *26*, 151–158.

3. McAllister, D.A., Liu, L., Shi, T., Chu, Y., Reed, C., Burrows, J., Adeloye, D., Rudan, I., Black, R.E., Campbell, H., et al. (2019). Global, regional, and national estimates of pneumonia morbidity and mortality in children younger than 5 years between 2000 and 2015: a systematic analysis. Lancet Glob Health *7*, e47–e57.

4. Obel, N., Christensen, K., Petersen, I., Sørensen, T.I.A., and Skytthe, A. (2010). Genetic and Environmental Influences on Risk of Death due to Infections Assessed in Danish Twins, 1943–2001. American Journal of Epidemiology *171*, 1007–1013.

5. Tian, C., Hromatka, B.S., Kiefer, A.K., Eriksson, N., Noble, S.M., Tung, J.Y., and Hinds, D.A. (2017). Genome-wide association and HLA region fine-mapping studies identify susceptibility loci for multiple common infections. Nat Commun *8*, 599.

6. Willer, C.J., Li, Y., and Abecasis, G.R. (2010). METAL: fast and efficient meta-analysis of genomewide association scans. Bioinformatics *26*, 2190–2191.

7. Watanabe, K., Taskesen, E., van Bochoven, A., and Posthuma, D. (2017). Functional mapping and annotation of genetic associations with FUMA. Nat Commun *8*, 1826.

8. Zhu, Z., Zheng, Z., Zhang, F., Wu, Y., Trzaskowski, M., Maier, R., Robinson, M.R., McGrath, J.J., Visscher, P.M., Wray, N.R., et al. (2018). Causal associations between risk factors and common diseases inferred from GWAS summary data. Nat Commun *9*, 224.

9. Liu, M., Jiang, Y., Wedow, R., Li, Y., Brazel, D.M., Chen, F., Datta, G., Davila-Velderrain, J., McGuire, D., Tian, C., et al. (2019). Association studies of up to 1.2 million individuals yield new insights into the genetic etiology of tobacco and alcohol use. Nat Genet *51*, 237–244.

10. Ferreira, M.A.R., Mathur, R., Vonk, J.M., Szwajda, A., Brumpton, B., Granell, R., Brew, B.K., Ullemar, V., Lu, Y., Jiang, Y., et al. (2019). Genetic Architectures of Childhood- and Adult-Onset Asthma Are Partly Distinct. Am J Hum Genet *104*, 665–684.

11. Giambartolomei, C., Vukcevic, D., Schadt, E.E., Franke, L., Hingorani, A.D., Wallace, C., and Plagnol, V. (2014). Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. PLoS Genet. *10*, e1004383.

12. Zhou, W., Nielsen, J.B., Fritsche, L.G., Dey, R., Gabrielsen, M.E., Wolford, B.N., LeFaive, J., VandeHaar, P., Gagliano, S.A., Gifford, A., et al. (2018). Efficiently controlling

for case-control imbalance and sample relatedness in large-scale genetic association studies. Nat. Genet. *50*, 1335–1341.

13. Bulik-Sullivan, B., Finucane, H.K., Anttila, V., Gusev, A., Day, F.R., Loh, P.-R., ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Duncan, L., et al. (2015). An atlas of genetic correlations across human diseases and traits. Nat. Genet. *47*, 1236–1241.

14. Wellcome Trust Case Control Consortium, Maller, J.B., McVean, G., Byrnes, J., Vukcevic, D., Palin, K., Su, Z., Howson, J.M.M., Auton, A., Myers, S., et al. (2012). Bayesian refinement of association signals for 14 loci in 3 common diseases. Nat. Genet. *44*, 1294–1301.

15. Benner, C., Havulinna, A.S., Järvelin, M.-R., Salomaa, V., Ripatti, S., and Pirinen, M. (2017). Prospects of Fine-Mapping Trait-Associated Genomic Regions by Using Summary Statistics from Genome-wide Association Studies. Am J Hum Genet *101*, 539–551.

16. de Leeuw, C.A., Mooij, J.M., Heskes, T., and Posthuma, D. (2015). MAGMA: generalized gene-set analysis of GWAS data. PLoS Comput. Biol. *11*, e1004219.

17. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P. (2015). The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell Syst *1*, 417–425.

18. Liu, Y., and Xie, J. (2020). Cauchy Combination Test: A Powerful Test With Analytic *p* -Value Calculation Under Arbitrary Dependency Structures. Journal of the American Statistical Association *115*, 393–402.

19. Liu, Y., Chen, S., Li, Z., Morrison, A.C., Boerwinkle, E., and Lin, X. (2019). ACAT: A Fast and Powerful p Value Combination Method for Rare-Variant Analysis in Sequencing Studies. Am. J. Hum. Genet. *104*, 410–421.

20. Turkmen, A., and Lin, S. (2017). Are rare variants really independent? Genet. Epidemiol. *41*, 363–371.

21. Talluri, R., and Shete, S. (2013). A linkage disequilibrium-based approach to selecting disease-associated rare variants. PLoS ONE *8*, e69226.

22. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W.J.H., Jansen, R., de Geus, E.J.C., Boomsma, D.I., Wright, F.A., et al. (2016). Integrative approaches for large-scale transcriptome-wide association studies. Nat. Genet. *48*, 245–252.

23. Mancuso, N., Freund, M.K., Johnson, R., Shi, H., Kichaev, G., Gusev, A., and Pasaniuc, B. (2019). Probabilistic fine-mapping of transcriptome-wide association studies. Nat. Genet. *51*, 675–682.

24. Zheng, J., Erzurumluoglu, A.M., Elsworth, B.L., Kemp, J.P., Howe, L., Haycock, P.C., Hemani, G., Tansey, K., Laurin, C., Early Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium, et al. (2017). LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. Bioinformatics *33*, 272–279.

25. O'Connor, L.J., and Price, A.L. (2018). Distinguishing genetic correlation from causation
across 52 diseases and complex traits. Nat. Genet. *50*, 1728–1734.

26. Reay, W.R., Kiltschewskij, D.J., Geaghan, M.P., Atkins, J.R., Carr, V.J., Green, M.J., and
Cairns, M.J. (2021). Genetic estimates of correlation and causality between blood-based
biomarkers and psychiatric disorders (Psychiatry and Clinical Psychology).

27. Reay, W.R., El Shair, S.I., Geaghan, M.P., Riveros, C., Holliday, E.G., McEvoy, M.A.,
Hancock, S., Peel, R., Scott, R.J., Attia, J.R., et al. (2021). Genetic association and causal
inference converge on hyperglycaemia as a modifiable factor to improve lung function. ELife
10, e63115.

28. Hemani, G., Zheng, J., Elsworth, B., Wade, K.H., Haberland, V., Baird, D., Laurin, C.,
Burgess, S., Bowden, J., Langdon, R., et al. (2018). The MR-Base platform supports
systematic causal inference across the human phenome. ELife *7*, e34408.

29. Willer, C.J., Schmidt, E.M., Sengupta, S., Peloso, G.M., Gustafsson, S., Kanoni, S.,
Ganna, A., Chen, J., Buchkovich, M.L., Mora, S., et al. (2013). Discovery and refinement of
loci associated with lipid levels. Nat Genet *45*, 1274–1283.

30. Bowden, J., Davey Smith, G., Haycock, P.C., and Burgess, S. (2016). Consistent
Estimation in Mendelian Randomization with Some Invalid Instruments Using a Weighted
Median Estimator. Genet. Epidemiol. *40*, 304–314.

31. Hartwig, F.P., Davey Smith, G., and Bowden, J. (2017). Robust inference in summary
data Mendelian randomization via the zero modal pleiotropy assumption. Int J Epidemiol *46*,
1985–1998.

32. Bowden, J., Davey Smith, G., and Burgess, S. (2015). Mendelian randomization with
invalid instruments: effect estimation and bias detection through Egger regression. Int J
Epidemiol *44*, 512–525.

33. Verbanck, M., Chen, C.-Y., Neale, B., and Do, R. (2018). Detection of widespread
horizontal pleiotropy in causal relationships inferred from Mendelian randomization between
complex traits and diseases. Nat. Genet. *50*, 693–698.

34. Thompson, S.G., and Sharp, S.J. (1999). Explaining heterogeneity in meta-analysis: a
comparison of methods. Stat Med *18*, 2693–2708.

35. Bowden, J., Del Greco M, F., Minelli, C., Davey Smith, G., Sheehan, N., and Thompson,
J. (2017). A framework for the investigation of pleiotropy in two-sample summary data
Mendelian randomization. Stat Med *36*, 1783–1802.

36. Reay, W.R., Shair, S.E., Geaghan, M.P., Riveros, C., Holiday, E.G., McEvoy, M.A.,
Hancock, S., Peel, R., Scott, R.J., Attia, J.R., et al. (2020). Genetically informed precision
drug repurposing for lung function and implications for respiratory infection (Respiratory
Medicine).

37. Reay, W.R., Atkins, J.R., Carr, V.J., Green, M.J., and Cairns, M.J. (2020).
Pharmacological enrichment of polygenic risk for precision medicine in complex disorders.
Sci Rep *10*, 879.

38. Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott,
P., Green, J., Landray, M., et al. (2015). UK biobank: an open access resource for identifying
the causes of a wide range of complex diseases of middle and old age. PLoS Med *12*,
e1001779.

39. Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A.,
Vukcevic, D., Delaneau, O., O'Connell, J., et al. (2018). The UK Biobank resource with deep
phenotyping and genomic data. Nature *562*, 203–209.

40. Choi, S.W., and O'Reilly, P.F. (2019). PRSice-2: Polygenic Risk Score software for
biobank-scale data. GigaScience *8*, giz082.

41. Mentzer, A.J., Brenner, N., Allen, N., Littlejohns, T.J., Chong, A.Y., Cortes, A., Almond,
R., Hill, M., Sheard, S., McVean, G., et al. (2019). Identification of host-pathogen-disease
relationships using a scalable Multiplex Serology platform in UK Biobank (Infectious
Diseases (except HIV/AIDS)).

42. Gu, Z., Eils, R., and Schlesner, M. (2016). Complex heatmaps reveal patterns and
correlations in multidimensional genomic data. Bioinformatics *32*, 2847–2849.

43. Linden, S.K., Sutton, P., Karlsson, N.G., Korolik, V., and McGuckin, M.A. (2008).
Mucins in the mucosal barrier to infection. Mucosal Immunol *1*, 183–197.

44. Chen, H.-H., Shaw, D.M., Petty, L.E., Graff, M., Bohlender, R.J., Polikowsky, H.G.,
Zhong, X., Kim, D., Buchanan, V.L., Preuss, M.H., et al. (2021). Host genetic effects in
pneumonia. The American Journal of Human Genetics *108*, 194–201.

45. Campos, A.I., Kho, P.F., Vazquez-Prada, K.X., García-Marín, L.M., Martin, N.G.,
Cuéllar-Partida, G., and Rentería, M.E. (2020). Genetic susceptibility to pneumonia: A
GWAS meta-analysis between UK Biobank and FinnGen (Respiratory Medicine).

46. Kim, K., Park, S., Park, S.Y., Kim, G., Park, S.M., Cho, J.-W., Kim, D.H., Park, Y.M.,
Koh, Y.W., Kim, H.R., et al. (2020). Single-cell transcriptome analysis reveals TOX as a
promoting factor for T cell exhaustion and a predictor for anti-PD-1 responses in human
cancer. Genome Med *12*, 22.

47. Sekine, T., Perez-Potti, A., Nguyen, S., Gorin, J.-B., Wu, V.H., Gostick, E., Llewellyn-
Lacey, S., Hammer, Q., Falck-Jones, S., Vangeti, S., et al. (2020). TOX is expressed by
exhausted and polyfunctional human effector memory CD8 [+] T cells. Sci. Immunol. *5*,
eaba7918.

48. Lin, C.-L., Liu, T.-C., Chung, C.-H., and Chien, W.-C. (2018). Risk of pneumonia in
patients with insomnia: A nationwide population-based retrospective cohort study. J Infect
Public Health *11*, 270–274.

49. Patel, S.R., Malhotra, A., Gao, X., Hu, F.B., Neuman, M.I., and Fawzi, W.W. (2012). A
prospective study of sleep duration and pneumonia risk in women. Sleep *35*, 97–101.

50. Burgess, S., and Thompson, S.G. (2015). Multivariable Mendelian randomization: the use
of pleiotropic genetic variants to estimate causal effects. Am J Epidemiol *181*, 251–260.

51. Parr, J.B. (2020). Time to Reassess Tocilizumab's Role in COVID-19 Pneumonia. JAMA
Intern Med.

52. Folkersen, L., Fauman, E., Sabater-Lleal, M., Strawbridge, R.J., Frånberg, M., Sennblad,
B., Baldassarre, D., Veglia, F., Humphries, S.E., Rauramaa, R., et al. (2017). Mapping of 79
loci for 83 plasma protein biomarkers in cardiovascular disease. PLoS Genet *13*, e1006706.

53. Ferreira, R.C., Freitag, D.F., Cutler, A.J., Howson, J.M.M., Rainbow, D.B., Smyth, D.J.,
Kaptoge, S., Clarke, P., Boreham, C., Coulson, R.M., et al. (2013). Functional IL6R 358Ala
allele impairs classical IL-6 receptor signaling and influences risk of diverse inflammatory
diseases. PLoS Genet *9*, e1003444.

54. Davis, W.A., Knuiman, M., Kendall, P., Grange, V., Davis, T.M.E., and Fremantle
Diabetes Study (2004). Glycemic exposure is associated with reduced pulmonary function in
type 2 diabetes: the Fremantle Diabetes Study. Diabetes Care *27*, 752–757.

55. King, J.B., West, M.B., Cook, P.F., and Hanigan, M.H. (2009). A novel, species-specific
class of uncompetitive inhibitors of gamma-glutamyl transpeptidase. J Biol Chem *284*, 9059–
9065.

56. Tuzova, M., Jean, J.-C., Hughey, R.P., Brown, L.A.S., Cruikshank, W.W., Hiratake, J.,
and Joyce-Brady, M. (2014). Inhibiting lung lining fluid glutathione metabolism with
GGsTop as a novel treatment for asthma. Front Pharmacol *5*, 179.

57. Smailhodzic, D., van Asten, F., Blom, A.M., Mohlin, F.C., den Hollander, A.I., van de
Ven, J.P.H., van Huet, R.A.C., Groenewoud, J.M.M., Tian, Y., Berendschot, T.T.J.M., et al.
(2014). Zinc Supplementation Inhibits Complement Activation in Age-Related Macular
Degeneration. PLoS ONE *9*, e112682.

58. Nan, R., Tetchner, S., Rodriguez, E., Pao, P.-J., Gor, J., Lengyel, I., and Perkins, S.J.
(2013). Zinc-induced self-association of complement C3b and Factor H: implications for
inflammation and age-related macular degeneration. J Biol Chem *288*, 19197–19210.

59. Bustamante-Marin, X.M., and Ostrowski, L.E. (2017). Cilia and Mucociliary Clearance.
Cold Spring Harb Perspect Biol *9*,.

60. Hewson, C.A., Haas, J.J., Bartlett, N.W., Message, S.D., Laza-Stanca, V., Kebadze, T.,
Caramori, G., Zhu, J., Edbrooke, M.R., Stanciu, L.A., et al. (2010). Rhinovirus induces
MUC5AC in a human infection model and in vitro via NF-κB and EGFR pathways. Eur
Respir J *36*, 1425–1435.

61. Barbier, D., Garcia-Verdugo, I., Pothlichet, J., Khazen, R., Descamps, D., Rousseau, K.,
Thornton, D., Si-Tahar, M., Touqui, L., Chignard, M., et al. (2012). Influenza A induces the
major secreted airway mucin MUC5AC in a protease-EGFR-extracellular regulated kinase-
Sp1-dependent pathway. Am J Respir Cell Mol Biol *47*, 149–157.

62. Singanayagam, A., Footitt, J., Kasdorf, B.T., Marczynski, M., Cross, M.T., Finney, L.J.,
Trujillo Torralbo, M.-B., Calderazzo, M., Zhu, J., Aniscenko, J., et al. (2019). MUC5AC
drives COPD exacerbation severity through amplification of virus-induced airway
inflammation (Immunology).

63. Lu, W., Liu, X., Wang, T., Liu, F., Zhu, A., Lin, Y., Luo, J., Ye, F., He, J., Zhao, J., et al.
(2020). Elevated MUC1 and MUC5AC mucin protein levels in airway mucus of critical ill
COVID-19 patients. J Med Virol.

64. He, J., Cai, S., Feng, H., Cai, B., Lin, L., Mai, Y., Fan, Y., Zhu, A., Huang, H., Shi, J., et
al. (2020). Single-cell analysis reveals bronchoalveolar epithelial dysfunction in COVID-19
patients. Protein Cell *11*, 680–687.

65. (2009). Essentials of Glycobiology (Cold Spring Harbor (NY): Cold Spring Harbor
Laboratory Press).

66. Wang, S.-S., del Solar, V., Yu, X., Antonopoulos, A., Friedman, A.E., Agarwal, K., Garg,
M., Ahmed, S.M., Addya, A., Nasirikenari, M., et al. (2020). Efficient Inhibition of O-glycan
biosynthesis using the hexosamine analog Ac_5 GalNTGc (Biochemistry).

67. Bae, S.S., Chang, L.C., Merkin, S.S., Elashoff, D., Ishigami, J., Matsushita, K., and
Charles-Schoeman, C. (2020). Major Lipids and Future Risk of Pneumonia: 20-Year
Observation of the Atherosclerosis Risk in Communities (ARIC) Study Cohort. The
American Journal of Medicine S0002934320306987.

68. Chien, Y.-F., Chen, C.-Y., Hsu, C.-L., Chen, K.-Y., and Yu, C.-J. (2015). Decreased
serum level of lipoprotein cholesterol is a poor prognostic factor for patients with severe
community-acquired pneumonia that required intensive care unit admission. J Crit Care *30*,
506–510.

69. Antcliffe, D., Jiménez, B., Veselkov, K., Holmes, E., and Gordon, A.C. (2017).
Metabolic Profiling in Patients with Pneumonia on Intensive Care. EBioMedicine *18*, 244–
253.

70. Nelson, M.R., Tipney, H., Painter, J.L., Shen, J., Nicoletti, P., Shen, Y., Floratos, A.,
Sham, P.C., Li, M.J., Wang, J., et al. (2015). The support of human genetic evidence for
approved drug indications. Nat Genet *47*, 856–860.

71. Asimit, J.L., Hatzikotoulas, K., McCarthy, M., Morris, A.P., and Zeggini, E. (2016).
Trans-ethnic study design approaches for fine-mapping. Eur J Hum Genet *24*, 1330–1336.

1236

1237

1238