1   **Colder and drier winter conditions are associated with greater SARS-CoV-2 transmission: a**
2   **regional study of the first epidemic wave in north-west hemisphere countries.**

4   Version 1, 26 January 2021

Jordi Landier[1*], Juliette Paireau[2,3], Stanislas Rebaudet[1,4], Eva Legendre[1], Laurent Lehot[1], Arnaud
Fontanet[5,6], Simon Cauchemez[2], Jean Gaudart[7]

[1] IRD, Aix Marseille Univ, INSERM, SESSTIM, Marseille, France.

[2] Mathematical Modelling of Infectious Diseases Unit, Institut Pasteur, UMR2000, CNRS, Paris, France.

[3] Santé publique France, French National Public Health Agency, Saint Maurice, France

[4] Hôpital Européen Marseille, France.

[5] Emerging Infectious Diseases Unit, Institut Pasteur, Paris, France.

[6] PACRI Unit, Conservatoire National des Arts et Métiers, Paris, France.

[7] Aix Marseille Univ, APHM, INSERM, IRD, SESSTIM, Hop Timone, BioSTIC, Marseille, France.

* corresponding author, jordi.landier@ird.fr

Abstract

Higher transmissibility of SARS-CoV-2 in cold and dry weather conditions has been hypothesized since
the onset of the COVID-19 pandemic but the level of epidemiological evidence remains low.

During the first wave of the pandemic for Spain, Italy, France, Portugal, Canada and USA presented an
early spread, a heavy COVID-19 burden, and low initial public health response until lockdowns. We
used regional death counts as a proxy for infections while diagnostic tests remained limited and
calculated a basic reproduction number (R0) in 63 regions. After adjusting for population density, early
spread of the epidemic, and aged population, temperature and humidity were negatively associated
to SARS-CoV-2 transmissibility. A -1g/m3 lower mean absolute humidity was associated with a 0.15-
unit higher R0. Below 10°C, a 1°C-higher temperature was associated with a 0.16-unit lower R0.

Our results confirm a strong dependency of SARS-CoV-2 transmissibility to weather conditions in the
absence of control measures. The initiation of the second wave in north-west hemisphere countries
was likely triggered by the transition from summer- to winter-like conditions. High levels of restrictions
to social activities should be maintained until spring to avoid or limit a third wave.

Introduction

The spread of SARS-CoV-2 in February-March 2020 caught the majority of European and North American countries unprepared. The spread of the virus was largely uncontrolled until movement restriction and social distancing policies were put in place at national or regional level with various degrees of intensity [1]. Most regions experienced a peak in hospital admissions, approximately 2 weeks after lockdown was imposed, corresponding to infections acquired around the date of lockdown [2].

The spread of this first wave was largely heterogeneous between countries. Various determinants were proposed to explain the differential spread of the virus. Asian countries (Japan, South Korea, Vietnam and Thailand) and isolated countries (Australia, New Zealand, Iceland) experienced only limited transmission and stood out for high levels of preparedness and efficient response strategies. Even within South West European and North American countries where no effective response was deployed before lockdowns, SARS-CoV-2 virus spread heterogeneously at the regional level, as observed from hospitalization, death counts, and confirmed by serological surveys [3, 4].

Epidemic spread is characterized by the basic reproduction number, or R0. R0 expresses the number of secondary cases resulting from a given case in the context of a naive population with uniform probability of contact.

R0 depends on individual susceptibility to infection when in contact with an infective individual, and on the probability of an infectious contact. Environmental parameters affect R0: population density increases the probability of contacts between individuals and weather conditions may affect the survival of the virus or the individual susceptibility to an infection. Most known respiratory viruses spread during the cold season in the temperate Northern hemisphere [5]. Weather conditions in winter can also affect individual susceptibility to infection through irritation of the nasal mucosa, but also influence the behaviour of individuals towards conditions prone to transmission (living or gathering in closed, heated spaces with a dry atmosphere) [5]. In addition, temperature, humidity, and UV, might directly affect the virus survival and modify infectiousness [6, 7]. Individual preventive behaviours (masks), collective strategies reducing mobility and contacts (lockdowns) or limiting the duration of the infectious period (detection and isolation) modify the number of secondary cases and the effective reproduction number can be calculated, which accounts for these alterations in the "natural" history of transmission.

In spite of a large number of studies, the evidence regarding the link between weather conditions and SARS-CoV-2 transmission remains limited. At date of 15 May 2020, a systematic review retained 61 studies analysing the relationship between COVID-19 epidemic and environmental factors [8]. Methodological issues included the lack of controlling for confounding factors such as population density [8]. Inappropriate epidemiological and statistical methods were also pointed out [8, 9]. Comparison between countries with different counter epidemic responses, testing strategy, or delayed onset of the epidemic might also have led to inconsistent results [8]. An earlier review retaining 17 studies highlighted the risk of bias and the low level of available evidence [10]. Between 15 May 2020 and 15 December 2020, our systematic search identified 82 research articles, of which only 15 (18%) analysed the growth rate or the reproduction number of SARS-CoV-2 (Appendix p2). Only four studies of climate and reproduction number included adjustments for confounding factors, and three of four included relevant covariates of population mobility, population density, and took into account interventions when necessary [11–13]. Of these, two were conducted over small geographical units: one study of 212 US counties identified a negative relationship between temperature increase and

SARS-CoV-2 growth, while the other study including 28 Japanese prefectures identified a positive relationship [11, 12]. The third study was at the global scale for 203 states (USA, Canada, Australia and China) or countries and identified a negative association between UV light exposure and SARS-CoV-2 growth. Temperature was negatively associated with growth only after adjustment for UV light exposure [13]. All three studies identified a positive association between population density and epidemic growth.

Overall, multiple studies described a negative relationship between temperature and COVID-19 outcomes but the majority were unadjusted ecological correlation studies with strong risk of bias bringing low quality evidence [8], and evidence from multivariable growth studies remains ambiguous.

A modelling study defined the range of possible dependency between SARS-CoV-2 transmission and absolute humidity, based on two known coronaviruses and influenza [14], and recent models still do not include specific SARS-CoV-2 data [15]. More precise estimations of the effect of meteorological conditions on the spread of SARS-CoV-2 are required to better anticipate and inform policies regarding seasonal adjustments [16].

The objective of this study was to evaluate the contribution of weather parameters in the transmission of SARS-CoV-2, by analysing their effect on SARS-CoV-2 basic reproduction number in a context of low public health response during the early phase of the first wave in 6 north-western hemisphere countries.

Material and methods

Study design

In order to compare the drivers of the epidemic dynamics between regions accurately, we calculated the basic reproduction number (R0) of the virus for each region affected by the first wave of COVID-19 epidemic in six countries, using the dynamics of the daily death counts. These six countries were located in the western part of the northern hemisphere, approximately between 25 and 50° of latitude (Figure S2). These countries experienced winter conditions and underwent significant SARS-CoV-2 transmission in a context of low public health response during the first wave of the epidemic.

This analysis was conducted at the first administrative subdivision country, here referred to as "region". Regions corresponded to States in USA and Canada, autonomous communities in Spain, and regions in Italy (regioni), Portugal (região) and France (regions).

Confirmed case counts were insufficiently reliable due to overall lack of tests and different testing strategies between countries, between regions of the same country and between periods for the same region. Hospitalization counts were not available consistently at regional level across countries. Overall, COVID-19 deaths were preferred as they were less likely to have different definitions within the same region or country and to undergo significant changes over the study period. R0 is an indicator of the speed of progression of the outbreak, and was therefore less likely to be biased compared to indicators based on cumulative counts or cumulative incidence rates. Estimating R0 values based on deaths counts relies on the minimal assumption that the infection-fatality rate was constant over the study period (~1 month) for a given region, which defined a proportional relationship between infections and deaths.

Study period

Daily reporting of COVID-19 deaths was not initiated immediately after the first recorded deaths, and often started in a staggered manner in the different regions of a country. Analyses started on the date when 10 cumulative deaths were reached in a given region to account for these limitations of available data and to limit stochasticity in the occurrence of the first deaths.

This study aimed at estimating SARS-CoV-2 R0 prior to implementation of major interventions in different regions. It was necessary to avoid the influence of the lockdown measures on the transmission. Reports indicated that symptoms occurred in median 5 days after infection (IQR=2-7), and hospitalization occurred in median 7 days after the onset of symptoms (IQR=3-9) [17, 18]. Reports of delay between hospitalization and death ranged from 5 days in New York to 12 days in Italy for patients admitted in ICU [19, 20]. In Spain and Italy, a median of 11 days (IQR=7-17), respectively 12 days (IQR 7-20), were reported between onset of symptoms and death [21, 22], while an early study in China indicated a median of 18 days from onset of symptoms to death [23]. These figures lead to an overall estimation that deaths occurred in median 17-18 days after infection, with a broad range of variations. In France, mortality peaked at week 14 when lockdown occurred at the beginning of week 12, i.e. for infections likely acquired during week 11. As a result, we initially assumed a 3-week delay between infection and death and excluded all data beyond 28 days from lockdown date to limit this study to deaths caused by pre-lockdown infections. We also conducted a more restrictive sensitivity analysis excluding all data beyond 18 days after lockdown date, which corresponded to the estimated median delay between infection and deaths.

Inclusion criteria

Regions where the smoothed daily death count did not rise above 5 deaths/day during the study period were not included in the analysis. Regions where the smoothed daily death count did not rise above 10 deaths/day during the linear growth phase were excluded. This limited the study to regions having displayed a clear exponential growth phase.

One Italian region was excluded because there was no weather station data available except for 2 stations >2000m altitude.

Data

Deaths from COVID-19

Regional level data on COVID-19 deaths were retrieved from data shared by national health ministries and/or public health agencies of Spain, Italy, Portugal, France, United States of America, and Canada (Table S1). Deaths were reported as daily new death counts or as a cumulative number.

Population and geographical data

Population structure by one- or five-year age groups, sex and region was retrieved from open access data shared by national institutes or administrations responsible for national statistics or demography (Table S1). The percentage of the region population aged $\geq$70 or $\geq$80 years was calculated in order to adjust for differences in way of life (e.g. rural regions have older population than metropolitan areas in Europe). The percentage of elderly inhabitants also represented an adjustment in case of a

differential spread of the epidemic among the elderly population (which would lead to a non-
proportional increase in deaths compared to cases.

Region shapefiles by country were obtained from national geographical authorities, open source
datasets or as provided in the coronavirus open data packages proposed by several national health
agencies (Table S1). The region surface was either obtained from the geographical layer (land surface
area), or calculated from the polygon extent. We estimated the percentage of each region surface with
$\geq$5 inhabitants per km^2 based on WorldPop 2020 raster dataset [24]. Population density was estimated
as the population in the region divided by the surface after excluding areas <5 inhabitants/km^2 in order
to limit the underestimation of population density when high heterogeneity existed between urban
centers and sparsely populated territories within the same region (deserts, mountains, polar regions).

Weather/climate

Weather station reports since 1 January 2020 were obtained from US National Oceanic and
Atmospheric Administration (NOAA) through the R-package {worldmet}. We extracted hourly
temperature, relative humidity (RH), dew point (DP), precipitation and windspeed observations for
each station. Hourly absolute humidity (AH, in g/m^3) was calculated from RH and temperature based
on the Clausius-Clapeyron formula [25]. Daily minimum, maximum and mean values were calculated
for temperature, AH, RH, dew point, as well as cumulative sum of precipitation and mean wind speed.
Days with $\geq$5 missing hourly record (no observation of any parameter) were excluded.

Each region was attributed stations based on geographic location. All available observations in weather
stations of the region contributed to the regional daily average. Weather stations can be located in
mountains or inhabited locations where they record weather conditions that differ strongly from
actual populated areas of the same region. To avoid this bias, observations were assigned population-
based weights: we estimated the population located within 10 km of each weather station using
WorldPop 2020 data; and for each day and each region, we calculated the total population within 10km
of any station reporting data for that day. Daily station observations were weighted in proportion of
the population around each station relative to the total population. Stations located within 10-km of
each other were included in a single buffer and each station was assigned equal weight within the
buffer.

For each weather parameter, the regional summary value was calculated over the assumed
transmission period. The assumed transmission period corresponded to the duration of the R0
calculation period, lagged by 3 weeks. Sensitivity analyses included lags ranging from 2 to 8 weeks from
R0 calculation period, i.e. -1 to +5 weeks from assumed transmission period. The longer lags were likely
irrelevant for actual transmission but aimed at identifying climate rather than weather aspects.

Using this approach, the weather parameters averaged over the assumed transmission period and over
each region included: minimum, mean and maximum average values for temperature, relative
humidity, absolute humidity, dew point, average cumulative rainfall per day and average wind speed.

Date of lockdown definition

Lockdown date was defined using Google mobility data as the date when a decrease >25% in workplace
localization was reported and sustained over 3 days in the region [26]. All references to a « date of
lockdown » hereafter refer to this definition. This definition matched national lockdowns in European

countries. This simplification was necessary due to the heterogeneity in social distancing measures
taken at regional (state) level in the USA and Canada. The objective was to exclude periods where
transmission would start to slow down due to these measures.

Distance to first region with 10 cumulative deaths

For each of the four European countries considered, the first region with 10 cumulative deaths was
defined. In Portugal, two regions reached $\geq$10 deaths on the same day, and the region with the highest
count (14 versus 12) was selected. Within each country, the euclidean distance in kilometers between
the main city in each region and the main city in the first region above 10 cumulative deaths was
calculated to reflect spatial autocorrelation due to proximity in the spread of the epidemic. In USA and
Canada, distance to the first region above 10 cumulative deaths were calculated separately for East
Coast and West Coast, using the limit between Central and Mountain time zones. This was necessary
due to the early start of the epidemic in the state of Washington, which reached 10 deaths by 2 March
2020, 16 days before the next state (New York on 18 March).

Statistical methods

Statistical analyses were performed using R version 4.0. Maps were produced using ArcGIS 10.7.1.

Daily death counts were smoothed using a 5-day moving average filter to account for irregularities in
data transmission and publication.

For each region, the exponential growth period was estimated using a log(deaths)=f(time)
representation and r (the exponential growth rate) was extracted as the coefficient of a Poisson
regression. R0 was calculated for each region using the generation time method assuming a gamma
distribution with parameters 7 and 5.2 for SARS-CoV-2 generation time [27, 28]. In order to improve
the adjustment of the regression, the start and end dates of calculation period were allowed to shift
by up to 2 days (+/-1 day or +1/+2 days for start date if calculation period began at the date of 10
cumulative deaths) using the built-in function sensitivity.analysis() of the package {R0} [29].

A directed acyclic graph (DAG) was constructed using Dagitty v3.0 web-based application
(http://www.dagitty.net/dags.html) in order to visualise the relationships between R0 and the
explanatory covariates (Figure S1). Dependence and independence assumptions were verified using
Spearman correlation coefficient.

Relative humidity values are temperature-dependent and absolute humidity or dew point
temperatures are also strongly correlated to temperature [30]. The different weather covariates
observed at the same 3-week lag from R0 calculation period were included separately in the models.
A generalized additive mixed model (gamm) regression was used to evaluate the effects of climate,
population and other determinants on the value of R0, using a Gaussian distribution and the identity
link function (package {mgcv}). A country-level random effect was included to account for within-
country correlations. Canada presented with only 2 regions and was grouped with USA for random
effect, while the single region included from Portugal was grouped with Spain. Univariate analyses
were conducted assuming linear and non-linear effects, using B-splines to model non-linear effect of
covariates. Models were compared using the percentage of deviance explained and Akaike's
Information Criterion.

We verified presence/absence of spatial autocorrelation in R0 values and in the final model residuals
by calculating Moran's I.

Sensitivity analyses

First, we assessed the importance of the 3-week delay for weather variables (corresponding to the
delay between transmission period and R0 calculation period based on death count exponential
growth). For this, we tested a variety of lags from -1 to 5 weeks from that estimated transmission
period (*i.e.* 2 to 8 weeks from the R0 calculation period) and included them as linear explanatory
variables in the univariable hierarchical model or as non-linear explanatory variables in the
multivariable hierarchical generalised additive model.

Second, we assessed the effect of the 28-day window to define the exponential growth period by using
a narrower window ending 18 days after date of lockdown. We recalculated R0 for regions where the
linear growth period retained for the main analysis extended beyond the 18-day limit, and followed
the same plan as the main analysis.

Third, we assessed possible continent specific effects by fitting continent-specific splines for weather
variables in the final multivariate models.

Results

Region selection

The six countries (United States, Canada, Spain, Italy, France and Portugal) included 128 regions/states.
Overseas regions (n=11) and regions which had experienced <10 cumulative deaths 28 days after
lockdown (n=15) were not included (Figure 1). Likewise, regions with a maximal daily mortality <5
deaths (n=19) within 28 days after lockdown were not included. Overall, 83 regions were assessed for
exponential growth period, and R0 was calculated for 64 regions with sufficient exponential growth
(Figure S2 in Appendix): 24 regions in the USA, 2 regions in Canada, 11 regions in France, 12 regions in
Spain, 13 regions in Italy and 1 region in Portugal.

Data description

For regions analysed, the maximal daily death counts was 39 deaths/day in median (interquartile range
(IQR)=18-83, max=779) and occurred in median 25 days (IQR=19-43) after the date of lockdown. Of
note, larger reductions in human mobility patterns after lockdown as assessed from google mobility
led to shorter delay (Spearman correlation coefficient = 0.63, p<0.0001, Figure S3). This delay was
reduced to 19 days in median (IQR=17-27) for South European countries with nationwide lockdowns
and strong mobility reductions (>60% in average, Figure S3).

R0 were estimated over a median exponential growth period of 11 days (IQR=9-14, range=5-19). In
median, R0 estimation period started 5 days (IQR=0-8.5) and ended 16 days after the date of lockdown
(IQR=11-21, range 1-27). Figure S4 presents details calculation periods.

The median R0 value was 2.58 (IQR=2.08-2.66). R0 estimates were lowest (<1.5) in two regions in
France and one in Spain, USA and Italy (respectively in Centre-Val de Loire, Nouvelle Aquitaine, La
Rioja, Alabama and Abruzzo). R0 were highest (>4.0) in New York (USA), Lombardia and Piemonte

(Italy), Castilla-La Mancha (Spain) and Ontario (Canada)(Figure 3A and B). Spain and Canada had overall
higher R0 values compared to Italy, France and USA (Figure 4A). R0 values exhibited significant spatial
autocorrelation (Moran's I=0.20, p=0.0087).

Covariates were heterogeneous between countries (Figure 3 and 4). Population density ranged
between 82 and 1056 inhabitants/km^2 (median=271, Figure 3C and D) and was overall higher in Italy
(Figure 4B). Mean absolute humidity during the estimated transmission period was 4.98 in median
(IQR=4.29-5.99, range=2.26-11.32, Figure 3E and F) corresponding to a median dew point of 3.7°C
(IQR=0.9-6.3, range=-7.7 – +15.3). Mean temperature was 9.8°C in median (IQR=7.1-11.5, range=-2.0-
19.9). Distance to the first region with 10 cumulative deaths was 406km in median (IQR=228-794) and
much larger in the USA compared to European countries (Figure 4F).

Factors associated with R0

Each variable was included in a univariate model assuming linear (Figure 5, Table S2) or non-linear
(spline smoothing, Table S3) relationships with R0. We identified significant relationships between R0
value and temperature, absolute humidity or dew point, average daily rainfall, but not with mean wind
speed over the transmission period.

Multivariable models were constructed according to the DAG. After adjustment for distance to the
nearest affected region, percentage of population over 80 and population density, there was a
consistent relationship between increasing temperature, AH or dew point temperature, and
decreasing R0. No relationship was found with average daily rainfall.

Mean temperature led to the model with the largest deviance explained compared to models including
minimum or maximum temperature, or any AH or dew point temperature (Table 1). In this model, a
10-fold (+1 log10 unit) increase in population density was associated with a +0.67 R0 unit increase
(95%CI=0.05-1.28). The proportion of population aged 80 years and older was not significantly
associated with R0. The relationship between mean temperature and R0 was not linear. A strong,
nearly linear drop of approximately 1.0 R0 unit was observed between 2.5 and 12°C, and a plateau
beyond 12°C (Figure 6 and 7). The residuals from this model did not exhibit significant spatial
autocorrelation (Moran's I=0.054, p=0.215).

Mean AH and mean dew point values exhibited similar profiles with increasing values associate with
decreasing R0 values (Table 1, Figure 6). In spite of the narrower range of values, it seemed that the
relationship between R0 and AH or dew point temperature did not reach a plateau (Figure 7). Assuming
a linear relationship, a 1 g/m3 higher absolute humidity translated in a 0.15 unit lower regional R0,
respectively a 1°C higher dew point temperature translated in a 0.08 unit lower R0 (Table S5).

Sensitivity analyses

First, using lagged weather summary values as linear univariate predictors, our statistical model found
similar relationships between R0 and temperature variables, but 0- , 1- and 5-week lags led to the
strongest effect for mean AH and mean DP (Figure S5). Using lagged weather summary values as non-
linear predictors in the multivariate model, the shapes of the relationships between temperature, AH
and DP remained similar, with a nearly linear drop reaching a plateau at values corresponding to milder
winter weather/climate (Figure S6). Overall, correlations were strong between weather summary
observations at the different lags (Figure S7). Correlations of lagged temperature and humidity
observations was highest between -1, 0, 1 and 5-week lagged observations compared to other lags.

Second, when limiting the R0 calculation window to 18-day after date of lockdown, 10 additional regions were excluded and analysis was conducted on 53 regions. After adjusting for population density, population over 80 and distance to the first region affected, the non-linear negative relationship between R0 and temperature remained unchanged, reaching a plateau around 10°C (spline p-value=0.0475). The shape of the relationship between AH, respectively DP, and R0 remained similar but did not reach statistical significance (p-value=0.19). Later AH or DP summary values, corresponding to 1 week later than the estimated transmission period (i.e. 2-week lag from R0 calculation period), restored the negative relationship with R0 (p=0.0661, respectively p=0.0667).

Third, fitting continent-specific splines for weather covariates, also resulted in low statistical power: only 37 regions in South Europe and 26 in North America. After adjusting for population density, population over 80 and distance to the first region affected, humidity variables (AH or dew point) retained a negative association with R0 (continent-specific spline p-values were 0.0915 (North America) and 0.0988 (South Europe) for AH, respectively 0.0506 and 0.1078 for dew point temperature). The relationship between R0 and mean temperature was markedly different between North America (negative association, p=0.005) and South Europe (p=0.17).

Discussion

Summary

In this study, we analysed SARS-CoV-2 propagation parameter R0 during the first wave of the pandemic in 63 regions of 6 north western countries. We showed that R0 values were influenced by population density (+0.6 for a 10-fold increase in density), by proximity with the first epidemic focus of the country or coast for USA (-0.3 for a 10-fold increase in distance to the first region to record 10 COVID-19 deaths), and by weather or climate conditions. For regions with mean temperatures below 10°C during the transmission period, a linear association was observed with R0 values: a difference between regions of +1°C was associated with a 0.16-unit decrease in R0. A difference in mean absolute humidity of +1 g/m^3 was associated with a 0.15-unit decrease in R0. Similar results were obtained with dew point temperatures, with a difference of 1°C was associated with a 0.08-unit decrease in R0 (Table S5). After adjusting for major confounders and spatial autocorrelation, our results indicate that weather conditions brought a significant contribution to drive the magnitude of the first wave, even if it was limited by an initial heterogeneous spread of the virus, which protected regions located furthest away from the first foci.

Strength and limits

This study relied on a regional scale analysis and accounted for different dynamics within the same country. The overall epidemic wave at country level was actually the sum of diverse dynamics, as is obvious for large countries but also true for Spain, Italy or France, where regions were heterogeneously affected, as confirmed by serological studies [3, 31]. Likewise, weather or climate heterogeneity between regions of a given country was large. By analyzing distinct spatial units with heterogeneous population density and weather, we were able to assess the effect of parameters that may be otherwise confounded by country-level parameters such as response strategy, timing of the analysis period compared to the progression of the epidemic, but also age structure of the population.

The 6 countries were selected due to their homogeneous location in the northern hemisphere between 25 and 50 degrees of latitude and the low efficacy of their counter epidemic responses until

a lockdown was decreed. We are therefore as close as possible of conditions allowing R0 estimation.
This 4-parameter hierarchical generalized additive model provides an explanation of up to 45% of R0
variability across 63 affected regions of the northern hemisphere, thus providing useful insights on the
drivers of the first wave, and allowing to estimate the contribution of population density and weather
conditions for the next waves, when the effect of local introduction is no longer relevant.

We acknowledge several limits. The first one is the necessity to rely on death counts to estimate R0
during this wave due to the lack of information on actual infections from limited testing and the lack
of consistency between countries for hospitalization counts. This assumption is similar to usual
assumptions for R0 estimation based on diagnosed cases, since true number of infections remains
unknown and detection occurs at variable delays after infection but it could have led to an
underestimation of the R0. Indeed, while the average delay between infection and death was 3 weeks,
deaths occurring from infections contracted the same day may in fact occur over a larger window
extending several weeks beyond this delay.

The second one is the use of a single summary weather observation over the assumed transmission
period, which may fail to express the dynamical aspects of weather. The choice of fixed, 1-week
increments to study the effect variations in the weather summary window may be too coarse to
capture effects for narrow exponential growth periods. The sensitivity analysis showed a slight increase
in effect when considering weather summary values calculated 1 week earlier than the estimated
transmission period, but the 18-day sensitivity analysis showed a stronger negative relationship
between R0 and AH summary values corresponding to 1-week later than the estimated transmission
period. The role of the weather conditions might be more important during the beginning of the
exponential growth period, until a sufficiently large number of persons becomes infected and
parameters such as population density become increasingly important. This study could also only
evaluate the effect of a limited range of weather conditions on SARS-CoV-2 transmission, since the
number of observations with a mean temperature below 2.5°C or above 15°C was low. This prevents
from analyzing the effect of higher temperatures such as during autumn. This restriction was however
necessary to achieve a minimal homogeneity of the studied regions in terms of their exposure and
response to the pandemic. Finally, this analysis includes only 63 regions and may lack statistical power.
The necessity to ensure that the epidemic growth of deaths was sufficient led to the exclusion of
regions that may be affected, but not enough for daily deaths counts to reach 10 deaths/day.

Finally, this study showed an association between SARS-CoV-2 R0 and temperature/absolute humidity,
but due to the strong correlation between absolute humidity and temperature in the seasonal
conditions analysed here, it is difficult to determine which parameter is more important and they could
not be analysed in combination [30]. The continent-specific analysis suggests that the relationship
between absolute humidity and R0 was more stable than that of temperature, which was strong in the
US but less so in South Europe. We could not conclude whether the relationship results from a direct
role of weather on individual susceptibility to viral invasion (*e.g.* dry nasal mucosa from indoors heating
and outdoors cold) or on viral persistence/survival, or from an indirect role of climate on human
behaviors (*e.g.* regions with cold winter favor more indoors living conditions and lead to bigger
infection opportunities).

Conclusion

Our study shows an important dependency of SARS-CoV-2 transmission to weather/climate, with a
0.16-unit increase in R0 for a 1°C difference in mean regional temperature below 10°C, or a 0.15-unit
increase for a -1g/m3 decrease in absolute humidity. Northern hemisphere countries experienced a
second wave of SARS-CoV-2 infections during autumnal transition from summer to winter, while still
actively maintaining control strategies. Public health strategies need to account for a further increase

in transmissibility when/where cold and dry winter conditions are reached. In the regions of the north-west hemisphere, lifting restrictions on social activities during the winter would likely lead to faster increases than observed during the second wave, when autumnal conditions were prevalent.

Authors' contribution

JL, JG, SC and AF designed the study. JL conducted the analysis with methodological contributions of JP, EL, and LL. JL JP EL LL AF SC JG contributed to interpreting the results and editing the manuscript.

Acknowledgements

The authors would like to thank Laurax Simgiane Ferbliegas from Hab' lab Marseille and L. Jae for help and support in retrieving multi-country regional data.

References

1. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. Nature. 2020;584:257–61.

2. Cauchemez S, Kiem CT, Paireau J, Rolland P, Fontanet A. Lockdown impact on COVID-19 epidemics in regions across metropolitan France. The Lancet. 2020;396:1068–9.

3. Pollán M, Pérez-Gómez B, Pastor-Barriuso R, Oteo J, Hernán MA, Pérez-Olmeda M, et al. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. Lancet. 2020;396:535–44.

4. Lai C, Wang J, Hsueh P. Population-based seroprevalence surveys of anti-SARS-CoV-2 antibody : An up-to-date review. Int J Infect Dis. 2020;101:314–22.

5. Moriyama M, Hugentobler WJ, Iwasaki A. Seasonality of Respiratory Viral Infections. Annu Rev Virol. 2020;7:83–101.

6. Matson MJ, Yinda CK, Seifert SN, Bushmaker T, Fischer RJ, Doremalen N Van, et al. Effect of Environmental Conditions on SARS-CoV-2 Stability in Human Nasal Mucus and Sputum. Emerg Infect Dis. 2020;26:9–11.

7. Morris DH, Claude Yinda K, Gamble A, Rossine FW, Bushmaker T, Fischer RJ, et al. Mechanistic theory predicts the effects of temperature and humidity on inactivation of SARS-CoV-2 and 2 other enveloped viruses. bioRxiv. 2020.

8. Briz-Redón Á, Serrano-Aroca Á. The effect of climate on the spread of the COVID-19 pandemic: A review of findings, and statistical and modelling techniques. Prog Phys Geogr. 2020;44:591–604.

9. Gillibert A, Jaureguiberry S, Hansmann Y, Argemi X, Landier J, Caumes E, et al. Comment on A. annua and A. afra infusions vs. Artesunate-amodiaquine (ASAQ) in treating Plasmodium falciparum malaria in a large scale, double blind, randomized clinical trial. Phytomedicine. 2019; in press:152981.

10. Mecenas P, da Rosa Moreira Bastos RT, Rosário Vallinoto AC, Normando D. Effects of temperature and humidity on the spread of COVID-19: A systematic review. PLoS One. 2020;15 9 September:1–21.

11. Azuma K, Kagi N, Kim H, Hayashi M. Impact of climate and ambient air pollution on the epidemic growth during COVID-19 outbreak in Japan. Environ Res. 2020;190:110042.

12. Rubin D, Huang J, Fisher BT, Gasparrini A, Tam V, Song L, et al. Association of Social Distancing, Population Density, and Temperature With the Instantaneous Reproduction Number of SARS-CoV-2 in Counties Across the United States. JAMA Netw open. 2020;3:e2016099.

13. Merow C, Urban MC. Seasonality and uncertainty in global COVID-19 growth rates. Proc Natl Acad Sci U S A. 2020;117:27456–64.

14. Baker RE, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. Susceptible supply limits the role of climate in the early SARS-CoV-2 pandemic. Science (80-). 2020;369:315–9.

15. Saad-roy CM, Wagner CE, Baker RE, Morris SE, Farrar J, Graham AL, et al. Immune life history, vaccination, and the dynamics of SARS-CoV-2 over the next 5 years. Science (80-). 2020;818 November:811–8.

16. Zaitchik BF, Sweijd N, Shumake-Guillemot J, Morse A, Gordon C, Marty A, et al. A framework for research linking weather, climate and COVID-19. Nat Commun. 2020;11:19–21.

17. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. JAMA - J Am Med Assoc. 2020;324:782–93.

18. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and transmission of COVID-19 in 391

cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. Lancet Infect Dis. 2020;20:911–9.

19. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes among 5700 Patients Hospitalized with COVID-19 in the New York City Area. JAMA - J Am Med Assoc. 2020;323:2052–9.

20. Grasselli G, Greco M, Zanella A, Albano G, Antonelli M, Bellani G, et al. Risk Factors Associated With Mortality Among Patients With COVID-19 in Intensive Care Units in Lombardy, Italy Supplemental content. JAMA Intern Med. 2020;180:1345–55.

21. ISCIII. Informe sobre la situación de COVID-19 en España. 2020.

22. Istituto Superiore di Sanità. Characteristics of SARS-CoV-2 patients dying in Italy Report based on available data on December 16th , 2020. 2020.

23. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet. 2020;395:1054–62.

24. Worldpop, Department of Geography and Geosciences University of Louisville, Department de Geographie Universite de Namur, CIESIN Columbia University. Global High Resolution Population Denominators Project. 2018. https://www.worldpop.org/doi/10.5258/SOTON/WP00647. Accessed 25 Jun 2020.

25. Lolli S, Chen YC, Wang SH, Vivone G. Impact of meteorological conditions and air pollution on COVID-19 pandemic transmission in Italy. Sci Rep. 2020;10:1–15.

26. Google LLC. Google COVID-19 Community Mobility Reports. https://www.google.com/covid19/mobility/data_documentation.html?hl=en. Accessed 4 Sep 2020.

27. Salje H, Kiem C, Lefrancq N, Courtejoie N, Paireau J, Andronico A, et al. Estimating the burden of SARS-CoV-2 in France To cite this version : HAL Id : pasteur-02548181. 2020.

28. Wallinga J, Lipsitch M. How generation intervals shape the relationship between growth rates and reproductive numbers. Proc R Soc B Biol Sci. 2007;274:599–604.

29. Obadia T, Haneef R, Boëlle P. The R0 package : a toolbox to estimate reproduction numbers for epidemic outbreaks. 2012.

30. Babin S. Use of Weather Variables in SARS-CoV-2 Transmission Studies. Int J Infect Dis. 2020;100:333–6.

31. Warszawski J, Bajos N, Meyer L, de Lamballerie X, Seng R, Beaumont A-L, et al. In May 2020, 4.5% of population in metropolitan France developed antibodies against SARS-CoV-2: first results from the national survey EpiCov [french]. 2020.

Tables

Table 1: Multivariable model results for the relationship between R0 and weather parameters,
obtained with the hierarchical generalized additive model. Weather parameters are temperature,
absolute humidity, and dew point temperature, adjusted for distance to the first region affected,
population density, and elderly population. Non-linear effects are presented in Figure 6. Models
assuming linear effects for weather covariates are presented in Table S5.

Model	Variable	Estimate	95%CI	p-value
Model 1	Intercept	0.78	[-0.88 - 2.45]	0.36152
	Population density (log10)	0.67	[0.07 - 1.26]	0.03218
	% population over 80	0.05	[-0.08 - 0.18]	0.44470
	Distance to first region affected in the country/coast	**spline**		0.08007
	Mean temperature	**spline**		0.00655
	Dev. explained: 41.5%			
Model 2	Intercept	1.28	[-0.36 - 2.92]	0.13239
	Population density (log10)	0.50	[-0.11 - 1.11]	0.11294
	% population over 80	0.03	[-0.1 - 0.16]	0.61862
	Distance to first region affected in the country/coast	**spline**		
	Mean AH	**spline**		0.03401
	Dev. explained: 33.6%			
Model 3	Intercept	1.20	[-0.47 - 2.88]	0.16427
	Population density (log10)	0.49	[-0.12 - 1.1]	0.12005
	% population over 80	0.05	[-0.08 - 0.18]	0.47092
	Distance to first region affected in the country/coast	**spline**		0.09756
	Mean Dew Point Temperature	**spline**		0.00494
	Dev. explained: 34.6%			

1   **Figures**

2   **Figure 1: Study flow chart**

1 **Figure 2: Deaths per day, by region and by country.** The thick red line figures the median date of
2 lockdown by each country, and the thin red line the median date +28 days.

1 **Figure 3: Map of regional values for R0 and selected covariates, panels are presented by continent.**
2 A,B: R0 ; C,D: population density (inhabitants per km2) ; D,E: mean temperature ; F,G: mean absolute
3 humidity.

1 **Figure 4: Distribution of R0 and selected covariates by country.** A: R0; B: population density (log10
2 inhabitants/km2); C: Mean absolute humidity (g/m3); D: Mean temperature (°C); E: Population over
3 80 years old (%); F: distance to the first region affected (km).

1   **Figure 5: Change in R0 on an additive scale estimated from the univariable model assuming a linear**
2   **relationship between R0 and the different variables.**

1 **Figure 6: Non-linear effects in the multivariable model for weather parameters (see Table 1 for linear**
2 **effects).** A: Temperature, model 1. B: Distance to first region affected, model 1. C: absolute humidity,
3 model 2. D: distance to first region affected, model 2. E: dew point temperature, model 3. F: distance
4 to first region affected, model 3.

Figure 7: Summary of estimated effect of temperature (A), absolute humidity (B) or dew point temperature (C) on R0 assuming a region with average population density (248 persons/km2) and percentage of inhabitants >80 years (5.6%), and corresponding to the first region first affected (distance=0km)**.**