

12-lead Electrocardiogram in Hospitalized COVID 19 Patients

Mohamed Shokr¹, Omar Chehab², Mustafa Ajam³, Manmohan Singh¹, Said Ashraf¹, John Dawdy¹, Mohit Pahuja¹, Vivek Reddy¹, Ahmed Subahi¹, M. Chadi Alraies¹, Luis Afonso¹, Randy Lieberman¹

1: Department of Cardiology, Wayne State University, Detroit, MI, USA

2: Department of Internal Medicine, Wayne State University, Detroit, MI, USA

3: Department of Pulmonary and Critical Care Medicine, Oregon Health and Science University, Portland, OR, USA

Corresponding Author Information:

Mohamed Shokr, MD

Phone: (313) 244-7083

Email: Mohamed.Shokr@nyulangone.org

Abstract

COVID-19 pandemic resulted in considerable morbidity and mortality. We analyzed 345 Electrocardiograms of 100 COVID-19 patients admitted to our tertiary care center in Detroit, during the initial month of Covid-19. Findings were correlated with mortality, cardiac injury and inflammatory markers. Our cohort included 61% males and 77% African Americans. The median age and BMI were 66 years (57-74) and 31 kg/m² (26.1-39), respectively. We observed atrial arrhythmias in 29% of the patients (17% new onset), First degree heart block in 12%, ST-T segment changes in 17%, S1Q3T3 pattern in 19%, premature ventricular complexes in 23%, premature atrial complexes in 13%, Q waves in 27%, T wave inversion in 42% of the cases. While presence of premature atrial complexes or left atrial abnormality correlated with mortality (P = 0.02 & 0.03, respectively), other findings did not show significant correlation in this small cohort of patients.

Manuscript

COVID-19 pandemic resulted in considerable morbidity and mortality (1). We analyzed 345 Electrocardiograms of 100 COVID-19 patients admitted to our tertiary care center in Detroit, during the initial month of Covid-19. Findings were correlated with mortality, cardiac injury and inflammatory markers (Tables 1, 2S & 3S supplementary index).

For our analysis, categorical and continuous variables were reported as frequency or percentage and mean \pm standard error (SE) or median \pm interquartile range, based on the normality of the data. Baseline demographics and comorbidities between groups were compared using the Pearson χ^2 test for categorical variables and one-way linear regression for continuous variables (Table 4S supplementary index)

Our cohort included 61% males and 77% African Americans. The median age and BMI were 66 years (57-74) and 31 kg/m² (26.1-39), respectively. History of Hypertension, Diabetes Mellitus, Cancer and lung disease was present in 71%, 40%, 15% and 26% of the patients, respectively. 12% had coronary artery disease, 10% had Systolic heart failure and 12% had history of atrial arrhythmias at baseline. The median hospital stay was 10.4 days (6.7-16) with 48% mortality.

We observed atrial arrhythmias in 29% of the patients (17% new onset), First degree heart block in 12% (6% were transient) , ST-T segment changes in 17% , S1Q3T3 pattern in 19%, Premature ventricular complexes (PVCs) in 23% , premature atrial complexes(PACs) in 13%, Q waves in 27%, T wave inversion in 42% of the cases . Long QTc (> 470 in males, >480 in females) was noted in 37 cases, of whom 20 patients were on Hydroxychloroquine and Azithromycin. There was no correlation between long QTc and mortality (P = 0.5).

Myocardial injury due to hypoxia and direct viral tissue invasion, sympathetic hyperactivity secondary to the cytokine storm and drug induced QTc prolongation are postulated mechanism of arrhythmias in COVID-19 cases (2)

Transient complete heart block was previously reported (3, 4). In our cohort, only first degree heart block was present without higher grade block.

S1Q3T3 pattern, possibly due to cor pulmonale, was reported in 2 Covid-19 cases (4). It was noted in 19% of patients in our cohort and did not correlate with mortality (P = 0.4)

ST elevation without obstructive coronary artery disease was reported in COVID-19 patients (5). We noted 8 cases of ST elevation, of which 5 were transient and 8 cases of ST depression. None of these patients underwent coronary angiogram during hospitalization. ST segment changes did not correlate with mortality (P = 0.5)

Premature complexes are possibly explained by the higher sympathetic tone, vasopressors use, electrolyte abnormalities and myocarditis. Right ventricular outflow tract was the most common site of origin (47% of cases). While PACs correlated with mortality (P = 0.02), PVCs correlation didn't reach

statistical significance ($P = 0.05$). EKG abnormalities are prevalent in this cohort of COVID-19 patients and can be a helpful diagnostic tool for underlying cardiac pathologies.

Acknowledgements:

None.

Financial Disclosures:

Authors have no financial disclosures to report.

Conflict of Interest:

The authors declare that they have no conflict of interest concerning this article.

Informed Consent:

Not applicable.

Authors Contributions:

Shokr M designed the research, collected the data and wrote the manuscript. Chehab O analyzed the data. All authors contributed to data collection. Shokr M and Lieberman R approved the final version of the manuscript.

References:

- 1- Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, Gong W, Liu X, Liang J, Zhao Q, Huang H. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA cardiology*. 2020 Mar 25.
- 2- Lazzarini PE, Boutjdir M, Capecchi PL. COVID-19, Arrhythmic Risk and Inflammation: Mind the Gap!. *Circulation*. 2020 Apr 14.

- 3- Azarkish M, Eslami M, Mollazadeh R. Transient complete heart block in a patient with critical COVID-19. *European Heart Journal*. 2020 Apr 14.
- 4- He J, Wu B, Chen Y, Tang J, Liu Q, Zhou S, Chen C, Qin Q, Huang K, Lv J, Chen Y, Peng D, Characteristic ECG Manifestations in Patients with COVID-19, *Canadian Journal of Cardiology* (2020), doi: <https://doi.org/10.1016/j.cjca.2020.03.028>.
- 5- Bangalore S, Sharma A, Slotwiner A, Yatskar L, Harari R, Shah B, Ibrahim H, Friedman GH, Thompson C, Alviar CL, Chadow HL. ST-segment elevation in patients with Covid-19—A case series. *New England Journal of Medicine*. 2020 Apr 17.

Table 1: Correlation of Electrocardiogram (EKG) findings with mortality, intensive care unit (ICU) admission, Cardiac injury (High sensitivity(Hs) Troponin > 17 ng/L) and Median C-reactive protein (175 mg/L)

EKG finding	Prevalence	Survival		P-value	ICU admission		P-value	Hs Troponin		P-value	Median CRP		P-value
		yes	No		No	Yes		≤17	>17		<175	≥175	
Transient 1 st °HB	6%	1.9%	10.4%	0.2	10.8%	3.2%	0.055	9.5%	6.4%	0.7	9.1%	2.3%	0.2
Persistent 1 st °HB	6%	5.8%	6.3%		0.0%	9.5%		4.8%	2.1%		4.5%	9.1%	
T wave inversion	42%	46.2%	37.5%	0.2	43.2%	41.3%	0.5	61.9%	36.2%	0.04	36.4%	52.3%	0.09
Premature ventricular complexes	23%	15.4%	31.3%	0.050	21.6%	23.8%	0.5	14.3%	23.4%	0.3	25.0%	18.2%	0.3
Premature atrial	13%	5.8%	20.8%	0.02	8.1%	15.9%	0.2	4.8%	12.8%	0.3	15.9%	9.1%	0.3

complexes													
Rhythm on admission				0.9			0.059			0.3			0.2
Sinus rhythm	56%	57.7%	54.2%		70.3%	47.6%		66.7%	46.8%		61.4%	45.5%	
Sinus Tachycardia	33%	30.8%	35.4%		18.9%	41.3%		28.6%	42.6%		27.3%	43.2%	
Atrial fibrillation	11%	11.5%	10.4%		10.8%	11.1%		4.8%	10.6%		11.4%	11.4%	
Rhythm during admission													
Sinus bradycardia	13%	17.3%	8.3%	0.1	24.3%	6.3%	0.01	9.5%	10.6%	0.6	18.2%	9.1%	0.1
Sinus tachycardia	81%	78.8%	83.3%	0.4	67.6%	88.9%	0.01	90.5%	80.9%	0.3	72.7%	90.9%	0.02
Left atrial abnormality	5.1%	9.8%	0%	0.03	5.6%	4.8%	0.6	0.0%	4.3%	0.5	2.4%	9.1%	0.1
Q waves				0.5			0.02			0.6			0.1
Q waves (Inferior)	22%	21.2%	22.9%		21.6%	22.2%		23.8%	25.5%		18.2%	29.5%	
(Anterior)	2%	3.8%	0%		5.4%	0.0%		0.0%	2.1%		2.3%	0.0%	
≥2 territories	3%	3.8%	2.1%		8.1%	0.0%		0.0%	6.4%		6.8%	0.0%	
Low voltage	16%	9.6%	22.9%	0.06	13.5%	17.5%	0.4	14.3%	14.9%	0.6	18.2%	11.4%	0.27
LVH	15%	17.3%	12.5%	0.3	16.2%	14.3%	0.5	9.5%	19.1%	0.3	6.8%	25.0%	0.01
Prolonged QTc	37%	36.5%	37.5%	0.5	29.7%	41.3%	0.2	42.9%	34.0%	0.3	31.8%	45.5%	0.1
S1Q3T3	19%	17.3%	20.8%	0.4	24.3%	15.9%	0.2	23.8%	25.5%	0.6	18.2%	22.7%	0.3

Table 2S: Correlation of EKG findings with Mortality, ICU admission, Cardiac injury and elevated Brain Natriuretic peptide (BNP) level (> 100 pg/ml)

EKG finding	N=100	Survivors		P-value	ICU admission		P-value	Cardiac injury		P-value	BNP		P-value
		yes	No		No	Yes		No	Yes		≤100	>100	
		1 st degree heart block (HB)	12%		7.7%	16.7%		0.1	10.8%		12.7%	0.5	
Transient 1 st degree HB	6%	1.9%	10.4%	0.2	10.8%	3.2%	0.055	9.5%	6.4%	0.7	5.9%	4.2%	0.9
Persistent 1 st degree HB	6%	5.8%	6.3%		0.0%	9.5%		4.8%	2.1%		5.9%	8.3%	
Transient ST elevation	5%	3.8%	6.3%	0.4	2.7%	6.3%	0.5	4.8%	6.4%	0.4	5.9%	4.25	0.7
Persistent ST elevation	3%	1.9%	4.2%		0.0%	4.8%		4.8%	2.1%		5.9%	0.0%	
ST depression	8%	11.5%	4.2%		8.1%	7.9%		14.3%	6.4%		11.8%	8.3%	

Early repolarization	1%	1.9%	0%		2.7%	0.0%		4.8%	0.0%		0.0%	4.2%	
T wave inversion	42%	46.2%	37.5%	0.2	43.2%	41.3%	0.5	61.9%	36.2%	0.04	41.2%	50.0%	0.4
T wave inversion (inferior)	5%	3.8%	6.3%	0.5	2.7%	6.3%	0.4	4.8%	6.4%	0.04	0.0%	12.5%	0.4
(anterior)	18%	21.2%	14.6%		16.2%	19.0%		14.3%	19.1%		23.5%	25.0%	
(Lateral)	2%	3.8%	0%		5.4%	0.0%		4.8%	2.1%		5.9%	0.0%	
(≥ 2 territories)	17%	17.3%	16.7%		18.9%	15.9%		38.1%	8.5%		11.8%	12.5%	
S1Q3T3	19%	17.3%	20.8%	0.4	24.3%	15.9%	0.2	23.8%	25.5%	0.6	29.4%	25.0%	0.5
Premature ventricular complexes	23%	15.4%	31.3%	0.050	21.6%	23.8%	0.5	14.3%	23.4%	0.3	29.4%	8.3%	0.1
Premature atrial complexes	13%	5.8%	20.8%	0.02	8.1%	15.9%	0.2	4.8%	12.8%	0.3	23.5%	4.2%	0.1
Poor R wave progression	14%	17.3%	10.4%	0.2	18.9%	11.1%	0.2	14.3%	14.9%	0.6	5.9%	16.7%	0.3
Rhythm on admission													
Sinus rhythm	56%	57.7%	54.2%	0.9	70.3%	47.6%	0.059	66.7%	46.8%	0.3	52.9%	58.3%	0.9
Sinus tachycardia	33%	30.8%	35.4%		18.9%	41.3%		28.6%	42.6%		35.3%	29.2%	
Atrial fibrillation	11%	11.5%	10.4%		10.8%	11.1%		4.8%	10.6%		11.8%	12.5%	
Rhythm during admission													
Atrial arrhythmias	29%	17.3%	27.1%	0.2	16.2%	25.4%	0.2	19.0%	25.5%	0.4	23.5%	25.0%	0.6
Atrial fibrillation	18%	15.4%	20.8%	0.3	13.5%	20.6%	0.3	14.3%	23.4%	0.3	23.5%	20.8%	0.6
Atrial flutter	6%	5.8%	6.3%	0.6	5.4%	6.3%	0.6	4.8%	2.1%	0.5	5.9%	8.3%	0.6
Atrial tachycardia	5%	1.9%	8.3%	0.5	2.7%	3.2%	0.7	4.8%	4.3%	0.7	0.0%	4.2%	0.6
Sinus bradycardia	13%	17.3%	8.3%	0.1	24.3%	6.3%	0.01	9.5%	10.6%	0.6	5.9%	16.7%	0.3
Sinus tachycardia	81%	78.8%	83.3%	0.4	67.6%	88.9%	0.01	90.5%	80.9%	0.3	82.4%	75.0%	0.4
Left atrial abnormality	5.1%	9.8%	0%	0.03	5.6%	4.8%	0.6	0.0%	4.3%	0.5	12.5%	4.2%	0.3
Right atrial abnormality	7.1%	5.9%	8.5%	0.4	5.6%	8.1%	0.5	5.0%	8.7%	0.5	6.3%	12.5%	0.5
Left axis deviation	24.2%	21.6%	27.1%	0.8	27.0%	22.6%	0.8	19.0%	27.7%	0.4	17.6%	33.3%	0.2
Right axis deviation	2.1%	2%	2.1%		2.7%	1.6%		0.0%	4.3%		0.0%	0.0%	
Right bundle branch block (RBBB)	4%	3.8%	4.2%	0.7	5.4%	3.2%	0.5	4.8%	4.3%	0.7	5.9%	0.0%	0.4
Incomplete RBBB	3%	3.8%	2.1%	0.5	0.0%	4.8%	0.2	0.0%	2.1%	0.7	0.0%	4.2%	0.6
Q waves	27%	28.8%	25%	0.4	35.1%	22.2%	0.1	23.8%	34.0%	0.3	35.3%	37.5%	0.6
Inferior Q waves	22%	21.2%	22.9%	0.5	21.6%	22.2%	0.02	23.8%	25.5%	0.6	29.4%	29.2%	0.8
Anterior Q waves	2%	3.8%	0%		5.4%	0.0%		0.0%	2.1%		5.9%	4.2%	
Q waves in ≥ 2 territories	3%	3.8%	2.1%		8.1%	0.0%		0.0%	6.4%		0.0%	4.2%	
Low voltage	16%	9.6%	22.9%	0.06	13.5%	17.5%	0.4	14.3%	14.9%	0.6	5.9%	29.2%	0.06
Left ventricular hypertrophy	15%	17.3%	12.5%	0.3	16.2%	14.3%	0.5	9.5%	19.1%	0.3	17.6%	12.5%	0.5
Left anterior fascicular block	12%	13.5%	10.4%	0.4	18.9%	7.9%	0.09	9.5%	17.0%	0.3	5.9%	20.8%	0.2
Bi-fascicular block	4%	3.8%	4.2%	0.7	5.4%	3.2%	0.5	4.8%	4.3%	0.7	5.9%	0.0%	0.4
Normal QRS width	95%	94.2%	95.8%	0.5	94.6%	95.2%	0.6	95.2%	95.7%	0.67	94.1%	100%	0.4

Table 3S: Correlation of EKG findings with median D-dimer of 1 mg/L, median C-reactive protein (CRP) of 175 mg/l, median Ferritin level of 900 ng/ml and median Length of stay (LOS) of 10 days

EKG finding	D-Dimer		P-value	CRP		P-value	Ferritin		P-value	LOS		P-value
	<1	≥1		<175	≥175		<900	≥900		<10	≥10	
	1 st degree heart block (HB)	6.3%		11.5%	0.47		13.6%	11.4%		0.5	13.6%	
Transient 1 st degree HB	0.0%	7.7%	0.4	9.1%	2.3%	0.2	9.1%	4.1%	0.5	6.4%	5.7%	0.7
Persistent 1 st degree HB	6.3%	3.8%		4.5%	9.1%		4.5%	8.2%		4.3%	7.5%	
Transient ST elevation	0.0%	7.7%	0.7	2.3%	6.8%	0.7	2.3%	8.2%	0.5	4.3%	5.7%	0.7
Persistent ST elevation	6.3%	3.8%		2.3%	2.3%		2.3%	4.1%		2.1%	3.8%	
ST depression	6.3%	9.6%		9.1%	9.1%		9.1%	8.2%		6.4%	9.4%	
Early repolarization	0.0%	1.9%		2.3%	0.0%		2.3%	0.0%		2.1%	0.0%	
T wave inversion	37.5%	48.1%	0.3	36.4%	52.3%	0.09	36.4%	51.0%	0.1	46.8%	37.7%	0.2
T wave inversion (Inferior)	0.0%	5.8%	0.7	2.3%	9.1%	0.2	2.3%	8.2%	0.3	4.3%	5.7%	0.1

(Anterior)	18.8%	19.2%		15.9%	20.5%		15.9%	20.4%		12.8%	22.6%	
(Lateral)	0.0%	3.8%		0.0%	4.5%		0.0%	4.1%		2.1%	1.9%	
(≥ two territories)	18.8%	19.2%		18.2%	18.2%		18.2%	18.4%		27.7%	7.5%	
S1 Q3T3	12.5%	19.2%	0.4	18.2%	22.7%	0.3	18.2%	22.4%	0.4	21.3%	17.0%	0.4
Premature ventricular complexes	12.5%	26.9%	0.2	25.0%	18.2%	0.3	25.0%	18.4%	0.3	25.5%	20.6%	0.4
Premature atrial complexes	6.3%	15.4%	0.3	15.9%	9.1%	0.3	15.9%	10.2%	0.3	12.8%	13.2%	0.9
Poor R wave progression	18.8%	11.5%	0.3	18.2%	11.4%	0.2	18.2%	10.2%	0.2	17.0%	11.3%	0.2
Rhythm on admission												
Sinus rhythm	56.3%	59.6%	0.1	61.4%	45.5%	0.2	61.4%	46.9%	0.34	59.6%	52.8%	0.7
Sinus tachycardia	25.0%	36.5%		27.3%	43.2%		27.3%	40.8%		29.8%	35.8%	
Atrial fibrillation	18.8%	3.8%		11.4%	11.4%		11.4%	12.2%		10.6%	11.3%	
Rhythm during admission												
Atrial arrhythmia	18.8%	17.3%	0.57	20.5%	20.5%	0.6	20.5%	22.4%	0.5	14.9%	28.3%	0.08
Atrial fibrillation	18.8%	13.5%	0.4	20.5%	18.2%	0.5	20.5%	16.3%	0.4	12.8%	22.6%	0.1
Atrial flutter	6.3%	1.9%	0.4	2.3%	6.8%	0.3	2.3%	8.2%	0.2	4.3%	7.5%	0.3
Atrial tachycardia	6.3%	3.8%	0.5	0.0%	4.5%	0.2	0.0%	6.1%	0.1	0.0%	5.7%	0.1
Sinus bradycardia	18.6%	9.6%	0.2	18.2%	9.1%	0.1	18.2%	8.2%	0.1	12.8%	13.2%	0.5
Sinus tachycardia	81.3%	92.3%	0.2	72.7%	90.9%	0.02	72.7%	89.8%	0.03	74.5%	86.8%	0.1
Left atrial abnormality	6.3%	2.0%	0.4	2.4%	9.1%	0.1	2.4%	8.2%	0.2	2.2%	7.5%	0.2
Right atrial abnormality	0.0%	7.8%	0.3	11.9%	4.5%	0.1	11.9%	4.1%	0.1	6.7%	7.5%	0.5
Left axis deviation	31.3%	23.1%	0.7	23.3%	31.8%	0.4	23.3%	28.6%	0.8	25.5%	23.1%	0.9
Right axis deviation	0.0%	1.9%		2.3%	0.0%		2.3%	2.0%		2.1%	1.9%	
Right bundle branch block (RBBB)	6.3%	1.9%	0.4	6.8%	2.3%	0.3	6.8%	2.0%	0.3	4.3%	3.8%	0.6
Incomplete RBBB	6.3%	1.9%	0.4	2.3%	4.5%	0.5	2.3%	4.1%	0.5	2.1%	3.8%	0.5
Q wave	18.8%	23.1%	0.5	27.3%	29.5%	0.5	27.3%	28.6%	0.5	25.5%	28.3%	0.4
Inferior Q wave	12.5%	23.1%	0.1	18.2%	29.5%	0.1	18.2%	28.6%	0.1	19.1%	24.5%	0.1
Anterior Q wave	0.0%	0.0%		2.3%	0.0%		2.3%	0.0%		0.0%	3.8%	
Q wave in ≥ 2 territories	6.3%	0.0%		6.8%	0.0%		6.8%	0.05		6.4%	0.0%	
Low voltage	12.5%	15.4%	0.56	18.2%	11.4%	0.3	18.2%	12.2%	0.3	23.4%	9.4%	0.051
Left ventricular hypertrophy	6.3%	11.5%	0.4	6.8%	25.0%	0.01	6.8%	22.4%	0.03	19.1%	11.3%	0.2
Left anterior fascicular block	25.0%	3.8%	0.02	13.6%	13.6%	0.6	13.6%	12.2%	0.5	17.0%	7.5%	0.1
Left posterior fascicular block	0.0%	1.9%	0.7	0.0%	0.0%	-	0.0%	2.0%	0.5	0.0%	1.9%	0.5
Bi-fascicular block	6.3%	1.9%	0.4	6.8%	2.3%	0.3	6.8%	2.0%	0.2	4.3%	3.8%	0.6
Normal QRS width	93.8%	98.1%	0.4	90.9%	97.7%	0.1	90.9%	98.0%	0.1	95.7%	94.3%	0.5

Table 4S : Baseline characteristics of our cohort of 100 patients diagnosed with COVID-19

Variable	N= 100	Survivors N=52	Non-survivors N=48	P-value
Age (median, IQR)	66 (57-74)	65 (47.5-71)	68.5 (60-75)	0.1
Race				
Male gender	61%	67.3%	54.2%	0.2
African Americans	77%	82.7%	70.8%	0.3
White	6%	3.8%	8.3%	
Other	17%	13.5%	20.8%	
Body mass index kg/m ²	31 (26.1 -39)	30.9 (25.9-36.8)	31.2 (26.6-41)	0.2
Hypertension	71%	71.2%	70.8%	1.0
Diabetes Mellitus	40%	30.8%	50%	0.05
Dyslipidemia	41%	40.4%	41.7%	0.9
Chronic/end-stage renal disease	19%	21.2%	16.7%	0.6
Lung disease	26%	23.1%	29.2%	0.5
Obstructive sleep apnea	11%	5.8%	16.7%	0.08
Cancer	15%	13.5%	16.7%	0.6
Immunological	7%	7.6%	6.2%	0.6
Mental disorder	12%	15.3%	8.3%	0.3
Neurological disorder	11%	11.5%	10.4%	0.9
Coronary artery disease	12%	17.3%	6.3%	0.08
Systolic heart failure	10%	15.4%	4.2%	0.06
Atrial fibrillation/Atrial flutter	10%	15.4%	4.2%	0.06
Troponin (Peak)	38.5(10-187.2)	23.5 (9- 185)	43.5 (18.75 -187.2)	0.2
Brain Natriuretic peptide (BNP)	130 (52.5-411.5)	123 (35.25 -261)	158 (56 -1034)	0.5
Ferritin	971 (418-1952.5)	884 (383-1345)	1035 (522-2802)	0.04
D-dimer	2.2 (1.12 -21.35)	1.34 (0.5-4.3)	4.9 (1.6 -35.2)	0.002
C-reactive protein (CRP)	175.0 (81.7-292.0)	171.5 (66.5-267.5)	207.5 (95.7-323.0)	0.2
Treatment				
No Hydroxychloroquine (HY)/ Azithromycin(AZ)/Doxycycline	12 (12.0%)	4 (7.7%)	8 (16.7%)	0.02
HY only	19 (19.0%)	9 (17.3%)	10 (20.8%)	
HY & AZ	40 (40.0%)	17 (32.7%)	23 (47.9%)	
HY & Doxycycline	29 (29.0%)	22 (42.3%)	7 (14.6%)	
Tocilizumab	5%	0%	10.4%	0.02
Steroids	53%	50%	56.3%	0.5
L-ascorbic acid	24%	23.1%	25%	0.8
Zinc	16%	15.4%	16.7%	0.9
Anticoagulation	32%	25%	39.6%	0.1
Single antiplatelet drug	33%	44.2%	20.8%	0.03
Dual antiplatelet therapy	7%	3.8%	10.4%	
Statins	38%	38.5%	37.5%	0.9
≥ Vasopressor	37%	5.8%	70.8%	<0.001
Beta blockers	41%	50%	31.3%	0.06
Calcium channel blocker	8%	7.7%	8.3%	0.6
Amiodarone	7%	3.8%	10.4%	0.2
Corrected QT(QTc) on admission (milliseconds)	443 (410.2 -472.7)	435 (404.5-476.7)	447 (416.2 -465.7)	0.8
Smoking	30%	36.5%	22.9%	0.1
Alcohol	20%	23.1%	16.7%	0.4
Illicit drugs	9%	15.4%	2.1%	0.02
Potassium trough	3.6 (3.3-4.0)	3.7 (3.3-3.9)	3.6 (3.4-4)	0.4
Potassium peak	4.75 (4.32-5.5)	4.75 (4.3-5.5)	4.75 (4.4-5.4)	0.7
Magnesium trough	1.9 (1.7-2.1)	1.9 (1.72-2.1)	2 (1.7-2.2)	0.5
Magnesium peak	2.4 (2.2-2.7)	2.3 (2.1-2.6)	2.6 (2.3- 2.87)	0.001
Sodium trough	134 (131-137)	134(132-136)	135 (130-138)	0.5
Sodium peak	143(140-147)	142 (140-145)	145 (140.2 -148.7)	0.03
Calcium trough	7.9(7.4-8.3)	8.1(7.6-8.4)	7.65 (7.1-8.2)	0.005
Calcium peak	9.0(8.5 -9.4)	9.2 (8.9-9.6)	8.7 (8.5-9.2)	0.001
Phosphorus trough	2.7 (2.2 -3.8)	2.6 (2.1-3.2)	3 (2.2-4.3)	0.2
Phosphorus peak	4.5(3.4-6.7)	3.7(3-4.7)	5.3(4.1-7.7)	<0.001
Intensive care unit (ICU) admission	48 (48.0%)	8 (21.6%)	40 (63.5%)	<0.001
ICU length of stay	3 (0- 8.9)	0 (0 - 3.75)	7 (2.2 -13.1)	< 0.001
Hospital length of stay	10.4 (6.7-16)	10.6 (7.07 -16.4)	9.4 (5.7-15.6)	0.5