Increased hazard of death in community-tested cases of SARS-CoV-2 Variant of Concern 202012/01

Nicholas G. Davies[1][†], Christopher I. Jarvis[1], CMMID COVID-19 Working Group, W. John Edmunds[1], Nicholas P. Jewell[2,3], Karla Diaz-Ordaz[2,3]*, Ruth H. Keogh[2,3]*

1. Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene and Tropical Medicine, London, UK.
2. Department of Medical Statistics, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK.
3. Centre for Statistical Methodology, London School of Hygiene and Tropical Medicine, London, UK.

[†]Corresponding author. E-mail: nicholas.davies@lshtm.ac.uk
*Equal contribution

VOC 202012/01, a SARS-CoV-2 variant first detected in the United Kingdom in September
2020, has spread to multiple countries worldwide. Several studies have established that
this novel variant is more transmissible than preexisting variants of SARS-CoV-2, but
have not identified whether the new variant leads to any change in disease severity. We
analyse a large database of SARS-CoV-2 community test results and COVID-19 deaths
for England, representing approximately 47% of all SARS-CoV-2 community tests and 7%
of COVID-19 deaths in England from 1 September 2020 to 22 January 2021. Fortuitously,
these SARS-CoV-2 tests can identify VOC 202012/01 because mutations in this lineage
prevent PCR amplification of the spike gene target (S gene target failure, SGTF). We
estimate that the hazard of death among SGTF cases is 30% (95% CI 9–56%) higher than
among non-SGTF cases after adjustment for age, sex, ethnicity, deprivation level, care
home residence, local authority of residence and date of test. In absolute terms, this
increased hazard of death corresponds to the risk of death for a male aged 55–69
increasing from 0.56% to 0.73% (95% CI 0.60–0.86%) over the 28 days following a positive
SARS-CoV-2 test in the community. Correcting for misclassification of SGTF, we
estimate a 35% (12–64%) higher hazard of death associated with VOC 202012/01. Our
analysis suggests that VOC 202012/01 is not only more transmissible than preexisting
SARS-CoV-2 variants but may also cause more severe illness.

SARS-CoV-2 Variant of Concern (VOC) 202012/01 (lineage B.1.1.7) carries several mutations,
including a 6-nucleotide deletion that prevents amplification of the S gene target by a
commercial PCR assay commonly used for community SARS-CoV-2 testing in England. By
linking individual records of positive community tests with and without S gene target failure
(SGTF) to COVID-19 deaths, we estimate the relative hazard of death from COVID-19
associated with infection by VOC 202012/01. We consider any PCR result with Ct < 30 for
ORF1ab, Ct < 30 for N, and no detectable S (Ct > 40) to be SGTF; any PCR result with cycle
threshold (Ct) < 30 for each of ORF1ab, N, and S to be non-SGTF; and exclude any other PCR
result from the analysis.

Characteristics of the study population

The study sample (**Table 1**) includes a total of 858,181 individuals who had a positive Pillar 2
(community) test between 1 November 2020 and 11 January 2021 processed by a laboratory
capable of producing an SGTF reading. Of these, 48% had SGTF. Females comprised 52.4% of
the sample; 45.3% were aged 1–34 years, 35.2% aged 35–54, 15.0% aged 55–69, 3.8% aged
70–85 and 0.7% aged 85 or older. The sex and age distributions were similar between those
with SGTF and without. The SGTF group had a slightly smaller percentage of females and
tended to be slightly younger (with 3.8% aged over 70 with SGTF versus 5.1% without SGTF).
The majority of individuals lived in standard residential accommodation, with 0.5% living in a
care or nursing home. Overall, 74.4% were White, 14.6% Asian, 3.9% Black and 7.0% of other,
mixed or unknown ethnicity. The SGTF group had a slightly greater percentage of individuals in
the Black and other, mixed or unknown ethnicity groups. More substantial differences were seen
between the SGTF groups when looking at index of multiple deprivation (IMD) decile and NHS
region. The SGTF group tended to have higher IMD (more deprived). The prevalence of SGTF

was low until the end of November 2020, with 95% of the observed SGTF cases appearing after
29 November 2020.

The highest prevalences of SGTF over the study period were observed in the East of England
(71.9%), South East (71.8%) and London (71.0%) NHS England regions. Prevalence of SGTF
was lowest in the North East and Yorkshire region. SGTF prevalence was similar in males and
females but lower in the older age groups: 48.6% in the 1-34 year olds compared with 36.3% in
those aged 85 and older. In keeping with these age patterns, SGTF prevalence was lower in
individuals living in a care or nursing home (30.7%, compared to 48.2% among those in
standard residential accommodation). SGTF prevalence by self-identified ethnicity was 46.6% in
the White group, 47.4% in the Asian group, 62.1% in the Black group, and 55.8% in the other,
mixed, or unknown ethnicity group. SGTF prevalence was lowest in the most deprived IMD
decile (34.1%) and highest in the least deprived decile (53.7%). The prevalence of SGTF
among those tested also increased steeply over time (**Fig. 1a**), ranging from 4.9% during 1–14
November 2020 to 78.7% during 27 December 2020–11 January 2021. **Table 2** presents
deaths within 28 days of a positive test among study subjects, **Table 3** presents crude death
rates within 28 days of a positive test per 10,000 person-days of follow-up, and **Table 4** for
unlimited follow-up (i.e. not restricted to 28 days); the maximum observed follow-up was 71
days. 2,967 individuals in the sample are known to have died, 2,091 of whom died within 28
days of their first positive test (**Fig. 1b**).

60-day survival assessed by Kaplan-Meier curves was higher in the SGTF group over the entire
study sample (**Fig. 1c**), but this apparently better survival for SGTF individuals is reversed when
stratifying the sample by age (**Fig. 1d,e**) indicating that the association between SGTF and
survival is confounded by the younger average age of SGTF cases. Stratifying by broad age
groups and by sex, place of residence, ethnicity, index of multiple deprivation, NHS England
region, and specimen date, crude death rates within 28 days of a positive SARS-CoV-2 test are
higher among SGTF than non-SGTF in 79 of the 104 stratifications assessed (76%; **Figs. 1f–k**).
Stratified death rates are considerably higher in older age groups and in individuals who live in a
care or nursing home, and generally, are higher in males than in females, higher in individuals
of Asian or Black ethnicity than in individuals of White or Mixed/unknown/other ethnicity, and
higher in the most deprived IMD deciles.

95
96

Fig. 1. Descriptive analyses. a The number of samples with and without SGTF by day from 1 November 2020 to 11 January 2021. **b** Number of deaths within 28 days of positive test by specimen date included in the analysis. **c, d, e** Kaplan-Meier plots showing survival in individuals tested in the community in England with SGTF versus without in (**c**) all individuals, (**d**) individuals under 70, and (**e**) individuals 70 years or older. The apparent better survival in the SGTF group in the whole study cohort (**c**) is due to confounding by age. Insets show the full *y*–axis range. **f–k** Crude death rates (with 95% confidence intervals) in SGTF versus other for deaths within 28 days of positive test stratified by broad age groups and (**f**) sex, (**g**) place of residence, (**h**) ethnicity, (**i**) index of multiple deprivation, (**j**) NHS England region, and (**k**) specimen date. Grey ribbons show the overall crude death rates by age group irrespective of SGTF status.

Cox regression analyses

To estimate the effect of SGTF on mortality while controlling for observed confounding, we fitted
a series of Cox proportional hazards models[1] to the data. In the Cox model with baseline hazard
stratified by lower tier local authority (LTLA) and specimen date and adjusted for the other
covariates, the estimated hazard ratio for SGTF was 1.30 (95% CI 1.09–1.56), indicating that
the hazard of death within 28 days of a positive test is 30% (9–56%) higher in those with SGTF
compared to non-SGTF (**Fig. 2a**). Without restricting the length of followup time, we estimated a
hazard ratio of 1.26 (1.07–1.50). Including age and IMD as restricted cubic splines rather than
linear terms did not change the results. An interaction between SGTF and time was included in
the model as an assessment of the proportional hazards assumption. There was strong
evidence of non-proportionality of hazards (likelihood ratio test P = 0.029; **Fig. 2a; Fig. S11**).
The hazard ratio is just below 1 initially, with confidence intervals including 1, and then
increases over time, crossing 1 at 3 days post positive test. The estimated time-varying hazard
ratio is 0.95 (0.69–1.33) 1 day after the positive test, 1.14 (0.92–1.42) on day 7, 1.41 (1.16–
1.70) on day 14, 1.73 (1.27–2.36) on day 21 and 2.13 (1.33–3.42) on day 28. There was no
evidence that adding higher order functions of time into the interaction terms improved model fit
(likelihood ratio test P = 0.371), and no evidence of a significant interaction between time and
age (P = 0.609), time and sex (P = 0.163), time and IMD (P = 0.703), time and ethnicity (P =
0.406), or time and residence type (P = 0.609).

We found no evidence of a significant interaction between SGTF and age group (likelihood ratio
test P = 0.804), sex (P = 0.543), IMD (P = 0.512), and ethnicity (P = 0.210). There was weak
evidence of an interaction between SGTF and residence type (P = 0.0574), with the associated
hazard ratio for SGTF being 1.26 (1.05–1.51) in residential settings (houses, flats, sheltered
accommodation and houses in multiple occupancy), 2.54 (1.38–4.68) in care/nursing homes,
and 1.64 (0.61–3.39) in "other" residence types (i.e. residential institutions including residential
education, prisons and detention centres, medical facilities, no fixed abode and other/unknown).

Fig. 2. Survival analyses. a, b Estimated hazard ratio of death within 28 days of positive test (**a**) using SGTF and (**b**) using variant of concern (VOC) 202012/01 as estimated by our misclassification analysis, in model stratified by LTLA and specimen date and adjusted for the other covariates. **c** Estimated hazard ratio of death within 28 days of positive test across each model tested. Model structures are coded as follows: VOC marker (special features) | stratification [date range]. VOC marker is either SGTF or p_voc. Special features: adjust for NHSE x week, NHS England region and specimen week included as fixed-effect covariates instead of in stratification of the baseline hazard; spline, age and IMD included as spline instead of as linear terms; symptom status, asymptomatic indicator included as a covariate; *X*-day followup, a followup length of *X* days is used instead of the default 28 days; Time-Y interaction, interaction terms included between followup time (optionally: also time^2) and covariate *Y*. Stratification of the baseline hazard in different models includes geographical level (NHSE: NHS England region; UTLA: upper-tier local authority; LTLA: lower-tier local authority). Date range: Specimens included from the indicated date (in 2020) forward; "by LTLA" signifies a start date chosen separately for each LTLA (see Methods), "1 Nov+" indicates no registration cutoff was used, and "1 Nov-20 Dec" uses an early cutoff so that all individuals have at least 28 days of followup time. Point estimates and 95% confidence intervals shown.

Misclassification analysis

Prior to the emergence of VOC 202012/01, a number of minor circulating SARS-CoV-2 lineages with spike mutations could also cause SGTF. We restrict our main analyses to specimens from 1 November 2020 onwards to minimise the number of these non-VOC 202012/01 lineages among SGTF-positive samples. However, the appearance of non-VOC 202012/01 samples in SGTF may dilute the estimated effect of VOC 202012/01 on the hazard of mortality, while the exclusion of any VOC 202012/01 samples taken from prior to 1 November 2020 may reduce the power of the analysis. We therefore undertook a misclassification analysis, modelling the relative frequency of SGTF over time for each NHS England region as a combination of a low, time-invariant frequency of non-VOC 202012/01 samples with SGTF plus a logistically growing frequency of VOC 202012/01 samples with SGTF, which allows us to assign to each SGTF sample a probability p_{VOC} that the sample is VOC 202012/01 based upon its specimen date and NHS England region (**Fig. S12**). Again restricting the analysis to specimens from 1 November 2020 onward, we find a hazard ratio associated with p_{VOC} of 1.35 (1.11–1.63), slightly higher than the hazard ratio associated with SGTF of 1.30 (1.09–1.56). Including all specimens from 1 September 2020 onward, the hazard ratio associated with p_{VOC} remains at 1.35 (1.11–1.63), while the hazard associated with SGTF for the same time period decreases to 1.25 (1.05–1.48).

Absolute risks

To estimate absolute risks for individuals with VOC 202012/01, we applied the 28- and 60-day hazard ratios obtained for SGTF and p_{VOC} to the baseline (i.e. not variant-specific) risk estimated from all specimens taken between 1 August and 31 October 2020 (**Table 5**). The absolute risk remains low amongst age groups less than 54, and the absolute risk was higher in males compared to females. For SGTF, in females aged 70–84 the risk of death within 28 days increased from 2.9% to 3.7% (95% CI 3.1–4.4%) and for females 85 or older increased from 12.8% to 16.4% (13.7–19.0%). For males aged 70–84 the risk of death within 28 days increased from 4.7% to 6.1% (5.0–7.1%) and for males 85 or older increased from 17.1% to 21.7% (18.3–25.1%). Estimates of the absolute risk at 60 days were higher than those for 28 days and show similar patterns, with the largest increase of 17.1% to 22.3% (18.6–26.1%) in males 85 years and older. Estimates based on p_{VOC} were marginally higher. These estimates reflect a substantial increase in absolute risk amongst older age groups. Note that these estimates do not reflect the infection fatality ratio, but the fatality ratio among people tested in the community, and are thus likely to be higher than the infection fatality rate as many infected individuals will not have been tested.

Further investigations

We conducted a number of sensitivity analyses to verify the robustness of our results. Our main analyses are stratified by LTLA (n = 316) and specimen date. Stratifying instead by upper-tier local authority (UTLA; n = 150) or by NHS England region (n = 7), or by week rather than by date of specimen, produced similar results, with the effect size of SGTF being greater for coarser strata; stratifying at the coarsest level—by NHS England region and specimen week—

yielded a hazard ratio for SGTF of 1.40 (1.23–1.59). Adjusting for, rather than stratifying by,
specimen week and NHSE region yielded a similar hazard ratio of 1.40 (1.23–1.60). We
interpret the decreased effect size of SGTF for finer stratifications as resulting from controlling
better for differences in heterogeneities in the availability of hospital services owing to increased
hospital demand as local SGTF prevalence increases. When restricting data to specimens
collected from September onwards, October onwards, November onwards, or December
onwards, the estimated hazard ratio of SGTF increased (**Fig. 2c**) as the data period became
more restricted, which we interpret as resulting from including fewer non-VOC 202012/01
specimens in the SGTF sample. Removing the administrative cutoff to followup time of 10 days
prior to data extraction (i.e. 11 January 2020) increased the hazard ratio of SGTF to 1.48 (1.28–
1.70), and restricting the analysis to individuals with at least 28 days' follow-up yielded a hazard
ratio of 1.34 (1.06–1.71). Pillar 2 testing data include an indicator for whether the subject was
asymptomatic at the time of requesting the test (or symptomatic, or unknown). Although
symptomatic status may lie on the causal pathway between SGTF status and death, we
adjusted for symptomatic status as a sensitivity analysis and found that it had no effect on the
relative hazard of SGTF (1.30 [1.09–1.56]).

Discussion
212
Our analysis focuses on deaths within the first 28 days following a positive test, which could
overestimate the change in mortality associated with SGTF if individuals infected by VOC
202012/01 die sooner than individuals infected with preexisting SARS-CoV-2 variants. However,
the consistency of results when analysing data with 60 days of followup or unlimited followup
(**Fig. 2c**) suggests that this is not the case. By stratifying on test time and region, we attempted
to control for the effects of pressure on health services, which cannot be adjusted for directly, as
these lie on the causal pathway between infection and mortality.
220
We do not identify the mechanism for an increased mortality rate in this analysis. There is some
evidence that infections with VOC 202012/01 may be associated with higher viral loads, as
measured by Ct values detected during PCR testing of specimens (**Fig. S13**), although Ct
values can be biased during the growth phase of an epidemic[2]. Higher viral loads resulting from
infection with VOC 202012/01 may be partly responsible for the observed increase in mortality,
partly because they may reduce the efficacy of standard antiviral treatments for COVID-19. The
impact of viral load on observed SGTF mortality could be assessed using a mediation analysis,
which is outside the remit of this study.
229
We previously identified that the novel SARS-CoV-2 lineage VOC 202012/01 appears to have a
substantially greater transmission rate than preexisting variants of SARS-CoV-2[3], but could not
robustly estimate any increase or decrease in associated disease severity from ecological
analysis. The individual-level linked community testing data analysed here suggest that the
fatality rate among individuals infected with VOC 202012/01 is higher than that associated with
infection by preexisting variants. Crucially, due to the nature of the data currently available, we
were only able to assess mortality among individuals who received a positive test for SARS-
CoV-2 in the community that was processed at one of the three national Lighthouse laboratories

capable of returning an SGTF positive or negative result. Indicators for VOC 202012/01 are not
currently available for the vast majority of individuals who die due to COVID-19, as they are first
tested in hospital. Accordingly, the evidence we provide here must be contextualised with
further study of a larger population sample. Our analysis is consistent with analyses by other
groups using different methods to verify the increased risk of death among community-tested
individuals with SGTF[4]. Estimates of increased mortality based upon Pillar 2 data will become
more robust as test results and mortality outcomes continue to accrue over time, although our
analysis using stratified Cox regression, which estimates hazard ratios for mortality by
comparing outcomes between individuals with and without SGTF who were tested in the same
place and at the same time, would no longer accrue additional information at the point when
SGTF becomes effectively fixed in England—which may occur as soon as February 2021 if
current trends continue[3].
250
251

253
254

Methods

Data sources — We linked three datasets provided by Public Health England: a linelist of all positive tests in England's "Pillar 2" (community) testing for SARS-CoV-2, containing specimen date and demographic information on the test subject; a linelist of cycle threshold (Ct) values for the ORF1ab, N (nucleocapsid), and S (spike) genes for positive tests that were processed in one of the three national laboratories (Alderley Park, Glasgow, or Milton Keynes) utilising the Thermo Fisher TaqPath COVID-19 assay; and a linelist of all deaths due to COVID-19 in England. We link these datasets using a numeric identifier common to all three datasets. We define S gene target failure (SGTF) as any test with Ct < 30 for ORF1ab and N targets but no detectable S gene; non-SGTF as any test with Ct < 30 for ORF1ab, N, and S targets; and all other tests as inconclusive and excluded from the analysis. Because cycle threshold values are not available for individuals who were not tested in the community, this linked dataset does not allow analysis of individuals who first tested positive in hospital, that is, those who presented to hospital after symptom onset without first being tested in the community. This is the reason why our linked dataset has SGTF status for 47% of all community tests (Table S1) from 1 November 2020–22 January 2021, but only 7.4% of all COVID-19 deaths in England following a positive test in either the community or hospital over the same time period (Table S2).

For our main analysis, we included only tests from after 1 November 2020 to avoid including an excess of tests with SGTF not resulting from infection by VOC 202012/01, and censored individuals who did not die within 28 days of their positive test at 28 days. For individuals with less than 28 days of followup after their positive test, we applied censoring at the date of data extraction minus 10 days to reduce any impact of late reporting of deaths. There were missing data for sex (*n* = 7, <0.01%), age (*n* = 56, <0.01%), and IMD and regional covariates (*n* = 1038, 0.1%). There were no missing specimen dates. Individuals with missing age, sex, or geographical location were excluded. We also excluded individuals from the dataset whose age was recorded as zero, as there were 15,400 age-0 individuals compared to 7,212 age-1 individuals in the dataset, suggesting that many of these age-0 individuals may have been miscoded. Categories for missing ethnicity and missing residence type were created.

We grouped residence types into three categories: Residential, which included the "Residential dwelling (including houses, flats, sheltered accommodation)" and "House in multiple occupancy (HMO)" groups; Care/Nursing home; and Other/Unknown, which included the "Medical facilities (including hospitals and hospices, and mental health)", "No fixed abode", "Other property classifications", "Overseas address", "Prisons, detention centres, secure units", "Residential institution (including residential education)", and "Undetermined" groups, as well as unspecified residence type. We grouped ethnicities into four categories according to the broad categories used in the 2011 UK Census: Asian, which included the "Bangladeshi (Asian or Asian British)", "Chinese (other ethnic group)", "Indian (Asian or Asian British)", "Pakistani (Asian or Asian British)", and "Any other Asian background" groups; Black, which included the "African (Black or Black British)", "Caribbean (Black or Black British)", and "Any other Black background" groups; White, which included the "British (White)", "Irish (White)", and "Any other White background" groups; and Other / Mixed / Unknown, which included the "Any other ethnic group", "White and

299   Asian (Mixed)", "White and Black African (Mixed)", "White and Black Caribbean (Mixed)", "Any
300   other Mixed background", and "Unknown" groups.
301
Statistical methods — There are several factors that we expect to be associated with both the
probability of SGTF and with risk of death, thus confounding the association between SGTF and
risk of death in those tested. Area of residence and specimen date were expected to be
potentially strong confounders. Area of residence is expected to be strongly associated with
SGTF status due to different virus variants circulating in different areas, and specimen date
because the prevalence of SGTF is known to have greatly increased over time. Area of
residence and specimen date are also expected to be associated with risk of death following a
test, including due to differential pressure on hospital resources by area and time. The following
variables were also identified as potential confounders: sex, age, place of residence
(Residential, Care/Nursing home, or Other/Unknown), ethnicity (White, Asian, Black, or
Other/Mixed/Unknown), index of multiple deprivation (IMD, in deciles).
313
Descriptive analyses were performed. We tabulated the association between SGTF and each of
the potential confounders (Table 1). Missing data in the exposure by levels of the confounders
were tabulated (Table S1). The unadjusted association between SGTF and mortality was also
assessed using a Kaplan-Meier plot and by tabulating mortality rates. Kaplan-Meier plots (**Figs.**
S1–S10) and mortality rates (**Tables 4–5**) are also presented separately according to categories
of the potential confounders. Exact Poisson CIs are used for mortality rates, assuming constant
rate.
321
322   We performed complete cases analysis. This assumes that for each analysis, the missing data
323   are independent from the outcome of interest, given the variables included in the models.
324   We also assumed that censoring is uninformative, which is plausible as all censoring is
325   administrative.
326
Cox regression was used to estimate the association between SGTF and the hazard for
mortality, conditioning on the potential confounders listed above. The baseline hazard in the
Cox model was stratified by both specimen date and LTLA, therefore finely controlling for these
variables. The remaining variables were included as covariates in the model (sex, age, place of
residence, ethnicity, IMD decile). Age and IMD were included either as linear terms, given an
observed log-linear effect of age on the infection fatality rate of SARS-CoV-2[5], or as restricted
cubic splines with 3 knots. The time origin for the analysis was specimen date and we
considered deaths up to 28 days after the specimen date. Individuals who did not die within 28
days were censored at the earlier of 28 days post specimen date and the administrative
censoring date, which we chose as the date of the most recent death linkable to SGTF status
minus 10 days in order to minimise any potential bias due to late reporting of deaths. We began
by assuming proportionality of hazards for SGTF and the covariates included in the model. The
proportional hazards assumption was assessed by including in the model an interaction
between each covariate and time, which was performed separately for SGTF and for each other
covariate. Schoenfeld residual plots were also obtained for each covariate. We assessed
whether the association between SGTF and the hazard was modified by age, sex, IMD,

ethnicity, and place of residence. Models with and without interactions were compared using
likelihood ratio tests.
345
Misclassification analysis — The exposure of SGTF is subject to misclassification, because a
number of minor circulating variants of SARS-CoV-2 in addition to VOC 202012/01 are also
associated with failure to amplify the spike gene target. Accordingly, a positive test with SGTF is
not necessarily indicative of infection with VOC 202012/01. Misclassification of an exposure can
result in bias in its estimated association with the outcome. We fitted a beta-binomial logistic
growth model to Pillar 2 data by NHS region to estimate a "background" rate of SGTF in the
absence of VOC 202012/01. This model is then used to estimate the probability that an
individual testing positive with SGTF is infected with VOC 202012/01, separately for individuals
in each NHS region. These probabilities can then be used in place of the binary SGTF exposure
in the Cox models (assuming proportional hazards, and allowing for a time-varying hazard ratio
where there is evidence of proportional hazards violations).
357
358   We fitted models of logistic growth accounting for false positives (modelled as regionally-varying
359   background rates of SGTF associated with non-VOC 202012/01 variants) to the SGTF data.
360   Our logistic beta-binomial model of VOC 202012/01 growth is as follows:
361

$$slope \sim normal(\mu = 0, \sigma = 1)$$
$$intercept \sim normal(\mu = 0, \sigma = 1000)$$
$$falsepos \sim beta(\alpha = 1.5, \beta = 15)$$
$$conc \sim normal(\mu = 0, \sigma = 500) \geq 2$$

366
367

$$f(t) = exp\ (slope \times (t - intercept))\ /\ (1 + exp\ (slope \times (t - intercept)))$$
$$s(t) = f(t) + (1 - f(t)) \times falsepos$$
$$k_t \sim betaBinomial(n = n_t, \alpha = s(t) \times (conc - 2) + 1, \beta = (1 - s(t)) \times (conc - 2) + 1)$$

371
372   Here, *f(t)* is the model-predicted frequency of VOC 202012/01 at time *t* based on the terms
373   *slope* and *intercept*, *s(t)* is the model-predicted frequency of S gene target failure at time *t* owing
374   to a background false positive rate *falsepos*, *conc* is the "concentration" parameter (= α + β) of a
375   beta distribution with mode *s(t)*, k_t is the number of S gene target failures detected at time *t* and
376   n_t is the total number of tests at time *t*.
377
Absolute risks — Estimates from the final Cox models were used to obtain estimates of absolute
risk of death for 28 and 60 days with SGTF and p_{VOC}. Given the strong influence of age on risk
of death, we present absolute risks by sex and age group (1-34, 35-54, 55-69, 70-84, 85+).
Absolute risks of death (case fatality rate) within 28 and 60 days were estimated by age group
and sex using data on individuals tested in August, September, and October 2020; this is
referred to as the baseline risk. The absolute risks of death for individuals with SGTF were then
estimated as follows. If the baseline absolute risk of death in a given age group is $(1 - A)$, then
the estimated absolute risk of death with SGTF is $(1 - A^{HR})$, where HR denotes the estimated
hazard ratio obtained from the Cox model with proportional hazards. We applied the hazard

ratio from 28 days to the baseline risk for 28 days, and the hazard ratio for 60 days to the
baseline risk for 60 days, to estimate absolute risks of death for individuals with SGTF and
uncertainty of these estimates. Standard errors are obtained via the delta method, and CIs
based on normal approximations.
391
Sensitivity analyses — Several sensitivity analyses were performed. After establishing the final
model through using the process outlined above we investigated the impact of using different
variables for stratification of the baseline hazard measuring region at a coarser level (UTLA, or
NHS England region), as well as coarser test specimen time (week rather than exact date).
Adjusting for these variables instead of using stratification was also explored. We also repeated
the main analysis restricting data to specimens collected from September onwards, October
onwards, November onwards, or December onwards.
To assess the impact of imposing an administrative cutoff to follow-up time of 10 days prior to
data extraction, we first reanalysed the data without this cutoff, as well as reanalysing the data
restricting the analysis to individuals with at least 28 days' follow-up.
Finally, we adjusted for symptomatic status at the time of requesting the test (asymptomatic,
symptomatic, or unknown).

Funding statement
NGD: UK Research and Innovation (UKRI) Research England; NIHR Health Protection
Research Unit in Immunisation (NIHR200929); UK Medical Research Council (MC_PC_19065).
CIJ: Global Challenges Research Fund project 'RECAP' managed through Research Councils
UK and the Economic and Social Research Council (ES/P010873/1). WJE: European
Commission (EpiPose 101003688), National Institutes of Health Research (NIHR200908). NPJ:
National Institutes of Health / National Institute of Allergy and Infectious Diseases
(R01AI148127). KDO: Royal Society-Wellcome Trust Sir Henry Dale Fellowship 218554/Z/19/Z.
RHK: UKRI Future Leaders Fellowship (MR/S017968/1).
413
Working group authors and acknowledgements
The CMMID COVID-19 working group is (randomized order) Kevin van Zandvoort, Samuel
Clifford, Fiona Yueqian Sun, Sebastian Funk, Graham Medley, Yalda Jafari, Sophie R Meakin,
Rachel Lowe, W John Edmunds, Matthew Quaife, Naomi R Waterlow, Rosalind M Eggo,
Nicholas G. Davies, Jiayao Lei, Mihaly Koltai, Fabienne Krauer, Damien C Tully, James D
Munday, Alicia Showering, Anna M Foss, Kiesha Prem, Stefan Flasche, Adam J Kucharski,
Sam Abbott, Billy J Quilty, Thibaut Jombart, Alicia Rosello, Gwenan M Knight, Mark Jit, Yang
Liu, Jack Williams, Joel Hellewell, Kathleen O'Reilly, Yung-Wai Desmond Chan, Timothy W
Russell, Christopher I Jarvis, Simon R Procter, Akira Endo, Emily S Nightingale, Nikos I Bosse ,
C Julian Villabona-Arenas, Frank G Sandmann, Amy Gimma, Kaja Abbas, William Waites,
Katherine E. Atkins, Rosanna C Barnard, Petra Klepac, Hamish P Gibbs, Carl A B Pearson, and
Oliver Brady.
426
Funding statements for the CMMID COVID-19 working group are as follows. KvZ: KvZ is
supported by the UK Foreign, Commonwealth and Development Office (FCDO)/Wellcome Trust

Epidemic Preparedness Coronavirus research programme (ref. 221303/Z/20/Z), and Elrha's
Research for Health in Humanitarian Crises (R2HC) Programme, which aims to improve health
outcomes by strengthening the evidence base for public health interventions in humanitarian
crises. The R2HC programme is funded by the UK Government (FCDO), the Wellcome Trust,
and the UK National Institute for Health Research (NIHR). SC: Wellcome Trust (grant:
208812/Z/17/Z). FYS: NIHR EPIC grant (16/137/109). SFunk: Wellcome Trust (grant:
210758/Z/18/Z), NIHR (NIHR200908). GFM: NTD Modelling Consortium by the Bill and Melinda
Gates Foundation (OPP1184344). YJ: LSHTM, DHSC/UKRI COVID-19 Rapid Response
Initiative. SRM: Wellcome Trust (grant: 210758/Z/18/Z). RL: Royal Society Dorothy Hodgkin
Fellowship. WJE: European Commission (EpiPose 101003688), NIHR (NIHR200908). MQ:
European Research Council Starting Grant (Action Number #757699); Bill and Melinda Gates
Foundation (INV-001754). NRW: Medical Research Council (grant number MR/N013638/1).
RME: HDR UK (grant: MR/S003975/1), MRC (grant: MC_PC 19065), NIHR (grant:
NIHR200908). NGD: UKRI Research England; NIHR Health Protection Research Unit in
Immunisation (NIHR200929); UK MRC (MC_PC_19065). JYL: Bill & Melinda Gates Foundation
(INV-003174). MK: Foreign, Commonwealth and Development Office / Wellcome Trust. FK:
Innovation Fund of the Joint Federal Committee (Grant number 01VSF18015), Wellcome Trust
(UNS110424). DCT: No funding declared. JDM: Wellcome Trust (grant: 210758/Z/18/Z). AS: No
funding declared. AMF: No funding declared. KP: Gates (INV-003174), European Commission
(101003688). SFlasche: Wellcome Trust (grant: 208812/Z/17/Z). AJK: Wellcome Trust (grant:
206250/Z/17/Z), NIHR (NIHR200908). SA: Wellcome Trust (grant: 210758/Z/18/Z). BJQ: This
research was partly funded by the National Institute for Health Research (NIHR) (16/137/109 &
16/136/46) using UK aid from the UK Government to support global health research. The views
expressed in this publication are those of the author(s) and not necessarily those of the NIHR or
the UK Department of Health and Social Care. BJQ is supported in part by a grant from the Bill
and Melinda Gates Foundation (OPP1139859). TJ: RCUK/ESRC (grant: ES/P010873/1); UK
PH RST; NIHR HPRU Modelling & Health Economics (NIHR200908). AR: NIHR (grant: PR-OD-
1017-20002). GMK: UK Medical Research Council (grant: MR/P014658/1). MJ: Gates (INV-
003174, INV-016832), NIHR (16/137/109, NIHR200929, NIHR200908), European Commission
(EpiPose 101003688). YL: Gates (INV-003174), NIHR (16/137/109), European Commission
(101003688). JW: NIHR Health Protection Research Unit and NIHR HTA. JH: Wellcome Trust
(grant: 210758/Z/18/Z). KO'R: Bill and Melinda Gates Foundation (OPP1191821). YWDC: No
funding declared. TWR: Wellcome Trust (grant: 206250/Z/17/Z). CIJ: Global Challenges
Research Fund (GCRF) project 'RECAP' managed through RCUK and ESRC (ES/P010873/1).
SRP: Bill and Melinda Gates Foundation (INV-016832). AE: The Nakajima Foundation. ESN:
Gates (OPP1183986). NIB: Health Protection Research Unit (grant code NIHR200908). CJVA:
European Research Council Starting Grant (Action number 757688). FGS: NIHR Health
Protection Research Unit in Modelling & Health Economics, and in Immunisation. AG: European
Commission (EpiPose 101003688). KA: Bill & Melinda Gates Foundation (OPP1157270, INV-
016832). WW: MRC (grant MR/V027956/1). KEA: European Research Council Starting Grant
(Action number 757688). RCB: European Commission (EpiPose 101003688). PK: This research
was partly funded by the Royal Society under award RP\EA\180004, European Commission
(101003688), Bill & Melinda Gates Foundation (INV-003174). HPG: This research was
produced by CSIGN which is part of the EDCTP2 programme supported by the European Union

(grant number RIA2020EF-2983-CSIGN). The views and opinions of authors expressed herein
do not necessarily state or reflect those of EDCTP. This research is funded by the Department
of Health and Social Care using UK Aid funding and is managed by the NIHR. The views
expressed in this publication are those of the author(s) and not necessarily those of the
Department of Health and SocialCare (PR-OD-1017-20001). CABP: CABP is supported by the
Bill & Melinda Gates Foundation (OPP1184344) and the UK Foreign, Commonwealth and
Development Office (FCDO)/Wellcome Trust Epidemic Preparedness Coronavirus research
programme (ref. 221303/Z/20/Z). OJB: Wellcome Trust (grant: 206471/Z/17/Z).

Ethical approval

Approved by the Observational / Interventions Research Ethics Committee at the London

School of Hygiene and Tropical Medicine (reference number 24020).

Code and data availability

Analysis code will be released at https://github.com/nicholasdavies/cfrvoc. Analysis data are

held by Public Health England. An anonymised data set which will be released with the code is

in preparation.

Tables
492
493   **Table 1. Characteristics of study subjects and prevalence of SGTF among study**
494   **subjects.**
495

	All	SGTF	Non-SGTF	SGTF prevalence
	858,181 (100%)	411,961 (100%)	446,220 (100%)	411,961 / 858,181 (48%)
Sex				
Female	449,688 (52.4%)	213,374 (51.8%)	236,314 (53%)	213,374 / 449,688 (47.4%)
Male	408,493 (47.6%)	198,587 (48.2%)	209,906 (47%)	198,587 / 408,493 (48.6%)
Age				
1–34	389,154 (45.3%)	188,967 (45.9%)	200,187 (44.9%)	188,967 / 389,154 (48.6%)
35–54	301,960 (35.2%)	147,747 (35.9%)	154,213 (34.6%)	147,747 / 301,960 (48.9%)
55–69	128,844 (15%)	59,664 (14.5%)	69,180 (15.5%)	59,664 / 128,844 (46.3%)
70–84	32,472 (3.8%)	13,495 (3.3%)	18,977 (4.3%)	13,495 / 32,472 (41.6%)
85 and older	5,751 (0.7%)	2,088 (0.5%)	3,663 (0.8%)	2,088 / 5,751 (36.3%)
Place of residence				
Residential	824,880 (96.1%)	397,203 (96.4%)	427,677 (95.8%)	397,203 / 824,880 (48.2%)
Care/Nursing home	4,669 (0.5%)	1,435 (0.3%)	3,234 (0.7%)	1,435 / 4,669 (30.7%)
Other/Unknown	28,632 (3.3%)	13,323 (3.2%)	15,309 (3.4%)	13,323 / 28,632 (46.5%)
Ethnicity				
White	638,520 (74.4%)	297,863 (72.3%)	340,657 (76.3%)	297,863 / 638,520 (46.6%)
Asian	125,676 (14.6%)	59,512 (14.4%)	66,164 (14.8%)	59,512 / 125,676 (47.4%)
Black	33,848 (3.9%)	21,028 (5.1%)	12,820 (2.9%)	21,028 / 33,848 (62.1%)
Other/Mixed/Unknown	60,137 (7%)	33,558 (8.1%)	26,579 (6%)	33,558 / 60,137 (55.8%)
Index of Multiple Deprivation decile				

1	105,211 (12.3%)	35,913 (8.7%)	69,298 (15.5%)	35,913 / 105,211 (34.1%)
2	103,249 (12%)	46,668 (11.3%)	56,581 (12.7%)	46,668 / 103,249 (45.2%)
3	100,018 (11.7%)	48,777 (11.8%)	51,241 (11.5%)	48,777 / 100,018 (48.8%)
4	90,559 (10.6%)	45,849 (11.1%)	44,710 (10%)	45,849 / 90,559 (50.6%)
5	84,802 (9.9%)	43,639 (10.6%)	41,163 (9.2%)	43,639 / 84,802 (51.5%)
6	78,761 (9.2%)	40,317 (9.8%)	38,444 (8.6%)	40,317 / 78,761 (51.2%)
7	77,367 (9%)	38,565 (9.4%)	38,802 (8.7%)	38,565 / 77,367 (49.8%)
8	77,017 (9%)	38,050 (9.2%)	38,967 (8.7%)	38,050 / 77,017 (49.4%)
9	74,305 (8.7%)	38,257 (9.3%)	36,048 (8.1%)	38,257 / 74,305 (51.5%)
10	66,892 (7.8%)	35,926 (8.7%)	30,966 (6.9%)	35,926 / 66,892 (53.7%)
NHS England region				
East of England	75,614 (8.8%)	54,339 (13.2%)	21,275 (4.8%)	54,339 / 75,614 (71.9%)
London	166,115 (19.4%)	117,899 (28.6%)	48,216 (10.8%)	117,899 / 166,115 (71%)
Midlands	169,911 (19.8%)	61,254 (14.9%)	108,657 (24.4%)	61,254 / 169,911 (36.1%)
North East and Yorkshire	157,965 (18.4%)	37,400 (9.1%)	120,565 (27%)	37,400 / 157,965 (23.7%)
North West	139,909 (16.3%)	43,112 (10.5%)	96,797 (21.7%)	43,112 / 139,909 (30.8%)
South East	120,712 (14.1%)	86,656 (21%)	34,056 (7.6%)	86,656 / 120,712 (71.8%)
South West	27,955 (3.3%)	11,301 (2.7%)	16,654 (3.7%)	11,301 / 27,955 (40.4%)
Specimen date				
1 Nov–14 Nov	164,495 (19.2%)	8,029 (1.9%)	156,466 (35.1%)	8,029 / 164,495 (4.9%)
15 Nov–28 Nov	111,198 (13%)	12,243 (3%)	98,955 (22.2%)	12,243 / 111,198 (11%)
29 Nov–12 Dec	104,865 (12.2%)	37,910 (9.2%)	66,955 (15%)	37,910 / 104,865 (36.2%)
13 Dec–26 Dec	169,491 (19.8%)	111,367 (27%)	58,124 (13%)	111,367 / 169,491 (65.7%)
27 Dec–11 Jan	308,132 (35.9%)	242,412 (58.8%)	65,720 (14.7%)	242,412 / 308,132 (78.7%)

Table 2. Deaths within 28 days of a positive test among study subjects.

	All	SGTF	Non-SGTF
	2,091 (100%)	687 (100%)	1,404 (100%)
Sex			
Female	895 (42.8%)	283 (41.2%)	612 (43.6%)
Male	1,196 (57.2%)	404 (58.8%)	792 (56.4%)
Age			
1–34	19 (0.9%)	6 (0.9%)	13 (0.9%)
35–54	141 (6.7%)	57 (8.3%)	84 (6%)
55–69	435 (20.8%)	173 (25.2%)	262 (18.7%)
70–84	869 (41.6%)	287 (41.8%)	582 (41.5%)
85 and older	627 (30%)	164 (23.9%)	463 (33%)
Place of residence			
Residential	1,569 (75%)	547 (79.6%)	1,022 (72.8%)
Care/Nursing home	475 (22.7%)	123 (17.9%)	352 (25.1%)
Other/Unknown	47 (2.2%)	17 (2.5%)	30 (2.1%)
Ethnicity			
White	1,705 (81.5%)	532 (77.4%)	1,173 (83.5%)
Asian	278 (13.3%)	109 (15.9%)	169 (12%)
Black	54 (2.6%)	28 (4.1%)	26 (1.9%)
Other/Mixed/Unknown	54 (2.6%)	18 (2.6%)	36 (2.6%)
Index of Multiple Deprivation decile			
1	341 (16.3%)	72 (10.5%)	269 (19.2%)
2	281 (13.4%)	88 (12.8%)	193 (13.7%)
3	225 (10.8%)	53 (7.7%)	172 (12.3%)
4	206 (9.9%)	81 (11.8%)	125 (8.9%)
5	176 (8.4%)	70 (10.2%)	106 (7.5%)

6	196 (9.4%)	81 (11.8%)	115 (8.2%)
7	157 (7.5%)	61 (8.9%)	96 (6.8%)
8	184 (8.8%)	73 (10.6%)	111 (7.9%)
9	171 (8.2%)	56 (8.2%)	115 (8.2%)
10	154 (7.4%)	52 (7.6%)	102 (7.3%)
NHS England region			
East of England	151 (7.2%)	85 (12.4%)	66 (4.7%)
London	229 (11%)	162 (23.6%)	67 (4.8%)
Midlands	445 (21.3%)	80 (11.6%)	365 (26%)
North East and Yorkshire	603 (28.8%)	91 (13.2%)	512 (36.5%)
North West	333 (15.9%)	64 (9.3%)	269 (19.2%)
South East	248 (11.9%)	187 (27.2%)	61 (4.3%)
South West	82 (3.9%)	18 (2.6%)	64 (4.6%)
Specimen date			
1 Nov–14 Nov	548 (26.2%)	20 (2.9%)	528 (37.6%)
15 Nov–28 Nov	339 (16.2%)	26 (3.8%)	313 (22.3%)
29 Nov–12 Dec	390 (18.7%)	124 (18%)	266 (18.9%)
13 Dec–26 Dec	475 (22.7%)	281 (40.9%)	194 (13.8%)
27 Dec–11 Jan	339 (16.2%)	236 (34.4%)	103 (7.3%)

498
499

500 **Table 3. Rates of death within 28 days of positive test among study subjects.** Total
501 number of deaths, number of days of followup, and deaths per 10,000 days of followup
502 reported.
503

	All	SGTF	Non-SGTF
	2,091 / 16,776,166 (1.25)	687 / 5,882,304 (1.17)	1,404 / 10,893,861 (1.29)
Sex			
Female	895 / 8,823,354 (1.01)	283 / 3,055,682 (0.93)	612 / 5,767,671 (1.06)
Male	1,196 / 7,952,812 (1.5)	404 / 2,826,622 (1.43)	792 / 5,126,190 (1.55)
Age			
1–34	19 / 7,693,867 (0.02)	6 / 2,767,823 (0.02)	13 / 4,926,044 (0.03)
35–54	141 / 5,907,410 (0.24)	57 / 2,133,536 (0.27)	84 / 3,773,873 (0.22)
55–69	435 / 2,458,532 (1.77)	173 / 789,704 (2.19)	262 / 1,668,828 (1.57)
70–84	869 / 615,306 (14.12)	287 / 168,523 (17.03)	582 / 446,784 (13.03)
85 and older	627 / 101,050 (62.05)	164 / 22,718 (72.19)	463 / 78,332 (59.11)
Place of residence			
Residential	1,569 / 16,108,070 (0.97)	547 / 5,668,687 (0.96)	1,022 / 10,439,383 (0.98)
Care/Nursing home	475 / 87,001 (54.6)	123 / 17,672 (69.6)	352 / 69,329 (50.77)
Other/Unknown	47 / 581,094 (0.81)	17 / 195,946 (0.87)	30 / 385,149 (0.78)
Ethnicity			
White	1,705 / 12,541,963 (1.36)	532 / 4,275,094 (1.24)	1,173 / 8,266,870 (1.42)
Asian	278 / 2,484,370 (1.12)	109 / 827,494 (1.32)	169 / 1,656,876 (1.02)
Black	54 / 598,024 (0.9)	28 / 286,850 (0.98)	26 / 311,174 (0.84)
Other/Mixed/Unknown	54 / 1,151,809 (0.47)	18 / 492,868 (0.37)	36 / 658,941 (0.55)
Index of Multiple Deprivation decile			
1	341 / 2,061,034 (1.65)	72 / 393,858 (1.83)	269 / 1,667,175 (1.61)
2	281 / 2,003,138 (1.4)	88 / 622,894 (1.41)	193 / 1,380,244 (1.4)
3	225 / 1,943,584 (1.16)	53 / 689,400 (0.77)	172 / 1,254,184 (1.37)
4	206 / 1,746,970 (1.18)	81 / 652,995 (1.24)	125 / 1,093,974 (1.14)
5	176 / 1,645,732 (1.07)	70 / 642,930 (1.09)	106 / 1,002,802 (1.06)
6	196 / 1,539,366 (1.27)	81 / 598,790 (1.35)	115 / 940,576 (1.22)

7	157 / 1,528,348 (1.03)	61 / 577,502 (1.06)	96 / 950,846 (1.01)
8	184 / 1,517,480 (1.21)	73 / 565,228 (1.29)	111 / 952,252 (1.17)
9	171 / 1,473,780 (1.16)	56 / 586,332 (0.96)	115 / 887,449 (1.3)
10	154 / 1,316,735 (1.17)	52 / 552,376 (0.94)	102 / 764,360 (1.33)
NHS England region			
East of England	151 / 1,389,416 (1.09)	85 / 865,050 (0.98)	66 / 524,366 (1.26)
London	229 / 3,208,122 (0.71)	162 / 1,976,046 (0.82)	67 / 1,232,076 (0.54)
Midlands	445 / 3,314,751 (1.34)	80 / 635,712 (1.26)	365 / 2,679,039 (1.36)
North East and Yorkshire	603 / 3,393,368 (1.78)	91 / 418,625 (2.17)	512 / 2,974,742 (1.72)
North West	333 / 2,603,784 (1.28)	64 / 372,149 (1.72)	269 / 2,231,635 (1.21)
South East	248 / 2,327,066 (1.07)	187 / 1,472,920 (1.27)	61 / 854,146 (0.71)
South West	82 / 539,660 (1.52)	18 / 141,804 (1.27)	64 / 397,856 (1.61)
Specimen date			
1 Nov–14 Nov	548 / 4,598,236 (1.19)	20 / 224,551 (0.89)	528 / 4,373,686 (1.21)
15 Nov–28 Nov	339 / 3,108,314 (1.09)	26 / 342,473 (0.76)	313 / 2,765,842 (1.13)
29 Nov–12 Dec	390 / 2,930,586 (1.33)	124 / 1,059,924 (1.17)	266 / 1,870,662 (1.42)
13 Dec–26 Dec	475 / 3,806,607 (1.25)	281 / 2,463,718 (1.14)	194 / 1,342,890 (1.44)
27 Dec–11 Jan	339 / 2,332,422 (1.45)	236 / 1,791,640 (1.32)	103 / 540,782 (1.9)

505 **Table 4. Rates of death within any time period following positive test among study**
506 **subjects.** Total number of deaths, number of days of followup, and deaths per 10,000 days of
507 followup reported.
508

	All	SGTF	Non-SGTF
	2,315 / 26,176,060 (0.88)	705 / 6,696,880 (1.05)	1,610 / 19,479,180 (0.83)
Sex			
Female	988 / 13,786,478 (0.72)	290 / 3,481,554 (0.83)	698 / 10,304,925 (0.68)
Male	1,327 / 12,389,581 (1.07)	415 / 3,215,326 (1.29)	912 / 9,174,255 (0.99)
Age			
1–34	23 / 11,944,753 (0.02)	6 / 3,162,130 (0.02)	17 / 8,782,623 (0.02)
35–54	164 / 9,153,070 (0.18)	60 / 2,421,846 (0.25)	104 / 6,731,224 (0.15)
55–69	510 / 3,918,577 (1.3)	178 / 895,650 (1.99)	332 / 3,022,926 (1.1)
70–84	942 / 1,000,954 (9.41)	293 / 191,878 (15.27)	649 / 809,076 (8.02)
85 and older	676 / 158,706 (42.59)	168 / 25,376 (66.21)	508 / 133,331 (38.1)
Place of residence			
Residential	1,758 / 25,106,189 (0.7)	560 / 6,451,174 (0.87)	1,198 / 18,655,015 (0.64)
Care/Nursing home	508 / 141,083 (36.01)	127 / 20,434 (62.15)	381 / 120,649 (31.58)
Other/Unknown	49 / 928,788 (0.53)	18 / 225,272 (0.8)	31 / 703,516 (0.44)
Ethnicity			
White	1,893 / 19,790,819 (0.96)	548 / 4,894,532 (1.12)	1,345 / 14,896,288 (0.9)
Asian	303 / 3,832,494 (0.79)	109 / 926,274 (1.18)	194 / 2,906,220 (0.67)
Black	61 / 851,538 (0.72)	30 / 322,224 (0.93)	31 / 529,314 (0.59)
Other/Mixed/Unknown	58 / 1,701,208 (0.34)	18 / 553,850 (0.32)	40 / 1,147,358 (0.35)
Index of Multiple Deprivation decile			
1	380 / 3,465,168 (1.1)	73 / 454,148 (1.61)	307 / 3,011,020 (1.02)
2	312 / 3,185,588 (0.98)	90 / 707,200 (1.27)	222 / 2,478,388 (0.9)
3	244 / 3,017,818 (0.81)	53 / 784,698 (0.68)	191 / 2,233,120 (0.86)
4	225 / 2,688,858 (0.84)	84 / 741,152 (1.13)	141 / 1,947,706 (0.72)
5	200 / 2,519,846 (0.79)	72 / 735,742 (0.98)	128 / 1,784,104 (0.72)
6	214 / 2,352,866 (0.91)	82 / 682,912 (1.2)	132 / 1,669,954 (0.79)

7	181 / 2,362,076 (0.77)	62 / 659,002 (0.94)	119 / 1,703,074 (0.7)
8	199 / 2,348,292 (0.85)	76 / 642,545 (1.18)	123 / 1,705,746 (0.72)
9	190 / 2,245,194 (0.85)	58 / 663,816 (0.87)	132 / 1,581,377 (0.83)
10	170 / 1,990,355 (0.85)	55 / 625,664 (0.88)	115 / 1,364,690 (0.84)
NHS England region			
East of England	168 / 1,842,747 (0.91)	87 / 960,092 (0.91)	81 / 882,656 (0.92)
London	244 / 4,322,423 (0.56)	167 / 2,219,614 (0.75)	77 / 2,102,810 (0.37)
Midlands	500 / 5,358,479 (0.93)	82 / 702,650 (1.17)	418 / 4,655,829 (0.9)
North East and Yorkshire	680 / 6,080,990 (1.12)	94 / 489,701 (1.92)	586 / 5,591,290 (1.05)
North West	372 / 4,457,542 (0.83)	64 / 399,443 (1.6)	308 / 4,058,099 (0.76)
South East	263 / 3,250,598 (0.81)	193 / 1,760,200 (1.1)	70 / 1,490,397 (0.47)
South West	88 / 863,280 (1.02)	18 / 165,180 (1.09)	70 / 698,100 (1)
Specimen date			
1 Nov–14 Nov	676 / 10,581,858 (0.64)	23 / 509,274 (0.45)	653 / 10,072,584 (0.65)
15 Nov–28 Nov	400 / 5,691,072 (0.7)	32 / 609,924 (0.52)	368 / 5,081,148 (0.72)
29 Nov–12 Dec	425 / 3,752,496 (1.13)	133 / 1,315,496 (1.01)	292 / 2,437,001 (1.2)
13 Dec–26 Dec	475 / 3,818,211 (1.24)	281 / 2,470,546 (1.14)	194 / 1,347,664 (1.44)
27 Dec–11 Jan	339 / 2,332,422 (1.45)	236 / 1,791,640 (1.32)	103 / 540,782 (1.9)

509
510

Table 5. Absolute risk associated with SGTF and p_{VOC}, as expressed by case fatality ratio (%) among individuals testing positive in the community. The baseline absolute risk after 28 days and 60 days post-test is derived using linked deaths for all individuals testing positive in the community from 1 August–31 October 2020.

Sex	Age	Baseline, 28 days	SGTF, 28 days	pVOC, 28 days	Baseline, 60 days	SGTF, 60 days	pVOC, 60 days
Female	0-34	0.00069 (0.00069-0.00069)	0.000900 (0.000741-0.00106)	0.000931 (0.000752-0.00111)	0.00138 (0.00138-0.00138)	0.00175 (0.00145-0.00205)	0.00182 (0.00148-0.00216)
Female	35-54	0.0326 (0.0326-0.0326)	0.0425 (0.035-0.0501)	0.0439 (0.0355-0.0524)	0.0396 (0.0396-0.0396)	0.0502 (0.0416-0.0588)	0.0521 (0.0424-0.0618)
Female	55-69	0.183 (0.183-0.183)	0.238 (0.196-0.28)	0.246 (0.199-0.293)	0.254 (0.254-0.254)	0.322 (0.267-0.377)	0.334 (0.272-0.396)
Female	70-84	2.88 (2.88-2.88)	3.74 (3.09-4.39)	3.86 (3.14-4.59)	3.43 (3.43-3.43)	4.33 (3.61-5.06)	4.49 (3.67-5.31)
Female	85 and older	12.8 (12.8-12.8)	16.4 (13.7-19)	16.8 (13.9-19.8)	15.9 (15.8-15.9)	19.7 (16.6-22.7)	20.3 (16.9-23.7)
Male	0-34	0.00306 (0.00306-0.00306)	0.00399 (0.00329-0.0047)	0.00413 (0.00333-0.00492)	0.00459 (0.00459-0.00459)	0.00582 (0.00483-0.00682)	0.00605 (0.00492-0.00717)
Male	35-54	0.0627 (0.0627-0.0627)	0.0818 (0.0673-0.0963)	0.0845 (0.0683-0.101)	0.0867 (0.0867-0.0868)	0.11 (0.0912-0.129)	0.114 (0.0928-0.135)
Male	55-69	0.559 (0.559-0.56)	0.729 (0.6-0.858)	0.753 (0.609-0.897)	0.734 (0.734-0.734)	0.93 (0.772-1.09)	0.965 (0.785-1.14)
Male	70-84	4.7 (4.7-4.7)	6.08 (5.04-7.12)	6.28 (5.11-7.44)	5.62 (5.62-5.62)	7.07 (5.91-8.24)	7.33 (6.02-8.65)
Male	85 and older	17.1 (17.1-17.1)	21.7 (18.3-25.1)	22.3 (18.6-26.1)	20.5 (20.4-20.5)	25.2 (21.5-28.9)	26 (21.9-30.2)

517   **Supplementary tables**

519   **Table S1. Characteristics of study subjects, including missing SGTF status.** Specimens
520   from 1 Nov 2020 to 11 January 2021.

	All	Missing	SGTF	Non-SGTF
	1,625,121 (100%)	766,940 (100%)	411,961 (100%)	446,220 (100%)
Sex				
Female	874,833 (53.8%)	425,145 (55.4%)	213,374 (51.8%)	236,314 (53%)
Male	750,288 (46.2%)	341,795 (44.6%)	198,587 (48.2%)	209,906 (47%)
Age				
1–34	729,865 (44.9%)	340,711 (44.4%)	188,967 (45.9%)	200,187 (44.9%)
35–54	556,849 (34.3%)	254,889 (33.2%)	147,747 (35.9%)	154,213 (34.6%)
55–69	240,984 (14.8%)	112,140 (14.6%)	59,664 (14.5%)	69,180 (15.5%)
70–84	69,560 (4.3%)	37,088 (4.8%)	13,495 (3.3%)	18,977 (4.3%)
85 and older	27,863 (1.7%)	22,112 (2.9%)	2,088 (0.5%)	3,663 (0.8%)
Place of residence				
Residential	1,518,839 (93.5%)	693,959 (90.5%)	397,203 (96.4%)	427,677 (95.8%)
Care/Nursing home	47,104 (2.9%)	42,435 (5.5%)	1,435 (0.3%)	3,234 (0.7%)
Other/Unknown	59,178 (3.6%)	30,546 (4%)	13,323 (3.2%)	15,309 (3.4%)
Ethnicity				
White	1,202,741 (74%)	564,221 (73.6%)	297,863 (72.3%)	340,657 (76.3%)
Asian	223,071 (13.7%)	97,395 (12.7%)	59,512 (14.4%)	66,164 (14.8%)
Black	73,628 (4.5%)	39,780 (5.2%)	21,028 (5.1%)	12,820 (2.9%)
Other/Mixed/Unknown	125,681 (7.7%)	65,544 (8.5%)	33,558 (8.1%)	26,579 (6%)
Index of Multiple Deprivation decile				
1	162,761 (10%)	57,550 (7.5%)	35,913 (8.7%)	69,298 (15.5%)
2	190,790 (11.7%)	87,541 (11.4%)	46,668 (11.3%)	56,581 (12.7%)
3	194,098 (11.9%)	94,080 (12.3%)	48,777 (11.8%)	51,241 (11.5%)
4	178,493 (11%)	87,934 (11.5%)	45,849 (11.1%)	44,710 (10%)

5	167,919 (10.3%)	83,117 (10.8%)	43,639 (10.6%)	41,163 (9.2%)
6	158,095 (9.7%)	79,334 (10.3%)	40,317 (9.8%)	38,444 (8.6%)
7	151,551 (9.3%)	74,184 (9.7%)	38,565 (9.4%)	38,802 (8.7%)
8	148,871 (9.2%)	71,854 (9.4%)	38,050 (9.2%)	38,967 (8.7%)
9	144,127 (8.9%)	69,822 (9.1%)	38,257 (9.3%)	36,048 (8.1%)
10	128,416 (7.9%)	61,524 (8%)	35,926 (8.7%)	30,966 (6.9%)
NHS England region				
East of England	205,900 (12.7%)	130,286 (17%)	54,339 (13.2%)	21,275 (4.8%)
London	388,515 (23.9%)	222,400 (29%)	117,899 (28.6%)	48,216 (10.8%)
Midlands	287,007 (17.7%)	117,096 (15.3%)	61,254 (14.9%)	108,657 (24.4%)
North East and Yorkshire	203,332 (12.5%)	45,367 (5.9%)	37,400 (9.1%)	120,565 (27%)
North West	182,175 (11.2%)	42,266 (5.5%)	43,112 (10.5%)	96,797 (21.7%)
South East	266,809 (16.4%)	146,097 (19%)	86,656 (21%)	34,056 (7.6%)
South West	91,383 (5.6%)	63,428 (8.3%)	11,301 (2.7%)	16,654 (3.7%)
Specimen date				
1 Nov–14 Nov	251,518 (15.5%)	87,023 (11.3%)	8,029 (1.9%)	156,466 (35.1%)
15 Nov–28 Nov	168,921 (10.4%)	57,723 (7.5%)	12,243 (3%)	98,955 (22.2%)
29 Nov–12 Dec	166,505 (10.2%)	61,640 (8%)	37,910 (9.2%)	66,955 (15%)
13 Dec–26 Dec	356,566 (21.9%)	187,075 (24.4%)	111,367 (27%)	58,124 (13%)
27 Dec–11 Jan	681,611 (41.9%)	373,479 (48.7%)	242,412 (58.8%)	65,720 (14.7%)

523   **Table S2. Deaths within 28 days of a positive test among study subjects, including**
524   **missing SGTF status**
525

	All	Missing	SGTF	Non-SGTF
	7,459 (100%)	5,368 (100%)	687 (100%)	1,404 (100%)
Sex				
Female	3,909 (52.4%)	3,014 (56.1%)	283 (41.2%)	612 (43.6%)
Male	3,550 (47.6%)	2,354 (43.9%)	404 (58.8%)	792 (56.4%)
Age				
1–34	32 (0.4%)	13 (0.2%)	6 (0.9%)	13 (0.9%)
35–54	272 (3.6%)	131 (2.4%)	57 (8.3%)	84 (6%)
55–69	814 (10.9%)	379 (7.1%)	173 (25.2%)	262 (18.7%)
70–84	2,566 (34.4%)	1,697 (31.6%)	287 (41.8%)	582 (41.5%)
85 and older	3,775 (50.6%)	3,148 (58.6%)	164 (23.9%)	463 (33%)
Place of residence				
Residential	2,702 (36.2%)	1,133 (21.1%)	547 (79.6%)	1,022 (72.8%)
Care/Nursing home	4,555 (61.1%)	4,080 (76%)	123 (17.9%)	352 (25.1%)
Other/Unknown	202 (2.7%)	155 (2.9%)	17 (2.5%)	30 (2.1%)
Ethnicity				
White	6,652 (89.2%)	4,947 (92.2%)	532 (77.4%)	1,173 (83.5%)
Asian	522 (7%)	244 (4.5%)	109 (15.9%)	169 (12%)
Black	116 (1.6%)	62 (1.2%)	28 (4.1%)	26 (1.9%)
Other/Mixed/Unknown	169 (2.3%)	115 (2.1%)	18 (2.6%)	36 (2.6%)
Index of Multiple Deprivation decile				
1	789 (10.6%)	448 (8.3%)	72 (10.5%)	269 (19.2%)
2	813 (10.9%)	532 (9.9%)	88 (12.8%)	193 (13.7%)
3	769 (10.3%)	544 (10.1%)	53 (7.7%)	172 (12.3%)
4	787 (10.6%)	581 (10.8%)	81 (11.8%)	125 (8.9%)

5	753 (10.1%)	577 (10.7%)	70 (10.2%)	106 (7.5%)
6	741 (9.9%)	545 (10.2%)	81 (11.8%)	115 (8.2%)
7	752 (10.1%)	595 (11.1%)	61 (8.9%)	96 (6.8%)
8	711 (9.5%)	527 (9.8%)	73 (10.6%)	111 (7.9%)
9	747 (10%)	576 (10.7%)	56 (8.2%)	115 (8.2%)
10	597 (8%)	443 (8.3%)	52 (7.6%)	102 (7.3%)
NHS England region				
East of England	848 (11.4%)	697 (13%)	85 (12.4%)	66 (4.7%)
London	744 (10%)	515 (9.6%)	162 (23.6%)	67 (4.8%)
Midlands	1,688 (22.6%)	1,243 (23.2%)	80 (11.6%)	365 (26%)
North East and Yorkshire	1,351 (18.1%)	748 (13.9%)	91 (13.2%)	512 (36.5%)
North West	797 (10.7%)	464 (8.6%)	64 (9.3%)	269 (19.2%)
South East	1,443 (19.3%)	1,195 (22.3%)	187 (27.2%)	61 (4.3%)
South West	588 (7.9%)	506 (9.4%)	18 (2.6%)	64 (4.6%)
Specimen date				
1 Nov–14 Nov	1,463 (19.6%)	915 (17%)	20 (2.9%)	528 (37.6%)
15 Nov–28 Nov	1,261 (16.9%)	922 (17.2%)	26 (3.8%)	313 (22.3%)
29 Nov–12 Dec	1,406 (18.8%)	1,016 (18.9%)	124 (18%)	266 (18.9%)
13 Dec–26 Dec	1,838 (24.6%)	1,363 (25.4%)	281 (40.9%)	194 (13.8%)
27 Dec–11 Jan	1,491 (20%)	1,152 (21.5%)	236 (34.4%)	103 (7.3%)

527 **Table S3. Rates of death within 28 days of positive test among study subjects, including**
528 **missing SGTF status.** Total number of deaths, number of days of followup, and deaths per
529 10,000 days of followup reported.
530

	All	Missing	SGTF	Non-SGTF
	7,459 / 29,601,000 (2.52)	5,368 / 12,824,834 (4.19)	687 / 5,882,304 (1.17)	1,404 / 10,893,861 (1.29)
Sex				
Female	3,909 / 15,961,968 (2.45)	3,014 / 7,138,615 (4.22)	283 / 3,055,682 (0.93)	612 / 5,767,671 (1.06)
Male	3,550 / 13,639,032 (2.6)	2,354 / 5,686,220 (4.14)	404 / 2,826,622 (1.43)	792 / 5,126,190 (1.55)
Age				
1–34	32 / 13,481,680 (0.02)	13 / 5,787,812 (0.02)	6 / 2,767,823 (0.02)	13 / 4,926,044 (0.03)
35–54	272 / 10,167,962 (0.27)	131 / 4,260,552 (0.31)	57 / 2,133,536 (0.27)	84 / 3,773,873 (0.22)
55–69	814 / 4,287,546 (1.9)	379 / 1,829,014 (2.07)	173 / 789,704 (2.19)	262 / 1,668,828 (1.57)
70–84	2,566 / 1,213,482 (21.15)	1,697 / 598,176 (28.37)	287 / 168,523 (17.03)	582 / 446,784 (13.03)
85 and older	3,775 / 450,330 (83.83)	3,148 / 349,280 (90.13)	164 / 22,718 (72.19)	463 / 78,332 (59.11)
Place of residence				
Residential	2,702 / 27,653,420 (0.98)	1,133 / 11,545,350 (0.98)	547 / 5,668,687 (0.96)	1,022 / 10,439,383 (0.98)
Care/Nursing home	4,555 / 844,984 (53.91)	4,080 / 757,984 (53.83)	123 / 17,672 (69.6)	352 / 69,329 (50.77)
Other/Unknown	202 / 1,102,596 (1.83)	155 / 521,502 (2.97)	17 / 195,946 (0.87)	30 / 385,149 (0.78)
Ethnicity				
White	6,652 / 22,097,241 (3.01)	4,947 / 9,555,278 (5.18)	532 / 4,275,094 (1.24)	1,173 / 8,266,870 (1.42)
Asian	522 / 4,102,698 (1.27)	244 / 1,618,328 (1.51)	109 / 827,494 (1.32)	169 / 1,656,876 (1.02)
Black	116 / 1,198,700 (0.97)	62 / 600,677 (1.03)	28 / 286,850 (0.98)	26 / 311,174 (0.84)
Other/Mixed/Unknown	169 / 2,202,360 (0.77)	115 / 1,050,551 (1.09)	18 / 492,868 (0.37)	36 / 658,941 (0.55)
Index of Multiple Deprivation decile				
1	789 / 3,104,612 (2.54)	448 / 1,043,579 (4.29)	72 / 393,858 (1.83)	269 / 1,667,175 (1.61)
2	813 / 3,461,175 (2.35)	532 / 1,458,037 (3.65)	88 / 622,894 (1.41)	193 / 1,380,244 (1.4)
3	769 / 3,481,612 (2.21)	544 / 1,538,028 (3.54)	53 / 689,400 (0.77)	172 / 1,254,184 (1.37)
4	787 / 3,180,420 (2.47)	581 / 1,433,450 (4.05)	81 / 652,995 (1.24)	125 / 1,093,974 (1.14)
5	753 / 3,015,647 (2.5)	577 / 1,369,916 (4.21)	70 / 642,930 (1.09)	106 / 1,002,802 (1.06)
6	741 / 2,847,643 (2.6)	545 / 1,308,278 (4.17)	81 / 598,790 (1.35)	115 / 940,576 (1.22)

7	752 / 2,777,872 (2.71)	595 / 1,249,524 (4.76)	61 / 577,502 (1.06)	96 / 950,846 (1.01)
8	711 / 2,726,232 (2.61)	527 / 1,208,752 (4.36)	73 / 565,228 (1.29)	111 / 952,252 (1.17)
9	747 / 2,650,394 (2.82)	576 / 1,176,614 (4.9)	56 / 586,332 (0.96)	115 / 887,449 (1.3)
10	597 / 2,355,392 (2.53)	443 / 1,038,658 (4.27)	52 / 552,376 (0.94)	102 / 764,360 (1.33)
NHS England region				
East of England	848 / 3,509,872 (2.42)	697 / 2,120,456 (3.29)	85 / 865,050 (0.98)	66 / 524,366 (1.26)
London	744 / 6,636,995 (1.12)	515 / 3,428,873 (1.5)	162 / 1,976,046 (0.82)	67 / 1,232,076 (0.54)
Midlands	1,688 / 5,506,792 (3.07)	1,243 / 2,192,040 (5.67)	80 / 635,712 (1.26)	365 / 2,679,039 (1.36)
North East and Yorkshire	1,351 / 4,309,108 (3.14)	748 / 915,740 (8.17)	91 / 418,625 (2.17)	512 / 2,974,742 (1.72)
North West	797 / 3,350,266 (2.38)	464 / 746,482 (6.22)	64 / 372,149 (1.72)	269 / 2,231,635 (1.21)
South East	1,443 / 4,675,060 (3.09)	1,195 / 2,347,994 (5.09)	187 / 1,472,920 (1.27)	61 / 854,146 (0.71)
South West	588 / 1,612,909 (3.65)	506 / 1,073,250 (4.71)	18 / 141,804 (1.27)	64 / 397,856 (1.61)
Specimen date				
1 Nov–14 Nov	1,463 / 7,020,658 (2.08)	915 / 2,422,422 (3.78)	20 / 224,551 (0.89)	528 / 4,373,686 (1.21)
15 Nov–28 Nov	1,261 / 4,709,996 (2.68)	922 / 1,601,682 (5.76)	26 / 342,473 (0.76)	313 / 2,765,842 (1.13)
29 Nov–12 Dec	1,406 / 4,640,583 (3.03)	1,016 / 1,709,998 (5.94)	124 / 1,059,924 (1.17)	266 / 1,870,662 (1.42)
13 Dec–26 Dec	1,838 / 7,889,946 (2.33)	1,363 / 4,083,340 (3.34)	281 / 2,463,718 (1.14)	194 / 1,342,890 (1.44)
27 Dec–11 Jan	1,491 / 5,339,816 (2.79)	1,152 / 3,007,394 (3.83)	236 / 1,791,640 (1.32)	103 / 540,782 (1.9)

Table S4. Rates of death within any time period following positive test among study
subjects, including missing SGTF status. Total number of deaths, number of days of
followup, and deaths per 10,000 days of followup reported.

	All	Missing	SGTF	Non-SGTF
	8,091 / 43,913,229 (1.84)	5,776 / 17,737,170 (3.26)	705 / 6,696,880 (1.05)	1,610 / 19,479,180 (0.83)
Sex				
Female	4,249 / 23,676,604 (1.79)	3,261 / 9,890,125 (3.3)	290 / 3,481,554 (0.83)	698 / 10,304,925 (0.68)
Male	3,842 / 20,236,626 (1.9)	2,515 / 7,847,044 (3.21)	415 / 3,215,326 (1.29)	912 / 9,174,255 (0.99)
Age				
1–34	36 / 19,921,946 (0.02)	13 / 7,977,194 (0.02)	6 / 3,162,130 (0.02)	17 / 8,782,623 (0.02)
35–54	304 / 14,986,480 (0.2)	140 / 5,833,410 (0.24)	60 / 2,421,846 (0.25)	104 / 6,731,224 (0.15)
55–69	920 / 6,489,470 (1.42)	410 / 2,570,893 (1.59)	178 / 895,650 (1.99)	332 / 3,022,926 (1.1)
70–84	2,773 / 1,855,836 (14.94)	1,831 / 854,882 (21.42)	293 / 191,878 (15.27)	649 / 809,076 (8.02)
85 and older	4,058 / 659,498 (61.53)	3,382 / 500,791 (67.53)	168 / 25,376 (66.21)	508 / 133,331 (38.1)
Place of residence				
Residential	2,965 / 40,917,878 (0.72)	1,207 / 15,811,690 (0.76)	560 / 6,451,174 (0.87)	1,198 / 18,655,015 (0.64)
Care/Nursing home	4,916 / 1,331,478 (36.92)	4,408 / 1,190,394 (37.03)	127 / 20,434 (62.15)	381 / 120,649 (31.58)
Other/Unknown	210 / 1,663,873 (1.26)	161 / 735,086 (2.19)	18 / 225,272 (0.8)	31 / 703,516 (0.44)
Ethnicity				
White	7,225 / 33,130,923 (2.18)	5,332 / 13,340,104 (4)	548 / 4,894,532 (1.12)	1,345 / 14,896,288 (0.9)
Asian	553 / 6,053,338 (0.91)	250 / 2,220,844 (1.13)	109 / 926,274 (1.18)	194 / 2,906,220 (0.67)
Black	126 / 1,635,050 (0.77)	65 / 783,512 (0.83)	30 / 322,224 (0.93)	31 / 529,314 (0.59)
Other/Mixed/Unknown	187 / 3,093,917 (0.6)	129 / 1,392,709 (0.93)	18 / 553,850 (0.32)	40 / 1,147,358 (0.35)
Index of Multiple Deprivation decile				
1	873 / 5,075,570 (1.72)	493 / 1,610,402 (3.06)	73 / 454,148 (1.61)	307 / 3,011,020 (1.02)
2	885 / 5,222,354 (1.69)	573 / 2,036,766 (2.81)	90 / 707,200 (1.27)	222 / 2,478,388 (0.9)
3	826 / 5,107,348 (1.62)	582 / 2,089,530 (2.79)	53 / 784,698 (0.68)	191 / 2,233,120 (0.86)
4	851 / 4,650,239 (1.83)	626 / 1,961,380 (3.19)	84 / 741,152 (1.13)	141 / 1,947,706 (0.72)
5	808 / 4,387,421 (1.84)	608 / 1,867,574 (3.26)	72 / 735,742 (0.98)	128 / 1,784,104 (0.72)
6	798 / 4,139,651 (1.93)	584 / 1,786,786 (3.27)	82 / 682,912 (1.2)	132 / 1,669,954 (0.79)

7	828 / 4,079,280 (2.03)	647 / 1,717,204 (3.77)	62 / 659,002 (0.94)	119 / 1,703,074 (0.7)
8	756 / 4,000,498 (1.89)	557 / 1,652,206 (3.37)	76 / 642,545 (1.18)	123 / 1,705,746 (0.72)
9	812 / 3,846,584 (2.11)	622 / 1,601,391 (3.88)	58 / 663,816 (0.87)	132 / 1,581,377 (0.83)
10	654 / 3,404,284 (1.92)	484 / 1,413,928 (3.42)	55 / 625,664 (0.88)	115 / 1,364,690 (0.84)
NHS England region				
East of England	889 / 4,510,888 (1.97)	721 / 2,668,140 (2.7)	87 / 960,092 (0.91)	81 / 882,656 (0.92)
London	776 / 8,484,137 (0.91)	532 / 4,161,714 (1.28)	167 / 2,219,614 (0.75)	77 / 2,102,810 (0.37)
Midlands	1,869 / 8,929,204 (2.09)	1,369 / 3,570,726 (3.83)	82 / 702,650 (1.17)	418 / 4,655,829 (0.9)
North East and Yorkshire	1,509 / 7,519,277 (2.01)	829 / 1,438,286 (5.76)	94 / 489,701 (1.92)	586 / 5,591,290 (1.05)
North West	893 / 5,612,248 (1.59)	521 / 1,154,706 (4.51)	64 / 399,443 (1.6)	308 / 4,058,099 (0.76)
South East	1,526 / 6,307,276 (2.42)	1,263 / 3,056,679 (4.13)	193 / 1,760,200 (1.1)	70 / 1,490,397 (0.47)
South West	629 / 2,550,199 (2.47)	541 / 1,686,918 (3.21)	18 / 165,180 (1.09)	70 / 698,100 (1)
Specimen date				
1 Nov–14 Nov	1,795 / 16,126,616 (1.11)	1,119 / 5,544,758 (2.02)	23 / 509,274 (0.45)	653 / 10,072,584 (0.65)
15 Nov–28 Nov	1,457 / 8,609,193 (1.69)	1,057 / 2,918,120 (3.62)	32 / 609,924 (0.52)	368 / 5,081,148 (0.72)
29 Nov–12 Dec	1,510 / 5,932,273 (2.55)	1,085 / 2,179,776 (4.98)	133 / 1,315,496 (1.01)	292 / 2,437,001 (1.2)
13 Dec–26 Dec	1,838 / 7,905,330 (2.33)	1,363 / 4,087,120 (3.33)	281 / 2,470,546 (1.14)	194 / 1,347,664 (1.44)
27 Dec–11 Jan	1,491 / 5,339,816 (2.79)	1,152 / 3,007,394 (3.83)	236 / 1,791,640 (1.32)	103 / 540,782 (1.9)

537 **Table S5. Hazard ratios for SGTF / VOC across models.**
538

Model	Parameter	Hazard ratio (95% CI)	P value
SGTF + lin age + lin IMD | NHSE + spec week	sgtf	1.40 (1.23–1.59)	< 0.001
SGTF + lin age + lin IMD | UTLA + spec week	sgtf	1.36 (1.17–1.57)	< 0.001
SGTF + lin age + lin IMD | LTLA + spec week	sgtf	1.34 (1.15–1.56)	< 0.001
SGTF + lin age + lin IMD | NHSE + spec date	sgtf	1.40 (1.23–1.60)	< 0.001
SGTF + lin age + lin IMD | UTLA + spec date	sgtf	1.33 (1.13–1.57)	< 0.001
SGTF + lin age + lin IMD | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003
SGTF + spl age + spl IMD | NHSE + spec week	sgtf	1.40 (1.23–1.59)	< 0.001
SGTF + spl age + spl IMD | UTLA + spec week	sgtf	1.36 (1.17–1.57)	< 0.001
SGTF + spl age + spl IMD | LTLA + spec week	sgtf	1.34 (1.15–1.56)	< 0.001
SGTF + spl age + spl IMD | NHSE + spec date	sgtf	1.40 (1.22–1.60)	< 0.001
SGTF + spl age + spl IMD | UTLA + spec date	sgtf	1.33 (1.13–1.57)	< 0.001
SGTF + spl age + spl IMD | LTLA + spec date	sgtf	1.31 (1.1–1.56)	0.003
Censoring: SGTF_07 + lin age + lin IMD | LTLA + spec week	sgtf	1.06 (0.81–1.38)	0.674
Censoring: SGTF_14 + lin age + lin IMD | LTLA + spec week	sgtf	1.19 (1.00–1.43)	0.052
Censoring: SGTF_21 + lin age + lin IMD | LTLA + spec week	sgtf	1.29 (1.10–1.51)	0.002
Censoring: SGTF_28 + lin age + lin IMD | LTLA + spec week	sgtf	1.34 (1.15–1.56)	< 0.001
Censoring: SGTF_60 + lin age + lin IMD | LTLA + spec week	sgtf	1.32 (1.14–1.53)	< 0.001
Censoring: SGTF999 + lin age + lin IMD | LTLA + spec week	sgtf	1.32 (1.14–1.52)	< 0.001
Time-age interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003
Time^2-age interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.55)	0.004
Time-sex interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003

Time^2-sex interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003
Time-IMD interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003
Time^2-IMD interaction term | LTLA + spec date	sgtf	1.3 (1.09–1.55)	0.004
Time-ethnicity interaction term | LTLA + spec date	sgtf	1.31 (1.09–1.56)	0.003
Time^2-ethnicity interaction term | LTLA + spec date	sgtf	1.31 (1.09–1.56)	0.003
Time-residence interaction term | LTLA + spec date	sgtf	1.30 (1.09–1.56)	0.003
Time^2-residence interaction term | LTLA + spec date	sgtf	1.3 (1.09–1.56)	0.003
Sensitivity: Sep onwards (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.25 (1.05–1.48)	0.011
Sensitivity: Oct onwards (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.25 (1.05–1.48)	0.011
Sensitivity: Nov onwards (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.30 (1.09–1.56)	0.003
Sensitivity: Dec onwards (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.32 (1.09–1.61)	0.005
Sensitivity: Prevalence cutoff (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.32 (1.09–1.61)	0.005
Sensitivity: no registration cutoff (SGTF + lin age + lin IMD | LTLA + spec date)	sgtf	1.48 (1.28–1.70)	< 0.001
Sensitivity: SGTF + lin age + lin IMD + NHSE * spec week	sgtf	1.40 (1.23–1.60)	< 0.001
p_voc + lin age + lin IMD | NHSE + spec week	p_voc	1.48 (1.28–1.70)	< 0.001
p_voc + lin age + lin IMD | UTLA + spec week	p_voc	1.43 (1.22–1.67)	< 0.001
p_voc + lin age + lin IMD | LTLA + spec week	p_voc	1.4 (1.19–1.65)	< 0.001
p_voc + lin age + lin IMD | NHSE + spec date	p_voc	1.48 (1.29–1.71)	< 0.001
p_voc + lin age + lin IMD | UTLA + spec date	p_voc	1.38 (1.16–1.65)	< 0.001
p_voc + lin age + lin IMD | LTLA + spec date	p_voc	1.35 (1.11–1.63)	0.002
p_voc + spl age + spl IMD | NHSE + spec week	p_voc	1.48 (1.28–1.70)	< 0.001
p_voc + spl age + spl IMD | UTLA + spec week	p_voc	1.43 (1.22–1.67)	< 0.001
p_voc + spl age + spl IMD | LTLA + spec week	p_voc	1.41 (1.19–1.65)	< 0.001

p_voc + spl age + spl IMD | NHSE + spec date	p_voc	1.48 (1.28–1.71)	< 0.001
p_voc + spl age + spl IMD | UTLA + spec date	p_voc	1.38 (1.16–1.65)	< 0.001
p_voc + spl age + spl IMD | LTLA + spec date	p_voc	1.35 (1.12–1.64)	0.002

   Supplementary Figures

542
543
544   **Fig. S1. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
545   **other positive SARS-CoV-2 tests by sex.**
546
547

548
549
Fig. S2. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all other positive SARS-CoV-2 tests by age group. Note that the Y axis differs for each panel.

552
553
554 **Fig. S3. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
555 **other positive SARS-CoV-2 tests by residence.** Note that the Y axis differs for each panel.

556
557
Fig. S4. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all
other positive SARS-CoV-2 tests by Ethnicity.

560
561
562 **Fig. S5. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
563 **other positive SARS-CoV-2 tests by Index of Multiple Deprivation decile (IMD).**

565 **Fig. S6. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
566 **other positive SARS-CoV-2 tests by NHS region.**

567
568
569 **Fig. S7. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
570 **other positive SARS-CoV-2 tests by specimen date.**
571

572
573 **Fig. S8. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
574 **other positive SARS-CoV-2 tests by specimen date for cases 35 years of age or older.**

585
586 **Fig. S9. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
587 **other positive SARS-CoV-2 tests by specimen date for cases 55 years of age or older.**
588
589
590
591

592
593 **Fig. S10. Kaplan Meier plots of survival within 60 days of positive test for SGTF versus all**
594 **other positive SARS-CoV-2 tests by specimen date for cases 70 years of age or older.**
595
596

597
Fig. S11. Schoenfeld residuals for survival model by SGTF stratified by LTLA and
specimen date. Model uses linear terms for age and IMD and a 28-day followup. Schoenfeld
residual tests give $P = 0.031$ for SGTF; $P = 0.425$ for age; $P = 0.170$ for sex; $P = 0.603$ for IMD
decile; $P = 0.410$ for ethnicity; $P = 0.728$ for residence type; and $P = 0.244$ globally.
602

Fig. S12. Misclassification model. For each NHS England region, we fit a beta-binomial model
(purple, Modelled SGTF) to the observed SGTF frequencies among Pillar 2 tests (black,
Observed SGTF), which estimates a constant proportion of "false positive" SGTF samples
among non-VOC 202012/01 specimens (orange, Modelled non-VOC SGTF) and a logistically
growing proportion of VOC 202012/01 specimens over time (blue, Modelled VOC). This allows
us to model the conditional probability that a specimen with SGTF represents VOC 202012/01
(teal, P(VOC|SGTF)). For our misclassification survival analysis, p_{VOC} = 0 for non-SGTF
specimens and p_{VOC} = P(VOC|SGTF) for SGTF specimens.

612
613 **Fig. S13. Ct values for SGTF versus other.** The distribution of Ct values for (**a**) ORF1ab and
614 (**b**) N gene targets among specimens collected between 1–11 January 2021.

References

1. Cox, D. R. Regression Models and Life-Tables. *Journal of the Royal Statistical Society: Series B (Methodological)* **34**, 187–202 (1972).

2. Sarah Walker, A. *et al.* Increased infections, but not viral burden, with a new SARS-CoV-2 variant. *medRxiv* 2021.01.13.21249721 (2021).

3. Davies, N. G. *et al.* Estimated transmissibility and severity of novel SARS-CoV-2 Variant of Concern 202012/01 in England. *bioRxiv* (2020) doi:10.1101/2020.12.24.20248822.

4. Department of Health and Social Care. NERVTAG paper on COVID-19 variant of concern B.1.1.7. https://www.gov.uk/government/publications/nervtag-paper-on-covid-19-variant-of-concern-b117 (2021).

5. Levin, A. T. *et al.* Assessing the age specificity of infection fatality rates for COVID-19: systematic review, meta-analysis, and public policy implications. *Eur. J. Epidemiol.* **35**, 1123–1138 (2020).