# HEALTHCARE WORKERS' PSYCHOLOGICAL DISTRESS At EARLY PHASE OF THE COVID-19 PANDEMIC IN MOROCCO

Jihane BELAYACHI, [1,4]; Sarah BENAMMI [2]; Hasnae CHIPPO [3]; Rhita Nechba BENNIS [1] ; Naoufel MADANI [1]; Abdelmalek HRORA[2,3];Redouane ABOUQAL[1,4]

1 Acute Medical Unit, Ibn Sina University Hospital, 10000, Rabat, Morocco
2 Surgical Departement, Ibn Sina University Hospital, 10000, Rabat, Morocco.
3 Hospital Administration, Ibn Sina University Hospital, 10000, Rabat, Morocco.
4 Laboratory of Biostatistics, Clincial, and Epidemiological Research, Faculté de Médecine et de Pharmacie - Université Mohamed V, 10000, Rabat, Morocco

**Corresponding author:**
Pr Redouane Abouqal at Medical Emergency Department, Ibn Sina University Hospital, 10000, Rabat, Morocco, <u>E-mail:</u> abouqal@invivo.edu, <u>Tel:</u> +21261224739; <u>Fax:</u> +21237672558

**Author's mail**
Jihane Belayachi: jihanebelayahi@gmail.com
Sarah Benammi: sara-benammi@hotmail.fr
Hasnae Chippo: chippohasna@gmail.com
Rhita Nechba Bennis: bennirhita@yahoo.fr
Naoufel Madani: naoufelmad@gmail.com
Abdelmalek Hrora: hroramalek@gmail.com
Redouane Abouqal: abouqal@invivo.edu

Conflict of Interest: none declared
Acknowledgments: none to declare
Ethical Consideration & Disclosure: Study protocol was approved by Rabat ethic committee, and informed consent was obtained from all participants
Running Head:  Psychological impact of COVID-19 Pandemic in Healthcare Workers
Abstract words 130

Manuscript Words Count: 849

*Keywords: COVID-19; Medical staff; Psychological distress;*

Pr Redouane Abouqal at Medical Emergency Department, Ibn Sina University Hospital, 10000, Rabat, Morocco, E-mail: abouqal@invivo.edu, Tel: +21261224739; Fax: +21237672558

**Author's mail**
Jihane Belayachi: jihanebelayahi@gmail.com
Sarah Benammi: sara-benammi@hotmail.fr
Hasnae Chippo: chippohasna@gmail.com
Rhita Nechba Bennis: bennirhita@yahoo.fr
Naoufel Madani: naoufelmad@gmail.com
Abdelmalek Hrora: hroramalek@gmail.com
Redouane Abouqal: abouqal@invivo.edu

Conflict of Interest: none declared

Acknowledgments, none declared

Ethical Considerations & Disclosure(s)

Running Head:  Psychological impact of COVID-19 Pandemic in Healthcare Workers

**Abstract words 130**

**MANUSCRIPT WORDS COUNT 995**
*KEYWORDS:*
*COVID-19; Medical staff; Psychological distress;*

## ABSTRACT

The mental being of healthcare workers with the COVID 19 pandemic is a determinant of their resilience. We investigated the psychological impact of healthcare workers during novel Coronavirus 2019-nCoV times in Morocco.

A cross-sectional study conducted among healthcare workers of an University Hospital. We evaluated the knowledge of the protective means against nCov-19, incidence and the degree of anxiety and depression according to health workers characteristic.

Among 287 health workers; 54% have been trained regarding protection procedures, and 94.8% declared that they are aware of individual protection measures.  The incidence of anxiety and depression was respectively 77.4% and 73.9%. High degree of anxiety and depression was associated with female gender. However, Higher degree of anxiety was also related to function, specialty of practice ,and knowledge of the protective measures against COVID-19.

**INTRODUCTION:**

Since its emergence in Wuhan China, the novel Coronavirus Disease 2019 (vCov-19 or COVID-19) has become a nationwide health crisis, indeed classified by the World Health Organization (WHO) as a pandemic since early March 2020(1). To date 27 April 2020, there have been over 2,99 M confirmed cases of nCov-19 or COVID-19 globally, and 207 k deaths worldwide. Amongst reported deaths over 100 healthcare professionals decease have been registered (2). In Morocco we report overall 4.065 confirmed cases,

and 161deaths. With the outbreak of the COVID-19 in China in December 2019 and in Europe in February 2020, the adaptive capacity of a health system appears to be more depending on the relevance and precaution of the preventive measures implemented than on maturity of health care system. On March 19, the Moroccan authorities declared a state of health emergency. In parallel to operations deployed to provide health system with necessary means to fight the Coronavirus, The Moroccan authorities deployed drastic measures including: Reinforcement of containment, and barrier measures, The obligation to wear the mask in public places; and expanded screening of contact; The initiation of mass treatment of cases confirmed by "the combination of chloroquine and Azithromicin. The increasing number of overwhelmed healthcare systems, reported global shortage of protective equipment, and clinical care challenges during these times, create a state of uncertainty regarding the situation progression among population in general and health care workers in particular. All these factors build up to create a high-risk environment to the physical and mental being of healthcare workers. Healthcare workers are one of the key resources of a health system. The mental being of healthcare workers with the COVID 19 pandemic is a determinant of their resilience. Indeed, we suppose that our global current context may become a source of psychological distress upon healthcare workers ready to face nCov-19 pandemic. Therefore, we sought to investigate the psychological burden of healthcare staff during novel Coronavirus 2019-nCoV (COVID-19) times.

## Methods

We conducted a cross-sectional study based on a survey of healthcare workers of the University Hospital Ibn-Sina in Morocco. Data were collected between 23th to 30th March 2020.Of each healthcare professional, sociodemographic, health characteristics

and professional characteristics were collected. We also evaluated the knowledge of the protective means against nCov-19 through 3 questions: 1 Do you know about individual protective measures? 2 What kind of individual protective measures do you know? 3 Did you have a training regarding protection procedures? Finally, we evaluated spiritual behaviour using prayers as a coping reaction to internal fear against novel Covid 19 Coronavirus. The mental health status of the healthcare staff was investigated using the Arabic validated version of HADS 14 items, evaluating hospital anxiety and depression (3). Anxiety or depression (defined as a subscale score>7) was the dependent variable. The univariate analysis was carried out using the students' t-test for the comparison of two groups. Analysis of variance for comparison of more than two groups and Pearson correlation for the link between two quantitative variables

## Résultats

Sociodemographic and health characteristics of 287 included health workers were described in table 1. Anxiety and Depression score showed good consistency with a Cronbach's alpha score was 0.81. The incidence of anxiety and depression was respectively 77.4% and 73.9%. Knowledge and training about individual protective measures against COVID-19 were presented in table 1. Comparison of anxiety and depression intensity according health care worker's characteristics was presented in table 2. Figure 1 showed Mean anxiety scale of health care workers according to protective procedures' knowledge.

## Discussion

We reported the result of the first evaluation of psychological burden of health worker during novel Coronavirus 2019-nCoV (COVID-19) times in Africa. The study showed high frequency of anxiety (77.4%) and depression (73.9%) among Moroccan health

worker in a hospital faced to COVID-19 patient management. High degree of anxiety and depression was associated with female gender. A previous study in a tertiary infectious disease hospital for COVID-19 in China, and concluded that female medical staff members were more vulnerable with high anxiety index (8).

However, High degree of anxiety was also related to function, specialty of practice, and knowledge of the protective measures against COVID-19. Higher anxiety was noted among emergency unit staff members, interventional units, laboratory units and surgical units. These results can be explained by the high risk of exposure proper to each department: direct manipulation of blood and high proximity to sharp instruments for surgical, laboratory and interventional units, and to frontline contact with all arriving patients to the hospital with unknown COVID-19 status regarding emergency units.Our study results go in accordance with previous studies evaluating anxiety and depression among medical staffdemonstrating both high anxiety and depression incidence (8).

## References

1- Ghebreyesus TD (2020) WHO Director-General's openin gremarks at the media briefing on COVID-19 (11 March 2020)
2- In Memoriam: HealthcareWorkersWho Have Died of COVID-19 ; Medscape April 01, 2020
3- Terkawi AS, Tsang S, AlKahtani GJ, Al-Mousa SH, Al Musaed S, AlZoraigi US, Alasfar EM, Doais KS, Abdulrahman A, Altirkawi KA.

Development and validation of Arabic version of theHospital Anxiety and Depress ion Scale. Saudi J Anaesth. 2017 May;11 (Suppl1): S11-S18.

4- Michopoulos, I., Douzenis, A., Kalkavoura, C., Christodoulou, C., Michalopoulou, P., Kalemi, G.,  Fineti K, Patapis P, Protopapas K, Lykouras, L. Hospital Anxiety and Depression Scale (HADS): validation in a Greek general hospital sample. Annals of General Psychiatry (2008). , 7(1), 4. doi:10.1186/1744-859x-7-4

5- Crow R, Gage H, Hampson S, Hart J, Kimber A, Storey L, Thomas H. The measurement of satisfaction withhealthcare: implications for practice from a systematicreview of the literature. Health Technol Assess 2002; 6:241-244.

6- El-Rufaie OE, Absood GH. Retesting the validity of the Arabic version of the Hospital Anxiety and Depression (HAD) scale in primary health care. Soc Psychiatr epidemiol 1995; 30:26-31.

7- Cronbach IJ. Coefficient alpha and the internal structure of tests. Psychometrika 1951;16:297-334. 26.

8- Zhonghua Lao Dong Wei Sheng ZhiYe Bing ZaZhi. Mental health survey of 230 medical staff in a tertiaryinfectiousdiseasehospital for COVID-19]. 2020 Mar 4;38(0):E001. doi: 10.3760/cma.j.cn121094-20200219-00063. [Epubahead of print]

## FIGURE LEDEND:

**Figure 1:** Mean anxiety scale of health care workers according to protective procedures' knowledge.

**Table.1 Characteristics of Healthcare Professionals**

|  | N (%) |
| --- | --- |
| Age Mean (± SD) | 34.4±12.18 |
| Gender |  |
| Female | 185(64.5%) |
| Male | 102(35.5%) |
| Marital Status |  |
| Married | 126(43.9%) |
| Single | 161(56.1%) |
| Years of Practice Median [IQR] | 4[1-16] |
| Comorbidities |  |
| None | 136 (47.4%) |
| Yes | 151 (52.6%) |

| Function | |
|---|---|
| Senior Doctor | 30 (10.6%) |
| Junior DOCTOR | 117 (40.8%) |
| Nurse | 72 (25.1%) |
| Health Technician | 50 (17.4) |
| Administration | 18 ( 6.3%) |
| Departments | |
| Emergency units | 49 (17.1%) |
| Interventional units | 31 (10.8%) |
| Laboratory units | 20 (7.1%) |
| Surgical departments | 60 (21.3%) |
| Medical departments | 65 (23%) |
| No Care departments | 17 (6%) |
| Intensive Care units | 31 (10.8%) |
| Acute Care unit | 9 (3.2%) |
| Training regarding protection procedures | |
| Yes | 155 (54%) |
| No | 132 (46%) |
| Knowledge about protection procedures | |
| Yes | 272 (94.8%) |
| No | 15 (5.2%) |
| Kind of individual protective measures | |
| Hand sanitizing | 236 (82.2%) |
| Mask | 214 (74.6%) |
| Social distancing | 170 (59.2%) |
| Eye Goggles | 109 (38%) |
| Hair Net | 124 (43.2%) |
| Gown | 144 (50.2%) |
| Gloves | 130 (45.3%) |
| No Surface Touching | 112 (39%) |
| Isolation | 107 (37.3%) |
| Shoe protection | 4 (1.4%) |
| Spiritual Behaviour | |
| Yes | 105 (36.6%) |
| No | 182 (63.4%) |
| Anxiety | |
| Yes | 222 (77.4%) |
| No | 65 (22.6%) |
| Depression | |
| Yes | 212 (73.9%) |
| No | 75 (26.1%) |

**Table.2 –Comparison of anxiety and depression intensity according health care worker's characteristics**

| | ANXIETY | | DEPRESSION | |
|---|---|---|---|---|
| | Mean± SD Or (r) | *P* | Mean± SD Or (r) | *P* |
| **Age (r)** | -0.05 | 0.41 | 0,06 | 0.2 |
| **Gender** | | <0.001 | | 0.001 |
| Female | 11.56 ±5. | | 9.46±3,9 | |
| Male | 8.83 ±4.5 | | 7.93±3,7 | |
| **Marital status** | | | | 0.20 |
| Married | 10.74 ±5.4 | 0.65 | 9.25 ±3.8 | |
| Single | 10.47 ±4.7 | | 8.65 ±3.9 | |

| | | | | |
|---|---|---|---|---|
| **Years of practice (r)** | -0.007 | 0.91 | 0.03 | 0.5 |
| **Comorbidities** | | 0.42 | | 0.87 |
| None | 10.84 ±5.2 | | 8.88 ±3.9 | |
| Yes | 10.37 ±4.8 | | 8.95 ±3.8 | |
| **Function** | | 0.04 | | 0.45 |
| Administration | 12.43 ±4.6 | | 8.77 ±4.3 | |
| Nurse | 11.36 ±4.8 | | 9.65 ±3.7 | |
| Health technician | 10.81 ±5.4 | | 8.75 ±3.6 | |
| Junior doctor | 10.26 ±4.7 | | 8.55 ±3.9 | |
| Senior doctor | 8.54 ±5.2 | | 8.92 ±4.2 | |
| **Training regarding protection procedures** | | 0.7 | | 0.21 |
| Yes | 10.67 ±4.8 | | 8.65 ±3.7 | |
| No | 10.49 ±5.2 | | 9.23 ±4.1 | |
| **Knowledge about protection procedures** | | 0.03 | | 0.27 |
| No | 13.25 ±5.3 | | 9.98 ±3.6 | |
| Yes | 10.44 ±4.9 | | 8.86 ±3.9 | |
| **Spiritual Behavior** | | 0.14 | | 0.07 |
| No | 10.67 ±5.2 | | 9.23 ±3.9 | |
| Yes | 10.44 ±4.6 | | 8.38 ±3.7 | |