**TITLE PAGE**

**Title:**

**Performance of a Computational Phenotyping Algorithm for Sarcoidosis Using**

**Diagnostic Codes in Electronic Medical Records: A Pilot Study from Two Veterans**

**Affairs Medical Centers**

**Authors:**

**Mohamed I Seedahmed, MD, MPH[1,2]\*, Izabella Mogilnicka, MD[1,5]\*, Siyang Zeng, MS[1,4],**

**Gang Luo, PhD[4], Charles McCulloch, PhD[3], Laura Koth, MD[2]γ, Mehrdad Arjomandi,**

**MD[1,2]γ**

[1] San Francisco Veterans Affairs Medical Center, San Francisco, California, USA

[2] Division of Pulmonary, Allergy, Critical Care and Sleep Medicine, Department of

Medicine, University of California San Francisco, California, USA

[3] Department of Epidemiology & Biostatistics, University of California San Francisco,

California, USA

[4] Department of Biomedical Informatics and Medical Education, School of Medicine,

University of Washington, Seattle, Washington, USA

[5] Department of Experimental Physiology and Pathophysiology, Laboratory of the Centre for

Preclinical Research, Medical University of Warsaw, Warsaw, Poland

* MIS and IM: These authors contributed equally to this work.

γ MA and LK: These authors share senior authorship.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**Email Addresses | ORCID iDs:**

MIS: mohamed.seedahmed@ucsf.edu | https://orcid.org/0000-0002-7446-7346

IM: izabella.mogilnicka@gmail.com | https://orcid.org/0000-0002-5735-8375

SZ: siyang.zeng@ucsf.edu | https://orcid.org/0000-0001-9346-301X

GL: luogang@uw.edu | https://orcid.org/0000-0001-7217-4008

CM: charles.mcculloch@ucsf.edu | https://orcid.org/0000-0002-1279-6179

LK: laura.koth@ucsf.edu | https://orcid.org/0000-0001-9541-3622

MA: mehrdad.arjomandi@ucsf.edu | https://orcid.org/0000-0002-0116-9217

**Word Count:** 3,829 words.

**Corresponding Author:**

**Mohamed I Seedahmed, MD, MPH**

Division of Pulmonary, Critical care, allergy and Immunology, and Sleep.

Department of Medicine, University of California, San Francisco

University of California San Francisco Helen Diller Medical Center

513 Parnassus Ave.

HSE 1314, Box 0111

San Francisco, CA 94143

Office: (415) 476-0735

Fax: (415) 502-2605

EMAIL: mohamed.seedahmed@ucsf.edu

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**Authors' Contributions:**

All authors read and approved the final manuscript.

Conceived and designed the study research: MIS, LK, MA

Developed study protocol: MIS, LK, MA

Worked on the methods: MIS, IM, SZ, CM, LK, MA

Analyzed and Interpreted data: MIS, IM, SZ, CM, LK, MA

Prepared and/or edit the manuscript: MIS, IM, GL, CM, LK, MA

**Keywords:**

Sarcoidosis, Electronic Medical Records (EMR), Computational Phenotyping, Diagnostic

Codes, Veterans Affairs (VA), ATS Practice Guidelines.

**Highlights**

• Identifying sarcoidosis cases using diagnostic codes in EMR has low accuracy.

• "Unstructured data" contain information useful in identifying cases of sarcoidosis.

• Computational algorithms could improve the accuracy and efficiency of

case identification in EMR.

• We introduce a new scoring system for assessing healthcare providers' compliance with

the American Thoracic Society (ATS) practice guideline.

• Compliance scoring could help automatically assess sarcoidosis patients' care delivery.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**ABSTRACT (250 words)**

**Background-**The accuracy of identifying sarcoidosis cases in electronic medical records

(EMR) using diagnostic codes is unknown.

**Methods-**To estimate the statistical performance of using diagnostic codes, ICD-9 and ICD-

10, to identify sarcoidosis cases in EMR, we searched the San Francisco and Palo Alto

Veterans Affairs (VA) medical centers EMR and randomly selected 200 patients coded as

sarcoidosis. To further improve diagnostic accuracy, we developed an "index of suspicion"

algorithm to identify probable sarcoidosis cases based on clinical and radiographic features.

We then determined the positive predictive value (PPV) of diagnosing sarcoidosis by two

computational methods using ICD only and ICD plus the "index of suspicion" against the gold

standard developed through manual chart review based on the American Thoracic Society

(ATS) practice guideline. Finally, we determined healthcare providers' adherence to the

guidelines using a new scoring system.

**Results-**The PPV of identifying sarcoidosis cases in VA EMR using ICD codes only was 71%

(95%CI=64.7%-77.3%). The inclusion of our construct of "index of suspicion" along with the

ICD codes significantly increased the PPV to 90% (95%CI=85.2%-94.6%). The care of

sarcoidosis patients was more likely to be classified as "Fully" or "Substantially" adherent with

the ATS practice guideline if their managing provider was a specialist (45% of primary care

providers vs. 74% of specialists; P=0.008).

**Conclusions-**Although ICD codes can be used as reasonable classifiers to identify

sarcoidosis cases within EMR, using computational algorithms to extract clinical and

radiographic information ("index of suspicion") from unstructured data could significantly

improve case identification accuracy.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**INTRODUCTION**

Sarcoidosis is a complex disease of unknown etiology that can involve multiple organs, with no universal and standardized measures that fully secure the final diagnosis.[1–3] In fact, only recently the American Thoracic Society (ATS) published its first practice guideline to provide recommendations for diagnosing sarcoidosis and the necessary screening tests.[3] The ATS practice guideline for diagnosis requires the presence of specific clinical and radiographic features, tissue biopsy revealing non-necrotizing granulomas, and exclusion of alternative conditions that can mimic sarcoidosis.[1,3,4]

Electronic medical record (EMR) data are commonly used in research and by healthcare systems, including the Department of Veterans Affairs, to predict outcomes or assess care quality.[5] EMR data are generally captured in two forms: 1) *structured data* such as billing codes like International Classification of Disease (ICD) codes, laboratory test results, procedural codes, and 2) *narrative or unstructured data* such as progress notes, pathology reports, and imaging reports. Unstructured data contain many more details of the clinical conditions, but extracting these details is challenging and time-consuming. In contrast, structured data are easier to search and offer the promise to identify cases computationally using diagnostic codes. But, diagnostic codes can be inaccurate and not easily verifiable, particularly for the case definition of complex diseases such as sarcoidosis.[6–8] Thus, there is a need to develop automated algorithms to verify the final diagnosis of sarcoidosis using unstructured data by incorporating the ATS diagnostic criteria.

The goals of this study are twofold. First, we sought to estimate the statistical performance of using diagnostic codes (ICD-9 and 10) to identify patients with sarcoidosis in the United States Veterans Affairs (VA) EMR through the VA Informatics and Computational Infrastructure (VINCI) via applying a recently published ATS practice guideline. Second, we

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

113 investigate healthcare providers' practice patterns and determine how adherent these patterns

114 are with the new practice guideline issued by the ATS. This study will help researchers and

115 healthcare systems understand the healthcare providers' compliance with these new

116 recommendations and conduct research using administrative databases.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

## METHODS

### Study Design

We determined the statistical performance of using diagnostic codes (the International

Classification of Diseases, Ninth and Tenth Revisions codes; ICD-9 and ICD-10) to identify

patients with sarcoidosis from the EMR. First, by identifying patients with ICD diagnosis code

of sarcoidosis, and then performing a comprehensive chart review of the entire unstructured

data to ascertain the true diagnosis of sarcoidosis. Furthermore, to improve the diagnostic

accuracy using ICD code alone, we generated an "index of suspicion" criteria from

unstructured clinical and radiographic data, but not histopathologic data, as a second decision

point along with ICD code. We then determined the positive predictive value of both the use

of ICD code and ICD code plus "index of suspicion" to determine whether the use of "index

of suspicion" could improve the accuracy of identifying sarcoidosis cases in EMR regardless

of tissue biopsy availability. Finally, we investigated the relevant healthcare utilization among

the identified patients to determine healthcare providers' adherence to the ATS practice

guideline.

### Data Source

This is a retrospective cohort study of electronic medical records (EMR) available from

the Veterans Affairs Informatics and Computational Infrastructure (VINCI). Developed by the

VA Health Services Research & Development (HSR&D) to improve veterans' healthcare,

VINCI provides access to comprehensive and integrated veterans' national datasets that are de-

identified and the necessary computational and analytical tools in a secure, high-performance

computing environment [9]. The University of California San Francisco Institutional Review

Board and the Veterans Health Administration Research and Development Committee

approved this study.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**Data Collection**

We searched the EMR data in VINCI from 1989 to 2019 and identified all patients

coded as sarcoidosis cases in the VA healthcare system, as defined by documentation of the

ICD-9 and 10 codes of 135 and D.86 (including all subcodes), respectively. Data were

extracted through executing Structured Query Language (SQL) queries in a SQL Server 2017

database.

We then performed a comprehensive and thorough manual chart review of the

unstructured data (available clinical notes, pathology reports, radiology reports, and laboratory

test results) to determine the accuracy of the ICD-9 or 10 code-based diagnosis of sarcoidosis

among cases. To limit the required chart review to a manageable level, we reviewed a total of

200 cases out of the 14,833 identified. Because our access to the detailed medical records was

limited to two VA medical centers (San Francisco VA [SFVA] and Palo Alto Medical Centers

[PAVA]), the 200 patients coded as sarcoidosis were selected from those two centers. We

stratified the list of sarcoidosis cases from the two centers by site and used the "lottery" method

to select 100 patients from each site randomly.[10]

Two independent reviewers (MIS, IM) confirmed the diagnoses of sarcoidosis by

performing a manual chart review of the 200 cases in the VA EMR, Computerized Patient

Record System (CPRS), and based on the ATS diagnostic criteria (clinical, radiographic,

pathological findings, and exclusion of other causes).[3] This approach was considered the

"gold standard" methodology for diagnostic accuracy.

Given that in clinical practice, not all suspected patients will have a tissue biopsy, we

generated an "index of suspicion" for sarcoidosis to identify patients with probable sarcoidosis

based on clinical and radiographic information and regardless of the availability of biopsy data.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

The "index of suspicion" was defined based on the interpretation of available unstructured data

from both clinical and radiographic findings without the inclusion of tissue biopsy results if

available and was used to determine the likelihood of an individual having sarcoidosis. Upon

completing the chart review, each patient was assigned to one of two groups of "high index of

suspicion" if the clinical and radiological presentations were supportive of sarcoidosis, or "low

index of suspicion" if clinical and radiological findings were not consistent with sarcoidosis.

We then classified the patients into three groups: (1) Patients with a high index of suspicion

and documented histopathological evidence of non-necrotizing granulomas were categorized

as "probable sarcoidosis with confirmed biopsy"; (2) Those with a high index of suspicion and

either no documented biopsy in the EMR or a biopsy showing no histopathological evidence

of non-necrotizing granulomas were categorized as "probable sarcoidosis without confirmed

biopsy"; and (3) those with a low index of suspicion were classified as "unlikely sarcoidosis."

This approach was used to compare the statistical performance of the two methods (ICD

code alone versus ICD code with "index of suspicion") in identifying sarcoidosis patients from

the EMR. As we started with a random sample of those with sarcoidosis diagnostic codes (ICD

codes 135 or D86), the further restriction of the sample to those with "high index of suspicion"

is still a random sample of the combination of both ICD codes and "index of suspicion."

**Disease-related Variables**

Organ involvement was assessed based on the clinical history obtained from

physicians' notes and imaging and biopsy reports available in CPRS. For this assessment, to

adjust for the variability in providers' documentation, we adapted a set of criteria previously

introduced in the NIH-sponsored Genomic Research in Alpha-1 Antitrypsin Deficiency and

Sarcoidosis (GRADS) Study.[11]

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

We collected the following data from chart review: clinical site (SFVA or PAVA),

gender, race, ICD-9 and 10 code for sarcoidosis (135 and D86), index date of ICD code (entry

date of the ICD code), specialty visits (specialty notes available), the pathological diagnosis

from any available biopsy, organ involvement (as described in **Figure 2**), Scadding staging of

chest x-ray (as described in radiology reports), history of bilateral hilar lymphadenopathy

(based on radiology reports and clinical notes), pulmonary function test (**PFT**) pattern (as

reported in PFT reports), vitamin D levels (25 [OH]D and 1,25[OH]D), and finally the

treatment status of sarcoidosis. Besides, we checked whether other imaging and laboratory

reports were performed and available in EMR as part of diagnostic workup and sarcoidosis

management; including echocardiography, 12-lead electrocardiogram (**ECG**), chest x-ray,

chest computed tomography (**CT**), cardiac magnetic resonance imaging (**MRI**) or positron

emission tomography (**PET**)/ CT, or abdominal CT scan or ultrasound.

Pathological diagnoses were categorized into "primary" histopathological if it was

available in the pathology report domains, or "secondary" if the data was only available in

clinical note domains due to either a remote history of biopsy or the biopsy being performed

outside of the VA. The PFT reports at SFVA and PAVA used Crapo reference equations to

calculate the Lower Limit of Normal (**LLN**) values for spirometry and lung volume

measurements.

Using the clinical data from chart abstraction, we classified the patients into the clinical

phenotypes proposed by the GRADS study with some modifications.[11] Because of the

variability of available data needed to confirm multiple organ involvement from EMR chart

review, we considered three or more organ involvement (instead of five or more as defined in

the GRADS study) to categorize "multiorgan phenotype."

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

Finally, we created a scoring system to estimate the physicians' compliance with the

recently published ATS clinical practice guideline.[3] For this analysis, four variables were

used in the scoring system: (1) having an ophthalmology visit, (2) 12-lead EKG, (3) chest X-

ray, and (4) vitamin D measurements (25-(OH) D and 1,25(OH) D) within one year after the

index date of ICD code. Each variable was given a score of three, and each patient could have

a total score of 12. We classified the physicians' compliance into four categories based on the

score: "Fully," "Substantially," "Partially," or "Non-compliant" for scores of 12, 9-11, 5-8, or

<5 points, respectively. This scoring system was applied to all probable sarcoidosis patients

with or without histopathology confirmation. Furthermore, the role of primary versus specialty

care evaluation and management was examined by comparing the healthcare providers'

compliance score.

### **Statistical Analyses**

All statistical analyses were performed with R software (RStudio, version 1.2.5,

Foundation for Statistical Computing, Vienna, Austria). Descriptive statistics were computed

to summarize the data. Categorical variables were presented as the frequency in percentages

and continuous data as means and standard deviations. We estimated the positive predictive

value (PPV) of the two computational diagnostic criteria for sarcoidosis described above (ICD

codes only and ICD codes along with "index of suspicion"). The PPV of the criterion of using

ICD code only was calculated as the number of patients verified to have sarcoidosis by chart

review ("gold standard" or true positives) divided by the total number of patients with an ICD-

9 or 10 diagnostic code of sarcoidosis. The PPV of the criterion of using ICD codes and "index

of suspicion" use was calculated as the number of patients verified to have sarcoidosis ("gold

standard" or true positives) divided by the total number of patients with a high index of

suspicion. We computed the 95% confidence intervals (CI) using the exact binomial method.

For our estimates, the significance was defined as $P < 0.05$.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

237    To determine whether it is worth interpreting the developed contingency table that

238 compares the healthcare providers' compliance score by primary versus specialty care, we

239 computed the chi-squared test of significance.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**RESULTS**

**Patients' Characteristics**

A total of 14,833 patients with at least one ICD-9 or 10 diagnostic code of sarcoidosis

were identified (**Figure 1**). The study cohort included 200 patients identified by ICD codes of

sarcoidosis: 169 with ICD-9 code of 135 and 39 with ICD-10 code of D.86 (**Table 1**). Of the

200 patients, 158 had a "high index of suspicion" for sarcoidosis based on clinical and

radiographic findings. Of those, 142 had confirmed sarcoidosis based on histopathological

evidence of non-necrotizing granuloma and were classified as "probable sarcoidosis with

confirmed biopsy." The remaining 16 patients with a "high index of suspicion" did not undergo

a biopsy and were classified as "probable sarcoidosis without confirmed biopsy" (**Figure 2**).

No patient had nondiagnostic biopsy results for sarcoidosis.

Among those patients with probable sarcoidosis (with and without confirmed biopsy),

90% (142/158) were males (**Table 1**), and there was a higher representation of African

Americans than non-Hispanic Whites (54% (85/158) vs. 33% (52/158), respectively). Overall,

90% (143/158) had a predominant pulmonary phenotype, and 29% (45/158) had a multi-organ

disease that included pulmonary. Among those with pulmonary phenotype, 28% (36/129), 22%

(28/129), and 19% (25/129) had restrictive, obstructive, and mixed lung function patterns,

respectively. The majority were in Scadding stage II (32%, 47/145), followed by Stage 0 and

Stage 1 (20% (29/145) and 18% (26/145), respectively). There was no significant difference in

age between those who did and those who did not have a biopsy performed to diagnose

sarcoidosis (mean age= 65.5 vs. 69.3, P= 0.18, respectively). In terms of clinical phenotypes,

the most common phenotype was multiorgan (33%, 52/158), followed by stage II or III treated

(29%, 45/158). Our study cohort did not include any individuals with acute presentation (acute,

untreated). Some patients with remitting phenotype overlapped with Groups 2 and 4 (**Table 1**)

(**Figure 3**).

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**Diagnostic Accuracy of ICD Codes**

We then calculated the positive predictive value (PPV) of using ICD codes to identify VA patients that met the ATS definition of sarcoidosis from the VINCI database. For this calculation, we used the curated dataset of 200 patients. The estimated PPV of using only ICD codes was 71% (95%CI= 64.7%-77.3%). The inclusion of our construct of "index of suspicion" along with the ICD codes into the identification of sarcoidosis increased the PPV estimate significantly to 90% (95%CI= 85.2%-94.6%) (**Table 2**), as the initial sarcoidosis cohort was restricted to the patients with "high index of suspicion."

**Providers' Compliance with the ATS Clinical Practice Guideline**

Among those with probable sarcoidosis (with and without biopsy), 13% were managed by primary care providers only, 51% (81/158) were managed by pulmonary physicians only, and pulmonary physicians managed 36% (57/158) along with other specialties such as ophthalmology or cardiology. The specialty care visits occurred in the context of diagnosis or management of the disease.

Within one year of the entry date of the ICD code of sarcoidosis, 91% (143/158) of patients had at least one chest x-ray, 83% (131/158) had at least one EKG, and 84% (133/158) had at least one visit to an ophthalmologist. In addition, 53% (82/158) had 25(OH)D, and 23% (36/158) had 1, 25(OH)D, including 20% (32/158) who had both measurements (**Table 1**).

Among the patients with probable sarcoidosis (with and without biopsy), 70% (111/158) had managing providers who were "Fully" or "Substantially" compliant with the ATS practice guideline based on our scoring system (**Table 3**). The majority of those patients, 92% (102/111), were managed by specialists, including pulmonary physicians, while primary

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

care providers managed 8% (9/111). Among the 30% (47/158) patients whose care only met

the "Partially compliant" or "Non-compliant" definition based on our scoring system, 23%

(11/47) were managed by primary care providers (**Figure 4**). This result indicates that the care

of sarcoidosis patients was more likely to be classified as "Fully" or "Substantially" adherent

with the ATS practice guideline if their managing provider was a specialist (45% [9/20] of

primary care providers vs. 74% [102/138] of specialists, P= .008) (**Table 4**).

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**DISCUSSION**

In this retrospective cohort study of Department of Veterans Affairs EMR, we evaluated 200 randomly selected patients diagnosed with sarcoidosis from the San Francisco and Palo Alto Veterans Affairs Medical Centers. We found that using ICD-9 and 10 codes to identify sarcoidosis patients within EMR has a relatively low accuracy (PPV of 71% [95%CI= 64.7%-77.3%]) of detecting patients with "probable sarcoidosis" as defined by the ATS clinical practice guideline. We also demonstrated that including an "index of suspicion" that incorporates only the clinical and radiographical data allows for a significant increase in diagnostic accuracy (PPV of 90% [85.2%-94.6%]), as the initial sarcoidosis cohort was restricted to the patients with "high index of suspicion." This "index of suspicion," which was developed based on the ATS clinical practice guideline [3], could be executed by manual chart review of the available clinical and radiographic data and could potentially be adapted for automated chart review algorithms using natural language processing and machine learning in EMR. Furthermore, we found that the care provided by specialists was more likely to be "Fully" or "Substantially" compliant with the ATS practice guideline compared to primary care providers.

This randomly selected cohort of 200 Veterans with sarcoidosis diagnosis consisted of 90% men and 10% women. While the sex distribution in our study was different from the ACCESS study[12], it is closely reflective of the veterans' population demographics.[13] This study further confirmed the higher prevalence of sarcoidosis in African-Americans (54%) compared to Caucasians (33%), which has been previously reported by many other sarcoidosis epidemiology studies [14–18]. At the same time, the study population was racially diverse, highlighting the potential utility of the VA EMR for studying sarcoidosis in populations of people of color.[19]

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

The novelty of this study is the development of a scoring system to objectively measure

physicians' compliance with the latest ATS practice guideline. Adherence to recommendations

might allow for earlier interventions in sarcoidosis management and ultimately improve

outcomes by delaying or preventing the development of morbid phenotypes. Variables used in

this scoring system included ophthalmology visit, 12-lead EKG, chest x-ray, and vitamin D

levels measurements within one year of the entry date of the ICD code. These variables were

chosen based on their utility and importance.[3] Not all tests and measurements discussed in

the guidelines have been included in our scoring system. Yet, we believe it can be easily

modified to include other clinical variables. Notably, a longitudinal analysis of the association

between physicians' compliance with the recommendations and clinical outcomes in

sarcoidosis could provide direction for future investigations and ultimately help better guide

clinical practice to be used as a metric to assess the improvement of the care delivery for

sarcoidosis patients.

Using ICD codes alone to extract health information is far more convenient than the

time-consuming manual review process of narrative datasets in unstructured data.[8] However,

using ICD codes to identify sarcoidosis cases in large datasets with thousands of patients poses

several practical challenges. First, given the heterogeneity of sarcoidosis, it is challenging to

confirm the disease's presence efficiently. The verification process requires careful analysis of

the available narrative data such as progress notes, imaging reports, and pathology reports to

establish the case definition based on the sarcoidosis diagnostic criteria.[3] Second, the precise

identification of the type of organ involvement through EMR is a complex process and requires

a thorough review of unstructured data. Although there are sub-codes for ICD diagnostic codes

that aim to capture various organ involvement, healthcare providers may or may not be familiar

with those sub-codes and may or may not use them correctly. Moreover, there are no specific

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

ICD codes for classifying some of the various organ involvement in sarcoidosis (such as central

nervous system or GI tract).[20] Third, ICD codes do not determine the extent of disease, such

as that described by chest x-ray stages [21], due to a lack of ICD codes for different stages of

pulmonary sarcoidosis.[20] Analysis of pulmonary features requires a manual review of every

patient's radiology reports and cannot be performed using only ICD codes alone. Finally, ICD

codes do not specify the various sarcoidosis presentations such as acute, remitting, or chronic

disease.[11,20] Thus, they cannot be used to classify patients into previously described

phenotype groups.

The definition of clinical phenotypes has become an essential goal for the sarcoidosis

scientific community, as genetic studies have identified different patterns of gene expression

associated with disease severity and disease course.[22,23] In 2015, the National Heart, Lung,

and Blood Institute (NHLBI) held a workshop to leverage current scientific knowledge and

defining platforms to address disease disparities, identify high-risk phenotypes, and improve

sarcoidosis outcomes [24]. Nine different steps and research strategies were recommended to

expand the scope of sarcoidosis research, including EMR-based research to provide a unified,

multidisciplinary approach to bring together stakeholders interested in reducing the burden and

severity of sarcoidosis. However, the major barrier in the efficient use of EMR data is the

accurate extraction of research-quality variables, case definitions, and outcomes.[25] Thus, the

rapid identification of cases and extraction of relevant clinical variables from EMR using

computable phenotype algorithms have emerged as an important next step in EMR-based

research. Computable phenotype definitions are also essential to conducting pragmatic clinical

trials and comparative effectiveness research, increasing the healthcare system's capacity to

deliver Precision Medicine effectively.[26]

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

The two most commonly applied approaches to defining computable phenotypes are:

(1) a "*high-throughput*" phenotype algorithm using only structured data (traditionally, the ICD

diagnosis codes). (2) a *"low-throughput"* phenotype algorithm that accesses structured and

unstructured data to develop a sequential flow chart that should end with a case definition.

Such a low-throughput approach employs high-performance computational tools (such as

natural language processing [**NLP**]) to process text and extract information utilizing linguistic

rules instead of labor-intensive manual review by researchers.[6] This approach should

streamline the development of registries and help enrich EMR-based research studies.[27] Our

study highlights the need for the development of such automated methods to improve the

computational case-definition of sarcoidosis and other high-quality sarcoidosis-related

research variables, such as determining the date of the diagnosis, organ involvements, Scadding

stages, and the clinical status (acute, chronic, acute on the chronic or remitting disease).

Our study has several limitations. First, we used the ICD diagnosis code's entered date

as the surrogate time point to establish the patients' care with active sarcoidosis or re-

establish the care for those with remitting disease. This approach is different from what

sarcoidosis researchers traditionally use as a surrogate to determine the date of diagnosis,

which is the date of performing the biopsy. However, because the ICD diagnosis code's date

is more relevant for assessing the managing healthcare providers' compliance, we chose to

take the above approach. Second, in the cases where the biopsy report was unavailable (either

due to a remote history of biopsy or biopsy performed outside the VA), we relied on the

"secondary" histopathological reports documented in the providers' narrative within the

clinical notes. This approach made the diagnosis of sarcoidosis less robust because

confirmatory biopsy reports in those patients were not directly verifiable. However, we used

the index of suspicion approach to define probable sarcoidosis cases regardless of whether

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

there was a confirmatory biopsy report available, which is consistent with the diagnostic

algorithm recommended by the ATS practice guideline.[3] Last, the generalizability of our

findings in VA EMR to other populations could be limited because the Veterans form a

special population with different demographics and exposures from the general population.

However, the US Veterans Affairs Healthcare System EMR data cover over 22 million

Veterans across the US and over 14,000 patients with sarcoidosis ICD diagnosis codes,

providing an enormous number of patients to study a rare disease.

**Conclusion**

Although ICD codes can be used as reasonable classifiers to identify sarcoidosis cases

within EMR, using computational algorithms to extract clinical and radiographic information

("index of suspicion") from unstructured data could significantly improve case identification

accuracy. Using automated emerging methods (such as NLP) to develop a novel sarcoidosis-

specific computational phenotype algorithm could increase the efficiency of identifying these

cases from large healthcare databases. Furthermore, we found that specialists are more likely

than primary care providers to be "Fully" or "Substantially" compliant with the ATS clinical

practice guideline. This finding calls attention to the importance of examining the association

between physicians' compliance with the recommendations and clinical outcomes

longitudinally.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

415 **TABLES:**

416 **Table 1.** Distribution of characteristics and clinical phenotype groups of patients with

417 probable sarcoidosis [a] (with and without confirmed biopsy)

| | Probable Sarcoidosis [a] | | |
|---|---|---|---|
| | Probable sarcoidosis with confirmed biopsy N (%) | Probable sarcoidosis without confirmed biopsy N (%) | p-value |
| Characteristics | 142 (90) | 16 (10) | |
| Age (mean years ±SD) | 65.5 ±10.8 | 69.3 ±10.3 | 0.18 |
| Gender | | | |
| M | 127 (89.4) | 15 (93.8) | 0.59 [c] |
| F | 15 (10.6) | 1 (6.2) | |
| Race | | | |
| African American | 74 (52.1) | 11 (68.8) | 0.62 [b] |
| Non-Hispanic White | 49 (34.5) | 3 (18.8) | |
| Hispanic White | 3 (2.1) | 0 (0) | |
| Unknown | 12 (8.5) | 2 (12.5) | |
| Other | 4 (2.8) | 0 (0) | |
| ICD codes for sarcoidosis | | | |
| ICD-9 | 98 (69) | 10 (62.5) | 0.60 [c] |
| ICD-10 | 44 (31) | 6 (37.5) | |
| Specialty visit | | | |
| Pulmonary only | 71 (50) | 10 (63) | 0.28 [b] |
| Other specialists (including multiple) | 54 (38) | 3 (18.5) | |
| Primary care only | 17 (12) | 3 (18.5) | |

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

| | | | |
|---|---|---|---|
| Ophthalmology visit | 120 (84.5) | 13 (81.2) | 0.93 [c] |
| Organ involvement | | | |
| Lung | 86 (60.5) | 12 (75) | |
| Multiorgan (pulmonary without cardiac involvement) | 39 (27.5) | 2 (12.5) | |
| Multiorgan (cardiac without pulmonary involvement) | 2 (1.4) | 0 (0) | |
| Multiorgan (not cardiac neither pulmonary involvement) | 11 (7.8) | 2 (0) | 0.38 [b] |
| Multiorgan (both cardiac and pulmonary involvement) | 4 (2.8) | 0 (0) | |
| EKG | 118 (83.1) | 13 (81.3) | 0.97 [c] |
| PFT pattern [d] | | | |
| Obstructive | 27 (19) | 1 (6.2) | |
| Restrictive | 30 (21.1) | 6 (37.5) | 0.03 [b] |
| Mixed | 20 (14) | 5 (31.3) | |
| Normal | 39 (27.5) | 1 (6.2) | |
| CXR | 129 (90.8) | 14 (87.5) | 0.93 [c] |
| 25(OH)D | 72 (50.7) | 10 (62.5) | 0.80 [c] |
| 1,25(OH)D | 31 (21.8) | 5 (31.3) | |
| Both (1,25 and 25 (OH)D) | 31(21.8) | 1(6.2) | |

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

| Scadding stage [e] | | | |
|---|---|---|---|
| Stage 0 | 24 (16.9) | 5 (31.2) | |
| Stage I | 23 (16.2) | 3 (18.8) | |
| Stage II | 45 (31.7) | 2 (12.5) | 0.06 [b] |
| Stage III | 17 (12) | 4 (25) | |
| Stage IV | 22 (15.5) | 0 (0) | |
| Clinical Phenotype Group [f] | | | |
| Group 1: Multiorgan | 50 (35.2) | 2 (12.5) | |
| Group 2: Nonacute, Stage I, untreated | 6 (4.2) | 2 (12.5) | |
| Group 3: Stage II-III, treated | 42 (29.6) | 3 (18.8) | |
| Group 4: Stage II-III, untreated | 14 (9.9) | 2 (12.5) | |
| Group 5: Stage IV, treated | 17 (12) | 0 (0) | 0.06 [b] |
| Group 6: Stage IV, untreated | 4 (2.8) | 2 (12.5) | |
| Group 7: Acute sarcoidosis, untreated | 0 (0) | 0 (0) | |
| Group 8: Remitting, untreated | 30 (21) | 5 (31) | |
| Group 9: Cardiac sarcoidosis, treated | 6 (4.2) | 0 (0.0) | |

418
419 [a] Based on the clinical diagnostic criteria recommended by the American Thoracic Society (clinical, radiographic,
420 and pathological findings) in the absence of other alternative diagnoses that could explain the presence of non-
421 necrotizing granulomas on histopathology.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

[b] Fisher's exact test.
[c] $X^2$ test.
[d] Evaluated based on pulmonary function tests reports available in CPRS.
[e] Scored based on reviewers' interpretation of imaging reports using Scadding staging. Stage 0: normal chest
radiograph, Stage I: hilar or mediastinal nodal enlargement only; Stage II: nodal enlargement and parenchymal
disease; Stage III: parenchymal disease only; Stage IV: end-stage lung disease (pulmonary fibrosis).
[g] Clinical Phenotype Groups [11]:
Group 1: Multiorgan involvement: Patients with more than 2 organs involved.
Group 2: Nonacute, Stage I, untreated: Patients with nonacute sarcoidosis, Stage I, never treated for sarcoidosis.
Group 3: Stage II-III, treated: Patients with nonacute sarcoidosis, Stage II or III, formerly treated for sarcoidosis
or treated within 3 months of data review.
Group 4: Stage II-III, untreated: Patients with nonacute sarcoidosis, Stage II or III, never treated for sarcoidosis.
Group 5: Stage IV, treated: Patients with nonacute sarcoidosis, Stage IV, formerly treated for sarcoidosis or treated
within 3 months of data review.
Group 6: Stage IV, untreated: Patients with nonacute sarcoidosis, Stage IV, never treated for sarcoidosis.
Group 7: Acute sarcoidosis, untreated: Patients with acute sarcoidosis (Löfgren syndrome).
Group 8: Remitting, untreated: Patients who have had no evidence of active clinical disease for more than one
year.
Group 9: Cardiac sarcoidosis, treated: Patients with cardiac manifestations of sarcoidosis, formerly treated for
sarcoidosis or treated within 3 months of data review.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

442  **Table 2.** Contingency 2x2 table of using ICD and index of suspicion for sarcoidosis cases

443  identification.

| | Index of Suspicion [a] | Sarcoidosis confirmed with a biopsy | | |
|---|---|---|---|---|
| | | Yes [b] | No [c] | Total |
| Patients with positive ICD 9 or ICD-10 for sarcoidosis | High index of suspicion | 142 [d] | 16 [e] | 158 |
| | Low index of suspicion | 0 [f] | 42 [g] | 42 |
| | Total | 142 | 58 | 200 |

[a] Index of suspicion for sarcoidosis based on both clinical and radiographic evidence but not biopsy.
[b] Available biopsies with primary or secondary histopathological reports.
[c] No biopsies were ordered or available in the electronic medical record.
[d] Probable sarcoidosis group with histopathological evidence of non-necrotizing granuloma (NNG) = true
positives
[e] Probable sarcoidosis group without histopathological evidence of NNG= false positives
[f] No sarcoidosis group due to lack of sufficient clinical and radiological features consistent with sarcoidosis even
in the presence of the histopathological evidence of NNG = false negatives
[g] No sarcoidosis group due to lack of sufficient clinical and radiological features consistent with sarcoidosis, in
addition to the absence of the histopathological evidence of NNG= true negatives

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

455 **Table 3.** Healthcare providers' compliance with the ATS clinical practice guideline.

| Healthcare providers' compliance with ATS practice guidelines [a] N (%) | Probable sarcoidosis with confirmed biopsy N=142 | Probable sarcoidosis without confirmed biopsy N= 16 |
|---|---|---|
| Fully compliant | 26 (18.3) | 1 (6.25) |
| Substantially compliant | 76 (53.5) | 8 (50) |
| Partially compliant | 31 (21.9) | 6 (37.5) |
| Non-compliant | 9 (6.3) | 1 (6.25) |

[a] Developed with modifications based on recently published ATS practice guideline. The scoring system has
been developed based on the availability of the following within one year of the diagnosis: ophthalmology visit
= 3 points, 12-leads EKG = 3 points, CXR = 3 points, vitamin D (25-OH + 1,25-OH = 3 points; 25(OH)D or
1,25(OH)D = 2 points, otherwise zero points). Each variable weight a score of 3, with a total score of 12. Each
patient was given a final score, and the purpose was to classify the degree of the providers' compliance into four
categories based on the score:
o *Fully compliant*: if the score = 12
o *Substantially compliant*: if the score = 9-11
o *Partially compliant*: if the score = 5-8
o *Non-compliant*: if the score <5.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

467    **Table 4.** Contingency 2x2 table of the healthcare providers' compliance score by primary

468    versus specialty care [a].

| Healthcare providers' compliance with ATS practice guideline scored by our classification | Healthcare providers | | |
|---|---|---|---|
| | Specialists | Primary care physicians | Total |
| "Fully" or "Substantially" compliant | 102 | 9 | 111 |
| "Partially" or "Non" compliant | 36 | 11 | 47 |
| Total | 138 | 20 | 158 |

470    [a] $X^2$ statistic= 6.9877, P-value= 0.008.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

471  **Ethics approval and consent to participate:** The University of California San Francisco

472  Institutional Review Board and the Veterans Health Administration Research and

473  Development Committee approved this study. [IRB Protocol #15-16660].

475  **Availability of data and material**: Due to the sensitive nature of health data analyzed in the

476  current study, data will remain confidential and are not publicly available.

477
**Competing interests:** The authors have no conflicts of interest to disclose relevant to the

present work.

**Funding**: This work was supported by funds from the Department of Veterans Affairs

Fellowship Award to MIS; the Flight Attendants Medical Research Institute (FAMRI)

(CIA190001 to MA); Department of Veterans Affairs Clinical Sciences Research and

Development (CSRD) (CXV-00125 to MA); the Tobacco-related Disease Research Program

of the University of California (T29IR0715 to MA).

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**Bibliography**

[1]    Judson MA. Advances in the diagnosis and treatment of sarcoidosis. F1000Prime Rep 2014;6:89. doi:10.12703/P6-89.

[2]    Keller AZ. Anatomic sites, age attributes, and rates of sarcoidosis in U. S. veterans. Am Rev Respir Dis 1973;107:615–20. doi:10.1164/arrd.1973.107.4.615.

[3]    Crouser ED, Maier LA, Wilson KC, Bonham CA, Morgenthau AS, Patterson KC, et al. Diagnosis and detection of sarcoidosis. an official american thoracic society clinical practice guideline. Am J Respir Crit Care Med 2020;201:e26–51. doi:10.1164/rccm.202002-0251ST.

[4]    Judson MA. The diagnosis of sarcoidosis. Curr Opin Pulm Med 2019;25:484–96. doi:10.1097/MCP.0000000000000596.

[5]    Szeto HC, Coleman RK, Gholami P, Hoffman BB, Goldstein MK. Accuracy of computerized outpatient diagnoses in a Veterans Affairs general medicine clinic. Am J Manag Care 2002;8:37–43.

[6]    Pendergrass SA, Crawford DC. Using electronic health records to generate phenotypes for research. Curr Protoc Hum Genet 2019;100:e80. doi:10.1002/cphg.80.

[7]    Horsky J, Drucker EA, Ramelson HZ. Accuracy and Completeness of Clinical Coding Using ICD-10 for Ambulatory Visits. AMIA Annu Symp Proc 2017;2017:912–20.

[8]    Ungprasert P, Matteson EL, Crowson CS. Accuracy of Diagnostic Coding for Sarcoidosis in Electronic Databases: A Population-Based Study. Lung 2017;195:713–5. doi:10.1007/s00408-017-0054-x.

[9]    Velarde KE, Romesser JM, Johnson MR, Clegg DO, Efimova O, Oostema SJ, et al. An initiative using informatics to facilitate clinical research planning and recruitment in the VA health care system. Contemp Clin Trials Commun 2018;11:107–12. doi:10.1016/j.conctc.2018.07.001.

[10]   Sampling Essentials: Practical Guidelines for Making Sampling Choices - SAGE Research Methods n.d. https://dx.doi.org/10.4135/9781452272047 (accessed December 23, 2020).

[11]   Moller DR, Koth LL, Maier LA, Morris A, Drake W, Rossman M, et al. Rationale and Design of the Genomic Research in Alpha-1 Antitrypsin Deficiency and Sarcoidosis (GRADS) Study. Sarcoidosis Protocol. Annals of the American Thoracic Society 2015;12:1561–71. doi:10.1513/AnnalsATS.201503-172OT.

[12]   Baughman RP, Teirstein AS, Judson MA, Rossman MD, Yeager H, Bresnitz EA, et al. Clinical characteristics of patients in a case control study of sarcoidosis. Am J Respir Crit Care Med 2001;164:1885–9. doi:10.1164/ajrccm.164.10.2104046.

[13]   National Center for Veteran Analysis and Statistics. Department of Veterans Affairs Statistics at A Glance (Updated 12/31/2019). February 2020. n.d. https://www.va.gov/vetdata/docs/Quickfacts/Stats_at_a_glance_4_6_20.PDF (accessed January 1, 2021).

[14]   Mirsaeidi M, Machado RF, Schraufnagel D, Sweiss NJ, Baughman RP. Racial difference in sarcoidosis mortality in the United States. Chest 2015;147:438–49. doi:10.1378/chest.14-1120.

[15]   Cozier YC, Berman JS, Palmer JR, Boggs DA, Serlin DM, Rosenberg L. Sarcoidosis in black women in the United States: data from the Black Women's Health Study. Chest 2011;139:144–50. doi:10.1378/chest.10-0413.

[16]   Rybicki BA, Major M, Popovich J, Maliarik MJ, Iannuzzi MC. Racial differences in sarcoidosis incidence: a 5-year study in a health maintenance organization. Am J Epidemiol 1997;145:234–41. doi:10.1093/oxfordjournals.aje.a009096.

[17]   Brito-Zerón P, Kostov B, Superville D, Baughman RP, Ramos-Casals M, Autoimmune Big Data Study Group. Geoepidemiological big data approach to sarcoidosis:

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

geographical and ethnic determinants. Clin Exp Rheumatol 2019;37:1052–64.

[18]   Arkema EV, Cozier YC. Sarcoidosis epidemiology: recent estimates of incidence,
prevalence and risk factors. Curr Opin Pulm Med 2020;26:527–34.
doi:10.1097/MCP.0000000000000715.

[19]  Minority Veteran Statistics n.d.
https://www.va.gov/HEALTHEQUITY/Race_Ethnicity.asp?utm_content&utm_mediu
m=email&utm_name&utm_source=govdelivery&utm_term (accessed January 1,
2021).

[20]   Organization WH. The ICD-10 Classification of Mental and Behavioural Disorders:
Diagnostic Criteria for Research. New Edition. Geneva: World Health Organization;
1993.

[21]   Scadding JG. Prognosis of intrathoracic sarcoidosis in england. BMJ 1961;2:1165–72.
doi:10.1136/bmj.2.5261.1165.

[22]   Su R, Li MM, Bhakta NR, Solberg OD, Darnell EPB, Ramstein J, et al. Longitudinal
analysis of sarcoidosis blood transcriptomic signatures and disease outcomes. Eur
Respir J 2014;44:985–93. doi:10.1183/09031936.00039714.

[23]   Zhou T, Zhang W, Sweiss NJ, Chen ES, Moller DR, Knox KS, et al. Peripheral blood
gene expression as a novel genomic biomarker in complicated sarcoidosis. PLoS One
2012;7:e44818. doi:10.1371/journal.pone.0044818.

[24]   Maier LA, Crouser ED, Martin WJ, Eu J. Executive summary of the NHLBI workshop
report: leveraging current scientific advancements to understand sarcoidosis variability
and improve outcomes. Annals of the American Thoracic Society 2017;14:S415–20.
doi:10.1513/AnnalsATS.201707-563OT.

[25]   Sharma H, Mao C, Zhang Y, Vatani H, Yao L, Zhong Y, et al. Developing a portable
natural language processing based phenotyping system. BMC Med Inform Decis Mak
2019;19:78. doi:10.1186/s12911-019-0786-z.

[26]   Richesson R. Electronic Health Records-Based Phenotyping n.d.

[27]   Paul DW, Neely NB, Clement M, Riley I, Al-Hegelan M, Phelan M, et al.
Development and validation of an electronic medical record (EMR)-based computed
phenotype of HIV-1 infection. J Am Med Inform Assoc 2018;25:150–7.
doi:10.1093/jamia/ocx061.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

**FIGURE LEGENDS**

**Figure 1. STROBE flow chart. Selection criteria for probable sarcoidosis cases.**

Abbreviations: ATS, American Thoracic Society; ICD, International Classification of

Diseases; PA, Palo Alto; SF, San Francisco; VA, Veterans Affairs

**Figure 2. Organ involvement assessment for probable sarcoidosis with and without**

**confirmed biopsy.** Abbreviations: AV, atrioventricular; CNS, central nervous system; CT,

computed tomography; CXR, chest radiograph; DLCO, diffusing capacity for carbon

monoxide; EKG, electrocardiogram; ENT, ears, nose and throat; MRI, magnetic resonance

imaging; PET, positron emission tomography; PFT, pulmonary function test.

**Figure 3. Distribution of characteristics among patients with probable sarcoidosis with**

**and without confirmed biopsy.** Abbreviations: PFT, pulmonary function test; ICD,

International Classification of Diseases.

**Figure 4.** Comparison of specialty visits across cases with probable sarcoidosis (with and

without confirmed biopsy) and with healthcare providers classified as "Fully" or

"Substantially" compliant vs. "Partially" or "Non-" compliant using our compliance scoring

system.

**Acknowledgment:** The Author(s) confirmed no figures or tables included from another

publication.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

593 **Figure 1.** STROBE flow chart. Selection criteria for probable sarcoidosis cases.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

| Organ involvement | |
|---|---|
| Lung | • Positive lung biopsy, positive mediastinal/hilar lymph node biopsy<br>• Chest X-ray/CT scan/PET scan demonstrating bilateral hilar lymphadenopathy, CT scan with perilymphatic nodules tracking the bronchovascular bundle, chest X-ray/CT scan/PET scan with diffuse infiltrates, chest X-ray with fibrosis<br>• PFT with restriction and low DLCO or isolated reduced DLCO |
| Skin | • Positive skin biopsy<br>• Lupus pernio and erythema nodosum [a] |
| Eye | • Positive conjunctival or scleral biopsy<br>• Optic neuritis, scleritis, uveitis, or retinitis |
| Cardiac | • Positive heart/pericardium biopsy<br>• 12-lead EKG showing Mobitz II or III-degree AV node block<br>• AV node block or cardiomyopathy responsive to treatment<br>• Cardiac MRI or PET-CT consistent with sarcoidosis |
| Liver | • Positive liver biopsy with a positive biopsy from another organ<br>• Evidence of hepatomegaly<br>• Abnormal liver enzymes |
| Multiorgan involvement | • More than two organs involved based on other criteria in this table |
| Neurosarcoidosis | • Positive brain/dura/peripheral nerve biopsy<br>• Clinical syndrome or symptoms consistent with CNS sarcoidosis along with a positive MRI |
| ENT | • Positive biopsy from ear, nose, or throat<br>• Direct laryngoscopy consistent with granulomatous disease |

**Figure 2.** Organ involvement assessment for probable sarcoidosis with and without confirmed biopsy [a].

598
599
[a] No biopsy is needed for acute skin sarcoidosis.

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

601
602

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

604
605

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

609
610

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

617 **Figure 3.** Distribution of characteristics among patients with probable sarcoidosis with and

618 without confirmed biopsy.

619 <u>Abbreviations:</u> PFT, pulmonary function test; ICD, International Classification of Diseases

Mohamed I Seedahmed (https://orcid.org/0000-0002-7446-7346)

623 **Figure 4.** Comparison of specialty visits across cases with probable sarcoidosis (with and

624 without confirmed biopsy) and with healthcare providers classified as "Fully" or

625 "Substantially" compliant vs. "Partially" or "Non-" compliant using our compliance scoring

626 system.