

1 West Nile virus in Portugal

2 Authors

3 José Lourenço ^{a*}, Sílvia C Barros ^b, Líbia Zé-Zé ^{c,d}, Daniel SC Damineli ^e, Marta Giovanetti ^{f,g}, Hugo
4 C Osório ^{c,h}, Fátima Amaro ^{c,h}, Ana M Henriques ^b, Fernanda Ramos ^b, Tiago Luís ^b, Margarida D
5 Duarte ^b, Teresa Fagulha ^b, Maria J Alves ^{c,h}, Uri Obolski ^{ij}

6 Affiliations

7 **a** Department of Zoology, University of Oxford, United Kingdom

8 **b** Instituto Nacional de Investigação Agrária e Veterinária, Virology Laboratory, Portugal

9 **c** Centro de Estudos de Vectores e Doenças Infecciosas, Instituto Nacional de Saúde Doutor Ricardo Jorge, Portugal

10 **d** Biosystems and Integrative Sciences Institute, Edifício TecLabs, Campus da FCUL, Portugal

11 **e** Department of Pediatrics, Faculdade de Medicina da Universidade de São Paulo, Brazil

12 **f** Laboratório de Flavivírus, Instituto Oswaldo Cruz Fiocruz, Brazil

13 **g** Laboratório de Genética Celular e Molecular, Universidade Federal de Minas Gerais, Brazil

14 **h** Instituto de Saúde Ambiental, Faculdade de Medicina da Universidade de Lisboa, Portugal

15 **i** School of Public Health, Faculty of Medicine, Tel Aviv University, Israel

16 **j** Porter School of the Environment and Earth Sciences, Faculty of Exact Sciences, Tel Aviv University, Israel

17 * Corresponding author(s): José Lourenço, jose.lourenco@zoo.ox.ac.uk

18 Competing interests

19 No competing interests.

20 **Abstract**

21 Background: West Nile virus (WNV) is a mosquito-borne pathogen with a globally expanding
22 geographical range. Many European countries regularly report outbreaks affecting both human
23 and equine populations, with an increase in epidemic activity in the past decade. Portugal, on
24 the far west of Europe, has not yet experienced large outbreaks of WNV, despite its suitable
25 Mediterranean climate.

26 Objectives: To provide a comprehensive, data-driven perspective on the past epidemiology,
27 surveillance and transmission suitability of WNV in Portugal.

28 Methods: Literature and database reviews were performed, covering the period between 1966
29 and 2020 on existing evidence for WNV circulation. Climatic data was used to estimate the
30 theoretical transmission potential of WNV, and to analyse long-term climate trends between
31 1981 and 2019.

32 Results: Existing data show that WNV-capable mosquito species are spread across the country,
33 while WNV molecular and serological evidence is vastly concentrated in the south. Our
34 analyses suggest that two different climate types, characteristic of the north and south of
35 Portugal, contribute to quantitative differences in WNV theoretical transmission potential. This
36 supports the north-south divide observed from previous molecular and serological evidence.
37 Due to long-term changes in local climate, suitability for WNV transmission is shown to have
38 been slowly increasing in the country.

39 Discussion: Data and theoretical analyses support the notion that the south of Portugal is more
40 suitable for WNV transmission, while not rejecting the possibility of transmission in the north.
41 Mosquito monitoring has been effective, but surveillance of key sentinel species remains
42 passive, creating important knowledge gaps for epidemic control and public health. To date,
43 evidence supports WNV circulation, but it remains unclear whether it is endemic or sporadically
44 transmitted. This study supports a timely change towards WNV active surveillance, to avoid
45 and mitigate potential public health emergencies as those recently observed in other European
46 countries.

47 **Introduction**

48 West Nile virus (WNV) is an RNA mosquito-borne virus of the Flaviviridae family, first identified
49 in 1937 in the West Nile district of Uganda. WNV ecology is characterised by a zoonotic
50 transmission cycle maintained between a large number of mosquito and wild avian species.
51 Occasional viral spillover can occur from this cycle to humans as well as domesticated and
52 wild mammals [1–3]. From such incidental hosts, humans and equines feature predominantly
53 in epidemiological data given a clear West Nile fever clinical profile, with equines often serving
54 as a sentinel species in many countries. Contrary to most avian species, mammals are
55 inefficient amplifier hosts, and hence can not establish mosquito-mammal transmission cycles
56 [3–5]. Approximately 80% of infections in humans are asymptomatic while the rest may
57 develop mild or severe disease of neuroinvasive nature and potentially death [6]. There are
58 currently no licensed vaccines nor antiviral treatments for humans [3,7], but four licensed

59 vaccines are on the market for use in equines [7].

60 Global trends in climate change and human factors continue to favour the dispersal of
61 mosquito species into new regions, and the long-distance movement of infectious hosts across
62 the globe. Both factors promote the introduction and epidemic activity of mosquito-borne
63 viruses such as WNV into previously unaffected areas [8–11]. Some of the most relevant
64 mosquito-borne examples of the past 10 years include the introduction of chikungunya and
65 Zika viruses into the Americas [12,13], as well as the resurgence of yellow fever virus in Brazil
66 and regions of Africa [14,15].

67 In the 20th century, WNV was mostly reported to circulate in Israel and some African countries
68 [16]. More recently, in 1999, it gained international notoriety after its introduction into the United
69 States of America (USA, New York), from where it quickly became endemic in most of the USA
70 and Canada [17]. Concurrently and in the following two decades, epidemic activity increased in
71 regions of Europe, the Middle East and Russia [11,16]. Presently, WNV epidemiology shows
72 large spatio-temporal heterogeneities across countries and continents, but much of that
73 variation is not well understood. However, given its well documented effects on the life-cycles
74 and dispersal of avian and mosquito species, climate has long been accepted as a major factor
75 influencing local WNV epidemic activity, dispersal and persistence.

76 The recent epidemiological history of WNV in Europe is characterised by increasing epidemic
77 activity in countries including France, Italy, Greece, Hungary, Romania, Bulgaria, Serbia and
78 Ukraine [18,19]. In 2018, the continent reported its largest epidemic ever across many
79 countries, exceeding cumulative infections since 2010 [20]. In 2020, the first autochthonous
80 human infections were reported in the Netherlands [21], and Spain had its largest epidemic to
81 date including regions previously not affected [22,23]. Compared to Spain and other countries
82 in the Mediterranean basin, Portugal remains an outlier in so far having reported only four,
83 serologically confirmed human infections in the past 40 years.

84 Routine laboratory diagnostics in Portugal were established in 1996, and are currently managed
85 by the Centro de Estudos de Vectors e Doenças Infecciosas (CEVDI, Center for Vector and
86 Infectious Disease Studies) of the Instituto Nacional de Saúde Dr. Ricardo Jorge (INSA,
87 National Health Institute Dr. Ricardo Jorge) and by the Instituto Nacional de Investigação
88 Agrária e Veterinária (National Agrarian and Veterinary Research Institute, INIAV) in human and
89 animals, respectively. Notification of WNV-related disease is currently mandatory in Portugal.
90 Being an European Union (EU) member state, Portugal reports outbreaks of equine WNV
91 neurological disorders to the European Animal Disease Notification System [24], and annual
92 reports for the EU region are published by the European Food Safety Authority [25] and
93 European Centre for Disease Prevention and Control (ECDC) [26].

94 Since 2008 the country has been performing extensive mosquito surveillance under REVIVE
95 (REde de Vigilância de VETores, surveillance network for vectors [27,28]). A multitude of
96 national agencies within the Ministry of Health are involved, including the General Directorate of
97 Health (DGS), INSA, the five Regional Health Administrations (Algarve, Alentejo, Lisboa e Vale
98 do Tejo, Centro and Norte), and the non-continental agencies the Institute of Health
99 Administration of Madeira and the Regional Health Directorate of the Azores. REVIVE is
100 responsible for the nationwide surveillance of the most significant hematophagous arthropods

101 for national public health (mosquitoes, ticks, and sandflies) including the cataloguing of extant
102 species and their population sizes. Surveillance of invasive mosquito species, such as *Aedes*
103 *aegypti* and *Aedes albopictus*, and screening of field-collected mosquitoes for arboviruses (e.g.
104 PCR) is also regularly performed. According to International Health Regulations, airports, ports,
105 storage areas, and specific border regions with Spain are regularly monitored throughout the
106 year according to the needs of local and regional authorities. Reports from REVIVE are provided
107 regularly to national agencies across the country [27], and to the European Network for
108 Arthropod Vector Surveillance for Human Public Health, a network for data sharing on the
109 geographic distribution of arthropod vectors, transmitting human and animal disease agents
110 (originally known as VBORNET, and VectorNet since 2014 [29]).

111 WNV surveillance in Portugal remains passive and mostly reactive to occasional reports of
112 clinically compatible events in equines and humans. However, many local clinicians and
113 veterinarians lack awareness of the importance of WNV screening in compatible disease
114 cases. As such, and since infections tend to be asymptomatic or mild, it is possible that only a
115 very small percentage of all infections end up being screened and therefore accounted for.
116 Detection of WNV RNA in serum, cerebrospinal fluid or urine offer significant technical
117 challenges in both humans and equines when assessed substantially after the start of the
118 symptomatic period. In Portugal, WNV diagnosis in humans is typically requested following the
119 failure of all other disease compatible diagnoses, commonly days or weeks after symptoms
120 onset and consequently after the viraemic phase. The reactive nature of the surveillance
121 system also implies that increased awareness and screening requests for humans and equines
122 tend to rise around areas with recent positive cases, but only for very short time windows which
123 remain insufficient for long-term risk and epidemiological assessment. These aforementioned
124 realities allow for systematic gaps in current knowledge regarding the local epidemiology of
125 WNV, hindering important conclusions such as whether or where the virus is endemic.

126 In this article, we review and describe the international and Portuguese literatures and official
127 data sources (1966-2020), further providing a theoretical geo-temporal assessment of WNV
128 transmission potential in Portugal (1981-2019). We find a clear north-south dichotomy in
129 existing evidence for the circulation of WNV in Portugal, although high resolution data on WNV-
130 capable mosquito species suggests transmission would be possible across the entire country.
131 Our theoretical assessments on transmission potential suggest that the south of the country
132 presents climate-driven seasonal signatures that justify WNV circulation to be restricted to that
133 region. Using climate data of the past four decades, we also quantify long-term trends in
134 climate and their possible historical effects on the local WNV transmission potential. In light of
135 our conclusions and the recent increase in epidemic behaviour of WNV in the European region,
136 we advocate for a more active surveillance of the virus in Portugal, in which ongoing circulation
137 may soon develop into a public health emergency.

138 **Methods**

139 *Literature reviews*

140 We queried PUBMED for articles related to WNV and Portugal using the query “((west nile virus)
141 AND (Portugal)) OR (vírus nilo AND Portugal)” in all fields; thus leaving the query non-specific
142 enough to capture a wide range of studies, and searching for articles both in Portuguese and

143 English. The query returned 42 hits as of 27th January 2020, from which 17 did not relate
144 directly to the circulation of WNV or its vectors in the country. A second literature search was
145 performed manually among Portuguese academic journals and PUBMED. References within the
146 studies found in our two searches were also scanned for potential studies related to WNV
147 circulation in Portugal. We also scanned the GenBank nucleotide database for existing WNV
148 sequences from Portugal, searched with the query “((west Nile virus) AND (Portugal)) OR
149 ((WNV) AND (Portugal))” OR (vírus nilo AND Portugal). The query returned 70 hits as of 27th
150 January 2021, of which 62 did not relate to WNV genomic regions.

151 We provide the results of our literature review in the form of tables in the main text. Studies
152 found in the PUBMED and manual searches are grouped by content type in **Table 1**. All
153 evidence related to the presence of WNV in Portugal across all studies found is summarized in
154 **Table 2** (also available in a permanent figshare repository [30]). Studies presenting time series
155 of relevant mosquito species in Portugal are summarized in **Table 3**. All publicly available WNV
156 genetic sequences obtained in Portugal are listed in **Table 4**.

157 *Mosquito data*

158 We used mosquito data collected by the REVIVE initiative in Portugal [27]. Data included
159 mosquito counts (adult and immature stages) for *Culex pipiens* per county in Portugal during
160 the period 2017-2019. The REVIVE data is in the public domain [27].

161 *Transmission suitability*

162 The basic reproduction number R_0 of WNV in the zoonotic reservoir can be summarised as the
163 product of two factors: the number of adult female mosquitoes per host M , and the
164 transmission potential of each adult female mosquito P (i.e. $R_0 = MP$). Both M and P oscillate
165 in time, driven by the influence of multiple eco-climatological factors. M , the ratio between the
166 population size of mosquitoes V and hosts N ($M = V/N$), depends on the local effects of
167 climate on mosquito survival and reproduction, affecting V [31], as well as on the reproduction
168 and migratory behaviours of the avian species, affecting N [32]. For most epidemiological
169 contexts and pathogens, high resolution data on either V or N is often lacking. The
170 transmission potential P depends on viral, mosquito and host factors (e.g. incubation periods,
171 lifespans, etc), some of which depend on local climate variations [31,33,34]. Since adequate
172 geo-temporal data for V and N is generally difficult to obtain, many recent theoretical
173 epidemiology studies resort to estimating the transmission potential of mosquito-borne viruses
174 using ‘suitability indices’. These indices are functions of the parameters that compose either M
175 or P or both, and generally depend on covariates such as climatic variables, vegetation indices,
176 host demography, mosquito and disease incidence reports, etc [34–36]. While a suitability
177 index is an incomplete measure of the real transmission potential of a mosquito-borne virus,
178 much work has been done to demonstrate the potential of an index to capture the geo-temporal
179 dynamics of the number of reported mosquito-borne infections in host species [34,37,38].

180 In this study, we assessed local WNV transmission potential by estimating one of such
181 suitability indices - termed the index P (the transmission potential of each adult female
182 mosquito from $R_0 = MP$). The theory and practice of P has been previously described in full by
183 Obolski et al., that also introduced the MVSE R-package capable of estimating P given some

184 epidemiological priors and climatic data [34]. The index P has been used in several studies
185 focusing on the population dynamics of the dengue, chikungunya, and Zika viruses (e.g.
186 [34,37,39,40]). Recently, the index was also successfully used to estimate the transmission
187 potential of WNV in Israel dependent on local relative humidity and temperature variables [41].
188 In that study, P was interpreted as a proxy for the risk of spillover to the human population
189 given that it measured the WNV transmission potential by single adult female mosquitoes in
190 the animal reservoir. In the current study, we use the same approach as applied to Israel,
191 informed by the epidemiological priors applied in that study (see Table in main text of [41]),
192 which relate to spp. *Culex*, WNV and a theoretical, average bird species.

193 To inform the estimation of the index P , climatic data was downloaded from the Copernicus
194 platform at www.copernicus.eu. We used the available dataset “essential climate variables for
195 assessment of climate variability” [42], which includes climatic variables at a time resolution of
196 1 month (1981-2019), and gridded spatial resolution of $0.25^\circ \times 0.25^\circ$. In the current study we
197 used MVSE R-package version 1.01r [43], which uses temperature, relative humidity and
198 precipitation as input data. For each latitude-longitude Portuguese data point in the Copernicus
199 dataset, we estimated the index P as described by Obolski et al. for Brazil, the Honduras and
200 South America [34]. The resulting geotemporal estimations, as presented in the main text, are
201 made available as a comma separated file in a permanent figshare repository [30].

202 *Analysis of peak suitability timing and size*

203 Peaks in the monthly index P time series 1981-2019 per county were found by the maximal
204 value and month where it occurred, using a circular mean to calculate the average time point of
205 the year for the north and south regions, independently. The distribution of values throughout
206 the years were then compared between regions with a circular ANOVA yielding a $p=0.005$.
207 Circular data was handled with the R-package 'circular' [44]. Mean peak values were calculated
208 across counties for each year, using Welch's t-test to compare north and south regions yielding
209 a $p=1 \times 10^{-16}$.

210 *Analysis of long-term climatic and suitability trends*

211 We applied linear quantile mixed models to the estimated index P , temperature, precipitation
212 and humidity between 1981 and 2019, for which the Copernicus climate dataset was available.
213 Our data contained yearly median values of the mentioned variables for each Portuguese
214 county. Since the data included highly irregular points, which might skew standard linear
215 regression models, we chose to use the more robust quantile regression using the median ($\tau=0.5$).
216 Furthermore, as different regions have repeated measures, we employed a mixed-
217 effects quantile regression (using the 'lqmm' R-package [45]). Our models contained a random
218 intercept term for each region, a fixed effect dummy variable coding for north versus south, and
219 a continuous variable representing the time unit (year). We tested the interaction between the
220 year and north-south terms and found it statistically significant ($p<0.01$) in all models except
221 when index P was the dependent variable ($p>0.7$) for which the term was dropped. All the
222 other fixed effects were statistically significant for all models ($p<0.001$) and are summarized in
223 **Table 5**.

224 *Analysis of long-term suitability periodicity*

225 Detrending and wavelet analysis followed the workflow of Damineli et al. [46]. Detrending with a
226 Maximal Overlap Discrete Wavelet Transform was performed in a multiresolution analysis with
227 the R-package 'wavelets' [47]. Significant periods of the continuous wavelets were assessed
228 considering the spectrum of an autoregressive model of order 1, as a null model using the R-
229 package 'biwavelet' [48].

230 *Equine WNV surveillance data 2016-2020*

231 Between 2016 and 2020 the virology laboratory of INIAV performed molecular and serology
232 testing in 165 horses from several regions of the country, reacting to local suspected
233 neuroinvasive disease. RNA was extracted by using the BioSprint 96 workstation with the
234 MagAttract 96 cadon pathogen kit, according to manufacturer's instructions (Qiagen, Hilden,
235 Germany). The samples were screened for WNV by real-time reverse transcriptase PCR (RT-
236 qPCR) targeting the NS2A gene [49]. Equine sera were tested for WNV-IgM antibodies by
237 capture ELISA (ID Screen West Nile IgM Capture ELISA, IDVET, Montpellier, France). Twenty
238 three horses were diagnosed as WNV-positive on the basis of IgM positivity, with one horse
239 being also positive by RT-qPCR (July 2020, Escalos de Cima-Castelo Branco, center Portugal).
240 Four of eight horses that were followed for clinical progression died or were euthanized. These
241 results are summarized in **Table 2**, including the time and region of sampling.

242 **Results**


243 *Evidence for West Nile virus circulation in Portugal*

244 The earliest evidence of WNV circulation in Portugal came from serological surveys in the
245 south of the country in 1966-1967, reporting high titres of hemagglutination-inhibiting
246 antibodies and neutralizing capacity against WNV in animal serum [50,51]. Such findings led to
247 several initiatives in 1969 to create institutional links between the national health and veterinary
248 medicine services with the purpose of seeding future studies and raising awareness to the
249 virus [52]. Some of the earlier initiatives included the capture and testing of animals (e.g.
250 equines, rodents, birds, mosquitoes) and serological testing of human serum [51–54],
251 eventually leading to the first viral isolation from the mosquito species *Anopheles maculipennis*
252 [53].

253 After a period of three decades with seemingly no research output in the country, further
254 evidence for the circulation of WNV in Portugal accrued from several studies. Between 1999
255 and 2002, Formosinho and colleagues reported serological data from several avian species
256 sampled across natural reserves and from equines sampled in the district of Santarém [55]. A
257 few studies in the period 2004-2010 also reported positive findings in several avian species
258 [56], mosquitoes [57,58], and equines [56]. More recently, between 2015 and 2020, following
259 reports of several clinically compatible equine outbreaks in the south of the country, a number
260 of equines were found serologically positive to the virus in the Alentejo and Algarve [59,60].
261 Some of this evidence is reported for the first time in this study (see **Methods** for details, and
262 **Table 2** for summary).

263 Although such evidence supports a long history of local circulation in animals and mosquitoes,

264 autochthonous WNV infections in humans in Portugal were only first reported in 2004. This was
 265 the case of two Irish bird-watchers found positive for both WNV IgG and IgM antibodies. They
 266 visited Ria Formosa, a marshland rich in endemic and migratory birds in the Algarve between
 267 the 26th June and 10th of July [61]. Two studies published thereafter were able to detect WNV
 268 by RT-PCR in pools of mosquitoes collected near the putative region of infection that year
 269 [57,58]. A third case was reported in July 2010 in the region of Setúbal, from a local rural
 270 resident with no history of international travel [62]. The resident tested positive for both WNV
 271 IgG and IgM antibodies in the weeks following the clinical diagnosis. Reactive mosquito
 272 collection performed around the patient's residence between the 13-18th of July revealed
 273 species suited for WNV transmission (e.g. *Culex pipiens*), although all mosquito pools tested
 274 negative by RT-PCR [62]. Two months later, two equine cases were notified about 4 kilometers
 275 away from the reported human case. The most recent human infection was reported in a rural
 276 area of Almancil (Algarve) in 2015 in a patient testing positive for both WNV IgG and IgM
 277 antibodies and viral neutralization, with no record of international travel or history of
 278 vaccination against flaviviruses [63]. In August and September of the same year several
 279 outbreaks in equines were reported in Loulé (Algarve) with some animals testing positive by
 280 serology [60].



281 **Figure 1 - Past evidence of WNV circulation in Portugal.** Summary of data sources (articles, national
 282 reports) on various types of evidence for the circulation of WNV in Portugal. Panel A presents the
 283 geographical region of each data source (latitude-longitude). Positioning is only approximate, given that

284 *some reports refer to regions of different areas. For data sources referring to large areas (e.g. districts), the*
285 *capital city of the region was used. Not all sources are mapped, e.g. not reporting where positive samples*
286 *were found. Mapped words represent the type of evidence reported, aggregated into two categories:*
287 *molecular, for PCR, isolation; and serology, for HI, VNT, ELISA. Panel B shows a timeline of the number of*
288 *data sources per year. The color scale (on the right) represents the different types of samples reported, and*
289 *refers to both panels. Not all years between 1966 and 2020 had data sources. Detailed summary of data*
290 *sources in **Table 2**.*


291 It is likely that the main reason why none of the human or equine reported infections in Portugal
292 have so far been confirmed by RT-PCR is due to diagnosis and screening being performed well
293 past the viraemic phase. This might also explain why obtaining sequencing data related to WNV
294 Portuguese infections has not been successful. To date, a single study has been published on
295 the local phylogenetics of WNV, with sequencing solely obtained from mosquito pools (**Table**
296 **4**) [58]. Such sequences, based on a short segment of the NS5 gene, demonstrated a close
297 genetic relationship with past WNV lineages circulating in the Mediterranean basin (France,
298 Morocco and Italy), but fell short in providing definite conclusions on viral diversity,
299 phylodynamics and persistence in the country.

300 When summarized, the past evidence for the circulation of WNV in Portugal is mostly restricted
301 to reports from the southern regions of the country, in particular the districts of Faro, Beja,
302 Setúbal, Évora, Lisboa, Portalegre and Santarém (**Figure 1, Table 2**). Such reports include
303 evidence from avian species, equines, mosquitoes and humans, achieved via different
304 confirmation methods. It is unknown whether the observed WNV circulation in the southern
305 regions, compared to the northern regions, stems from sampling bias, or from disparate
306 ecological conditions dictating WNV epidemic activity. Sampling bias could be due to large
307 differences in regional biodiversity and population sizes of sentinel species (mosquitoes, birds,
308 equines) across the country. Indeed, there is evidence that the equine population is larger in the
309 south [64], while the population sizes and diversity of birds in the north and south are largely
310 unknown for many resident and migratory species. To the ecological point, climate is a major
311 driver of the population dynamics of mosquito and avian species participating in the zoonotic
312 cycle of WNV. According to the Köppen climate classification, Portugal presents a north-south
313 divide in climate types, with the northern region classified as Temperate Mediterranean, and the
314 southern region as Warm Mediterranean. A lack of past studies exploring the effects of climate
315 on WNV epidemiology in Portugal has, however, not allowed for much speculation on its role in
316 the seemingly restricted circulation of the virus to the south of the country. Considering the
317 possible sampling bias and lack of the above mentioned ecological and genetic information,
318 and with other factors still to be researched in the national context, there is currently no
319 consensus on where and whether WNV circulates in Portugal sporadically or endemically.

320 *West Nile virus vectors in Portugal*

321 To date, very few articles in the research literature have provided geo-temporal data on the
322 population size and dynamics of mosquito species relevant for the transmission of WNV in
323 Portugal (**Table 3**). We obtained REVIVE *Culex pipiens* data collected between 2017 and 2019
324 across Portuguese counties (**Figure 2**). The methods of collection (e.g. type of trap used), as
325 well as resources and regional objectives, varied by year, as described in detail in the published
326 reports [27]. Summarizing the data revealed that both sampling coverage and the total number
327 of yearly collections per county varied in time, both for adult (**Figures 2A-C**) and immature
328 (**Figures 2E-G**) forms of this mosquito species. Moreover, we noticed that even in the presence

329 of many project-related factors affecting sampling, *Culex pipiens* was found to be present
330 across the country (**Figures 2DH**). Since *Culex pipiens* is considered the main vector for the
331 possible transmission of WNV in Portugal, these findings stand in contrast with the north-south
332 divide of WNV circulation found in the data review from the literature and public databases
333 (**Figure 1, Table 2**).




334 **Figure 2 - REVIVE *Culex pipiens* counts in Portugal (2017-2019).** Borders represent Portuguese counties,
335 the level at which *Culex pipiens* data are reported. In panels A-C and E-G, counties are colored according to
336 the yearly observations (transformed to $\log_{10}(\text{counts}+1)$) of adult and immature mosquitoes
337 (respectively). In panels D and H, counties are colored by the yearly mean observations (transformed to
338 $\log_{10}(\text{mean}(\text{counts})+1)$) of adult and immature mosquitoes (respectively). Counties for which there was
339 zero counts are coloured in white. Counties for which collections were not performed are in grey.

340 West Nile virus transmission potential in Portugal

341 In this study we use a climate-driven suitability index (termed index *P*) to estimate the geo-
342 temporal transmission potential of WNV in Portugal (see **Methods** for details). This index is the
343 theoretical transmission potential *P* of each adult female mosquito in the zoonotic reservoir
344 (derived from the decomposition of $R_0=MP$). *P* can be interpreted as a proxy for the risk of
345 spillover to the human and equine populations (as demonstrated recently for Israel [41]).
346 Typically, the estimated seasonal variations in index *P* would be validated by quantifying geo-

347 temporal correlations between time series of the index and suspected or confirmed infections
 348 in humans or equines. However, WNV infection time series do not exist for Portugal given the
 349 absence of reported chains of transmission (**Figure 1, Table 2**).




350 **Figure 3 - Monthly REVIVE *Culex pipiens* counts in Portugal versus estimated transmission potential**
 351 ***P* (2017-2019).** Panels A-D present the monthly time series for $\log_{10}[1+\text{mosquito population size}]$ (grey
 352 bars) and mean (colored lines) and standard deviation (coloured areas) for four districts: two from the south
 353 (A for Beja in purple, C for Faro in purple) and two from the north (B for Bragança in red, D for Viana do
 354 Castelo in blue). The map panel presents all districts, coloured by their location. Panel E shows the mean
 355 index *P* for all districts 2009-2019 (North in grey and South in black). Panel F shows the seasonal index *P*
 356 per month from panel E independently of year. Panel G is the distribution of index *P* across the period 2009-
 357 2019. For the estimation of the index *P*, the monthly mean of the climate variables was used per district.

358 We thus first compared the only available time series data related to WNV transmission in
 359 Portugal, which included the newly estimated index *P* and regional *Culex pipiens* population
 360 sizes from the REVIVE initiative (**Figure 3**). Given the north-south disparity in past evidence for
 361 the circulation of WNV (**Figure 1**) and the widespread geographical distribution of *Culex pipiens*
 362 (**Figure 2**), we compared directly two of the most northern and southern districts of the country.
 363 Although the index *P* measures transmission per mosquito and not population size, we found
 364 the index to present peaks and troughs coinciding in time with those of the mosquito
 365 population sizes, both in the south (**Figures 3AC**) and north (**Figures 3BD**). Apart from the
 366 previously demonstrated potential for the index *P* to be correlated with WNV human infections
 367 [41], these results thus suggest that (at least in Portugal) the index can also be used to
 368 estimate the seasonal timing of variation in *Culex pipiens* population sizes.

369 The patterns in **Figures 3A-D** further suggest several characteristics of the typical season of
 370 WNV transmission in Portugal, as informed by both mosquito population size and estimated
 371 transmission potential. Seasons appeared to occur between the months of March and
 372 November of each year, often peaking in the summer and reaching minimum transmission
 373 potential values during the winter months. We calculated the peak timing for each region using
 374 the index *P* time series of all counties, and found that historically peaks occur in July, although
 375 on average ≈ 12 days later in the south compared to the north (t-test $p < 0.01$, see **Methods** for
 376 details). When comparing the northern and southern districts for a longer time period (2009-
 377 2019, **Figures 3EF**), we found that seasonal variation in index *P* typically presented higher

378 values during the summer months in the north compared to the south ($\approx 35\%$ higher peaks, t -
379 test $p < 0.001$). However, compared to the northern districts, the southern districts presented
380 wider seasonal waves, with transmission potential increasing earlier in the spring and
381 remaining non-zero well into late autumn (**Figure 3F**). The higher values of P in the north,
382 suggesting higher summer transmission potential compared to the south, contrasted with the
383 existing evidence for WNV circulation (**Figure 1**). However, the long-term time distribution of
384 index P seasonal waves in the north was skewed, with lower medians than in the south (**Figure**
385 **3G**), potentially supporting the north-south divide found in past evidence for WNV circulation in
386 the country (**Figure 1**).




387 **Figure 4 - Estimated WNV transmission potential across Portugal (2016-2019).** Maps present summary
388 statistics of estimated index P (WNV transmission potential) at the spatial resolution available in the
389 climate dataset. Panels A, C, E and G present the yearly median index P per coordinate according to the
390 presented color scale on the right. Panels B, D, F and H present the number of months during each year for
391 which the estimated index P per coordinate was above zero, according to the presented color scale on the
392 right. Boundaries in dark grey represent districts.

393 To explore in detail the regional differences in transmission potential across the country, we
394 next estimated the index P per available latitude-longitude coordinate in our climate dataset for
395 the time period between 2016 and 2019 (**Figure 4**). Yearly summary statistics per latitude-
396 longitude coordinate revealed a north-south divide similar to the one presented by past
397 evidence of WNV circulation in the country (**Figure 1**). In particular, we found the southern
398 region of Portugal to universally present higher median yearly index P when compared to the

399 north (**Figures 4ACEG**). However, the geographical ‘boundary’ between north and south, in
 400 terms of WNV transmission potential, changed over the years (compare e.g. **Figures 4A to 4C**).
 401 This output showed that, in accordance to previously described geo-temporal patterns for other
 402 mosquito-borne viruses [34] and WNV [41] in different countries, natural climate variations
 403 strongly influence both inter- and intra-yearly variation in transmission potential. When
 404 quantifying the number of months within a year for which the estimated transmission potential
 405 was not zero, we again found a clear division between the north and south (**Figures 4BDFH**).
 406 Regions in the south and south-west typically spent nine or more months of the year with non-
 407 zero transmission potential, contrasting to regions in the north and north-east which spent only
 408 half or less of the year with non-zero transmission potential. Both the higher yearly median and
 409 longer seasonal waves in the south compared to the north, could thus be seen to support the
 410 mentioned north-south divide in past evidence for WNV circulation in Portugal (**Figure 1**),
 411 hinting that the south may provide particularly favourable conditions for the virus.

412 *Long-term WNV suitability and climatic trends in Portugal*

413 Estimating transmission potential of a pathogen via the index P , while informed by climatic
 414 variables, provides means to reconstruct historical trends when long-term climatic data is
 415 available. We thus used the past 40 years of annual climate data per Portuguese county to
 416 explore long-time trends in climate variables and estimated WNV transmission potential, using
 417 linear quantile mixed models (see **Methods** for technical details).



418 **Figure 5 - Long-term trends and periodicity of transmission potential of WNV and climate across different**
 419 **regions of Portugal.** Panels A, B, and C present the median of climatic variables among counties (points)
 420 within the north (blue) and south (orange) regions defined by the set of districts listed in the map subpanel
 421 (on the right). Panel D presents the same as A-C but for the index P (transmission potential). Panel E
 422 presents the cross-wavelet transform of the median yearly transmission potential between north and south
 423 time-series. Significant periodic components are circled in white (p -value < 0.01), where the heat map
 424 denotes power with peaks indicated by black dots (wavelet ridges) and mean power throughout time shown
 425 in the right subplot.

426 We found that all four explored variables presented significant differences between the north
 427 and south regions of the country (**Figure 5**). Among these, precipitation had the smallest

428 change over time, with a yearly decrease of $\approx -1.2 \times 10^{-6}$ in the north, and increase of
429 $\approx 4.8 \times 10^{-6}$ in the south (**Figure 5A**). Following the Köppen climate classifications, temperature
430 was historically warmer in the south compared to the north. There were similar yearly increases
431 of ≈ 0.041 (north) and ≈ 0.039 (south) Celsius per year between 1981 and 2019, equating to
432 cumulative changes of ≈ 1.55 and ≈ 1.44 degrees for the north and south, respectively (**Figure**
433 **5B**). As the country warmed, it also became dryer, with an accumulated decrease in relative
434 humidity of $\approx 2.8\%$ in the north and $\approx 1.6\%$ in the south (**Figure 5C**). Such historical climate
435 trends translated into an yearly increase in estimated transmission potential for both regions
436 (**Figure 5D**). The index P had an yearly increase of ≈ 0.00097 , accumulating a change of ≈ 0.038
437 between 1981 and 2019. During this period, the south had an absolute difference of ≈ 0.083 in
438 estimated transmission potential compared to the north.

439 As described in **Figure 3**, the northern and southern regions presented an evident yearly
440 periodicity in climate-driven transmission potential, with cycles being stable throughout the
441 years and in phase between the regions. To explore the possibility of periodicity beyond the 12
442 month cycles, we performed wavelet analysis on the north and south yearly median of
443 transmission potential (see **Methods** for details). The analysis revealed longer periodic
444 components in both regions, which were seen to change between 1981 and 2019 (**Figure 5E**).
445 Before 2005, periodicity was mostly found in ≈ 8 year cycles, although a strong 2-3 year
446 component was present around 1995. From 2005 onward, periodicity shifted to become
447 strongly dominated not by one but two components at 2-3 and 4-5 years. The presence of such
448 quasi-periods indicates a Moran effect [65], i.e. that climate synchronizes transmission
449 potential across large geographical areas in Portugal beyond the classic intra-year, summer
450 winter cycles. Finally, the significant shifts in the periodic components over the years do not
451 support an historically stable influence of climate, but are rather suggestive that inter-year
452 instability is an expected feature of WNV transmission potential in Portugal.

453 **Discussion**

454 Recently, several European countries in the Mediterranean basin have quickly progressed from
455 sporadically reporting WNV infections to reporting yearly epidemic activity affecting both
456 humans and equines (e.g. Spain, France, Greece [20]). To date, sufficient evidence has
457 accumulated to support ongoing circulation of WNV in Portugal. However, despite Portugal
458 sharing similar climate types and bird migration routes with nearby countries that have
459 experienced WNV introduction and local dissemination, no events of sustained epidemic
460 activity have been reported. Without such reports, the perception of the public health
461 importance of WNV in Portugal remains minimal. Although several studies have focused on
462 describing WNV human, animal and mosquito infections in the country, its epidemiology
463 remains largely uncharacterised and poorly understood. In this study we have generated an
464 historical, comprehensive perspective of the virus in Portugal using a combination of existing
465 data and climate-informed theoretical approaches.

466 We collated existing data from the research literature, national databases and recent
467 serological samples in equines representative of the time period between 1966 and 2020.
468 Recent data revealed that *Culex pipiens*, considered to be the most suitable species for WNV
469 transmission in Portugal, is present across the entire country. In contrast, we found that the

470 vast majority of WNV-related evidence has been reported in the districts of Faro, Beja, Setúbal,
471 Évora, Lisboa, Portalegre, and Santarém. This group of districts creates a clear north-south
472 divide at the ≈ 39.5 latitude axis of the country, suggesting that past or ongoing WNV circulation
473 is restricted to the south of Portugal. Coincidentally, such a divide is representative of the known
474 geographical separation of the two Köppen climate classifications that exist in the country, with
475 the north being classified as Temperate Mediterranean and the south as Warm Mediterranean.

476 Climate has long been accepted as a major driver of WNV epidemic activity and dissemination,
477 given its effects on both the mosquito and avian species' population dynamics. To consolidate
478 and complement existing data, we explored the effects of local climate variation on theoretical
479 estimations of WNV transmission potential over the years. We estimated that, similarly to other
480 countries in the northern hemisphere, WNV transmission seasons across the country tended to
481 occur between March and November (Spring, Autumn), often peaking in July (Summer). Our
482 analyses identified quantitative differences between the north and the south of Portugal,
483 supporting the existing north-south divide in evidence of WNV circulation. Namely, we found
484 that inter- and intra-year median transmission potential was substantially higher in the south
485 when compared to the north. This implies that the south presents longer seasons suitable for
486 WNV transmission, while most of the year is unsuitable for WNV transmission in the north of
487 Portugal.

488 In light of the ongoing discussion of the effects of climate change on the long-term
489 epidemiology of mosquito-borne viruses (including WNV), we explored historical climate
490 variations and the corresponding WNV transmission potential in Portugal. Differences between
491 the north and south were clear and remained similar over the past 40 years. In general, we
492 presented evidence that Portugal has slowly changed towards a warmer and dryer climate over
493 this period. Temperature presented a especially large change across the years, with a median
494 increase of +1 degree Celsius. Temperature remains a critical factor in mosquito-borne
495 epidemiology, often positively affecting a large number of viral and entomological traits that
496 favour transmission potential. Key examples are the positive relationship of higher
497 temperatures with a shorter viral extrinsic incubation period, longer adult mosquito life-span
498 and shorter aquatic mosquito development [31]. Following the aforementioned historical trends
499 in climate, we also found a small but continuous increase in transmission potential over the
500 years. Our analyses also suggest that recent changes in climate, since 2005, appear to have
501 introduced unstable inter-year variations in transmission potential, the practical consequences
502 of which are difficult to assess.

503 The data and modelling outputs presented include some limitations. For example, the
504 geotemporal patterns of transmission potential reflect solely the contribution of natural climate
505 variation. Our estimations do not take into consideration other factors that could affect WNV
506 transmission suitability in Portugal. These may include the geographical distribution of
507 adequate terrain and vegetation (e.g. wetland, marshland) that favour avian-vector mixing and
508 their proximity to human populations, or geo-temporal hotspots for inbound migratory avian
509 species, vector and avian population sizes, biotype composition of the *Culex pipiens*
510 populations, etc. Such factors should be the basis of future research extending the methods
511 and analyses presented in this study, when such data becomes available at adequate
512 resolutions. The original resolution of the climate data has also restricted some of our outputs.
513 Namely, the resolution of our maps may have missed small-area regions of importance, and we

514 were only able to estimate the duration of seasons (non-zero transmission potential) in the
515 scale of months. The collated empirical data of past WNV circulation may be another source of
516 limitations. For instance, some evidence was impossible to map geographically, given
517 deficiencies in reporting (e.g. no region, or non-specific regions e.g. “south” or “litoral”). It also
518 remains unknown if the data includes a sampling bias towards the south of the country, which
519 could partially explain the seemingly north-south divide in WNV circulation. This bias could
520 arise from the reactive nature of surveillance initiatives responding to reports of symptomatic
521 disease in sentinel species which may be distributed differently across the country, or
522 alternatively, due to differences in WNV awareness among farmers and veterinarians.

523 The large European WNV epidemic of 2018 provided key opportunities to assess existing WNV
524 surveillance and public health mechanisms. Countries implementing cross-sectoral and cross-
525 disciplinary early warning systems based on One Health surveillance (vector, avian, equine and
526 human) consistently reported positive impacts of such infrastructures [19]. Common areas for
527 possible improvement included investment and sustainability of mosquito surveillance, and
528 improved information and media management. In Portugal, mosquito surveillance remains an
529 exception in an otherwise fragmented and insufficient WNV surveillance infrastructure. Key
530 improvements would include sustainability of laboratory capacities and their timely responses,
531 active One Health surveillance beyond the mosquito, and investment in awareness and training
532 across public health domains and the general public, in particular in the south of the country.

533 Our study provides a comprehensive data-driven WNV perspective, and a first spatio-temporal
534 assessment of the contribution of local climate to the theoretical transmission potential of the
535 virus. The ubiquitous countrywide oscillations in transmission potential between summer and
536 winter months demonstrate that climate can couple WNV seasons throughout Portugal,
537 independently of the absolute transmission potential of each region. Consequently, concerted
538 surveillance and control actions spanning the entire territory will be required in the future to
539 better understand and tackle WNV spread. Portugal is slowly accruing suitability for WNV
540 transmission, and hence a shift from passive to active One Health surveillance will be
541 necessary to prevent and manage future epidemics. Our results carry implications for future
542 research, surveillance and public health policy.

543 **Conclusion**

544 Past data suggests that WNV circulates mostly in the south of Portugal, infecting birds,
545 equines, mosquitoes, and occasionally humans. In contrast, mosquito data shows that the
546 spatial distribution of WNV-capable species includes all districts, from north to south. Natural
547 differences in climate types between the north and the south contribute to relevant quantitative
548 differences in WNV theoretical transmission potential, supporting the geographical divide
549 observed in data. It is still uncertain if WNV circulates endemically in any region of the country,
550 but sufficient evidence exists to support ongoing seasonal circulation. The south of Portugal
551 may be more suitable for WNV transmission, but the possibility of the north supporting
552 transmission in shorter periods of each year can not be rejected. A shift towards active
553 surveillance in sentinel species, and awareness and training initiatives are required to timely
554 detect and avert future sustained transmission events that may result in human public health
555 emergencies, as reported recently for other countries in Europe.

556 Tables

Group	PUBMED [in English]	Manual [in English & Portuguese]
WNV in animals, in Portugal (serology / viral isolation / VNT)	[50,51,56,59,66]	[50,52,55,67]
WNV in vectors, in Portugal (viral isolation)	[53,57,58,66,68–70]	[55]
WNV in humans, in Portugal (serology / viral isolation / VNT)	[63,66]	[54,62]
WNV vectors in Portugal but no WNV research	[28,71–78]	[79]
WNV modelling including regions of Portugal	[80,81]	
WNV phylogenetics in Portugal	[70]	

557 **Table 1 - Studies including information on WNV or its vectors in Portugal.**

Sample	Type	N	Region	Period	Study
Equine	VNT	7/24	Aljustrel	Summer 1970	[52]
Human	HI	38/1649	Country wide	1969 - 1973	[54]◆
Mammal	HI	16%	Portalegre, Évora, Beja	Sep 1966 - Feb 1967	[51]◆
Mammal	HI	3%	Portalegre, Évora, Beja	Sep 1966 - Feb 1967	[51]◆
Vector	PCR	pool	Ancão, Almancil	Jul 2004	[57]◆
Vector	Isolation	pool	Barragem do Roxo	Sept 1969	[53]
Human	Serology	2	Faro	Jun - Jul 2004	[61]
Human	Serology	1	Setúbal	Jul 2010	[62]
Equine	VNT	40/1313	Santarém, Aveiro, Portalegre, Évora, Setúbal, Algarve	2004 - 2010	[56]▶
Avian	VNT	23/116	Vila Real, Lisboa, Setúbal, Algarve	2004 - 2010	[56]▶
Equine	VNT	22/23	Loulé	Aug - Nov 2015	[59]▶
Equine	VNT	1/1	Faro	Aug - Oct 2015	[59]▶
Equine	VNT	1/1	Olhão	Sept 2015	[59]▶
Equine	VNT	8/9	Lagos	Sept - Oct 2015	[59]▶
Equine	VNT	4/4	Alcácer do Sal	Sept - Oct 2015	[59]▶
Equine	VNT	1/1	Arronches	Oct 2015	[59]▶
Equine	VNT	1/1	Alpiarça	Oct 2015	[59]▶

Equine	VNT	2/2	Loulé	Feb - Oct 2016	[59] ▶
Equine	VNT	1/1	Lagos	Jul - Dec 2016	[59] ▶
Equine	VNT	3/3	Silves	Oct - Dec 2016	[59] ▶
Equine	VNT	1/1	Odemira	May 2016	[59] ▶
Equine	VNT	1/1	Salvaterra de Magos	Jun - Dec 2016	[59] ▶
Equine	VNT	2/2	Portalegre	Sep - Nov 2016	[59] ▶
Equine	VNT	1/1	Alter-do-chão	Oct 2016	[59] ▶
Equine	VNT	1/1	Benavente	Nov 2016	[59] ▶
Equine	VNT	1/1	Elvas	Nov - Dec 2016	[59] ▶
Equine	VNT	1/1	Beja	Nov - Dec 2016	[59] ▶
Vector	PCR		Ria Formosa	Jul 2004	[58]
Avian	HI	1/4	Parque Nacional da Peneda-Gerês	1999 - 2002	[55] ◆
Avian	HI	1/19	Parque Natural do Vale do Guadiana	1999 - 2002	[55] ◆
Avian	HI	3/44	Parque Ecológico de Monsanto	1999 - 2002	[55] ◆
Avian	HI	2/12	Parque Ecológico de Monsanto	1999 - 2002	[55] ◆
Avian	HI	1/4	Parque Nacional da Peneda-Gerês	1999 - 2002	[55] ◆
Avian	HI	1/4	Parque Ecológico de Monsanto	1999 - 2002	[55] ◆
Avian	HI	2/6	Parque Nacional da Peneda-Gerês	1999 - 2002	[55] ◆
Avian	HI	1/3	Reserva Natural de Santo André	1999 - 2002	[55] ◆
Avian	HI	1/3	Reserva Natural de Santo André	1999 - 2002	[55] ◆
Avian	HI	3/13	Parque Nacional da Peneda-Gerês	1999 - 2002	[55] ◆
Equine	Serology	3/41	Santarém	1999 - 2002	[55] ◆
Human	VNT	1	Almancil	Jul 2015	[63]
Equine	Serology	1/18	Loulé	Aug 2015	[60]
Equine	Serology	1/54	Faro	Aug 2015	[60]
Equine	Serology	2/10	Loulé	Aug 2015	[60]
Avian	ELISA	8/123	Reserva Natural de Santo André, Tagus estuary, Sado estuary	2006, 2008 - 2010	[67] ◆

Avian	ELISA	5/46	Reserva Natural de Santo André, Tagus estuary, Sado estuary	2006, 2008 - 2010	[67]♦
Avian	ELISA	9/213	Reserva Natural de Santo André, Tagus estuary, Sado estuary	2006, 2008 - 2010	[67]♦
Equine	ELISA / PCR	23/165	Loulé, Portalegre, Silves, Alardo-Chão, Lagoa, Elvas, Lagos, Benavente, Alcácer do Sal, Alcáçovas, Samora Correia, Albufeira, Escalos de Cima-Castelo Branco, Comporta	Aug 2016 - Sep 2020	This study

558 ▶: Also IgM and prE-IgG serology, VNT only used when confirmed serology; ♦: positive, implying Flavivirus cross-
559 reactivity, but no WNV confirmed by other means; HI: hemagglutinin inhibition; VNT: virus neutralization test; PCR:
560 polymerase chain reaction; ELISA: enzyme-linked immunosorbent assay.

561 **Table 2 - Sources including past evidence for WNV circulation in Portugal.**

Source	Region, period	Study
<i>Aedes berlandi</i> , <i>Aedes caspius</i> , <i>Anopheles atroparvus</i> , <i>Anopheles claviger</i> , <i>Culex hortensis</i> , <i>Culex laticinctus</i> , <i>Culex mimeticus</i> , <i>Culex pipiens</i> , <i>Culex theileri</i> , <i>Culex univittatus</i> , <i>Culiseta annulata</i> , <i>Culiseta longiareolata</i> , <i>Culiseta sunbochrea</i>	Alqueva, monthly, 2004-2005, 2007	[68]
<i>Aedes caspius</i> , <i>Aedes detritus</i> , <i>Anopheles atroparvus</i> , <i>Culex pipiens</i> , <i>Culex theileri</i> , <i>Culex univittatus</i> , <i>Culiseta annulata</i> , <i>Culiseta longiareolata</i> , <i>Culiseta subochrea</i>	Comporta, monthly, 2004-2005	[68]
<i>Aedes caspius</i> , <i>Aedes detritus</i> , <i>Anopheles atroparvus</i> , <i>Anopheles algerienis</i> , <i>Coquillettidia richiardii</i> , <i>Culex modestus</i> , <i>Culex pipiens</i> , <i>Culex theileri</i> , <i>Culex univittatus</i> , <i>Culiseta longiareolata</i>	Algarve, monthly, Jun Sep 2007	[79]
<i>Culex pipiens</i> , <i>Ochlerotatus caspius</i>	Sado estuary, Ria Formosa, monthly, May Oct 2005, 2006	[79]
<i>Culex pipiens</i>	Entire country, 2016 - 2019	REVIVE [27]

562 **Table 3 - Articles including time series of mosquito population dynamics in Portugal.**

Location	Host	Genomic region	Year	Accession ID
Algarve	Mosquitoes	NS5	2004	AJ965631
Algarve	Mosquitoes	NS5	2004	AJ965627
Algarve	Mosquitoes	Polyprotein precursor	2004	AJ965630
Algarve	Mosquitoes	Polyprotein	2004	AJ965629

		precursor		
Algarve	Mosquitoes	Polyprotein precursor	2004	AJ965628
Algarve	Mosquitoes	Polyprotein precursor	2004	AJ965626
Alentejo	Mosquitoes	Polyprotein precursor	1969	AM404308
Alentejo	Mosquitoes	Env	1969	AY727828

563 **Table 4 - Existing WNV genetic sequences from Portugal.**

Dependent variable	Year slope (95% CI)	South intercept (95%CI)	South*year slope (95% CI):
Index P	9.8×10^{-4} ($9.4 \times 10^{-4}, 10^{-3}$)	0.082 (0.077,0.088)	-----
Humidity (relative, %)	-0.07 (-0.08, -0.06)	-65 (-94, -36)	0.03 (0.015,0.045)
Temperature (Celsius)	0.04 (0.04,0.04)	9.6 (5.3,13.9)	-0.003 (-0.005, -0.001)
Precipitation (m day ⁻¹)	-1.2×10^{-6} ($-1.8 \times 10^{-6}, -6.6 \times 10^{-7}$)	-0.013 (-0.015, -0.014)	6×10^{-6} ($5 \times 10^{-6}, 7 \times 10^{-6}$)

564 **Table 5 - Statistical output from the linear quantile mixed modelling.**

565 Acknowledgements

566 DSCD was funded by the grant 19/23343-7 and 20/06160-3 within the scope of 15/22308-2
567 from the São Paulo Research Foundation (FAPESP). JL was supported by a research
568 lectureship by the Department of Zoology, University of Oxford. MG was supported by
569 Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ). We are grateful to The
570 Ministry of Health / National Institute of Health (INSA) under the National Vector Surveillance
571 Network - REVIVE for supporting this research. We are also grateful to the REVIVE team for the
572 mosquito collection nationwide, and to Dr Patrícia Rosa Ramos Rodrigues for the collection of
573 the reported CSF equine sample.

574 References

- 575 1. Granwehr BP, Lillibridge KM, Higgs S, Mason PW, Aronson JF, Campbell GA, et al. West Nile
576 virus: where are we now? *Lancet Infect Dis.* 2004;4: 547–556.
- 577 2. Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet Infect Dis.* 2002;2:
578 519–529.
- 579 3. Petersen LR, Brault AC, Nasci RS. West Nile virus: review of the literature. *JAMA.* 2013;310.
580 doi:10.1001/jama.2013.8042

- 581 4. Gamino V, Höfle U. Pathology and tissue tropism of natural West Nile virus infection in
582 birds: a review. *Veterinary Research*. 2013. p. 39. doi:10.1186/1297-9716-44-39
- 583 5. Bunning ML, Bowen RA, Cropp CB, Sullivan KG, Davis BS, Komar N, et al. Experimental
584 infection of horses with West Nile virus. *Emerg Infect Dis*. 2002;8.
585 doi:10.3201/eid0804.010239
- 586 6. Hayes EB, Sejvar JJ, Zaki SR, Lanciotti RS, Bode AV, Campbell GL. Virology, pathology, and
587 clinical manifestations of West Nile virus disease. *Emerg Infect Dis*. 2005;11: 1174–1179.
- 588 7. Saiz J-C. Animal and Human Vaccines against West Nile Virus. *Pathogens*. 2020;9.
589 doi:10.3390/pathogens9121073
- 590 8. Rizzoli A, Tagliapietra V, Cagnacci F, Marini G, Arnoldi D, Rosso F, et al. Parasites and
591 wildlife in a changing world: The vector-host- pathogen interaction as a learning case.
592 *International Journal for Parasitology: Parasites and Wildlife*. 2019. pp. 394–401.
593 doi:10.1016/j.ijppaw.2019.05.011
- 594 9. Wang Y, Yim SHL, Yang Y, Morin CW. The effect of urbanization and climate change on the
595 mosquito population in the Pearl River Delta region of China. *Int J Biometeorol*. 2020;64:
596 501–512.
- 597 10. Braack L, Gouveia de Almeida AP, Cornel AJ, Swanepoel R, de Jager C. Mosquito-borne
598 arboviruses of African origin: review of key viruses and vectors. *Parasites and Vectors*.
599 2018;11: 29.
- 600 11. Johnson N, Fernández de Marco M, Giovannini A, Ippoliti C, Danzetta ML, Svartz G, et al.
601 Emerging Mosquito-Borne Threats and the Response from European and Eastern
602 Mediterranean Countries. *Int J Environ Res Public Health*. 2018;15.
603 doi:10.3390/ijerph15122775
- 604 12. Lourenço J, Maia de Lima M, Faria NR, Walker A, Kraemer MU, Villabona-Arenas CJ, et al.
605 Epidemiological and ecological determinants of Zika virus transmission in an urban
606 setting. *Elife*. 2017;6. doi:10.7554/eLife.29820
- 607 13. Giovanetti M, Faria NR, Lourenço J, Goes de Jesus J, Xavier J, Claro IM, et al. Genomic and
608 Epidemiological Surveillance of Zika Virus in the Amazon Region. *Cell Rep*. 2020;30: 2275–
609 2283.e7.
- 610 14. Faria NR, Kraemer MUG, Hill SC, Goes de Jesus J, Aguiar RS, Iani FCM, et al. Genomic and
611 epidemiological monitoring of yellow fever virus transmission potential. *Science*. 2018;361:
612 894–899.
- 613 15. Wu JT, Peak CM, Leung GM, Lipsitch M. Fractional dosing of yellow fever vaccine to extend
614 supply: a modelling study. *Lancet*. 2016;388: 2904–2911.
- 615 16. Murgue B, Zeller H, Deubel V. The ecology and epidemiology of West Nile virus in Africa,
616 Europe and Asia. *Curr Top Microbiol Immunol*. 2002;267: 195–221.
- 617 17. Pybus OG, Suchard MA, Lemey P, Bernardin FJ, Rambaut A, Crawford FW, et al. Unifying the
618 spatial epidemiology and molecular evolution of emerging epidemics. *Proc Natl Acad Sci U*
619 *S A*. 2012;109: 15066–15071.
- 620 18. Haussig JM, Young JJ, Gossner CM, Mezei E, Bella A, Sirbu A, et al. Early start of the West
621 Nile fever transmission season 2018 in Europe. *Euro Surveill*. 2018;23. doi:10.2807/1560-
622 7917.ES.2018.23.32.1800428

- 623 19. Riccardo F, Bolici F, Fafangel M, Jovanovic V, Socan M, Klepac P, et al. West Nile virus in
624 Europe: after action reviews of preparedness and response to the 2018 transmission
625 season in Italy, Slovenia, Serbia and Greece. *Global Health*. 2020;16: 47.
- 626 20. Bakonyi T, Haussig JM. West Nile virus keeps on moving up in Europe. *Eurosurveillance*.
627 2020;25. doi:10.2807/1560-7917.ES.2020.25.46.2001938
- 628 21. Vlaskamp DRM, Thijsen SFT, Reimerink J, Hilkens P, Bouvy WH, Bantjes SE, et al. First
629 autochthonous human West Nile virus infections in the Netherlands, July to August 2020.
630 *Eurosurveillance*. 2020;25: 2001904.
- 631 22. West Nile virus in Europe in 2020 - human cases compared to previous seasons, updated 8
632 October 2020. 9 Oct 2020 [cited 27 Jan 2021]. Available:
633 [https://www.ecdc.europa.eu/en/publications-data/west-nile-virus-europe-2020-human-](https://www.ecdc.europa.eu/en/publications-data/west-nile-virus-europe-2020-human-cases-compared-previous-seasons-updated-8)
634 [cases-compared-previous-seasons-updated-8](https://www.ecdc.europa.eu/en/publications-data/west-nile-virus-europe-2020-human-cases-compared-previous-seasons-updated-8)
- 635 23. Weekly updates: 2020 West Nile virus transmission season. [cited 27 Jan 2021]. Available:
636 [https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data/disease-](https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data/disease-data-ecdc)
637 [data-ecdc](https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data/disease-data-ecdc)
- 638 24. Council Directive 82/894/EEC of 21 December 1982 on the notification of animal diseases
639 within the Community. In: EUR-Lex [Internet]. [cited 11 Jan 2021]. Available: [https://eur-](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:31982L0894)
640 [lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:31982L0894](https://eur-lex.europa.eu/legal-content/EN/ALL/?uri=CELEX:31982L0894)
- 641 25. European Food Safety Authority. [cited 11 Jan 2021]. Available:
642 <https://www.efsa.europa.eu/en>
- 643 26. European Centre for Disease Prevention and Control - West Nile virus. [cited 11 Jan 2021].
644 Available: <https://www.ecdc.europa.eu/en/west-nile-virus-infection>
- 645 27. REVIVE - Rede de Vigilância de Vetores. [cited 15 Jan 2021]. Available:
646 [http://www2.insa.pt/sites/INSA/Portugues/AreasCientificas/DoencasInfecciosas/](http://www2.insa.pt/sites/INSA/Portugues/AreasCientificas/DoencasInfecciosas/AreasTrabalho/EstVectDoencasInfecciosas/Paginas/Revive.aspx)
647 [AreasTrabalho/EstVectDoencasInfecciosas/Paginas/Revive.aspx](http://www2.insa.pt/sites/INSA/Portugues/AreasCientificas/DoencasInfecciosas/AreasTrabalho/EstVectDoencasInfecciosas/Paginas/Revive.aspx)
- 648 28. Osório HC, Zé-Zé L, Amaro F, Alves MJ. Mosquito surveillance for prevention and control of
649 emerging mosquito-borne diseases in Portugal - 2008-2014. *Int J Environ Res Public*
650 *Health*. 2014;11: 11583–11596.
- 651 29. European network for sharing data on the geographic distribution of arthropod vectors,
652 transmitting human and animal disease agents (VectorNet). [cited 1 Feb 2021]. Available:
653 [https://www.ecdc.europa.eu/en/about-us/partnerships-and-networks/disease-and-](https://www.ecdc.europa.eu/en/about-us/partnerships-and-networks/disease-and-laboratory-networks/vector-net)
654 [laboratory-networks/vector-net](https://www.ecdc.europa.eu/en/about-us/partnerships-and-networks/disease-and-laboratory-networks/vector-net)
- 655 30. Lourenco J. MVSE - WNV related files for Portugal. 2021 [cited 26 Jan 2021].
656 doi:10.6084/m9.figshare.c.5281664.v1
- 657 31. Mordecai EA, Cohen JM, Evans MV, Gudapati P, Johnson LR, Lippi CA, et al. Detecting the
658 impact of temperature on transmission of Zika, dengue, and chikungunya using
659 mechanistic models. *PLoS Negl Trop Dis*. 2017;11: e0005568.
- 660 32. Jiguet F, Devictor V, Ottvall R, Van Turnhout C, Van der Jeugd H, Lindström A. Bird
661 population trends are linearly affected by climate change along species thermal ranges.
662 *Proc Biol Sci*. 2010;277: 3601–3608.
- 663 33. Cator LJ, Johnson LR, Mordecai EA, Moustaid FE, Smallwood TRC, LaDeau SL, et al. The
664 Role of Vector Trait Variation in Vector-Borne Disease Dynamics. *Front Ecol Evol*. 2020;8.

- 665 doi:10.3389/fevo.2020.00189
- 666 34. Obolski U, Perez PN, Villabona-Arenas CJ, Thézé J, Faria NR, Lourenço J. MVSE: An R-
667 package that estimates a climate-driven mosquito-borne viral suitability index. *Methods*
668 *Ecol Evol.* 2019;10: 1357–1370.
- 669 35. Kraemer MUG, Sinka ME, Duda KA, Mylne AQN, Shearer FM, Barker CM, et al. The global
670 distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Elife.* 2015;4:
671 e08347.
- 672 36. Hamlet A, Jean K, Perea W, Yactayo S, Biey J, Van Kerkhove M, et al. The seasonal
673 influence of climate and environment on yellow fever transmission across Africa. *PLoS*
674 *Negl Trop Dis.* 2018;12: e0006284.
- 675 37. Petrone ME, Earnest R, Lourenço J, Kraemer MUG, Paulino-Ramirez R, Grubaugh ND, et al.
676 Asynchronicity of endemic and emerging mosquito-borne disease outbreaks in the
677 Dominican Republic. *Nat Commun.* 2021;12: 151.
- 678 38. Thézé J, Li T, du Plessis L, Bouquet J, Kraemer MUG, Somasekar S, et al. Genomic
679 Epidemiology Reconstructs the Introduction and Spread of Zika Virus in Central America
680 and Mexico. *Cell Host Microbe.* 2018;23: 855–864.e7.
- 681 39. Perez-Guzman PN, Carlos Junior Alcantara L, Obolski U, de Lima MM, Ashley EA, Smithuis
682 F, et al. Measuring Mosquito-borne Viral Suitability in Myanmar and Implications for Local
683 Zika Virus Transmission. *PLoS Curr.* 2018;10.
684 doi:10.1371/currents.outbreaks.7a6c64436a3085ebba37e5329ba169e6
- 685 40. Pereira Gusmão Maia Z, Mota Pereira F, do Carmo Said RF, Fonseca V, Gräf T, de Bruycker
686 Nogueira F, et al. Return of the founder Chikungunya virus to its place of introduction into
687 Brazil is revealed by genomic characterization of exanthematic disease cases. *Emerg*
688 *Microbes Infect.* 2020;9: 53–57.
- 689 41. Lourenço J, Thompson RN, Thézé J, Obolski U. Characterising West Nile virus
690 epidemiology in Israel using a transmission suitability index. *Euro Surveill.* 2020;25.
691 doi:10.2807/1560-7917.ES.2020.25.46.1900629
- 692 42. Copernicus Climate Data Store. [cited 13 Jan 2021]. Available:
693 <https://cds.climate.copernicus.eu/cdsapp#!/dataset/ecv-for-climate-change?tab=overview>
- 694 43. Lourenço J, Obolski U. MVSE R-package official page. [cited 21 Jan 2021]. Available:
695 <https://sourceforge.net/projects/mvse/>
- 696 44. R-Forge: Circular Statistics: Project Home. [cited 25 Jan 2021]. Available: [https://r-forge.r-](https://r-forge.r-project.org/projects/circular/)
697 [project.org/projects/circular/](https://r-forge.r-project.org/projects/circular/)
- 698 45. Geraci M. Linear Quantile Mixed Models: The lqmm Package for Laplace Quantile
699 Regression. *J Stat Softw.* 2014;57: 1–29.
- 700 46. Damineli DSC, Portes MT, Feijó JA. Oscillatory signatures underlie growth regimes in
701 *Arabidopsis* pollen tubes: computational methods to estimate tip location, periodicity, and
702 synchronization in growing cells. *J Exp Bot.* 2017;68: 3267–3281.
- 703 47. wavelets: Functions for Computing Wavelet Filters, Wavelet Transforms and
704 Multiresolution Analyses. [cited 21 Jan 2021]. Available:
705 <https://CRAN.R-project.org/package=wavelets>

- 706 48. biwavelet GitHub repository. [cited 21 Jan 2021]. Available:
707 <https://github.com/tgouhier/biwavelet>
- 708 49. Barros SC, Ramos F, Zé-Zé L, Alves MJ, Fagulha T, Duarte M, et al. Simultaneous detection
709 of West Nile and Japanese encephalitis virus RNA by duplex TaqMan RT-PCR. *J Virol*
710 *Methods*. 2013;193: 554–557.
- 711 50. Filipe AR. Anticorpos contra virus transmitidos por artrópodos-arbovirus do grupo B em
712 animais do Sul de Portugal: inquérito serológico preliminar com o vírus West Nile, estirpe
713 Egypt 101. *Ann Escola Nacional de Saúde Pública de Medicina Tropical*. 1967;1: 197–204.
- 714 51. Filipe AR, Pinto MR. Survey for antibodies to arboviruses in serum of animals from
715 southern Portugal. *Am J Trop Med Hyg*. 1969;18: 423–426.
- 716 52. Filipe AR, Campaniço M. Encefalomielite equina por arbovírus. A propósito de uma
717 epizootia presuntiva causada pelo vírus West Nile. *Revista Portuguesa de Ciências*
718 *Veterinárias*. 1973;LXVIII.
- 719 53. Filipe AR. Isolation in Portugal of West Nile virus from *Anopheles maculipennis*
720 mosquitoes. *Acta Virol*. 1972;16. Available: <https://pubmed.ncbi.nlm.nih.gov/4403183/>
- 721 54. Filipe AR. Anticorpos contra arbovírus na população de Portugal. *Separata de O Médico*.
722 1973;LXVII: 731–732.
- 723 55. Formosinho P, Santos-Silva MM, Santos A, Melo P, Encarnação V, Santos N, et al. O vírus
724 West Nile em Portugal – estudos de vigilância epidemiológica. *Revista Portuguesa de*
725 *Ciências Veterinárias*. 2006;101: 61–68.
- 726 56. Barros SC, Ramos F, Fagulha T, Duarte M, Henriques M, Luís T, et al. Serological evidence
727 of West Nile virus circulation in Portugal. *Vet Microbiol*. 2011;152: 407–410.
- 728 57. Almeida APG, Galão RP, Sousa CA, Novo MT, Parreira R, Pinto J, et al. Potential mosquito
729 vectors of arboviruses in Portugal: species, distribution, abundance and West Nile
730 infection. *Trans R Soc Trop Med Hyg*. 2008;102: 823–832.
- 731 58. Esteves A, Almeida AP, Galão RP, Parreira R, Piedade J, Rodrigues JC, et al. West Nile virus
732 in Southern Portugal, 2004. *Vector Borne Zoonotic Dis*. 2005;5.
733 doi:10.1089/vbz.2005.5.410
- 734 59. Barros SC, Ramos F, Fagulha T, Duarte M, Henriques AM, Waap H, et al. West Nile virus in
735 horses during the summer and autumn seasons of 2015 and 2016, Portugal. *Vet Microbiol*.
736 2017;212: 75–79.
- 737 60. World Organization for Animal Health (OIE) - West Nile reports. In: Information received on
738 03/09/2015 from Prof. Dr Álvaro Mendonça, Director General, Direcção Geral de
739 Alimentação e Veterinária, Ministério da Agricultura E do Mar, Lisboa, Portugal (2015)
740 [Internet]. [cited 18 Jan 2021]. Available:
741 [https://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?](https://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?page_refer=MapFullEventReport&reportid=18585)
742 [page_refer=MapFullEventReport&reportid=18585](https://www.oie.int/wahis_2/public/wahid.php/Reviewreport/Review?page_refer=MapFullEventReport&reportid=18585)
- 743 61. Connell J, Mckeown P, Garvey P, Cotter S, Conway A, O’Flanagan D, et al. Two linked cases
744 of West Nile virus (WNV) acquired by Irish tourists in the Algarve, Portugal. *Weekly*
745 *releases (1997–2007)*. 2004;8: 2517.
- 746 62. Alves MJ, Poças JM, Luz T, Amaro F, Zé-Zé L, Osório H. Infecção por vírus West Nile
747 [Flavivírus] em Portugal. Considerações acerca de um caso clínico de síndrome febril com

- 748 exantema. 2012;8: 46–51.
- 749 63. Zé-Zé L, Proença P, Osório HC, Gomes S, Luz T, Parreira P, et al. Human case of West Nile
750 neuroinvasive disease in Portugal, summer 2015. *Euro Surveill.* 2015;20.
751 doi:10.2807/1560-7917.ES.2015.20.38.30024
- 752 64. Direcção-Geral de Veterinária (Directorate-General of Veterinary). In: National statistics on
753 official number of equines in subregions of Portugal. [Internet]. [cited 13 Jan 2021].
754 Available: [http://srvbamid.dgv.min-agricultura.pt/portal/page/portal/DGV/genericos?](http://srvbamid.dgv.min-agricultura.pt/portal/page/portal/DGV/genericos?actualmenu=23555&generico=33698230&cboui=33698230)
755 [actualmenu=23555&generico=33698230&cboui=33698230](http://srvbamid.dgv.min-agricultura.pt/portal/page/portal/DGV/genericos?actualmenu=23555&generico=33698230&cboui=33698230)
- 756 65. Hansen BB, Grøtan V, Herfindal I, Lee AM. The Moran effect revisited: spatial population
757 synchrony under global warming. *Ecography.* 2020;43: 1591–1602.
- 758 66. Filipe AR, de Andrade HR. Arboviruses in the Iberian Peninsula. *Acta Virol.* 1990;34: 582–
759 591.
- 760 67. Pardal S, Alves JA, Zé-Zé L, Osório H, Rocha A, Lopes RJ, et al. Shorebird low spillover risk
761 of mosquito-borne pathogens on Iberian wetlands. *J Ornithol.* 2013;155: 549–554.
- 762 68. Almeida APG, Freitas FB, Novo MT, Sousa CA, Rodrigues JC, Alves R, et al. Mosquito
763 surveys and West Nile virus screening in two different areas of southern Portugal, 2004-
764 2007. *Vector Borne Zoonotic Dis.* 2010;10: 673–680.
- 765 69. Freitas FB, Novo MT, Esteves A, de Almeida AP. Species Composition and WNV Screening
766 of Mosquitoes from Lagoons in a Wetland Area of the Algarve, Portugal. *Front Physiol.*
767 2012;2. doi:10.3389/fphys.2011.00122
- 768 70. Parreira R, Severino P, Freitas F, Piedade J, Almeida APG, Esteves A. Two distinct
769 introductions of the West Nile virus in Portugal disclosed by phylogenetic analysis of
770 genomic sequences. *Vector Borne Zoonotic Dis.* 2007;7: 344–352.
- 771 71. Fotakis EA, Mastrantonio V, Grigoraki L, Porretta D, Puggioli A, Chaskopoulou A, et al.
772 Identification and detection of a novel point mutation in the Chitin Synthase gene of *Culex*
773 *pipiens* associated with diflubenzuron resistance. *PLoS Negl Trop Dis.* 2020;14: e0008284.
- 774 72. Mixão V, Bravo Barriga D, Parreira R, Novo MT, Sousa CA, Frontera E, et al. Comparative
775 morphological and molecular analysis confirms the presence of the West Nile virus
776 mosquito vector, *Culex univittatus*, in the Iberian Peninsula. *Parasit Vectors.* 2016;9: 601.
- 777 73. Osório HC, Zé-Zé L, Alves MJ. Host-feeding patterns of *Culex pipiens* and other potential
778 mosquito vectors (Diptera: Culicidae) of West Nile virus (Flaviviridae) collected in Portugal.
779 *J Med Entomol.* 2012;49: 717–721.
- 780 74. Gomes B, Parreira R, Sousa CA, Novo MT, Almeida APG, Donnelly MJ, et al. The *Culex*
781 *pipiens* complex in continental Portugal: distribution and genetic structure. *J Am Mosq*
782 *Control Assoc.* 2012;28: 75–80.
- 783 75. Gomes B, Wilding CS, Weetman D, Sousa CA, Novo MT, Savage HM, et al. Limited genomic
784 divergence between intraspecific forms of *Culex pipiens* under different ecological
785 pressures. *BMC Evol Biol.* 2015;15: 197.
- 786 76. Calzolari M, Zé-Zé L, Růžek D, Vázquez A, Jeffries C, Defilippo F, et al. Detection of
787 mosquito-only flaviviruses in Europe. *J Gen Virol.* 2012;93: 1215–1225.
- 788 77. Hernández-Triana LM, Brugman VA, Pramual P, Barrero E, Nikolova NI, Ruiz-Arrondo I, et al.

- 789 Genetic diversity and population structure of *Culex modestus* across Europe: does recent
790 appearance in the United Kingdom reveal a tendency for geographical spread? *Med Vet*
791 *Entomol.* 2020;34: 86–96.
- 792 78. Osório HC, Zé-Zé L, Amaro F, Nunes A, Alves MJ. Sympatric occurrence of *Culex pipiens*
793 (Diptera, Culicidae) biotypes *pipiens*, *modestus* and their hybrids in Portugal, Western
794 Europe: feeding patterns and habitat determinants. *Med Vet Entomol.* 2014;28: 103–109.
- 795 79. Alves JM, Osório H, Zé-Zé L, Amaro F, Luz T, Encarnação V, et al. Flavivírus transmitidos
796 por mosquitos: um risco potencial para Portugal. *Investigação em ambiente e saúde -*
797 *desafios e estratégias* (Universidade de Aveiro). 2009.
- 798 80. Conte A, Candeloro L, Ippoliti C, Monaco F, De Massis F, Bruno R, et al. Spatio-Temporal
799 Identification of Areas Suitable for West Nile Disease in the Mediterranean Basin and
800 Central Europe. *PLoS One.* 2015;10: e0146024.
- 801 81. García-Carrasco J-M, Muñoz A-R, Olivero J, Segura M, Real R. Predicting the spatio-
802 temporal spread of West Nile virus in Europe. *PLoS Negl Trop Dis.* 2021;15: e0009022.