Title: **Status of hypertension control in urban slums of Central India : A Community**

health worker based two-year follow up

Authors

1. Pakhare Abhijit, MD (Community Medicine), Associate Professor, AIIMS Bhopal

2. Lahiri Anuja, PhD Scholar (Public Health), AIIMS Bhopal

3. Shrivastava Neelesh, PhD Scholar, Barkatullah University Bhopal

4. N Subba K, PG (Community Medicine), AIIMS Bhopal

5. Kurra Veera Durga Vaishnavi, (Intern), AIIMS Bhopal

6. Joshi Ankur, MD (Community Medicine), Assistant Professor, AIIMS Bhopal

7. Atal Shubham, MD (Pharmacology), Assistant Professor, AIIMS Bhopal

8. Khadanga Sagar, MD (Medicine), Associate Professor, AIIMS Bhopal

9. Joshi Rajnish, PhD (Public Health), MD (Medicine), HOD and Professor, AIIMS Bhopal

Corresponding Author:

Name: Rajnish Joshi

Address: Baghsewania, AIIMS campus, Saket Nagar, Bhopal (462020), MP

E-mail address: rajnish.genmed@aiimsbhopal.edu.in

Total number of pages: 12 (without references)

Total number of photographs: 2 Figures

Word counts

for abstract: 250

for the text: 2856 (excluding tables, references)

Source(s) of support: ICMR extramural project grant (Grant – PI- Dr Rajnish Joshi, IRIS-

2014-0976)

Conflicting Interest (If present, give more details): Nil

Acknowledgement: Investigators would like acknowledge efforts of all Community Health

Workers (ASHAs) and their officials affiliated to Urban Public Health system in

implementing this study.

Status of hypertension control in urban slums of Central India : A Community health worker based

two-year follow up

Abstract

Background: Hypertension is a leading cause of cardiovascular diseases and its control is poor. There

is heterogeneity in levels of blood-pressure control among various population sub-groups. The present

study was conducted within the framework of National Program for prevention and control of cancer,

diabetes, cardiovascular diseases and stroke (NPCDCS) in India. It aims to estimate proportion of

optimal blood pressure control and identify potential risk factors pertaining to uncontrolled

hypertension consequent to initial screening.

Methods: We assembled a cohort of individuals with hypertension confirmed in a baseline screening

in sixteen urban slum clusters of Bhopal (2017-2018). Sixteen Accredited Social Health Activists

(ASHAs) were trained from within these slums. Individuals with hypertension were linked to primary

care providers and followed-up for next two years. Obtaining optimal blood-pressure control (defined

as SBP< 140 and DBP<90 mm of Hg) was a key outcome.

Results: Of a total of 6174 individuals, 1571 (25.4%) had hypertension, of which 813 were

previously known and 758 were newly detected during baseline survey. Two year follow up was

completed for 1177 (74.9%). Blood-pressure was optimally controlled in 301 (26%) at baseline, and

in 442 (38%) individuals at two years (absolute increase of 12%; 95% CI 10.2-13.9). Older age,

physical-inactivity, higher BMI and newly diagnosed hypertension were significantly associated with

uncontrolled blood-pressure.

Conclusions: We found about six of every ten individuals with hypertension were on-treatment, and

about four were optimally controlled. These findings provide a benchmark for NPCDCS, in terms of

achievable goals within short periods of follow-up.

Key words: hypertension, optimal blood pressure control, community health worker,

physical inactivity

Status of hypertension control in urban slums of Central India : A Community health worker based two-year follow up

Introduction

Hypertension (HTN) is a leading risk factor of cardiovascular diseases and its control is poor. There occurs a loss of numbers at screening-awareness-treatment-control pathway. Among those with hypertension, in urban India less than half are aware of their elevated blood pressure, and only one in five individuals are controlled.(1) Studies from central India found that 22.3% of all individuals in the age group of 18-69 years had hypertension, and only 11.8% of those who had hypertension were being treated.(2) Better control of hypertension depends upon availability and access to quality health-care.(3) Once diagnosed, management of untreated and uncontrolled HTN is challenging. There needs to be a focus on improved adherence.(4) Various engagement strategies have been used, including but not limited to mobile-health applications, (5) telemedicine services, and community health worker (CHW) based engagement strategies.(6) While some population based studies have had a limited impact on blood-pressure levels, (7,8) other facility based studies have shown benefits.(9,10) Clusters where more members have uncontrolled hypertension at baseline, with a higher baseline blood-pressure, have reported greater and more significant reductions.(11) Even amongst a homogenous health-system, variation in blood pressure control exists, attributable to better patient engagement.(12) National Program for prevention and control of cancer, diabetes, cardiovascular diseases and stroke (NPCDCS) in India recommends annual hypertension screening for all adults, and proportion of optimally controlled individuals as a facility level indicator. Aim of the current study is to identify proportion of individuals with optimal blood pressure control and to identify risk-factors for uncontrolled hypertension.

Methods

Design and ethics: We established a community based cohort. The study design was approved by the Institutional Human Ethics Committee (Ref: IHEC-LOP/2017/EF00045) and funded by Indian Council of Medical Research. All participants provided written informed consent prior to initiation of any study procedures.

Setting: The study was conducted in 16 urban slum clusters of Bhopal, a city with population of about

18 Lakhs, located in central India. A baseline door-to-door survey was conducted in the urban slum

clusters between November 2017 and March 2018, details of which are provided elsewhere.(13)

Briefly, all consenting non-pregnant adults above the age of 30 years formed a cross-section, in whom

cardiovascular disease (CVD) risk assessment was performed by Accredited social health activists

(ASHAs), who are CHWs in these communities. Study supervisors provided oversight, and also

confirmed key variables, before stratifying individuals as high risk. Individuals identified at a high

CVD risk, especially those with hypertension and diabetes were sought to be linked to urban primary

health centres (UPHCs) which are primary-care public-health facilities in these communities. All

participants were however free to seek care from alternate private-providers of their choice. Limited

generic anti-hypertensive drugs (Amlodipine, Losartan, and Hydrochlorothiazide) are available at no-

cost from UPHCs with a prescription refill duration varying from two to four weeks. Participants who

visit private primary-care providers usually incur out-of-pocket expenditure for their drug therapy,

with a refill duration dependent on their capacity to pay.

Participants: All individuals residing in selected urban slum clusters of age 30 years or more were

screened for hypertension as part of baseline assessment. Individuals with hypertension (either a

previously known or a newly detected hypertension), were identified in a baseline survey. Individuals

who reported themselves as diagnosed to have hypertension in the past, were defined as having

previously known-hypertension. ASHAs obtained a first-set of blood-pressure values using a digital

sphygmomanometer (Omron digital apparatus, Model 7200, Kyoto, Japan) at home of the

participants. All the blood pressure readings were then confirmed in a second set of readings obtained

by the study supervisor, using the same device. Each set of blood-pressure readings consisted of an

average of three values, obtained one-minute apart, using a standardized technique. A person was

defined as a *newly detected hypertensive*, if the blood pressure values were above Systolic blood

pressure (SBP) of 140 mm Hg or a Diastolic blood pressure (DBP) of 90mm Hg on both the

occasions. This is the definition for diagnosis of HTN under NPCDCS. All individuals with

hypertension (previously known or newly detected) were provided with referrals for physicians

available at UPHC, who initiated them with or optimized their anti-hypertensive medication. All

participants, who completed the two-year follow-up assessment in November-December 2019 are included in the current study.

Procedures: We followed individuals with hypertension (previously known or newly detected) for a period of two years. Follow-up was performed using the existing human resources in the public health-delivery system, facilitated by cloud-based data-management tools, and study supervisors. We prepared a list of all individuals with HTN in a database, and stratified them by cluster. We provided these lists of each cluster to respective ASHAs, who were responsible for visiting each member at-least once in every three-months. Purpose of these visits was to promote adherence to drug therapies, measurement of blood pressure, and to provide focussed CVD-prevention health education. Monetary incentives were given to ASHAs for every completed follow-up (rupees 50 INR about one USD). In each visit, ASHAs recorded linkage to UPHC or a private care provider, and continuation of drug therapy. We defined *treatment interruption* as failure to obtain a prescription-refill for two-weeks or more after exhaustion of previously obtained drug supplies.

We designed a cloud-based digital tool using CommCare based application to facilitate follow-up. This tool was available on mobile phones of all ASHAs, and in addition to facilitate follow-up, it provided access to baseline values of the participants, and also was to be used for data-entry of each subsequent visit. Study supervisors monitored cohort follow-up using this CommCare based application, and performed additional site-visits for troubleshooting. We also conducted a quarterly-meeting of all ASHAs and study supervisors to monitor follow-ups. A community clinic based one-year adherence assessment was performed by a team of pharmacologists who were trained to assess pill-burden and adherence to previously prescribed therapies. This assessment was limited to those who were on any anti-hypertensive drug therapy in the previous month.

Outcome assessment: Blood pressure control at two-year assessment visit was the key outcome. It was defined as *optimally controlled* if average SBP was less than 140mm Hg and DBP was less than 90mm Hg. This two-year assessment was performed by independent field investigators, who visited households of all participants and measured their blood pressures. Multiple mop-up visits were conducted, and facilitation by ASHAs and study supervisors was solicited to improve proportion of follow-ups.

Statistical Analysis: ASHAs and study supervisors entered all study-variables obtained in baseline survey, confirmatory-visits, UPHC-visits, or on follow-up using mobile-phone based data collection system. All the variables were stored on cloud-based CommCare data-management system. The demographic and cardiovascular-risk variables (age, gender, education level, wealth quintiles, tobacco use, alcohol consumption, physical activity levels, body-mass index, waist circumference) were obtained from entries in the baseline questionnaire. History of hypertension obtained at baseline and subsequently obtained SBP and DBP values were used for classification of hypertension status. Primary-care physician consultation decisions culminating in drug escalation, de-escalation or no action, were taken from UPHC visit logs. The distribution of these explanatory variables was compared with respect to outcome, using appropriate tests of significance. The difference was considered as significant if p-value for null hypothesis was less than 0.05.

We performed a multivariable logistic regression, to evaluate independent risk factors of uncontrolled hypertension. All explanatory variables that were marginally significant on univariate analysis (p<0.25) were included in the full model. Assumptions for using logistic regression were tested and Hosmer-Lemeshow Goodness-of-fit was also used. All data analysis and visualizations were done using Statistical software R (14) and *arsenal* ,(15) *finalfit* ,(16) *ggplot2* ,(17) *ggpubr* ,(18) *ggstatplot* ,(19) *gtsummary* ,(20) *performance* (21) and *tidyverse* (22) packages.

Results

In November 2017 we initiated a baseline survey in 16 urban slum clusters of Bhopal. Of 6174 individuals surveyed at baseline, we identified a total of 1571 individuals (25.4%) with hypertension; and all of these were sought to be visited by CHWs once in every two months. A total of 1177 (74.9%) individuals completed two-year follow up visit done in November-December 2019. Individuals who were lost to follow up were more likely to be men, who were older, and were newly detected with hypertension. (Table S1) Of 1177 individuals who completed follow up, 623 (53%) knew about their hypertension status at baseline, and 554 (47%) were newly detected during screening. At baseline, 301 (26%) individuals were optimally controlled, and at the end of two years

their proportion increased to 442 (38%), an absolute increase of 12% (95%CI 10.2-13.9). When we

split these by status of hypertension diagnosis, of the 623 previously known hypertensives, 48% were

optimally controlled at baseline, which became 46% at the end of two years, and of the 554 newly

detected hypertensives, and 29% were optimally controlled at the end of two years. There was a

significant decline of 1.7 (95%CI -2.9 to -0.5) and 1.7mm Hg (95%CI (-2.4 to -1.1) in mean SBP and

DBP respectively from baseline to follow up. This change can be largely attributed to reduction in

SBP among those with uncontrolled hypertension. (Table 1, Figure 1)

The proportion of individuals with uncontrolled blood pressure was progressively higher in each age

band (55%, 63%, 67% and 73% in those with baseline age of 30-44, 45-59, 60-74, and above 75 years

respectively). We observed a significant decline in SBP and DBP, across all age-bands of individuals

with uncontrolled hypertension at baseline. About 65% of individuals who were sedentary at baseline,

developed uncontrolled hypertension, as compared to only 54% of those who were non-sedentary i.e.

physically active. This difference was statistically significant. Individuals who were newly detected to

have hypertension, and those who had elevated blood-pressure at baseline, were more likely to be

uncontrolled on follow-up. (Table 2, Figure 2)

A total of 699 (59.4%) individuals reported to be on anti-hypertensive drug therapy, 533 of these were

linked to UPHC, and 166 to private-care providers. All these individuals received follow up visits by

ASHAs, and during these visits self-reported adherence to prescribed therapies was similar in both

optimally controlled and uncontrolled individuals. (84.4% vs 85.8%; p=0.842) An independent

adherence assessment was also done at the end of one-year in 263 of the UPHC attendees, and

perceived adherence was lower in those subsequently found to be uncontrolled (69.5% vs 79.1%;

p=0.006). The proportion of individuals who were uncontrolled after two years was significantly

higher amongst those who were linked to UPHC as compared to those linked to private-providers

(67% vs 61%; p=0.009). Of those linked to UPHC, median number of visits to the facility was similar

in optimally controlled and uncontrolled subgroups (p=0.436) and 116 (21.7%) individuals had one or

more visit culminating in a drug-escalation advise. While 24 (20.6%) of individuals who were advised

drug-escalation were optimally controlled, remaining 92(79%) were not. Of the 478 of 1177

individuals (40.6%), defaulted for initiation of drug-therapy, and 276 (58%) had their blood pressures

above 140/90 mm Hg at two-year follow up.

We performed multivariable logistic regression to identify risk-factors for uncontrolled hypertension.

Hosmer-Lemeshow *goodness-of-fit* test was non-significant, there was no multi-collinearity,

assumptions of homogeneity of variance, residual distribution and absence of influential observations

were met, however model's explanatory power was weak (Tjur's R2=0.06, RMSE=0.46). After

adjusting for age, and gender newly detected hypertension (OR 2.42 (1.78-3.31)), and BMI (OR 1.04

(1.01-1.07)) were significant independent risk-factors for uncontrolled hypertension. (Table 3)

Discussion

In the current study we found that after initial screening and two-year follow-up, about six of every

ten individuals with hypertension were on-treatment, and about four were optimally controlled. While

overall proportion of individuals with optimally controlled blood-pressure rose over the two-year

period, this modest increase was due to blood-pressure control in less than a-third of newly detected

hypertensives. More than two-thirds of the remaining newly detected and about half of previously

known hypertensives remained uncontrolled. Those who were uncontrolled, were more likely to be

older, had a higher BMI, and were newly detected to have hypertension. These findings demonstrate

that obtaining blood-pressure control in vulnerable communities, though feasible is challenging.

These findings provide a benchmark for NPCDCS, in terms of achievable goals within short periods

of follow-up.

A recent systematic review (23) reported that in India only 13% of all individuals with hypertension

are treated, and a meagre eight percent are controlled. Our baseline screening bridged the awareness

gap, and CHW based adherence reinforcement improved proportion of on-treatment and controlled

individuals from 43% and 29% to 60% and 38% respectively. While these numbers are encouraging,

and approach and even exceed proportions for control achieved in previous studies in high-income

settings,(24,25) they bring-forth challenges of initiating anti-hypertensive drugs in all who need them,

improving adherence in those initiated, and appropriate dosing requirements.(26) In the current study

education, wealth, or gender did not increase risk of uncontrolled hypertension. These socioeconomic

characteristics are perceived as proxy-indicators for awareness, access or empowerment. It is likely

that more complex behavioural phenomenon such as fear, reluctance and adverse impact of being

labelled with an illness, affected both initiation and adherence to prescribed drug therapies.(27) We

found that risk of being uncontrolled was consistently higher with increasing age, a factor that

contributes to therapeutic inertia.(28) Further, presence of fixed-dose combinations in the drug

formulary in UPHCs could have helped reduce inertia,(29) as use of combination treatment in newly

detected hypertensives is likely to achieve a faster blood-pressure control.(24)

Similar findings were reported in recent studies on predictors of hypertension control in primary

health care settings.(30–32)   These studies have reported age, gender, race, Waist-Hip Ratio, BMI,

presence of diabetes mellitus, intake of multiple drugs and lack of knowledge as predictors. (30–32)

Physical activity is important to achieve blood-pressure control, and is an important component of

various population based programs.(33) Self-reported physical activity levels correlate well with

reduced blood pressure levels.(34) In the current study, hypertensive individuals who reported

themselves to be sedentary at baseline were more-likely to be uncontrolled two years later. While this

effect observed on univariate analysis, was not significant when adjusted for age and gender, it is an

important life-style measure that has a strong legacy effect. Improved physical activity achieves blood

pressure reduction through diverse mechanisms such as reduction in vascular resistance, sympathetic

activity, inflammation and stress.(34) It also leads to lower BMI, which was an independent predictor

of blood-pressure control in our study. While some studies, such as SPRINT trial have found no

relationship between BMI and blood-pressure control, (35) another large study from China found a

significant lowering effect.(36) While mean BMI of participants in SPRINT trial was 29 kg/m2, it was

24.4 kg/m2 in the Chinese study. Mean BMI of our participants was 25.8 kg/m2, closer to that of the

Chinese study. It is likely that blood-pressure control in population sub-groups with a lower BMI is

more feasible, as compared to those with obesity.

Improving blood pressure control is challenging, and regular follow up with intensive

pharmacotherapy is most useful to achieve this goal. In a Cochrane review of 56 randomized control

trials, other measures such as self-monitoring, appointment reminders, and health-education had little

independent effects and were less useful.(37) Feasibility of achieving even lower targets than

envisaged in NPCDCS, was demonstrated in intensive blood-pressure control arm in SPRINT

trial.(38) Similar intensive management is also possible through CHWs in low-income settings. Two

recent trials have demonstrated this effect. First, HOPE4 study demonstrated that CHW based

intensive blood pressure control can improve proportion of optimally controlled individuals to 69% as

compared to 30% in standard care arm.(39) Second, in another randomized control trial from

Argentina, CHW led multi-dimensional care that involved intensive health-coaching, home BP

monitoring, audits and feedbacks optimally controlled 72% individuals.(9) When CHWs are provided

with intensive supportive supervision, empowered to obtain quick feedbacks on treatment decisions, it

can help improve therapeutic inertia and achieve better control levels.(40)

Conclusion

We have established a cohort of individuals with hypertension, with an aim to improve CHW based

care delivery in vulnerable populations and generate evidence that may help implementation of

NPCDCS. We used efficient data-management systems, within the programmatic settings to establish

a system of community-based follow-up. Our study however has had some limitations. We could

follow up only three-fourths of our intended cohort, mostly due to migration, and limited availability

at home due to long working hours of many participants. Proportion of individuals on-treatment was

modest, reflecting the real-world apprehensions many individuals with hypertension have, regarding

initiation of drug therapy. Further, the process of obtaining prescription refills from a health-facility

often requires missing work on that day, which may not be feasible for the working populations. Last,

our follow up duration was only two years. While this is short, but we believe it is sufficient to

demonstrate improvements within program settings.

To conclude, in the current study we found about six of every ten individuals with hypertension were

on-treatment, and about four of them were optimally controlled. Our study has implications for

implementation and practice of NPCDCS. These findings truly provide a benchmark for NPCDCS

which are in terms of achievable goals within short periods of follow-up. Newly detected individuals

with hypertension, and those who are obese, are two such subgroups that need more intensive

monitoring and follow-up to improve optimal control levels in the community. CHWs based follow-

ups are feasible, but more intensive treatments and follow-up systems are needed for better blood-

pressure control outcomes.

What was already known?

In India only around 13% of all individuals with hypertension were on anti-hypertensive medications,

and a meagre eight percent were controlled. NPCDCS program is rolled out to improve this situation.

What this study adds?

In the current study we found that after initial screening and two-year follow-up, about six of every

ten individuals with hypertension were on-treatment, and about four were optimally controlled.

Those who were uncontrolled, were more likely to be older, had a higher BMI, and were newly

detected to have hypertension.

These findings demonstrate that obtaining blood-pressure control in vulnerable communities, though

feasible is challenging.

References

References

1.   Anchala R, Kannuri NK, Pant H, Khan H, Franco OH, Di Angelantonio E, et al. Hypertension in
India: a systematic review and meta-analysis of prevalence, awareness, and control of
hypertension. J Hypertens. 2014 Jun;32(6):1170–7.

2.   Kokane AM, Joshi R, Kotnis A, Chatterjee A, Yadav K, Revadi G, et al. Descriptive profile of
risk factors for cardiovascular diseases using WHO STEP wise approach in Madhya Pradesh.
PeerJ. 2020;8:e9568.

3.   Gupta R, Gaur K, S Ram CV. Emerging trends in hypertension epidemiology in India. J Hum
Hypertens. 2019 Aug;33(8):575–87.

4.   Stafford RS. New High Blood Pressure Guidelines: Back on Track With Lower Treatment Goals,
but Implementation Challenges Abound. Am J Prev Med. 2018 Oct;55(4):575–8.

5.   Thangada ND, Garg N, Pandey A, Kumar N. The Emerging Role of Mobile-Health Applications
in the Management of Hypertension. Curr Cardiol Rep. 2018 Jul 26;20(9):78.

6.   Vedanthan R, Bernabe-Ortiz A, Herasme OI, Joshi R, Lopez-Jaramillo P, Thrift AG, et al.
Innovative Approaches to Hypertension Control in Low- and Middle-Income Countries. Cardiol
Clin. 2017 Feb;35(1):99–115.

7.   R V, Jh K, Ak D, V N, Ca B, Gs B, et al. Community Health Workers Improve Linkage to
Hypertension Care in Western Kenya. J Am Coll Cardiol. 2019 Sep 2;74(15):1897–906.

8.   Joshi R, Agrawal T, Fathima F, Usha T, Thomas T, Misquith D, et al. Cardiovascular risk factor
reduction by community health workers in rural India: A cluster randomized trial. Am Heart J.
2019 Oct;216:9–19.

9.   He J, Irazola V, Mills KT, Poggio R, Beratarrechea A, Dolan J, et al. Effect of a Community
Health Worker-Led Multicomponent Intervention on Blood Pressure Control in Low-income
Patients in Argentina: A Randomized Clinical Trial. JAMA. 2017 Sep 19;318(11):1016–25.

10.   Jafar TH, Gandhi M, de Silva HA, Jehan I, Naheed A, Finkelstein EA, et al. A Community-Based
Intervention for Managing Hypertension in Rural South Asia. N Engl J Med. 2020 Feb
20;382(8):717–26.

11.   Neupane D, McLachlan CS, Mishra SR, Olsen MH, Perry HB, Karki A, et al. Effectiveness of a
lifestyle intervention led by female community health volunteers versus usual care in blood
pressure reduction (COBIN): an open-label, cluster-randomised trial. Lancet Glob Health.
2018;6(1):e66–73.

12.   Smolen JR, Wang JJ, Anane SP. Challenges and strategies for hypertension management:
qualitative analysis of small primary care practices with varying blood pressure control. Fam
Pract. 2019 Oct 8;36(5):644–9.

13.   Pakhare AP, Joshi A, Khadanga S, Kumar S, Atal S, Ingle V, et al. Feasibility of Community
Health Worker based cardiovascular risk reduction strategies in urban slums of Bhopal: Rationale
and design of community based study. medRxiv. 2020 Sep 18;2020.09.18.20189639.

14.   R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2019. Available from: https://www.R-
project.org/

15. Heinzen E, Sinnwell J, Atkinson E, Gunderson T, Dougherty G, Votruba P, et al. arsenal: An
Arsenal of "R" Functions for Large-Scale Statistical Summaries [Internet]. 2020 [cited 2020 Dec
24]. Available from: https://CRAN.R-project.org/package=arsenal

16. Harrison E, Drake T, Ots R. finalfit: Quickly Create Elegant Regression Results Tables and Plots
when Modelling [Internet]. 2020 [cited 2020 Oct 17]. Available from: https://CRAN.R-
project.org/package=finalfit

17. Wickham H, Chang W, Henry L, Pedersen TL, Takahashi K, Wilke C, et al. ggplot2: Create
Elegant Data Visualisations Using the Grammar of Graphics [Internet]. 2020 [cited 2020 Oct 17].
Available from: https://CRAN.R-project.org/package=ggplot2

18. Kassambara A. ggpubr: "ggplot2" Based Publication Ready Plots [Internet]. 2020 [cited 2020
Dec 24]. Available from: https://CRAN.R-project.org/package=ggpubr

19. Patil I. ggstatsplot: "ggplot2" Based Plots with Statistical Details [Internet]. 2020 [cited 2020 Dec
24]. Available from: https://CRAN.R-project.org/package=ggstatsplot

20. Sjoberg DD, Curry M, Hannum M, Whiting K, Zabor EC, Drill E, et al. gtsummary: Presentation-
Ready Data Summary and Analytic Result Tables [Internet]. 2020 [cited 2020 Oct 17]. Available
from: https://CRAN.R-project.org/package=gtsummary

21. easystats/performance [Internet]. easystats; 2020 [cited 2020 Dec 24]. Available from:
https://github.com/easystats/performance

22. Wickham H, RStudio. tidyverse: Easily Install and Load the "Tidyverse" [Internet]. 2019 [cited
2020 Oct 19]. Available from: https://CRAN.R-project.org/package=tidyverse

23. Prenissl J, Manne-Goehler J, Jaacks LM, Prabhakaran D, Awasthi A, Bischops AC, et al.
Hypertension screening, awareness, treatment, and control in India: A nationally representative
cross-sectional study among individuals aged 15 to 49 years. PLOS Med. 2019 May
3;16(5):e1002801.

24. Jb B, C Z, Hm T, Dj M, Pj O, Kl M, et al. Combination therapy as initial treatment for newly
diagnosed hypertension. Am Heart J. 2011 Jul 18;162(2):340–6.

25. Krousel-Wood M, Muntner P, Carson A, Anderson AH, Delaune E, Cushman WC, et al.
Hypertension control among newly treated patients before and after publication of the main
ALLHAT results and JNC 7 guidelines. J Clin Hypertens Greenwich Conn. 2012 May;14(5):277–
83.

26. Chia YC, Kario K, Turana Y, Nailes J, Tay JC, Siddique S, et al. Target blood pressure and
control status in Asia. J Clin Hypertens. 2020 Mar 1;22(3):344–50.

27. Marx G, Witte N, Himmel W, Kühnel S, Simmenroth-Nayda A, Koschack J. Accepting the
unacceptable: medication adherence and different types of action patterns among patients with
high blood pressure. Patient Educ Couns. 2011 Dec;85(3):468–74.

28. Gil-Guillén V, Orozco-Beltrán D, Carratalá-Munuera C, Márquez-Contreras E, Durazo-Arvizu R,
Cooper R, et al. Clinical inertia in poorly controlled elderly hypertensive patients: a cross-
sectional study in Spanish physicians to ascertain reasons for not intensifying treatment. Am J
Cardiovasc Drugs Drugs Devices Interv. 2013 Jun;13(3):213–9.

29. Mancia G, Rea F, Cuspidi C, Grassi G, Corrao G. Blood pressure control in hypertension. Pros
and cons of available treatment strategies. J Hypertens. 2017 Feb;35(2):225–33.

30. Ware LJ, Chidumwa G, Charlton K, Schutte AE, Kowal P. Predictors of hypertension awareness,
treatment and control in South Africa: results from the WHO-SAGE population survey (Wave 2).
J Hum Hypertens. 2019 Feb;33(2):157–66.

31. Shelley D, Tseng TY, Andrews H, Ravenell J, Wu D, Ferrari P, et al. Predictors of blood pressure
control among hypertensives in community health centers. Am J Hypertens. 2011
Dec;24(12):1318–23.

32. Knight EL, Bohn RL, Wang PS, Glynn RJ, Mogun H, Avorn J. Predictors of Uncontrolled
Hypertension in Ambulatory Patients. Hypertension. 2001 Oct 1;38(4):809–14.

33. Carey RM, Muntner P, Bosworth HB, Whelton PK. Prevention and Control of Hypertension:
JACC Health Promotion Series. J Am Coll Cardiol. 2018 Sep 11;72(11):1278–93.

34. Diaz KM, Shimbo D. Physical Activity and the Prevention of Hypertension. Curr Hypertens Rep.
2013 Dec;15(6):659–68.

35. Oxlund CS, Pareek M, Rasmussen BSB, Vaduganathan M, Biering-Sørensen T, Byrne C, et al.
Body Mass Index, Intensive Blood Pressure Management, and Cardiovascular Events in the
SPRINT Trial. Am J Med. 2019 Jul;132(7):840–6.

36. Zhang YQ, Li Y, Dong YG, Wu YH, Bian R, Li JH, et al. A nationwide assessment of blood
pressure control and the associated factors in Chinese type 2 diabetes mellitus patients. J Clin
Hypertens. 2019;21(11):1654–63.

37. Fahey T, Schroeder K, Ebrahim S. Interventions used to improve control of blood pressure in
patients with hypertension. Cochrane Database Syst Rev. 2006 Oct 18;(4):CD005182.

38. SPRINT Research Group, Wright JT, Williamson JD, Whelton PK, Snyder JK, Sink KM, et al. A
Randomized Trial of Intensive versus Standard Blood-Pressure Control. N Engl J Med. 2015 Nov
26;373(22):2103–16.

39. Schwalm JD, McCready T, Lopez-Jaramillo P, Yusoff K, Attaran A, Lamelas P, et al. A
community-based comprehensive intervention to reduce cardiovascular risk in hypertension
(HOPE 4): a cluster-randomised controlled trial. Lancet Lond Engl. 2019 Oct
5;394(10205):1231–42.

40. Long H, Huang W, Zheng P, Li J, Tao S, Tang S, et al. Barriers and Facilitators of Engaging
Community Health Workers in Non-Communicable Disease (NCD) Prevention and Control in
China: A Systematic Review (2006‾2016). Int J Environ Res Public Health. 2018 Oct 26;15(11).

   Table-1: Change in cardiovascular risk factors over two-year follow-up in the cohort (n=1177)

Variables	Baseline (2017)	Follow Up (2019)	Absolute Difference (CI)	p-value
All participants (N=1177)				
SBP (mean, SD)	146.6 (20.6)	145.0 (21.2)	-1.7 (-2.9 to -0.5)	0.007
DBP(mean, SD)	88.2 (12.6)	86.5 (12.6)	-1.7 (-2.4 to -1.1)	<0.001
Optimal BP control	301 (26%)	442 (38%)	+141 (12%)	<0.001
BMI(mean, SD)	25.7 (4.9)	25.8 (5.2)	+0.1 (-0.05 to 0.18)	0.257
Obese or overweight	536 (54%)	533 (54%)	-3 (1%)	1.000
Waist Circum. (mean, SD)	89.4 (14.4)	92.2 (12.2)	2.9 (2.3 to 3.5)	<0.001
Men	89.4 (19.5)	92.5 (11.5)	3.2 (1.1 to 5.2)	0.002
Women	88.6 (16.8)	92.1 (12.7)	3.4 (1.9 to 5.0)	<0.001
Abdominal Obesity	817 (79%)	868 (82%)	51 (3%)	0.0008
Previously known hypertension at baseline (N=623)				
SBP (mean, SD)	141.8 (22.9)	141.6 (21.0)	-0.2 (-1.9 to 1.5)	0.825
DBP(mean, SD)	84.3 (13.4)	83.8 (12.3)	-0.5 (-1.4 to 0.5)	0.316
Optimal BP Control	301 (48%)	284 (46%)	-17 (2%)	0.277
BMI(mean, SD)	26.2 (4.9)	26.3 (5.0)	0.1 (-0.05 to 0.26)	0.178
Obese or overweight	320 (59%)	313 (57%)	-7 (2%)	0.371
Waist Circum. (mean, SD)	89.9 (14.8)	93.1 (12.3)	3.2 (2.3 to 4.0)	<0.001
Men	89.7 (16.3)	93.8 (11.8)	4.0 (1.2 to 6.9)	0.005
Women	89.2 (18.5)	92.7 (12.4)	3.6 (1.4 to 5.7)	0.001
Abdominal Obesity	464 (85%)	488 (87%)	+22 (2%)	0.021
Newly detected hypertension at baseline (N=554)				
SBP (mean, SD)	152.0 (16.1)	148.7 (20.7)	-3.3 (-5.0 to -1.6)	<0.001
DBP(mean, SD)	92.7 (10.1)	89.5 (12.3)	-3.2 (-4.14 to -2.2)	<0.001
Optimal BP control	0 (0%)	158 (29%)	+158 (29%)	<0.001
BMI(mean, SD)	25.2 (4.9)	25.2 (5.4)	0.0 (-0.15 to 0.18)	0.865
Obese or overweight	216 (48%)	220 (49%)	+4(1%)	0.429
Waist Circum. (mean, SD)	88.7 (14.0)	91.3 (12.0)	2.6 (1.7 to 3.5)	<0.001
Men	89.1 (21.4)	91.7 (11.2)	2.6 (0.26 to 5.4)	0.075
Women	87.8 (13.8)	91.0 (13.0)	3.2 (1.0 to 5.5)	0.005
Abdominal Obesity	353 (72%)	380 (76%)	+27(4%)	0.161

409 **Table-2: Distribution of status of BP control among various socio-demographic, behavioral and**
410 **biological variables**

| Baseline Characteristics | All N = 1,177 | Blood-pressure outcome after follow-up | | p-value |
		Optimal control N = 442 (38%)	Not controlled N = 735 (62%)	
Mean age	51.7 (12.7)	49.9 (12.3)	52.9 (12.8)	<0.001
Age Group				<0.001
31-40	284	134 (47%)	150 (53%)	
41-50	347	124 (36%)	223 (64%)	
51-60	272	101 (37%)	171 (63%)	
60+	274	83 (30%)	191 (70%)	
Gender				0.669
Men	482	185 (38%)	297 (62%)	
Women	695	257 (37%)	438 (63%)	
Education				0.047
Literate	685	274 (40%)	411 (60%)	
Illiterate	492	168 (34%)	324 (66%)	
Wealth Quintiles				0.641
Q1	148	49 (33%)	99 (67%)	
Q2	179	70 (39%)	109 (61%)	
Q3	218	81 (37%)	137 (63%)	
Q4	230	93 (40%)	137 (60%)	
Q5	332	120 (36%)	212 (64%)	
Tobacco				0.688
Non-user	725	276 (38%)	449 (62%)	
User	452	166 (37%)	286 (63%)	
Alcohol				0.688
Non-user	951	354 (37%)	597 (63%)	
User	226	88 (39%)	138 (61%)	
Physical Activity				<0.001
Non-sedentary	285	131 (46%)	154 (54%)	
Sedentary	892	311 (35%)	581 (65%)	
Fruits and Vegetables (Median servings)	1.0 (0.5, 2.0)	1.0 (1.0, 2.0)	1.0 (0.5, 2.0)	0.081
Mean Salt Intake (gm/person/day)	8.3 (6.7, 11.1)	8.3 (6.7, 11.1)	8.3 (6.7, 11.1)	0.803

Mean Edible-oil intake (ml/person/day)	30.0 (25.0, 42.0)	30.0 (25.7, 42.0)	30.0 (25.0, 42.0)	0.808
Abdominal Obesity				0.418
Non-obese	360	142 (39%)	218 (61%)	
Obese	816	300 (37%)	516 (63%)	
HTN History				<0.001
Newly Detected	554	158 (29%)	396 (71%)	
Past HTN	623	284 (46%)	339 (54%)	
BP Status at Baseline				<0.001
Not controlled	876	245 (28%)	631 (72%)	
Controlled	301	197 (65%)	104 (35%)	
Dysglycemia				0.417
Absent	822	302 (37%)	520 (63%)	
Present	355	140 (39%)	215 (61%)	
Weight (mean ; SD)	62.5 (13.6)	61.7 (13.6)	62.9 (13.6)	0.169
BMI (mean ; SD)	25.8 (5.2)	25.5 (5.3)	26.0 (5.2)	0.172
Waist Circumference (mean ; SD)	88.9 (17.9)	88.6 (16.5)	89.1 (18.8)	0.376
Random Blood Sugar (mean ; SD)	138.2 (61.0)	138.1 (60.9)	138.3 (61.2)	0.696

414   **Table-3: Logistic regression analysis for determinants of uncontrolled hypertension**

Baseline variables	OR (univariable)	OR (multivariable)
Age	1.02 (1.01-1.03, p<0.001)	1.02 (1.01-1.04, p<0.001)
Women vs Men	1.06 (0.84-1.35, p=0.625)	0.95 (0.69-1.30, p=0.744)
Illiterate vs Literate	1.29 (1.01-1.64, p=0.041)	1.21 (0.88-1.65, p=0.241)
Tobacco User vs Non-user	1.06 (0.83-1.35, p=0.643)	-
Alcohol User vs Non-user	0.93 (0.69-1.26, p=0.632)	-
Physical Inactivity vs Non-sedentary	1.59 (1.21-2.08, p=0.001)	1.44 (1.05-1.98, p=0.024)
BMI	1.02 (0.99-1.04, p=0.211)	1.04 (1.01-1.07, p=0.007)
Waist Circumference	1.00 (0.99-1.01, p=0.650)	-
Newly Detected HTN at baseline vs Previously known HTN	2.10 (1.65-2.68, p<0.001)	2.38 (1.80-3.17, p<0.001)
Dysglycemia at baseline vs No Dysglycemia	0.89 (0.69-1.15, p=0.381)	-

Number in data frame = 1177, Number in model = 997, Missing = 180, AIC = 1247.8, C-statistic = 0.65, H&L = Chi-sq(8) 5.81 (p=0.669)

OR- Odds Ratio, BMI- Body Maas Index, HTN- Hypertension

Change in SBP among newly detected and previously known hypertension groups stratified by BP control status at baseline

Grp: Newly Detected : Uncontrolled

$t_{\text{Student}}(552) = 3.30$, $p = 0.001$, $\widehat{d}_{\text{Cohen}} = 0.14$, $CI_{95\%}$ [0.06, 0.22], $n_{\text{pairs}} = 553$

Grp: Past HTN : Controlled

$t_{\text{Student}}(300) = -8.43$, $p = 1.43e{-}15$, $\widehat{d}_{\text{Cohen}} = -0.49$, $CI_{95\%}$ [-0.61, -0.37], $n_{\text{pairs}} = 301$

Grp: Past HTN : Uncontrolled

$t_{\text{Student}}(320) = 6.91$, $p = 2.7e{-}11$, $\widehat{d}_{\text{Cohen}} = 0.39$, $CI_{95\%}$ [0.27, 0.50], $n_{\text{pairs}} = 321$