Recruitment location influences bias and uncertainty in SARS-CoV-2 seroprevalence estimates

Tyler S. Brown[1,2,*], Pablo Martinez de Salazar Munoz[2], Abhishek Bhatia[2], Bridget Bunda[1], Ellen K.
Williams, David Bor[3], James S. Miller[1], Amir M. Mohareb[1], Julia Thierauf[1], Wenxin Yang[1], Julian
Villalba[1], Vivek Naranbai[1], Wilfredo Garcia Beltran[1], Tyler E. Miller[1], Doug Kress[4], Kristen Stelljes[5],
Keith Johnson[5], Daniel B. Larremore[6], Jochen Lennerz[1], A. John Iafrate[1], Satchit Balsari[2], Caroline O.
Buckee[2,**], Yonatan H. Grad[2,**]
7

[1] Massachusetts General Hospital, Boston, MA
[2] Harvard T.H. Chan School of Public Health, Boston, MA
[3] Cambridge Health Alliance, Cambridge, MA
[4] Board of Health, City of Somerville, Massachusetts
[5] SomerStat, City of Somerville, Massachusetts
[6] University of Colorado, Boulder
* Corresponding author: tsbrown@mgh.harvard.edu
** Joint senior authors

Abstract: 147 words

Main text: 3,225

Abstract

The initial phase of the COVID-19 pandemic in the US was marked by limited diagnostic testing,

resulting in the need for seroprevalence studies to estimate cumulative incidence and define epidemic

dynamics. In lieu of systematic representational surveillance, venue-based sampling was often used to

rapidly estimate a community's seroprevalence. However, biases and uncertainty due to site selection

and use of convenience samples are poorly understood. Using data from a SARS-CoV-2

serosurveillance study we performed in Somerville, Massachusetts, we found that the uncertainty in

seroprevalence estimates depends on how well sampling intensity matches the known or expected

geographic distribution of seropositive individuals in the study area. We use GPS-estimated foot traffic

to measure and account for these sources of bias. Our results demonstrated that study-site selection

informed by mobility patterns can markedly improve seroprevalence estimates. Such data should be

used in the design and interpretation of venue-based serosurveillance studies.

Keywords: SARS-CoV-2, serosurveillance, study design, human mobility

Introduction

Studies estimating SARS-CoV-2 seroprevalence have been critical to our understanding of COVID-19 pandemic dynamics, particularly in times when diagnostic testing was limited and the extent of community spread was unknown [1–4]. However, many of these estimates were derived from non-representational convenience sampling [4–6], thus making these estimates subject to multiple sources of bias and uncertainty that remain poorly understood [7]. Low precision can obscure differences in estimated seroprevalence between populations, limiting efforts to understand heterogeneity in epidemic intensity or vaccination coverage and reducing the effectiveness of public health interventions. As cost, speed, and logistical concerns continue to motivate the use of convenience sampling [8], identifying and accounting for bias and uncertainty are therefore critical to improving the design and interpretability of these studies.

Convenience sampling involves inherently non-uniform sampling across demographic or geographic subgroups. For example, in venue-based (or "walk-up") studies, in which participants are recruited from among visitors to a central or highly trafficked location [4,9], the geographic distribution of participants is expected to be skewed toward individuals living closer to the study location. A similar concern applies to studies using discarded blood samples, where the catchment area of a given hospital or clinical laboratory may strongly constrain the geographic distribution of samples available for analysis [5].

Multiple recent studies have underscored the utility of GPS- and mobile phone-associated human mobility data in understanding COVID-19 epidemiology. These studies have clarified the role of highly-visited "super-spreader" locations [10,11] in local COVID-19 epidemics and identified higher aggregate mobility as a predictor of neighborhood-level COVID-19 risk [3]. Prospective applications of these data sources, including their use in the design and implementation of epidemiological studies, are limited.

Here, we examine seroprevalence estimates obtained via venue-based sampling [4,9], with direct

applications to other forms of convenience sampling [5,12,13]. We performed a venue-based

serosurveillance study in Somerville, Massachusetts and used these data to analyze bias and uncertainty

arising from inherent geographic variation in sampling intensity. This analysis showed that substantial

loss of precision can occur if the distribution of sampling intensity poorly matches the geographic

distribution of true seropositive individuals in the study area. As GPS-estimated foot traffic offers a

proxy measurement for the geographic distribution of visitors to different locations, we evaluated the

extent to which informed selection of locations for venue sampling reduces uncertainty and bias

introduced by geographic heterogeneity in underlying seropositivity. Our results thus offer an approach

to significantly improve the design and interpretation of seroprevalence studies that use venue-based

convenience sampling with little if any impact on cost and speed.

Methods

SARS-CoV-2 seroprevalence study design and participant information

We obtained serological and participant demographic data for 398 asymptomatic adults tested at a

temporary study site near an essential business location in Somerville, Massachusetts. The study was

conducted over 4 days (June 4th, 5th, 8th, and 9th, 2020), approximately 6 weeks after the first wave of the

COVID-19 epidemic peaked in Massachusetts [14]. The study was designated minimal risk human

subjects research and approved by institutional review boards at Massachusetts General Hospital and the

Harvard T.H. Chan School of Public Health (Protocol number: 2020P001081). The study recorded

participant demographic information including age, gender, and self-reported home locations (by ZIP

code and electoral ward). We also collected information on how participants learned about the study in

order to distinguish participants directly recruited on site at the study location from those who learned

about the study from friends, family, or social media. We did not advertise or announce enrollment for

the study prior to its implementation, with the goal of increasing the proportion of individuals recruited

78   on site at the study location. We used this data to calculate P_j^{direct}, the proportion of all directly recruited

79   participants from Somerville with self-reported home locations in electoral ward $j \in \{1,\dots,7\}$. We refer

80   to $\{P_1^{\text{direct}},\dots,P_7^{\text{direct}}\}$ as the "survey participant catchment distribution."

Public health acute infection data

We obtained data on 916 PCR-confirmed COVID-19 cases with documented home addresses in

Somerville (collected from the onset of the epidemic through June, 2020) from the Massachusetts

Virtual Epidemiologic Network (MAVEN). COVID-19 cases are reported to MAVEN by state and local

health agencies, and cases are designated as "confirmed" if they have a positive result for SARS-CoV-2

RNA detection using an FDA-approved molecular amplification detection test, for example RT-PCR.

The total number of Somerville residents tested for SARS-COV-2 via RT-PCR was also obtained from

MAVEN. Data were anonymized and aggregated by electoral ward prior to analysis. We calculated the

cumulative incidence of PCR-confirmed infections by ward (θ_j^{PCR}) and the proportion of all PCR tests

with positive results ("PCR positivity").

GPS-estimated business foot traffic

We used GPS-estimated foot traffic (SafeGraph, *safegraph.com*) to approximate the distribution of

home locations for daytime visitors for different locations of interest, which we refer to as the "GPS-

estimated catchment area." We use these data to estimate home and work locations at the level of census

block group (CBG) for visitors to designated points-of-interest, such as businesses, and specific CBGs.

We used CBG-level visitor data for June 2020 as the primary data source for visitors in our analysis.

These data were filtered for CBGs with low visitor counts and re-aggregated from CBGs to electoral

wards prior to analysis (Supplementary Material, Figures S1 and S2). We used the filtered, re-

aggregated data to obtain GPS-estimated visitor catchment distributions for two locations: (1) the actual

study venue, located in Somerville electoral Ward 2 (denoted V_j^{site}) and (2) a hypothetical alternative

study venue located in Somerville electoral Ward 1 (denoted V_j^{alt}).

 Simulations

We used numerical simulation to examine bias and uncertainty in estimated seroprevalence ($\hat{\theta}_{pop}$) as a

function of subgroup sizes, true subgroup seropositivity, and sample allocation. Here, "subgroup" is

used to describe any potential stratification of the overall population, including by demographic and/or

geographic characteristics. Briefly, we used demographic [15] and SARS-CoV-2 acute infection data

from Somerville, MA to generate a simulated population with varying true seropositivity $\theta_{j,k}$ across

subgroups stratified by age group j and location k (where locations are electoral wards in Somerville).

Briefly, the simulation randomly draws $n_{j,k}$ individuals from each subgroup, calculates weighted

population-level seroprevalence (adjusted for serological test performance), and repeats this process

10000 times to generate distributions of $\hat{\theta}_{pop}$ values. We report W, the width of the 95th percentile

interval for each distribution, as an approximate measure of uncertainty for $\hat{\theta}_{pop}$ for a given set of

simulation parameters. Additional details are available in the Supplementary Material. *R* code for the

numerical simulations is available at *https://github.com/svsero/COVID19serosurveillance-Somerville*.

Results

Study participant catchment distributions and geographic heterogeneity in COVID-19 epidemic intensity

We first examined how participant catchment distributions align with, or mismatch, the geographic

distribution of seropositive individuals in a given study area. We observed that the survey participant

catchment distribution in our serosurveillance study was skewed strongly toward locations near the

study site (Figure 1A). Among directly recruited participants with home locations in Somerville, 43%

(43/100) reported home locations in Somerville Ward 2 (where the study site was located) compared to

4% in Ward 1 and 4% in Ward 4. In contrast, the cumulative incidence of PCR-confirmed SARS-CoV-2

infections (θ_j^{PCR}) was approximately three-fold higher in electoral Ward 1 compared to Wards 2 and 6

(Figure 1B) and the proportion of SARS-CoV-2 PCR tests with positive results was approximately five-

fold higher (Supplementary Figure S3A). Both of these proxy measures of epidemic intensity (θ_j^{PCR} and

PCR test positivity) are limited by potential biases, some of which are likely to still be present even if

PCR testing rates are relatively equal by ward [16]. Nonetheless, these measures suggest substantial

heterogeneity in the underlying epidemic intensity, with an apparent higher rate of previously infected

individuals (as a proportion of the population) in Somerville Wards 1 and 4. Thus, we observed that the

venue location chosen for this study, and its associated survey participant catchment distribution,

resulted in relative under-sampling of wards with expected higher seropositivity and oversampling of

those with lower expected seropositivity (Figure 1C).

GPS-estimated visitor catchment distributions

Recognizing that survey participant catchment distributions can be poorly matched to the underlying

geographic distribution of seropositivity, we explored the use of GPS-estimated foot traffic data as a tool

for evaluating actual or candidate locations for venue-based sampling. We evaluated correlations

between the observed participant catchment distribution for the actual Somerville study location, GPS-

estimated visitor catchment distributions for this site and a hypothetical alternative site in Somerville

Ward 1, and the cumulative incidence of PCR-confirmed infections by ward. The participant catchment

distribution at the actual study site (P_j^{direct}) closely matched its corresponding GPS-estimated visitor

catchment distribution, V_j^{site} (Pearson's $r = 0.90$, $p = 0.0131$, Figure S4). However, the GPS-estimated

visitor catchment distribution for the actual study site was poorly correlated with the cumulative

incidence of PCR-confirmed infections ($r= -0.11$, $p = 0.55$, Figure 2C).

We evaluated whether choosing an alternative study site could improve the correlation between sample

allocation and cumulative incidence of PCR-confirmed infections by ward, with the goal of reducing

uncertainty in estimated seroprevalence. We observed that the GPS-estimated visitor catchment

distribution at the alternative site (V_j^{alt}) is strongly correlated with ward-level cumulative incidence of

PCR-confirmed infections ($r=0.93$, $p=0.0072$, Figure 2D). If subgroup sizes are known and differences

in subgroup-level seropositivity can be inferred or assumed, allocating more samples to larger subgroups

and those with higher expected seropositivity will improve precision for weighted population-level

seroprevalence estimates (Figure S5 and Equation 2) [17]. This suggests that, if the GPS-estimated

visitor catchment distribution reliably predicts the survey participant catchment at the alternative study

site, this location would yield improved sample allocation across geographic subgroups that more

closely approximates optimal sample allocation.

Venue location and uncertainty in SARS-CoV-2 seroprevalence estimates

We used numerical simulation to quantify uncertainty in estimated SARS-CoV-2 seroprevalence under

different survey participant catchment distributions. Specifically, we constructed a simple synthetic

population with seven geographic subgroups, corresponding to the seven electoral wards in Somerville,

each divided into three age-based subgroups. We specified the size of each subgroup using local census

data [15] and specified the true underlying seropositivity for each age-location subgroup by assuming

these values are proportional to the observed cumulative incidence of PCR-confirmed infections for

each subgroup. Using this model, we compared three sample allocation scenarios: (1) optimal allocation,

in which the number of individuals sampled from each age-location subgroup is specified to optimally

reduce uncertainty in the resulting seroprevalence estimates (per Equation 2 in the Supplementary

Information); (2) allocation according to the observed survey participant catchment distribution for the

actual study site; (3) allocation according to the GPS-estimated visitor distribution at the hypothetical

alternative study site. (Additional details on model specification and sensitivity testing for model

parameters are available in the Supplementary Information.)

We observed 1.5- to 2-fold higher uncertainty when sampling effort was allocated according to the

participant catchment distribution at the study site compared to the alternative site or optimal allocation

(Figure 3). This observation suggests that choice of recruitment location can result in suboptimal sample

allocation and higher uncertainty.

Bias due to unappreciated heterogeneity in seropositivity across geographic subgroups

Biased seroprevalence estimates can result if geographic heterogeneity in sample allocation results in

substantial over- or under-sampling in locations with higher (or lower) seropositivity, and if procedures

for generating weighted prevalence estimates do not appropriately account for geographic heterogeneity

in underlying seropositivity. We compared estimated seroprevalence versus true seroprevalence for

numerical simulations in which the final seroprevalence estimates were weighted by the sampling

probability for each age-location group or by the sampling probability of each age subgroup alone

(Figure 4). The first weighting procedure accounts for heterogeneity across age and location subgroups,

whereas the second procedure accounts only for heterogeneity across age subgroups. Using the second

procedure resulted in over- or under-estimation of seroprevalence, depending on whether sample

allocation enriches for participants from areas with high or low underlying seropositivity, respectively.

Discussion

Convenience sampling, despite its inherent limitations, may have continued utility in the public health

response to the COVID-19 pandemic and for infectious disease outbreaks generally. Cost and logistical

considerations may limit the feasibility of randomized structured sampling, particularly in resource-

constrained contexts or in situations where census data, population rosters, or household mapping data

are unavailable or unreliable. Certain forms of convenience sampling are better suited for reaching

important subgroups compared to structured approaches. Lower-wage or frontline workers who are at

higher risk of SARS-CoV-2 exposure [18-20], including undocumented workers [20], may be less likely

to participate if recruited using conventional survey outreach methods (e.g., mail or phone contact) due

to constraints on their time [21-23] and lack of incentives [21]. Convenience sampling at highly visited

community locations such as essential businesses may be an attractive alternative to structured sampling

in this important population, similar to sampling approaches developed to study so-called "hidden

populations" [24].

Geographic heterogeneity in SARS-CoV-2 epidemic intensity within cities has been a repeatedly observed feature of the pandemic [1–3]. This phenomenon poses unique challenges for seroprevalence studies that employ venue-based and other convenience sampling strategies, in which sample allocation across subgroups cannot be pre-specified and is non-uniform across geographic space.

Examining data from our seroprevalence study in Somerville, MA, we observed that venue-based sampling resulted in substantial undersampling of areas where proxy measures (cumulative incidence of PCR-confirmed infections and PCR test positivity rates) suggest higher epidemic intensity. This mismatch between the survey participant catchment distribution and the geographic distribution of seropositive individuals can result in suboptimal sample allocation and higher variance in resultant seroprevalence estimates.

Study locations can have widely divergent participant catchment distributions. These distributions can be heavily enriched for participants living in the immediate vicinity of the study location. This limitation has important implications for bias and uncertainty of resulting seroprevalence estimates (as detailed here) and raises questions about potential undersampling or exclusion of important subgroups in venue-based studies. Multiple studies have identified geographic location as a strong surrogate for multiple risk factors associated with severe infection, hospitalization, and/or death due to COVID-19 [25] and undersampling in neighborhoods where these risk factors co-localize together can compromise the reliability and interpretability of seroprevalence estimates. Recruiting participants directly from such communities, where rates of COVID-19 related hospitalization and deaths are often higher, has yielded seroprevalence estimates that are substantially higher than city-level or state-level estimates [4,9]. Notably, the areas that were most undersampled in our study strongly overlap neighborhoods with lower socioeconomic status, larger proportions of non-white residents, lower proportions of English-speaking households (Figure 1A, Supplementary Figure S3C).

Our work has three practical findings that are applicable to the design, implementation, and interpretation of convenience-based seroprevalence studies.

(1) Uncertainty in population-level seroprevalence estimates is minimized when sample allocation is proportional to the size and underlying seropositivity of individual subgroups in the population (Equation 2 and Figure S5). The practical application of this finding may be limited because of challenges in reliably ascertaining differences in the underlying seroprevalence between subgroups *a priori*. For example, this may arise when access to diagnostic testing for acute infections is limited or disparate across subgroups. However, even in this situation, allocating sampling effort proportional to subgroup sizes alone can substantially reduce uncertainty (Figure S1). Applying this finding may also be difficult because an inherent feature of venue-based sampling is that allocation of sampling effort is not pre-specified, but instead results from a stochastic process that depends on the location of the study venue. Thus, although optimal sample allocation is likely not achievable via venue-based sampling, careful selection of venue locations, with the objective of enriching for participants from geographic subgroups with larger populations and/or higher expected seroprevalence, can at least improve sample allocation and help reduce uncertainty.

(2) GPS-estimated foot traffic can inform the selection of venue-based recruitment locations. The GPS-estimated visitor catchment distribution at our study location correlated closely with the survey participant catchment distribution. Validation against other data sources that directly measure the geographic distributions of visitors to locations of interest (for example, aggregated geographic and registration data from COVID-19 mobile testing programs) can help further evaluate this potentially important data source.

(3) Convenience sampling can produce biased seroprevalence estimates if geographic heterogeneity in underlying subgroup-level seropositivity is not properly accounted for (Figure 5). Consistent with prior studies [2, 3], we observed that COVID-19 incidence can vary widely, even over a relatively small geographic area. To avoid this problem, studies that employ convenience sampling should collect geographic data on participants' home locations that is granular enough to capture potential geographic heterogeneity in seroprevalence within the study area. This information, combined with data on the

catchment distributions of individual recruitment sites (including GPS-estimated foot traffic data, as

examined here), can be used to quantify what would otherwise be an unmeasured source of bias in

resulting seroprevalence estimates.

Multiple considerations are important for contextualizing our findings and the recommendations above.

These include participation bias that may result in exclusion of individuals with disabilities or others

who are less likely to leave their homes. Methods designed to account for participation bias, including

those developed for use with time-location sampling [26], may be applicable here. Likewise, collecting

information on non-respondents in venue-based sampling—for example, brief demographic surveys

collected before recruitment for serological testing—can help measure and account for potential sources

of participation bias.

The GPS-estimated foot traffic data used in our study have several important limitations. Identification

of visitors and their home locations in this data may be biased by differences in mobile device usage

between demographic groups, potentially under-sampling visitors from important populations or

oversampling others. In addition, this data can be sparse and thus more subject to stochastic variation

when only small numbers of users are captured in either the point of interest or home location. Different

forms of participation bias (described above) are expected to skew the geographic distribution of study

participants away from GPS-estimated catchment distributions, potentially making it difficult to use this

data source in real-world public health practice.

Lastly, we make several simplifying assumptions in our numerical model. The numerical model assumes

that the true seropositivity in each age-location subgroup is proportional to its observed cumulative

incidence of PCR-confirmed SARS-CoV-2 infections (per local public health data from Somerville,

MA). However, wards with higher PCR positivity rates (an indicator of greater epidemic intensity) have

relatively the same rates of overall PCR testing per capita (Supplementary Figure S3B), indicating that

there were gaps in testing effort in areas of Somerville with more incident infections overall [16]. The

assumed true underlying seroprevalence of each age-location group, which is specified using the

observed cumulative incidence of PCR-confirmed infection and does not account for the testing gap

described above, are less dispersed across age-location groups than what would be expected if PCR

testing effort better matched epidemic intensity by ward (i.e., greater PCR testing effort in heavily

impacted areas would likely reveal even larger differences in cumulative incidence between wards). This

misspecification, and resultant smaller dispersion in assumed true cumulative incidence by ward, is

expected to result in more conservative values for the uncertainty in estimated population-level

seroprevalence; otherwise, this limitation is not expected to change our primary findings from the

numerical model.

In summary, we have examined how geographic heterogeneity in sample allocation, combined with

underlying heterogeneity in geographic distribution of seropositive individuals, can influence

seroprevalence estimates derived from venue-based sampling. Our findings are relevant to studies

employing venue-based recruitment and are also applicable to other kinds of convenience sampling, for

example, studies using a hospital's discarded blood specimens from patients drawn from the hospital's

geographic catchment area.

References

1.   Feehan AK, Fort D, Garcia-Diaz J, et al. Seroprevalence of SARS-CoV-2 and Infection Fatality

Ratio, Orleans and Jefferson Parishes, Louisiana, USA, May 2020. Emerging Infectious Diseases **2020**;

26(11).

2.   Kim SJ, Bostwick W. Social Vulnerability and Racial Inequality in COVID-19 Deaths in

Chicago. Health Education & Behavior **2020**; 47(4):509–513.

3.   Kissler S, Kishore N, Prabhu M, et al. Reductions in commuting mobility predict geographic

differences in SARS-CoV-2 prevalence in New York City. Nature Communications **2020**; 11(1):4674.

4.   Rosenberg ES, Tesoriero JM, Rosenthal EM, et al. Cumulative incidence and diagnosis of

SARS-CoV-2 infection in New York. Annals of Epidemiology. **2020**; 48:23–29

5.   Havers FP, Reed C, Lim T, et al. Seroprevalence of Antibodies to SARS-CoV-2 in 10 Sites in

the United States, March 23-May 12, 2020. JAMA Internal Medicine **2020;** 180(12):1576–1586

6.   Bendavid E, Mulaney B, Sood N, et al. COVID-19 Antibody Seroprevalence in Santa Clara

County, California. International Journal of Epidemiology **2021**; 50(2):410-419.

7.   Shook-Sa BE, Boyce RM, Aiello AE. Estimation Without Representation: Early Severe Acute

Respiratory Syndrome Coronavirus 2 Seroprevalence Studies and the Path Forward. J Inf Dis. **2020**;

222(7):1086–1089.

8.   Kelly H. A random cluster survey and a convenience sample give comparable estimates of

immunity to vaccine preventable diseases in children of school age in Victoria, Australia. Vaccine.

2002; 20(25-26):3130–3136.

9.   Naranbhai V, Chang CC, Beltran WFG, et al. High seroprevalence of anti-SARS-CoV-2

antibodies in Chelsea, Massachusetts. J Inf Dis **2020**; 222(12):1955–1959.

10.   Chang, S., Pierson, E., Koh, P.W. et al. Mobility network models of COVID-19 explain

inequities and inform reopening. Nature **2021**; 589**:** 82–87.

11.   Ma, KC and Lipsitch M. Big data and simple models used to track the spread of COVID-19 in

cities. Nature **2021**; 589: 26-28.

12.   Hobbs CV, Drobeniuc J, Kittle T, et al. Estimated SARS-CoV-2 seroprevalence among persons

aged <18 years - Mississippi, May-September 2020. MMWR Morbidity and Mortality Weekly Report.

2021; 70(9):312–315.

13.   Sutton M, Cieslak P, Linder M. Notes from the field: Seroprevalence estimates of SARS-CoV-2

infection in convenience sample - Oregon, May 11-June 15, 2020. MMWR Morbidity and Mortality

Weekly Report. **2020**; 69(32):1100–1101.

14.   Massachusetts State Department of Health. COVID-19 Response Reporting. Available from:

https://www.mass.gov/info-details/covid-19-response-reporting

15.   City of Somerville, Massachusetts and Cambridge Health Alliance. The Wellbeing of Somerville

Report. Available at: https://www.somervillema.gov/sites/default/files/wellbeing-of-somerville-report-

2017.pdf

16.   Dryden-Peterson S, Velásquez GE, Stopka TJ, Davey S, Lockman S, Ojikutu B. Disparities in

SARS-CoV-2 Testing in Massachusetts During the COVID-19 Pandemic. JAMA Network Open **2020**;

4(2):e2037067

17.   Larremore DB, Fosdick BK, Bubar KM, et al. Estimating SARS-CoV-2 seroprevalence and

epidemiological parameters with uncertainty from serological surveys. Elife **2021;** 10:e64206

18.   Hawkins D. Social Determinants of COVID-19 in Massachusetts, United States: An Ecological

Study. Journal of Preventive Medicine and Public Health. Korean Society for Preventive Medicine;

2020; 53(4):220–227.

19.   Baker MG, Peckham TK, Seixas NS. Estimating the burden of United States workers exposed to

infection or disease: A key factor in containing risk of COVID-19 infection. PLOS ONE **2020**;

15(4):e0232452.

20.   Feehan AK, Velasco C, Fort D, et al. Racial and workplace disparities in seroprevalence of

SARS-CoV-2 in Baton Rouge, Louisiana, July 15-31, 2020. Emerging Infectious Diseases **2021**; 27(1):

314-317

21.   Hernández MG, Nguyen J, Casanova S, Suárez-Orozco C, Saetermoe CL. Doing no harm and

getting it right: Guidelines for ethical research with immigrant communities. New Directions for Child

and Adolescent Development. **2013**; 2013(141):43–60.

22.   Corbie-Smith DM. Minority recruitment and participation in health research. North Carolina

Medical Journal. **2004**; 65(6):385-7

23.   Keyzer JF, Melnikow J, Kuppermann M, et al. Recruitment strategies for minority participation:

challenges and cost lessons from the POWER interview. Ethn Dis **2005**; 15(3):395–406.

24.   Muhib FB, Lin LS, Stueve A, et al. A Venue-Based Method for Sampling Hard-to-Reach

Populations. Public Health Reports **2001**; 116:216–222.

25.   Maroko AR, Nash D, Pavilonis BT. COVID-19 and inequity: A comparative spatial analysis of

New York City and Chicago hot spots. Journal of Urban Health **2020**; 97(4):461–470.

26.   Leon L, Jauffret-Roustide M, Le Strat Y. Design-based inference in time-location sampling.

Biostatistics **2015**; 16(3):565–579.

Figures

Figure 1. Sample allocation and geographic heterogeneity in proxy measures of epidemic intensity. (A) Survey participant catchment distribution. Wards are shaded by P_j^{direct}, the proportion of all directly recruited participants from each of Somerville Wards 1-7; (B) Cumulative incidence of prior PCR-confirmed SARS-CoV-2 infections by Ward as of June 8th, 2020 (θ_j^{PCR}); (C) Correlation between P_j^{direct} and θ_j^{PCR}. Significance of the correlation is calculated via permutation testing, as described in the Supplementary Material.

Figure 2. GPS-estimated visitor catchment distributions for actual and hypothetical alternative study sites. GPS-estimated visitor catchment distributions for (A) the actual study location V_j^{site} or (B) a hypothetical alternative study site in Somerville Ward 1 V_j^{alt}. Correlations between V_j^{site}, V_j^{alt}, and the cumulative incidence of PCR-confirmed SARS-CoV-2 infection θ_j^{PCR} are shown in (C) and (D).

Figure 3. Uncertainty in estimated SARS-CoV-2 seroprevalence obtained using different

sample allocation strategies. The uncertainty (W, the width of the 95th percentile interval for

10000 estimated seroprevalence values) versus mean estimated seroprevalence for different values

of n (the total number of individuals sampled) when individuals are sampled using (1) in the left

panel, the optimal sample allocation according to Equation 2 in the Supplementary Material (S_{op});

(2), in the center panel, the sampling distribution of participants in the Somerville seroprevalence

survey (S_{site}); (3) the sampling distribution at the proposed alternative study site (S_{alt}).

Figure 4. Bias in estimated seroprevalence by sampling strategy and weighting procedures. Left: Estimated seroprevalence, weighted only by age subgroups. Right: Estimates weighted by age-location subgroups are shown in the right panel. Blue: Sample allocation specified by the observed participant distribution catchment distribution in the Somerville study. Red: Sample allocation specified by the catchment distribution of GPS-estimated visitors to the proposed alternate study site. Dotted line indicates where estimated equals true seroprevalence.

Funding

This work was supported by the Andrew and Corey Morris-Singer Foundation, National Cancer Institute at the National Institutes of Health [U01CA261277] and the National Institute of Allergy and Infectious Diseases at the National Institutes of Health [T32AI007061 to TSB]. This project has been funded in part by contract 200-2016-91779 with the Centers for Disease Control and Prevention. The findings, conclusions, and views expressed in this presentation are those of the author(s) and do not necessarily represent the official position of the Centers for Disease Control and Prevention (CDC).